

# Claim A in 3D: Proof Attempt via Caccioppoli–Gehring

WORKING DOCUMENT — T-First Programme

Route B (Gehring) + Route C ( $\theta$ -bootstrap) analysis

Pratik Menon  
T-First Programme

May 2026 — v0.1

## Abstract

This working document attempts to prove Claim A in three space dimensions:

Claim A:  $\nu|\nabla u|^2 \in L^{1+\delta}(Q_T)$  for some  $\delta > 0$ ,

for Leray–Hopf solutions of the 3D incompressible Navier–Stokes equations at  $\varepsilon = 0$ . We systematically pursue two routes and identify the precise gap in each.

**Route B (Caccioppoli–Gehring):** CKN local energy inequality  $\Rightarrow$  Caccioppoli for  $|\nabla u|^2 \Rightarrow$  reverse Hölder  $\Rightarrow$  Gehring  $\Rightarrow \delta > 0$ . **Status:** Steps 1–2 complete. Step 3 (reverse Hölder with controlled pressure) identified as the key task.

**Route C ( $\theta$ -bootstrap at  $\varepsilon = 0$ ):** Identity  $\nu|\nabla u|^2 = \partial_t \theta + u \cdot \nabla \theta - \nu \Delta \theta \Rightarrow$  bootstrap via parabolic regularity. **Status:** Bootstrap closes for any  $\delta_0 > 0$  (proved below). Base case  $\delta_0 > 0$  supplied by §20 vorticity recursion for  $u_0 \in H^1$ .

**Key finding:** Routes B and C are *complementary*: Route B provides the base case  $\delta_0 > 0$  (quantitative, from Gehring); Route C then iterates  $\delta_0 \rightarrow \infty$  (smooth). **WITHDRAWN (S31 audit — see §25):** the combined argument was claimed complete for  $u_0 \in H^1(\mathbb{T}^3)$  via §20, Cor. 20.6, but the independent audit in §25 finds the §20 induction arithmetically impossible for general data (Defect 1) and two invalid analytic steps (Defects 2–3); the Clay Prize claim for  $u_0 \in H^1$  is withdrawn. Global regularity for general  $H^1$  data remains **open**.

## Contents

|          |                                |          |
|----------|--------------------------------|----------|
| <b>1</b> | <b>Setup and Goal</b>          | <b>9</b> |
| 1.1      | The System . . . . .           | 9        |
| 1.2      | The Auxiliary Scalar . . . . . | 9        |
| 1.3      | Claim A Precisely . . . . .    | 9        |

|          |                                                                               |           |
|----------|-------------------------------------------------------------------------------|-----------|
| <b>2</b> | <b>Route B: Caccioppoli–Gehring</b>                                           | <b>9</b>  |
| 2.1      | Step 1: Suitable Weak Solutions and Local Energy . . . . .                    | 9         |
| 2.2      | Step 2: Caccioppoli Inequality . . . . .                                      | 10        |
| 2.3      | Step 3: Controlling the Pressure . . . . .                                    | 10        |
| 2.4      | Step 4: Reverse Hölder Inequality . . . . .                                   | 11        |
| 2.5      | Step 5: Gehring’s Lemma Gives $\delta_0 > 0$ . . . . .                        | 12        |
| <b>3</b> | <b>Route C: <math>\theta</math>-Bootstrap at <math>\varepsilon = 0</math></b> | <b>13</b> |
| 3.1      | The Bootstrap Map . . . . .                                                   | 13        |
| 3.2      | Convergence of the Bootstrap . . . . .                                        | 14        |
| 3.3      | From $\delta_n \rightarrow \infty$ to Smoothness . . . . .                    | 14        |
| <b>4</b> | <b>The Single Gap: Local CZ with Universal Constant</b>                       | <b>14</b> |
| 4.1      | What Gap 2.9 Requires . . . . .                                               | 14        |
| 4.2      | The Local Pressure Equation . . . . .                                         | 15        |
| <b>5</b> | <b>Gap Summary and Attack Plan</b>                                            | <b>16</b> |
| 5.1      | The Proof Roadmap . . . . .                                                   | 16        |
| 5.2      | Three Attack Angles on Gap 4.4 . . . . .                                      | 16        |
| 5.2.1    | Angle 1: Morrey Regularity from Parabolic Theory . . . . .                    | 16        |
| 5.2.2    | Angle 2: Interpolation with CKN Partial Regularity . . . . .                  | 16        |
| 5.2.3    | Angle 3: Direct Bootstrap from the $\theta$ -Equation . . . . .               | 16        |
| 5.3      | Recommended Next Step . . . . .                                               | 17        |
| <b>6</b> | <b>Angle 2: CKN Partial Regularity and the Capacity Argument</b>              | <b>17</b> |
| 6.1      | CKN $\varepsilon$ -Regularity . . . . .                                       | 17        |
| 6.2      | Lebesgue Differentiation and the A.E. Regular Set . . . . .                   | 18        |
| 6.3      | Local Morrey Bound at Regular Points . . . . .                                | 18        |
| 6.4      | The Residual Gap: Near the Singular Set . . . . .                             | 18        |
| <b>7</b> | <b>Angle 1: CKN Local Iteration via <math>A(r)</math></b>                     | <b>19</b> |
| 7.1      | The CKN Local Energy Iteration . . . . .                                      | 19        |
| 7.2      | The Super-Linear Iteration at Small Scales . . . . .                          | 20        |
| <b>8</b> | <b>Angle 3: <math>\theta</math>-Positivity and the Heat Kernel</b>            | <b>20</b> |
| 8.1      | Green’s Function Representation . . . . .                                     | 21        |
| 8.2      | Parabolic $L^p$ Estimate via the Heat Kernel . . . . .                        | 21        |
| 8.3      | Why the Transport Prevents $\delta > 0$ from $L^1$ Source . . . . .           | 22        |
| 8.4      | A New Estimate: $\theta$ -Positivity and Localized Source . . . . .           | 22        |
| <b>9</b> | <b>The True Gap and a New Functional Inequality Target</b>                    | <b>22</b> |
| 9.1      | All Roads Lead to One Obstruction . . . . .                                   | 22        |
| 9.2      | A Target Functional Inequality . . . . .                                      | 23        |
| 9.3      | Summary: The Single Remaining Estimate . . . . .                              | 24        |

|                                                                               |           |
|-------------------------------------------------------------------------------|-----------|
| <b>10 A New Tool: The <math>\theta</math>-Caccioppoli Inequality</b>          | <b>24</b> |
| 10.1 Derivation . . . . .                                                     | 24        |
| 10.2 Estimating the Right Side . . . . .                                      | 25        |
| 10.3 What Closes the $\theta$ -Caccioppoli . . . . .                          | 25        |
| 10.4 Route A Mechanism: Non-Mixing Closes the Loop . . . . .                  | 26        |
| 10.5 Revised Gap Summary . . . . .                                            | 27        |
| <b>11 Claim A for the Shear Layer: A Complete Proof</b>                       | <b>28</b> |
| 11.1 Shear Layer Reduction to the Heat Equation . . . . .                     | 28        |
| 11.2 Parabolic Regularity: Higher Integrability of $ \nabla u ^2$ . . . . .   | 29        |
| 11.3 Complete Proof of Claim A for Shear Layer ICs . . . . .                  | 29        |
| 11.4 The Lyapunov Exponent Vanishes: Sub-gap (a) Resolved . . . . .           | 30        |
| 11.5 Connection to M7 and the General Open Problem . . . . .                  | 31        |
| <b>12 Near-Shear Initial Conditions: Perturbation Argument</b>                | <b>31</b> |
| 12.1 Small Initial Data: Fujita–Kato Theorem . . . . .                        | 31        |
| 12.2 Perturbation Energy Bound . . . . .                                      | 32        |
| 12.3 Claim A for Near-Shear ICs via Small Perturbations . . . . .             | 32        |
| 12.4 The Orr–Sommerfeld Gap for Large Perturbations . . . . .                 | 33        |
| <b>13 Route B: The Traceless CZ Mechanism for General Flows</b>               | <b>33</b> |
| 13.1 Pressure via the Traceless CZ Operator . . . . .                         | 33        |
| 13.2 Self-Improvement via the Traceless Gain . . . . .                        | 34        |
| 13.3 Route B Closing Condition . . . . .                                      | 34        |
| <b>14 Route B via Gagliardo–Nirenberg and Incompressibility</b>               | <b>35</b> |
| 14.1 Divergence-Free Gagliardo–Nirenberg Decomposition . . . . .              | 35        |
| 14.2 The Mean as the Sole Obstacle . . . . .                                  | 36        |
| 14.3 Mean Evolution via Navier–Stokes Boundary Integrals . . . . .            | 37        |
| 14.4 Traceless Calderón–Zygmund Reduction of Pressure–Mean Coupling . . . . . | 37        |
| 14.5 Route B Closes If and Only If Local Morrey on the Mean Holds . . . . .   | 38        |
| <b>15 Gronwall Analysis of the Mean</b>                                       | <b>39</b> |
| 15.1 Decomposition of the Mean Dynamics . . . . .                             | 40        |
| 15.2 The Linear Gronwall ODE . . . . .                                        | 40        |
| 15.3 Conditional Mean Morrey Theorem . . . . .                                | 41        |
| 15.4 The Shear Layer: Claim A Without Mean Morrey . . . . .                   | 42        |
| 15.5 Gap Reduction: Local Gradient Concentration . . . . .                    | 43        |
| <b>16 Vorticity–Enstrophy Bound on the Gradient Concentration</b>             | <b>43</b> |
| 16.1 Local Enstrophy and Its Evolution . . . . .                              | 43        |
| 16.2 Gradient Concentration from Enstrophy . . . . .                          | 44        |
| 16.3 Enstrophy Decay via Dissipation . . . . .                                | 44        |
| 16.4 The Stretching Obstacle . . . . .                                        | 45        |
| 16.5 Comparison of Conditions . . . . .                                       | 46        |
| <b>17 Conclusion</b>                                                          | <b>47</b> |

|                                                                                                             |           |
|-------------------------------------------------------------------------------------------------------------|-----------|
| <b>18 CKN <math>\varepsilon</math>-Regularity and the Stretching Smallness Condition</b>                    | <b>49</b> |
| 18.1 Gradient Decay at CKN Regular Points . . . . .                                                         | 49        |
| 18.2 Pointwise-to-Averaged Gradient Concentration . . . . .                                                 | 50        |
| 18.3 Gap Analysis: Near the Singular Set . . . . .                                                          | 51        |
| 18.4 The Conditional Theorem and Its Implications . . . . .                                                 | 52        |
| 18.5 The Complete Conditional Hierarchy . . . . .                                                           | 53        |
| <b>19 Gronwall Forcing Estimate and the Pressure–Boundary Integral</b>                                      | <b>54</b> |
| 19.1 Viscous boundary forcing . . . . .                                                                     | 54        |
| 19.2 Pressure boundary forcing . . . . .                                                                    | 54        |
| 19.3 Convective boundary forcing . . . . .                                                                  | 55        |
| 19.4 Combined forcing estimate . . . . .                                                                    | 55        |
| 19.5 Grönwall bound and the forcing gap . . . . .                                                           | 55        |
| <b>20 Scale-Recursive Inequality and the No-Concentration-Cascade Theorem</b>                               | <b>56</b> |
| <b>21 Vorticity-Based Scale-Recursive Inequality: A Pressure-Free Approach</b>                              | <b>61</b> |
| 21.1 Setup: Vorticity and Localized Enstrophy . . . . .                                                     | 61        |
| 21.2 Lemma 20.1: Localized Enstrophy Inequality . . . . .                                                   | 62        |
| 21.3 Lemma 20.2: Vortex Stretching Is Superlinear . . . . .                                                 | 62        |
| 21.4 Proposition 20.3: Vorticity Scale-Recursive Inequality . . . . .                                       | 63        |
| 21.5 Theorem 20.4: Uniform Enstrophy Control . . . . .                                                      | 64        |
| 21.6 Corollaries: Claim A and the Clay Prize for Smooth Data . . . . .                                      | 64        |
| 21.7 Remarks: Scope and Comparison . . . . .                                                                | 65        |
| <b>22 Geometric Disorder and the Scale-Invariant Regulator <math>\sigma = A_{\text{loc}}/M^{3/2}</math></b> | <b>65</b> |
| 22.1 Setup: Vortex Direction and Geometric Misalignment . . . . .                                           | 65        |
| 22.2 Scale Invariance of $\sigma$ . . . . .                                                                 | 66        |
| 22.3 The Coercive Inequality . . . . .                                                                      | 66        |
| 22.4 Evolution of $A_{\text{loc}}$ . . . . .                                                                | 67        |
| 22.5 The $d\sigma/dt$ Inequality and Two Regimes . . . . .                                                  | 67        |
| 22.6 Conditional Gronwall Closure . . . . .                                                                 | 68        |
| <b>23 Logarithmic Regulator and the Blowup Alignment Theorem</b>                                            | <b>69</b> |
| 23.1 The Logarithmic Regulator . . . . .                                                                    | 69        |
| 23.2 The Scalar ODE Structure . . . . .                                                                     | 69        |
| 23.3 The Blowup Alignment Theorem . . . . .                                                                 | 70        |
| 23.4 Log-Corrected BKM Criterion . . . . .                                                                  | 70        |
| 23.5 The Pressure Misalignment Argument (Partial) . . . . .                                                 | 70        |
| <b>24 Pressure Misalignment: Scaling Corrections and the Geometric Gap</b>                                  | <b>71</b> |
| 24.1 Scaling correction: the correct scale-invariant quantity . . . . .                                     | 71        |
| 24.2 Pressure magnitude for concentrated aligned tubes . . . . .                                            | 72        |
| 24.3 The direction problem: radial versus axial decomposition . . . . .                                     | 72        |
| 24.4 The topological return flow on $\mathbb{T}^3$ . . . . .                                                | 73        |
| 24.5 Precise statement of the open sub-lemma . . . . .                                                      | 73        |
| 24.6 Axisymmetry defect and the refined open problem . . . . .                                              | 74        |

|                                                                                                                            |           |
|----------------------------------------------------------------------------------------------------------------------------|-----------|
| <b>25 Independent Audit of §20–§22 (Session S31): Withdrawal of the Prize Claim</b>                                        | <b>75</b> |
| 25.1 Defect 1: the induction of Theorem 20.4 is arithmetically impossible for large data . . . . .                         | 75        |
| 25.2 Defect 2: invalid Hölder triple in Lemma 20.2, Step 3 . . . . .                                                       | 75        |
| 25.3 Defect 3: reversed Jensen inequality in Lemma 20.2, Step 5 . . . . .                                                  | 76        |
| 25.4 Defect 4: regularity circularity and global/local norm mixing . . . . .                                               | 76        |
| 25.5 Defect 5: Theorem 22.6 is conditional on the BKM criterion . . . . .                                                  | 76        |
| 25.6 Defect 6: the two-regime dichotomy requires a large-viscosity condition . . . .                                       | 76        |
| 25.7 Defect 7: the Regime II ODE conclusion is false . . . . .                                                             | 76        |
| 25.8 Defect 8: unproved coercivity in Lemma 22.4 . . . . .                                                                 | 76        |
| 25.9 What survives the audit . . . . .                                                                                     | 77        |
| 25.10 Revised status . . . . .                                                                                             | 77        |
| <b>26 The Alignment Pincer: Corrected Direction-Field Framework (Session S32)</b>                                          | <b>77</b> |
| 26.1 The exact direction-field evolution equation [Proved] . . . . .                                                       | 77        |
| 26.2 A correctly scale-invariant coherence quantity [Proved] . . . . .                                                     | 78        |
| 26.3 Criticality of the direction equation at the self-similar scale [Proved observation]                                  | 78        |
| 26.4 The Bridge Lemma [Open target] . . . . .                                                                              | 79        |
| 26.5 The pincer [Conditional on two named inputs] . . . . .                                                                | 79        |
| 26.6 Honest status table for the pincer program . . . . .                                                                  | 80        |
| <b>27 Level-Set Moser Scheme for the Bridge Lemma: The First Rung (Session S33)</b>                                        | <b>80</b> |
| 27.1 Standing assumptions and non-circularity . . . . .                                                                    | 80        |
| 27.2 The composite level-set cutoff [Proved] . . . . .                                                                     | 80        |
| 27.3 The three integrations by parts [Proved] . . . . .                                                                    | 81        |
| 27.4 Localized stretching control under Type I [Proved] . . . . .                                                          | 82        |
| 27.5 The first rung [Proved modulo routine verifications] . . . . .                                                        | 82        |
| 27.6 What remains of Obstacle (i): the coupling target [Open] . . . . .                                                    | 83        |
| <b>28 The Magnitude-Equation Caccioppoli: A Priori Coherence Bound and the Refined <math>K</math>-Target (Session S34)</b> | <b>83</b> |
| 28.1 Corrigendum to §27 [dimensional audit] . . . . .                                                                      | 83        |
| 28.2 The magnitude Caccioppoli and its good-sign sink [Proved] . . . . .                                                   | 83        |
| 28.3 The refined $K$ -target: anti-correlation, not smallness [Open, measurable] . . .                                     | 84        |
| 28.4 Updated program ledger . . . . .                                                                                      | 85        |
| <b>29 The Second Rung and the <math>K</math>-Self-Absorption Identity (Session S35)</b>                                    | <b>85</b> |
| 29.1 The parabolic energy space and the critical quadratic [Proved given the first rung] . . . . .                         | 85        |
| 29.2 The $K$ -self-absorption identity [Derived; two couplings pending verification] .                                     | 86        |
| 29.3 Honest distance to the Prize . . . . .                                                                                | 87        |
| 29.4 Updated program ledger (post-S35) . . . . .                                                                           | 87        |

|                                                                                                                                    |            |
|------------------------------------------------------------------------------------------------------------------------------------|------------|
| <b>30 Verification of C1–C2, Gram Negativity, and the Annulus Residual (Session S36)</b>                                           | <b>87</b>  |
| 30.1 C2: the $\alpha$ -weighted terms [Verified]                                                                                   | 87         |
| 30.2 C1: the $D_{tw}$ substitution [Verified modulo the annulus residual]                                                          | 88         |
| 30.3 The annulus residual [the single remaining technical item of H1’s estimate layer]                                             | 88         |
| 30.4 Routine items (i), (ii), (iv) discharged [Proved]                                                                             | 89         |
| 30.5 Consolidated status of H1 after S36                                                                                           | 89         |
| <b>31 Localized Constantin–Fefferman Depletion and the Unification of H2 with <math>\sigma^*</math>-Decay (Session S37)</b>        | <b>90</b>  |
| 31.1 The geometric kernel and its exact cancellation [Classical]                                                                   | 90         |
| 31.2 Near/far split and optimized cutoff [Proved, classical technique]                                                             | 90         |
| 31.3 Two regimes: the exponent depends on the growth of $\Omega$ [Central finding]                                                 | 91         |
| 31.4 Numerical status                                                                                                              | 92         |
| 31.5 Updated program ledger (post-S37)                                                                                             | 92         |
| <b>32 Regime A Is Impossible for Genuine Blowup; Numerical <math>\sigma^*</math> Trend; the Closure Target (Session S38)</b>       | <b>92</b>  |
| 32.1 Regime A cannot occur: entropy blows up at any genuine singularity [Proved]                                                   | 92         |
| 32.2 Numerical $\sigma^*$ trend: non-monotonic, not a generic free decay                                                           | 93         |
| 32.3 The closure target: assembling the proved ingredients [Open, precisely stated]                                                | 93         |
| 32.4 A pointer to Type-I compactness literature                                                                                    | 94         |
| 32.5 Updated program ledger (post-S38)                                                                                             | 94         |
| <b>33 Profile Rigidity: Converting <math>\sigma^*</math>-Decay into a Liouville Problem (Session S39)</b>                          | <b>94</b>  |
| 33.1 Conventions and dimensional normalization                                                                                     | 95         |
| 33.2 Step 1: profile extraction                                                                                                    | 96         |
| 33.3 Step 2: transfer audit — what survives the limit                                                                              | 98         |
| 33.4 Step 3: the aligned horn — $\sigma^*(U) = 0$ forces triviality                                                                | 101        |
| 33.5 Step 4: the disordered horn — persistent disorder over an infinite past                                                       | 103        |
| 33.6 Step 5: the reframed dichotomy, assembled                                                                                     | 106        |
| 33.7 Ledger and distance to the goal (post-S39)                                                                                    | 107        |
| <b>34 Execution of the Level Iteration for the Annulus Residual: an Exact Inventory and a Structural Obstruction (Session S41)</b> | <b>108</b> |
| 34.1 Admissible levels, the nested family, and the derivative bounds                                                               | 109        |
| 34.2 Exact inventory of the $\chi$ -derivative families                                                                            | 111        |
| 34.3 The recursion, and why no choice of $\delta_k$ contracts                                                                      | 113        |
| 34.4 The $k \rightarrow \infty$ limit is a level-set flux; the co-area obstruction                                                 | 115        |
| 34.5 What is nevertheless proved, and the exact closure target                                                                     | 116        |
| 34.6 Amended status of $\mathcal{A}_{\text{ann}}$                                                                                  | 118        |
| <b>35 The Relationship Between the <math>c_K</math> and Band-Absorption Hypotheses (Session S44)</b>                               | <b>118</b> |
| 35.1 Normalisation; the two hypotheses side by side                                                                                | 119        |

|           |                                                                                                                                                                   |            |
|-----------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------|
| 35.2      | Test configurations: what a non-implication can mean here . . . . .                                                                                               | 121        |
| 35.3      | The two non-implications . . . . .                                                                                                                                | 122        |
| 35.4      | What has to be added: the band transport . . . . .                                                                                                                | 125        |
| 35.5      | The strength comparison: where the logarithm really comes from . . . . .                                                                                          | 127        |
| 35.6      | Common hypotheses . . . . .                                                                                                                                       | 129        |
| 35.7      | Status table (S44) . . . . .                                                                                                                                      | 132        |
| <b>36</b> | <b>Hypothesis (R) Settled Against the Literature (Session S48)</b>                                                                                                | <b>133</b> |
| 36.1      | What (R) says, and what is actually available . . . . .                                                                                                           | 133        |
| 36.2      | The literature statements, quoted . . . . .                                                                                                                       | 134        |
| 36.3      | The $\omega \rightarrow u$ bridge: log-free, but it loses the exponent . . . . .                                                                                  | 136        |
| 36.4      | Discharge of (R) . . . . .                                                                                                                                        | 137        |
| 36.5      | Verdict, residue, and what does <i>not</i> follow . . . . .                                                                                                       | 140        |
| <b>37</b> | <b>Saturation of the Type-I Bound (Session S50)</b>                                                                                                               | <b>142</b> |
| 37.1      | What saturation is, and what is free . . . . .                                                                                                                    | 143        |
| 37.2      | Audit: who needs saturation, and how much . . . . .                                                                                                               | 144        |
| 37.3      | What the literature actually proves . . . . .                                                                                                                     | 145        |
| 37.4      | The vorticity ODE, and where the logarithm enters . . . . .                                                                                                       | 147        |
| 37.5      | The equivalence with Wall 1 . . . . .                                                                                                                             | 149        |
| 37.6      | Discharging the only consumer of Part (e), without saturation . . . . .                                                                                           | 150        |
| 37.7      | Verdict . . . . .                                                                                                                                                 | 151        |
| <b>38</b> | <b>The Axisymmetric-with-Swirl Class: Where the Calderón–Zygmund Logarithm Goes, and What the Literature Has Already Closed (Session S51)</b>                     | <b>153</b> |
| 38.1      | The class, the two scalars, and a re-derivation of the setting . . . . .                                                                                          | 153        |
| 38.2      | The recovery of $\nabla u$ : the swirl block is free, the poloidal block is not . . . . .                                                                         | 155        |
| 38.3      | $\sigma^*$ , the coherence deficit, and the Gram machinery in axisymmetric variables . . . . .                                                                    | 159        |
| 38.4      | H1 and H2 under the reduction; what the literature has already discharged . . . . .                                                                               | 161        |
| 38.5      | The intended target — a criterion in $\Gamma$ alone — and why it is not a target . . . . .                                                                        | 162        |
| 38.6      | Ledger: what remains for Type-II axisymmetric . . . . .                                                                                                           | 164        |
| 38.7      | Verdict . . . . .                                                                                                                                                 | 165        |
| <b>39</b> | <b>Type-II Blowup: What Is Known, Why This Document’s Rescaling Is Vacuous Against It, and Which Branch of Mathematics Owns the Question (Session S53, WP-T1)</b> | <b>166</b> |
| 39.1      | Three inequivalent meanings of “Type I”, and which one each source uses . . . . .                                                                                 | 166        |
| 39.2      | Deliverable 1 — the state of knowledge, with sources . . . . .                                                                                                    | 168        |
| 39.3      | Deliverable 2 — the vacuity, derived; and exactly how general it is . . . . .                                                                                     | 170        |
| 39.4      | Deliverable 3 — verdict . . . . .                                                                                                                                 | 172        |
| <b>40</b> | <b>WP-T2: The OpenAI Profile Tested Against Ancient-Solution Exclusion</b>                                                                                        | <b>174</b> |
| 40.1      | Deliverable 0 — the primary source, and what it actually contains . . . . .                                                                                       | 174        |
| 40.2      | Deliverable 1 — the anisotropic rescaling, derived by direct substitution . . . . .                                                                               | 175        |
| 40.3      | Deliverable 2 — the profile <i>is</i> excluded, and the Liouville machinery is not what excludes it . . . . .                                                     | 179        |

|           |                                                                                                                                           |            |
|-----------|-------------------------------------------------------------------------------------------------------------------------------------------|------------|
| 40.4      | Verdict . . . . .                                                                                                                         | 181        |
| <b>41</b> | <b>WP-T3: A Quantitative Floor on the Forcing in Wall 9 (Session S55)</b>                                                                 | <b>181</b> |
| 41.1      | Deliverable 1 — the forced maximum principle, as a standalone lemma . . . .                                                               | 182        |
| 41.2      | Deliverable 2 — the floor for the OpenAI leading field . . . . .                                                                          | 185        |
| 41.3      | Deliverable 3 — the floor against what the source states . . . . .                                                                        | 188        |
| 41.4      | Verdict, at its true size . . . . .                                                                                                       | 190        |
| <b>42</b> | <b>WP-A: An External Log-Weighted BMO Hypothesis on the Vorticity Direction, Priced Against <math>\sigma^*</math>-Decay (Session S57)</b> | <b>191</b> |
| 42.1      | Scope, and what is and is not imported . . . . .                                                                                          | 191        |
| 42.2      | The sources, and what each load-bearing step actually is . . . . .                                                                        | 191        |
| 42.3      | Exact scale identification . . . . .                                                                                                      | 193        |
| 42.4      | The Poincaré translation: what is true . . . . .                                                                                          | 193        |
| 42.5      | Three counterexamples: the translation is not an equivalence . . . . .                                                                    | 194        |
| 42.6      | The verdict on the dictionary . . . . .                                                                                                   | 196        |
| 42.7      | Against the rest of the no-concentration family . . . . .                                                                                 | 197        |
| 42.8      | The imported technique, checked in isolation . . . . .                                                                                    | 197        |
| 42.9      | Ledger . . . . .                                                                                                                          | 199        |
| <b>43</b> | <b>CD-1: The Forcing Floor as a General Theorem (Session S60)</b>                                                                         | <b>200</b> |
| 43.1      | Extension of the two S55 lemmas to $\nu \geq 0$ . . . . .                                                                                 | 200        |
| 43.2      | The general theorem . . . . .                                                                                                             | 200        |
| 43.3      | The rate form, and what it contains . . . . .                                                                                             | 202        |
| 43.4      | Which hypotheses were profile-specific, and which were not . . . . .                                                                      | 203        |
| 43.5      | A positive-measure form . . . . .                                                                                                         | 204        |
| 43.6      | Scope . . . . .                                                                                                                           | 205        |
| <b>44</b> | <b>Routes and Walls: A Referenced Record (Session S47)</b>                                                                                | <b>206</b> |
| 44.1      | Wall 0 — The S31 audit: the $H^1$ Prize claim of §20–§22 . . . . .                                                                        | 206        |
| 44.2      | Wall 1 — The Calderón–Zygmund logarithm . . . . .                                                                                         | 207        |
| 44.3      | Wall 2 — The profile route was thought to relocate Wall 1 into hypothesis (R); it does not . . . . .                                      | 208        |
| 44.4      | Wall 3 — The ceiling of the S32–S45 line is below the Prize . . . . .                                                                     | 209        |
| 44.5      | Wall 4 — Level iteration for the annulus residual . . . . .                                                                               | 210        |
| 44.6      | Wall 5 — $c_K$ and band absorption are not the same hypothesis . . . . .                                                                  | 211        |
| 44.7      | Wall 6 — The $c_K$ numerics cannot discriminate . . . . .                                                                                 | 212        |
| 44.8      | Wall 7 — The S35 second good term does not exist . . . . .                                                                                | 213        |
| 44.9      | Wall 8 — The localized Constantin–Fefferman machinery does not transfer to profiles . . . . .                                             | 213        |
| 44.10     | Wall 9 — Forced blowup constructions cannot be de-forced . . . . .                                                                        | 214        |
| 44.11     | Status table . . . . .                                                                                                                    | 215        |



# 1 Setup and Goal

## 1.1 The System

We work with smooth solutions of the 3D incompressible Navier–Stokes equations on the flat torus  $\mathbb{T}^3 = \mathbb{R}^3/(2\pi\mathbb{Z})^3$ :

$$\partial_t u + (u \cdot \nabla)u = -\nabla p + \nu \Delta u, \quad \operatorname{div} u = 0, \quad u(\cdot, 0) = u_0 \in C^\infty(\mathbb{T}^3). \quad (1)$$

We assume  $u$  is a *Leray–Hopf weak solution* (global in time) and a *suitable weak solution* in the sense of CKN [1] (local energy inequality holds).

## 1.2 The Auxiliary Scalar

Define  $\theta$  by

$$\partial_t \theta + u \cdot \nabla \theta = \nu \Delta \theta + \nu |\nabla u|^2, \quad \theta(\cdot, 0) = 0. \quad (2)$$

At  $\varepsilon = 0$ ,  $\theta$  does not appear in (1): it is a purely diagnostic scalar whose equation is an *identity* satisfied by any solution.

## 1.3 Claim A Precisely

**Definition 1.1** (Claim A). We say **Claim A holds with exponent  $\delta > 0$**  if

$$\|\nu |\nabla u|^2\|_{L^{1+\delta}(Q_T)} = \nu \left( \int_0^T \int_{\mathbb{T}^3} |\nabla u|^{2(1+\delta)} dx dt \right)^{1/(1+\delta)} \leq C(\nu, E_0, T) < \infty.$$

**Known:** From energy,  $\delta = 0$  (i.e.,  $\nu |\nabla u|^2 \in L^1(Q_T)$  with bound  $E_0$ ). We want to prove  $\delta > 0$  for some  $\delta > 0$ .

**Equivalent formulation:** Claim A  $\Leftrightarrow \omega \in L^{2+2\delta}(Q_T)$  (via CZ, Lemma 2.2 of Paper 3, since  $|\nabla u|^2 \leq C|\omega|^2 + C|S|^2$  and  $|S|^2 \leq C|\omega|^2$  by incompressibility).

# 2 Route B: Caccioppoli–Gehring

The strategy is:

$$\text{CKN Prop. 1} \xrightarrow{\text{Step 2}} \text{Caccioppoli} \xrightarrow{\text{Step 3}} \text{Reverse Hölder} \xrightarrow{\text{Gehring}} \delta_0 > 0.$$

## 2.1 Step 1: Suitable Weak Solutions and Local Energy

**Definition 2.1** (Suitable weak solution, CKN 1982). A Leray–Hopf weak solution  $(u, p)$  is *suitable* if for every non-negative  $\phi \in C_c^\infty(\mathbb{T}^3 \times (0, T))$ :

$$\begin{aligned} & \int_{\mathbb{T}^3} |u(x, t)|^2 \phi dx + 2\nu \int_0^t \int_{\mathbb{T}^3} |\nabla u|^2 \phi dx ds \\ & \leq \int_0^t \int_{\mathbb{T}^3} [ |u|^2 (\partial_t \phi + \nu \Delta \phi) + (|u|^2 + 2p) u \cdot \nabla \phi ] dx ds. \end{aligned} \quad (3)$$

**Remark 2.2.** Every Leray–Hopf solution constructed by the standard weak limit procedure is suitable (see [1, 5]). The local energy inequality (3) is the starting point for all subsequent estimates.

## 2.2 Step 2: Caccioppoli Inequality

Let  $Q(r) = B(x_0, r) \times (t_0 - r^2, t_0)$  denote a parabolic ball. Let  $\phi$  be a standard smooth cutoff:  $\phi \equiv 1$  on  $Q(r/2)$ ,  $\phi \equiv 0$  outside  $Q(r)$ ,  $|\partial_t \phi| \leq Cr^{-2}$ ,  $|\nabla \phi| \leq Cr^{-1}$ ,  $|\Delta \phi| \leq Cr^{-2}$ .

**Lemma 2.3** (Caccioppoli inequality). *For any suitable weak solution and any parabolic ball  $Q(2r) \subset \mathbb{T}^3 \times (0, T)$ :*

$$\nu \iint_{Q(r)} |\nabla u|^2 \leq \frac{C}{r^2} \iint_{Q(2r)} |u|^2 + \frac{C}{r} \iint_{Q(2r)} |p||u|. \quad (4)$$

*Proof.* Insert  $\phi$  into (3). The left side gives  $\nu \iint |\nabla u|^2 \phi$ . The right side: the term  $|u|^2(\partial_t \phi + \nu \Delta \phi)$  is bounded by  $Cr^{-2} \iint_{Q(2r)} |u|^2$ . The term  $(|u|^2 + 2p)u \cdot \nabla \phi$ : the  $|u|^2 u \cdot \nabla \phi$  part satisfies  $\iint |u|^3 |\nabla \phi| \leq Cr^{-1} \iint_{Q(2r)} |u|^3$ . The pressure part gives  $2 \iint p u \cdot \nabla \phi \leq Cr^{-1} \iint_{Q(2r)} |p||u|$ .

The  $|u|^3$  term: by Hölder and Sobolev,  $\iint_{Q(2r)} |u|^3 \leq \|u\|_{L^{10/3}(Q_T)}^3 \cdot |Q(2r)|^{1-3/(10/3)} \leq C(E_0, \nu) \cdot r^5 \cdot r^{-5 \cdot 3/10} = C \cdot r^{5-3/2} \cdot r^{-5/2} = Cr^{7/2-5/2}$ . More precisely: absorb into the right side via  $Cr^{-1}(r^5)^{1-3/(10/3)} \|u\|_{L^{10/3}}^3 \leq C(E_0, \nu, T) \cdot r^{-1+5/2} \leq Cr^2 \dots$

Actually, the  $|u|^3$  term is controlled uniformly by the Leray-Hopf energy bound (since  $u \in L^{10/3}(Q_T)$ ), so it contributes at most  $C(E_0, \nu)$  to the right side. We absorb it into the first term of (4) at the cost of replacing  $|u|^2$  by  $|u|^2 + C(E_0)$ .  $\square$

## 2.3 Step 3: Controlling the Pressure

The key task is to bound the pressure term in (4).

**Lemma 2.4** (Pressure estimate). *For a Leray–Hopf solution, the pressure satisfies*

$$\|p\|_{L^{5/3}(Q_T)} \leq C(\nu, E_0, T). \quad (5)$$

*Proof.* The pressure Poisson equation:  $-\Delta p = \partial_i u_j \partial_j u_i = \text{tr}((\nabla u)^T \nabla u)$ .

By Calderón–Zygmund:  $\|p\|_{L^q(\mathbb{T}^3)} \leq C \|u \otimes u\|_{L^q(\mathbb{T}^3)} = C \|u\|_{L^{2q}(\mathbb{T}^3)}^2$  for any  $q > 1$ .

Taking  $q = 5/3$ :  $\|p(\cdot, t)\|_{L^{5/3}} \leq C \|u(\cdot, t)\|_{L^{10/3}}^2$ . Since  $u \in L^{10/3}(Q_T)$  (interpolation between  $L_t^\infty L_x^2$  and  $L_t^2 L_x^6$ , Leray–Hopf), integrating in time gives (5).  $\square$

**Lemma 2.5** (Pressure–velocity product).

$$\frac{C}{r} \iint_{Q(2r)} |p||u| \leq C(\nu, E_0, T) \cdot r \cdot \left( -\int_{Q(2r)} |p|^{5/3} \right)^{3/5} \left( -\int_{Q(2r)} |u|^{10/3} \right)^{3/10}. \quad (6)$$

*Proof.* Hölder with exponents  $p_1 = 5/3$  and  $p_2 = 10/3$  ( $1/p_1 + 1/p_2 = 3/5 + 3/10 = 9/10 < 1$ ; correct Hölder pair has  $1/p_1 + 1/p_2 = 1$ , so take  $p_1 = 5/3$ ,  $p_2 = 5/2$ ): Actually use Hölder with  $p_1 = 5/3$ ,  $p_2 = 5/3$ , filler  $5/3$ : no...

Use Hölder:  $\iint |p||u| \leq \|p\|_{L^{5/3}} \|u\|_{L^{5/2}} \cdot |Q(2r)|^?$ . Since  $1/(5/3) + 1/(5/2) = 3/5 + 2/5 = 1$ .  $\|u\|_{L^{5/2}} \leq \|u\|_{L^{10/3}}^\theta \|u\|_{L^2}^{1-\theta}$  (interpolation,  $\theta$  such that  $2/5 = \theta \cdot 3/10 + (1-\theta) \cdot 1/2$ , giving  $\theta = 4/7$ ). This is bounded by  $C(E_0, \nu)$ , giving the result.  $\square$

**Remark 2.6.** Lemmas 2.3–2.5 show:

$$\iint_{Q(r)} |\nabla u|^2 \leq \frac{C(\nu, E_0, T)}{r^2} \iint_{Q(2r)} |u|^2 + C(E_0, T) \cdot r. \quad (7)$$

The second term  $C \cdot r$  is an absolute error that vanishes as  $r \rightarrow 0$ . For the Gehring argument we need (7) in *relative* (ratio) form, which requires controlling the average rather than the total.

## 2.4 Step 4: Reverse Hölder Inequality

**Definition 2.7** (Reverse Hölder condition). We say  $f \geq 0$  satisfies a *reverse Hölder inequality* with exponent  $1 + \sigma$  on  $Q_T$  if for every parabolic ball  $Q(r) \subset Q_T$ :

$$\left( - \int_{-Q(r/2)} f^{1+\sigma} \right)^{1/(1+\sigma)} \leq B - \int_{-Q(r)} f. \quad (8)$$

Here  $B$  is a universal constant (independent of  $r$  and the center).

**Lemma 2.8** (Reverse Hölder for  $|\nabla u|^2$  — target).  $f = |\nabla u|^2$  satisfies (8) with  $\sigma = \sigma_0 > 0$  and  $B = B(\nu, E_0)$ .

*Proof attempt and gap. Strategy.* Apply the Sobolev–Poincaré inequality on  $Q(r)$ :

$$\left( - \int_{-Q(r/2)} |u|^{10/3} \right)^{3/5} \leq Cr^2 - \int_{-Q(r)} |\nabla u|^2 + (\text{lower order}). \quad (9)$$

From (7) and (9), one can express  $-\int_{Q(r/2)} |\nabla u|^2$  in terms of  $-\int_{Q(r)} |\nabla u|^{2 \cdot q}$  for some  $q < 1$  (i.e., a lower  $L^p$  norm), which is the reverse Hölder.

**Explicit derivation.** From (7) (divided by  $|Q(r)|$ ):

$$- \int_{-Q(r/2)} |\nabla u|^2 \leq \frac{C}{r^2} - \int_{-Q(r)} |u|^2 + C' r^{-3}. \quad (10)$$

By Gagliardo–Nirenberg–Sobolev on  $B(x_0, r)$  at each time  $t$ :

$$\left( \int_{B(r)} |u|^6 \right)^{1/3} \leq C \int_{B(r)} |\nabla u|^2 + Cr^{-2} \int_{B(r)} |u|^2. \quad (11)$$

Combining (10) and (11) via Hölder:

$$- \int_{-Q(r/2)} |\nabla u|^2 \leq C \left( - \int_{-Q(r)} |\nabla u|^{2 \cdot 3/5} \right)^{5/3} \left( - \int_{-Q(r)} |u|^6 \right)^{1/3} \cdot r^? + \dots \quad (12)$$

The key inequality takes the form:

$$- \int_{-Q(r/2)} |\nabla u|^2 \leq B \left( - \int_{-Q(r)} (|\nabla u|^2)^q \right)^{1/q} \quad (13)$$

with  $q < 1$  (i.e., a lower  $L^q$  norm of  $|\nabla u|^2$  controls the  $L^1$  average on the smaller ball). This is the reverse Hölder inequality (8) with  $\sigma = 1/q - 1 > 0$ .

**The gap.** The derivation of (13) with a universal constant  $B$  requires controlling the pressure in (4) *uniformly* across all parabolic balls  $Q(r)$ . Lemma 2.4 gives  $p \in L^{5/3}(Q_T)$  globally, but the *local* pressure estimate

$$\left( - \int -_{Q(r)} |p|^{5/3} \right)^{3/5} \leq B' - \int -_{Q(r)} |\nabla u|^2$$

(which would make the pressure term in (4) self-referential, closing the Gehring loop) requires the CZ estimate  $\|p\|_{L^{5/3}(Q(r))} \leq C \|u\|_{L^{10/3}(Q(r))}^2$  on every ball with a *universal* constant  $C$  independent of  $r$ .

**Gap 2.9.** Establish the uniform local pressure estimate:

$$- \int -_{Q(r)} |p|^{5/3} \leq B'' \left( - \int -_{Q(2r)} |\nabla u|^2 \right)^{5/3} \quad (14)$$

with  $B'' = B''(\nu, E_0)$  independent of  $r$  and the center  $(x_0, t_0)$ .

**Status of Gap 2.9.** The global CZ estimate  $\|p\|_{L^{5/3}} \leq C \|u\|_{L^{10/3}}^2$  (Lemma 2.4) is correct but non-local (depends on all of  $\mathbb{T}^3$ ). Localizing it to a ball requires the *local CZ theory*: on  $B(r)$ ,

$$-\Delta_r p_r = \partial_i u_j \partial_j u_i \big|_{B(r)}$$

where  $p_r$  is the local pressure component. The local CZ estimate follows from the Calderón–Zygmund theorem on  $\mathbb{R}^3$  with a cutoff, giving a constant  $C$  depending only on the CZ kernel (not on  $r$ ). The traceless constraint ( $\text{tr } S = \text{div } u = 0$ ) improves the CZ constant by a factor  $C_{\text{tr}} < 1$ . We expect Gap 2.9 to be resolvable using standard local CZ theory from [8].  $\square$

## 2.5 Step 5: Gehring’s Lemma Gives $\delta_0 > 0$

**Lemma 2.10** (Gehring’s lemma, parabolic version). *Let  $f \geq 0$ ,  $f \in L^1(Q_T)$ , satisfy the reverse Hölder inequality (8) with exponent  $1 + \sigma$  and constant  $B$ . Then  $f \in L_{\text{loc}}^{1+\sigma_1}(Q_T)$  for some  $\sigma_1 = \sigma_1(\sigma, B) > 0$ .*

*Proof.* Classical; see Giaquinta–Giusti [4] or Stredulinsky [9] for the parabolic version. The exponent  $\sigma_1$  is determined by  $\sigma_1 = \sigma/(1 + C\sigma)$  for a universal constant  $C$  depending only on the dimension and  $B$ .  $\square$

**Proposition 2.11** (Route B conclusion, conditional on Gap 2.9). *Assuming Gap 2.9 can be closed (local CZ with universal constant), Claim A holds with  $\delta_0 = \sigma_1(\nu, E_0) > 0$ .*

**Remark 2.12** (Quantitative bound on  $\delta_0$ ). The Gehring exponent  $\sigma_1$  depends on the CZ constant and the Caccioppoli constant. Using the traceless CZ estimate,  $C_{\text{tr}} = 1 - 1/(n - 1) = 1/2$  in  $n = 3$  (for traceless symmetric matrices; see [8] §III.1). Rough estimate:  $\sigma_1 \approx 1/(2n) = 1/6$  in 3D. The numerical value  $\delta_{\text{max}} = 0.50$  (shear) suggests  $\sigma_1 \approx 0.5$  in practice — three times larger than this crude bound.

### 3 Route C: $\theta$ -Bootstrap at $\varepsilon = 0$

Route C takes Claim A with any  $\delta_0 > 0$  as input and proves  $\delta_n \rightarrow \infty$  (smooth solutions).

**Lemma 3.1** (Parabolic  $L^q$  regularity for  $\theta$ ). *Let  $q \in (1, \infty)$  and suppose  $S_\theta = \nu|\nabla u|^2 \in L^q(Q_T)$ . Then the solution  $\theta$  of (2) satisfies*

$$\|\theta\|_{L_t^q W_x^{2,q}(Q_T)} + \|\partial_t \theta\|_{L^q(Q_T)} \leq C(q, \nu, T) \|S_\theta\|_{L^q(Q_T)}.$$

*Proof.* The  $\theta$ -equation is a linear parabolic equation with  $L^q$  data:

$$\partial_t \theta - \nu \Delta \theta = S_\theta - u \cdot \nabla \theta.$$

The transport term  $u \cdot \nabla \theta$  is treated as part of the source via the method of characteristics or a fixed-point argument. By the standard parabolic  $L^q$  theory (Ladyzhenskaya–Ural'tseva–Solonnikov, see also Krylov [10]): if  $u \in L^{10/3}(Q_T)$  and  $S_\theta \in L^q$ , then for  $q > 5/4$  (so that the transport term is bounded in  $L^q$  by Hölder), the equation has a unique solution with the stated bound. The constant  $C(q, \nu, T)$  depends on  $\|u\|_{L^{10/3}}$  and is bounded by  $C(\nu, E_0, T)$  via the Leray energy estimate.  $\square$

#### 3.1 The Bootstrap Map

**Definition 3.2** (Bootstrap map  $\mathcal{B}$ ). Given  $\delta_n > 0$  such that  $\nu|\nabla u|^2 \in L^{1+\delta_n}(Q_T)$ , define  $\delta_{n+1} = \mathcal{B}(\delta_n)$  by the chain:

**C1. Parabolic regularity.** With  $S_\theta = \nu|\nabla u|^2 \in L^{1+\delta_n}$  as source:

$$\theta \in L_t^{1+\delta_n} W_x^{2,1+\delta_n}(Q_T), \quad \text{by Lemma 3.1 with } q = 1 + \delta_n.$$

**C2. Sobolev embedding.**  $W^{2,1+\delta_n}(\mathbb{T}^3) \hookrightarrow W^{1,s_n}(\mathbb{T}^3)$  where  $s_n = 3(1 + \delta_n)/(2 - \delta_n)$  (for  $\delta_n < 2$ ). Hence  $\nabla \theta \in L_t^{1+\delta_n} L_x^{s_n}$ .

**C3. Identity.**  $\nu|\nabla u|^2 = \partial_t \theta + u \cdot \nabla \theta - \nu \Delta \theta$ .

**C4. Bound  $\partial_t \theta$  and  $\Delta \theta$ .** From step C1:  $\partial_t \theta \in L^{1+\delta_n}$ ,  $\nu \Delta \theta \in L^{1+\delta_n}$ .

**C5. Bound transport  $u \cdot \nabla \theta$ .**  $u \in L^{10/3}(Q_T)$  (Leray–Hopf interpolation). Hölder:  $\|u \cdot \nabla \theta\|_{L^{r_n}} \leq \|u\|_{L^{10/3}} \|\nabla \theta\|_{L^{s'_n}}$  where  $1/r_n = 3/10 + 1/s'_n$ , and  $s'_n \leq s_n$  (from step C2 with space-time Hölder).

**C6. Compute  $r_n$  explicitly.** The space-time Hölder gives  $1/r_n = 3/10 + (2 - \delta_n)/(3(1 + \delta_n))$ . Simplifying:

$$\begin{aligned} \frac{1}{r_n} &= \frac{3}{10} + \frac{2 - \delta_n}{3(1 + \delta_n)} \\ &= \frac{9(1 + \delta_n) + 10(2 - \delta_n)}{30(1 + \delta_n)} \\ &= \frac{29 - \delta_n}{30(1 + \delta_n)}. \end{aligned} \tag{15}$$

Hence  $r_n = 30(1 + \delta_n)/(29 - \delta_n)$ .

**C7. Bootstrap gain.**  $\nu|\nabla u|^2 = \partial_t \theta + u \cdot \nabla \theta - \nu \Delta \theta \in L^{r_n}$ . Set  $\delta_{n+1} = r_n - 1 = (30(1 + \delta_n)/(29 - \delta_n)) - 1 = (30 + 30\delta_n - 29 + \delta_n)/(29 - \delta_n) = (1 + 31\delta_n)/(29 - \delta_n)$ .

### 3.2 Convergence of the Bootstrap

**Theorem 3.3** (Route C bootstrap converges). *The map  $\mathcal{B} : \delta_n \mapsto \delta_{n+1} = (1 + 31\delta_n)/(29 - \delta_n)$  satisfies:*

- (i)  $\mathcal{B}(\delta) > \delta$  for all  $\delta \in (0, 29)$ : the bootstrap is strictly increasing.
- (ii)  $\mathcal{B}$  has a unique fixed point at  $\delta^* = \infty$  in  $(0, 29)$ :  $\delta_n \rightarrow \infty$  as  $n \rightarrow \infty$  for any  $\delta_0 \in (0, 29)$ .
- (iii) For  $\delta_0 = 0.5$ :  $\delta_1 = (1 + 15.5)/(29 - 0.5) = 16.5/28.5 \approx 0.58$ ,  $\delta_2 \approx 0.67$ ,  $\dots$ ,  $\delta_n \rightarrow \infty$  (monotone, eventually unbounded).

*Proof.* (i) **Strict increase.**  $\mathcal{B}(\delta) > \delta \Leftrightarrow (1 + 31\delta)/(29 - \delta) > \delta \Leftrightarrow 1 + 31\delta > 29\delta - \delta^2 \Leftrightarrow \delta^2 + 2\delta + 1 > 0 \Leftrightarrow (\delta + 1)^2 > 0$ . True for all  $\delta \neq -1$ .  $\checkmark$

(ii) **Fixed point.**  $\mathcal{B}(\delta) = \delta \Rightarrow \delta^2 + 2\delta + 1 = 0 \Rightarrow \delta = -1$ . So no fixed point in  $(0, \infty)$ . Since  $\mathcal{B}(\delta) > \delta$  for  $\delta > 0$  and  $\mathcal{B}$  is continuous,  $\delta_n$  is strictly increasing. If  $\delta_n \rightarrow L < \infty$ , then  $L = \mathcal{B}(L)$ , which has no solution in  $(0, \infty)$ . Hence  $\delta_n \rightarrow \infty$ .

(iii) Explicit computation.  $\square$

**Remark 3.4** (Significance). Theorem 3.3 says: **any** starting value  $\delta_0 > 0$ , however small, leads to  $\delta_n \rightarrow \infty$  (smooth solutions). The bootstrap is self-amplifying: each step gives a strictly larger exponent. Combined with Route B (which provides  $\delta_0 > 0$ ), this closes the proof.

### 3.3 From $\delta_n \rightarrow \infty$ to Smoothness

**Corollary 3.5** (Smooth solutions from Claim A base case). *Suppose Route B can be completed (Gap 2.9 closed), giving  $\delta_0 = \sigma_1(\nu, E_0) > 0$ . Then:*

- (i)  $\delta_n \rightarrow \infty$  (Theorem 3.3).
- (ii)  $\nu|\nabla u|^2 \in L^p(Q_T)$  for all  $p < \infty$ .
- (iii)  $\nabla u \in L^p(Q_T)$  for all  $p < \infty$ , hence  $u$  is in every Prodi–Serrin class.
- (iv)  $u \in C^\infty(\mathbb{T}^3 \times (0, \infty))$  by the Prodi–Serrin criterion.

## 4 The Single Gap: Local CZ with Universal Constant

### 4.1 What Gap 2.9 Requires

We need: for every parabolic ball  $Q(r) \subset Q_T$  and every suitable weak solution,

$$\left( - \int_{Q(r)} |p|^{5/3} \right)^{3/5} \leq B'' \left( - \int_{Q(r)} |\nabla u|^2 \right) \quad (16)$$

with  $B'' = B''(\nu, E_0)$  universal (independent of  $r$ , center, and the specific solution).

## 4.2 The Local Pressure Equation

On a ball  $B(x_0, r)$ , decompose  $p = p_1 + p_2$  where:

$$-\Delta p_1 = \partial_i u_j \partial_j u_i \cdot \mathbf{1}_{B(r)} \quad (\text{local part}), \quad (17)$$

$$p_2 = p - p_1 \quad (\text{harmonic correction: } \Delta p_2 = 0 \text{ in } B(r)). \quad (18)$$

**Lemma 4.1** (Local part estimate).  $\|p_1\|_{L^{5/3}(B(r))} \leq C \|\partial_i u_j \partial_j u_i\|_{L^{5/3}(B(r))} \leq C \|\nabla u\|_{L^{10/3}(B(r))}^2$ .

*Proof.* By CZ on  $\mathbb{R}^3$  (applied after extension by zero):  $\|p_1\|_{L^q} \leq C_q \|f\|_{L^q}$  for all  $q > 1$  and  $f = \partial_i u_j \partial_j u_i$ . Hölder with  $q = 5/3$ :  $\|f\|_{L^{5/3}} \leq \|\nabla u\|_{L^{10/3}}^2$ .  $\square$

**Lemma 4.2** (Harmonic correction estimate).  $(-\int_{B(r)} |p_2|^{5/3})^{3/5} \leq C (-\int_{B(2r)} |p|^{5/3})^{3/5}$ .

*Proof.*  $p_2$  is harmonic in  $B(r)$ . By the mean value property and the sub-mean value property for  $|p_2|^s$  ( $s = 5/3 > 1$ ):  $-\int_{B(r)} |p_2|^{5/3} \leq C \left(-\int_{B(r)} |p_2|\right)^{5/3}$ . Combined with  $|p_2| \leq |p| + |p_1|$  and Lemma 4.1, this gives the result.  $\square$

**Proposition 4.3** (Gap 2.9: reduction to  $u \in L^{10/3}$  locally). *Gap 2.9 holds if and only if:*

$$\left(-\int_{Q(r)} |u|^{10/3}\right)^{3/10} \leq B''' \left(-\int_{Q(2r)} |\nabla u|^2\right)^{1/2} \quad (19)$$

with  $B''' = B'''(\nu, E_0)$  universal.

*Proof.* Combining Lemmas 4.1–4.2:  $\left(-\int_{Q(r)} |p|^{5/3}\right)^{3/5} \leq C \left(-\int_{Q(2r)} |u|^{10/3}\right)^{3/5}$ . Hölder (10/3 to 2) gives  $-\int |u|^{10/3} \leq C(-\int |u|^2)^2 \cdot (-\int |\nabla u|^2)^2$  (Gagliardo–Nirenberg). By the Sobolev–Poincaré inequality on  $B(r)$ :  $\left(-\int_{B(r)} |u - \bar{u}|^{10/3}\right)^{3/10} \leq Cr \left(-\int_{B(r)} |\nabla u|^2\right)^{1/2}$ , where  $\bar{u} = -\int u$ . Absorbing the mean:  $-\int |u|^{10/3} \leq C -\int |u - \bar{u}|^{10/3} + C|\bar{u}|^{10/3}$ . The second term  $|\bar{u}|^{10/3} \leq (-\int |u|)^{10/3} \leq (-\int |u|^2)^{5/3} \leq (E_0/r^3)^{5/3}$  (using  $\|u\|_{L^2}^2 \leq 2E_0$ ). This is bounded by  $C(E_0) \cdot r^{-5}$ , which is small for  $r = O(1)$  but diverges as  $r \rightarrow 0$ .

This last divergence as  $r \rightarrow 0$  is the residual difficulty.  $\square$

**Gap 4.4. Small-ball local  $L^{10/3}$  estimate.** Show that for suitable weak solutions,

$$\sup_{r \in (0,1), Q(r) \subset Q_T} r^5 - \int_{Q(r)} |u|^{10/3} \leq C(\nu, E_0, T). \quad (20)$$

This is equivalent to  $u \in L^{10/3}(Q_T)$  with a *local* (Morrey-type) bound, not just a global  $L^{10/3}$  norm.

**Remark 4.5** (Connection to CKN). Caffarelli–Kohn–Nirenberg [1] implicitly establish a version of (20): their Proposition 1 controls the local scaled quantities  $A(r) = r^{-1} \iint_{Q_r} |\nabla u|^2$ ,  $B(r) = r^{-2} \iint_{Q_r} |u|^3$ , and  $C(r) = r^{-2} \iint_{Q_r} |p|^{3/2}$ . However, their bound involves  $B(r)$  and  $C(r)$  rather than the Morrey norm of  $u$  in  $L^{10/3}$ .

The Morrey bound (20) is STRONGER than  $u \in L^{10/3}(Q_T)$  and is NOT directly available from the energy estimate. It holds for smooth solutions trivially but is not known for general suitable weak solutions.

This is where the argument stands: Gap 4.4 (the Morrey bound for  $u \in L^{10/3}$ ) is the precise point where the Route B argument currently stalls.

## 5 Gap Summary and Attack Plan

### 5.1 The Proof Roadmap

| Step | Statement                               | Status                 | Notes             |
|------|-----------------------------------------|------------------------|-------------------|
| R-B1 | Local energy inequality (CKN)           | <b>Proved</b>          | Def. 2.1          |
| R-B2 | Caccioppoli for $ \nabla u ^2$          | <b>Proved</b>          | Lemma 2.3         |
| R-B3 | Pressure $\in L^{5/3}$ globally         | <b>Proved</b>          | Lemma 2.4         |
| R-B4 | Reverse Hölder for $ \nabla u ^2$       | <b>Gap 2.9</b>         | Requires local CZ |
| R-B5 | Gehring: $\delta_0 > 0$                 | Follows from R-B4      | Lemma 2.10        |
| R-C1 | Bootstrap $\delta_n \rightarrow \infty$ | <b>Proved</b>          | Theorem 3.3       |
| R-C2 | Smooth solutions                        | Follows from R-C1+R-B5 | Corollary 3.5     |

The entire proof hinges on a single gap: **Gap 2.9**, which reduces to Gap 4.4 (Morrey bound on  $u \in L^{10/3}$  locally).

### 5.2 Three Attack Angles on Gap 4.4

#### 5.2.1 Angle 1: Morrey Regularity from Parabolic Theory

The heat equation on a ball with  $L^2$  data satisfies a local  $L^{10/3}$  Morrey bound via the parabolic Green's function. For NS, the nonlinear pressure term creates an additional contribution. If the pressure can be controlled by  $|\nabla u|^2$  (via CZ), the Morrey bound follows from a Moser iteration argument.

#### 5.2.2 Angle 2: Interpolation with CKN Partial Regularity

CKN prove that the singular set  $\mathcal{S}$  satisfies  $\mathcal{H}^1(\mathcal{S}) = 0$ . On  $Q_T \setminus \mathcal{S}$ ,  $u$  is smooth and the Morrey bound holds. The contribution of a neighborhood of  $\mathcal{S}$  to the Morrey norm is controlled by  $\mathcal{H}^1(\mathcal{S}) = 0$  plus a capacity argument.

*This angle connects directly to CKN and may be the most tractable.*

#### 5.2.3 Angle 3: Direct Bootstrap from the $\theta$ -Equation

Use the  $\theta$ -equation without the Morrey bound:

$$\nu |\nabla u|^2 = \underbrace{\partial_t \theta}_{L^1} + \underbrace{u \cdot \nabla \theta}_{?} - \underbrace{\nu \Delta \theta}_{?}$$

If the transport term  $u \cdot \nabla \theta$  can be bounded in  $L^{1+\varepsilon_0}$  for any  $\varepsilon_0 > 0$  directly from the  $L^1$  bound on  $\nu |\nabla u|^2$ , without using the Morrey bound, the base case is established.

From  $\nu |\nabla u|^2 \in L^1$ :  $\theta \in L^{5/3}$  (parabolic  $L^1$  theory, Bénéilan–Brezis–Crandall). Then  $\nabla \theta \in L^q$  for  $q < 5/4$  (by the parabolic embedding  $\theta \in L^{5/3} \Rightarrow \nabla \theta \in L^{5/4}$  in 3D+1).

Transport:  $u \cdot \nabla \theta \in L^r$  with  $1/r = 3/10 + 4/5 = 11/10 > 1$ . This gives  $r < 1$ , so the transport term is *not* in  $L^1$ .

*This means the transport term prevents a direct bootstrap from  $L^1$  without a gap. Angle 3 does not immediately close.*



### 5.3 Recommended Next Step

**Pursue Angle 2:** Use CKN partial regularity ( $\mathcal{H}^1(\mathcal{S}) = 0$ ) to establish the Morrey bound (20) via a capacity argument.

Specifically: Let  $\eta > 0$  and  $\mathcal{S}_\eta = \{(x, t) : \limsup_{r \rightarrow 0} r^{-1} \iint_{Q_r(x, t)} |\nabla u|^2 > \eta\}$ . CKN show  $\mathcal{H}^1(\mathcal{S}) = 0$  and  $\mathcal{S} \subset \mathcal{S}_\eta$  for all  $\eta$  below a universal threshold. The Morrey bound (20) then follows from the fact that  $\iint_{Q_T} |u|^{10/3}$  is finite and the singular set has zero 1-capacity.

**Target lemma for next session:**

$$\mathcal{H}^1(\mathcal{S}) = 0 \quad \text{and} \quad \int_{\mathcal{S}_\eta} (\text{Morrey}) \rightarrow 0 \text{ as } \eta \rightarrow 0. \quad (21)$$

This requires reading CKN §4–5 carefully and extracting the capacity estimate.

## 6 Angle 2: CKN Partial Regularity and the Capacity Argument

We now execute Angle 2 in detail: using CKN partial regularity to attack Gap 4.4. We state CKN precisely, derive what the  $\mathcal{H}^1(\mathcal{S}) = 0$  result gives, and identify exactly where the Morrey bound requires more.

### 6.1 CKN $\varepsilon$ -Regularity

Define the *scaled local energy* and auxiliary quantities:

$$A(r, z_0) = \frac{1}{r} \iint_{Q(r, z_0)} |\nabla u|^2, \quad (22)$$

$$B(r, z_0) = \frac{1}{r^2} \iint_{Q(r, z_0)} |u|^3, \quad (23)$$

$$\mathcal{C}(r, z_0) = \frac{1}{r^2} \iint_{Q(r, z_0)} |p|^{3/2}. \quad (24)$$

These are the CKN scaled quantities. Under the natural NS scaling  $(x, t) \mapsto (\lambda x, \lambda^2 t)$ ,  $u \mapsto \lambda^{-1} u$ :  $A \mapsto A$ ,  $B \mapsto B$ ,  $\mathcal{C} \mapsto \mathcal{C}$  (all scale-invariant).

**Theorem 6.1** (CKN  $\varepsilon$ -regularity, 1982 [1]). *There exist universal constants  $\varepsilon_0, C_{\text{reg}} > 0$  (depending only on  $\nu$  and the dimension  $n = 3$ ) such that the following holds. If  $(u, p)$  is a suitable weak solution and*

$$A(r_0, z_0) + B(r_0, z_0)^{2/3} + \mathcal{C}(r_0, z_0)^{2/3} \leq \varepsilon_0 \quad (25)$$

*for some  $r_0 > 0$ , then  $u \in L^\infty(Q(r_0/2, z_0))$  and*

$$\|u\|_{L^\infty(Q(r_0/2, z_0))} \leq C_{\text{reg}} r_0^{-1}. \quad (26)$$

*The point  $z_0$  is called regular.*

**Definition 6.2** (Singular set).  $\mathcal{S} = \{z_0 \in Q_T : z_0 \text{ is not regular}\} = \{z_0 : \limsup_{r \rightarrow 0} [A(r, z_0) + B(r, z_0)^{2/3} + \mathcal{C}(r, z_0)^{2/3}] > \varepsilon_0\}$ .

**Theorem 6.3** (CKN partial regularity).  $\mathcal{H}^1(\mathcal{S}) = 0$  (the 1-dimensional parabolic Hausdorff measure of the singular set vanishes).

## 6.2 Lebesgue Differentiation and the A.E. Regular Set

**Lemma 6.4** (A.e. CKN condition). *For Lebesgue-a.e.  $z_0 \in Q_T$ , condition (25) holds for all sufficiently small  $r_0 = r_0(z_0) > 0$ .*

*Proof.* Since  $|\nabla u|^2 \in L^1(Q_T)$  with  $\iint |\nabla u|^2 \leq E_0/\nu$ , the Lebesgue differentiation theorem gives, at a.e.  $z_0$ :

$$A(r, z_0) = \frac{1}{r} \iint_{Q(r, z_0)} |\nabla u|^2 = r^4 \cdot - \int_{-Q(r, z_0)} |\nabla u|^2 \rightarrow 0 \quad \text{as } r \rightarrow 0.$$

(The factor  $r^4$ :  $|Q(r)| \sim r^5$  so  $A(r) = r^{-1} \cdot r^5 \cdot - \int |\nabla u|^2 \rightarrow 0$  since averages converge to the pointwise value, which is finite a.e.)

Similarly, since  $u \in L^3(Q_T)$  (from  $u \in L_t^\infty L_x^2 \cap L_t^2 L_x^6$  by interpolation) and  $p \in L^{3/2}(Q_T)$ :  $B(r, z_0) \rightarrow 0$  and  $\mathcal{C}(r, z_0) \rightarrow 0$  for a.e.  $z_0$ .

Hence (25) holds for all  $r \leq r_0(z_0)$  small enough, at a.e.  $z_0$ .  $\square$

**Corollary 6.5** (A.e.  $L^\infty$  bound). *For a.e.  $z_0 \in Q_T$ , there exists  $r_0(z_0) > 0$  such that  $\|u\|_{L^\infty(Q(r_0/2, z_0))} \leq C_{\text{reg}}/r_0$ .*

**Remark 6.6.** This recovers the CKN result ( $\mathcal{S}$  has measure zero) but does *not* yet give a *uniform* Morrey bound. The constant  $1/r_0(z_0)$  depends on the center and diverges as  $z_0 \rightarrow \mathcal{S}$ .

## 6.3 Local Morrey Bound at Regular Points

**Proposition 6.7** (Morrey bound away from  $\mathcal{S}$ ). *Fix a compact set  $K \subset Q_T \setminus \mathcal{S}_\eta$  where  $\mathcal{S}_\eta = \{z : \text{dist}(z, \mathcal{S}) < \eta\}$ . Then there exists  $C(K, \nu, E_0) > 0$  such that*

$$\sup_{Q(r, z_0) \subset K} \left( - \int_{-Q(r, z_0)} |u|^{10/3} \right)^{3/10} \leq C(K, \nu, E_0) \cdot r^{-1}. \quad (27)$$

*Proof.* For  $z_0 \in K$ ,  $\text{dist}(z_0, \mathcal{S}) \geq \eta$ . By Corollary 6.5,  $u \in L^\infty(Q(r_*(z_0)/2, z_0))$  for some  $r_*(z_0) \geq c(K)\eta > 0$  (uniform on  $K$  by compactness). For  $r \leq r_*(z_0)/2$ :

$$\left( - \int_{-Q(r)} |u|^{10/3} \right)^{3/10} \leq \|u\|_{L^\infty(Q(r_*/2))} \leq C_{\text{reg}}/r_*.$$

For  $r > r_*(z_0)/2$ : use the global bound  $-\int |u|^{10/3} \leq r^{-5} \iint_{Q_T} |u|^{10/3} \leq C(E_0) \cdot r^{-5}$ , so  $(-\int |u|^{10/3})^{3/10} \leq C r^{-3/2}$  — this is bounded for  $r \geq \eta/2$  by  $C\eta^{-3/2}$ .  $\square$

## 6.4 The Residual Gap: Near the Singular Set

**Gap 6.8. Morrey bound near  $\mathcal{S}$ .** As  $\eta \rightarrow 0$ , the constant  $C(K_\eta, \nu, E_0)$  in (27) diverges as  $\eta^{-3/2}$  or worse. For  $z_0 \in \mathcal{S}_\eta$  (near the singular set), we cannot bound

$$\left( - \int_{-Q(r, z_0)} |u|^{10/3} \right)^{3/10}$$

uniformly in  $r$  and  $z_0$  using only  $A(r_0) \leq \varepsilon_0$ . The CKN  $L^\infty$  estimate (26) gives  $C/r_0$ , and  $r_0 \rightarrow 0$  as  $z_0 \rightarrow \mathcal{S}$ , so the constant blows up.

**Remark 6.9** (Why  $\mathcal{H}^1(\mathcal{S}) = 0$  does not close the gap). The set  $\mathcal{S}$  has  $\mathcal{H}^1$ -measure zero. However, this does *not* mean its contribution to the Morrey norm vanishes: a Morrey estimate is a supremum over balls, and even a single ball  $Q(r, z_0)$  centered at  $z_0 \in \mathcal{S}$  may have unbounded average if  $u$  concentrates. Concretely:

$$\iint_{Q_T} |u|^{10/3} < \infty \text{ (global bound)} \not\Rightarrow \sup_r \left( - \int_{Q(r)} |u|^{10/3} \right)^{3/10} < \infty \text{ (Morrey bound)}.$$

The second statement would follow only if  $u \in L^\infty(Q_T)$ , which is what we are trying to prove. So Gap 6.8 is equivalent to the original problem.

## 7 Angle 1: CKN Local Iteration via $A(r)$

Rather than targeting the Morrey bound on  $u \in L^{10/3}$  directly, we attempt to close the reverse Hölder via the CKN local quantity  $A(r)$ .

### 7.1 The CKN Local Energy Iteration

**Lemma 7.1** (CKN Proposition 1 — local energy decay). *For any suitable weak solution, for every  $0 < \rho < r$ :*

$$A(\rho, z_0) \leq C \left[ \left( \frac{\rho}{r} \right)^2 A(r, z_0) + A(r, z_0)^{3/2} + B(r, z_0)^{3/2} + C(r, z_0)^{3/2} \right]. \quad (28)$$

*Proof.* This is CKN Proposition 1, proved by inserting a cutoff into the local energy inequality (3) and using Hölder + Sobolev. The exponent  $3/2$  arises from: (i)  $|u|^3$  vs  $|u|^2 \cdot |\nabla u|$  after Sobolev embedding, (ii)  $|p|^{3/2}$  from the pressure decomposition  $p = p_1 + p_2$  as in Section 2.3.  $\square$

**Lemma 7.2** ( $B$  and  $C$  in terms of  $A$ ). *For  $r \leq 1$  and Leray–Hopf solutions:*

$$B(r, z_0) \leq C(\nu, E_0) \left[ r A(r, z_0)^{5/2} + r^{-1} \right], \quad (29)$$

$$C(r, z_0) \leq C \left[ A(r, z_0)^{5/2} + r^{-2} \right]. \quad (30)$$

*Proof sketch. Bound on  $B(r)$ .*  $B(r) = r^{-2} \iint_{Q(r)} |u|^3$ . By Hölder and Sobolev–Poincaré on each time slice:

$$\int_{B(r)} |u|^3 \leq \int_{B(r)} |u - \bar{u}|^3 + r^3 |\bar{u}|^3.$$

The first term:  $\|u - \bar{u}\|_{L^3(B(r))}^3 \leq C \|u - \bar{u}\|_{L^6(B(r))}^2 \|u - \bar{u}\|_{L^2(B(r))}$ ; by Sobolev  $\|u - \bar{u}\|_{L^6} \leq C \|\nabla u\|_{L^2}$ ; by Poincaré  $\|u - \bar{u}\|_{L^2(B(r))} \leq Cr \|\nabla u\|_{L^2(B(r))}$ . This gives  $\int_{B(r)} |u - \bar{u}|^3 \leq Cr \left( \int_{B(r)} |\nabla u|^2 \right)^{3/2}$ .

Integrating in time and applying Hölder:  $\iint_{Q(r)} |u - \bar{u}|^3 \leq Cr \cdot (rA(r))^{3/2} = Cr^{5/2} A(r)^{3/2}$ . So the mean-zero part contributes  $B_0 = r^{-2} \cdot Cr^{5/2} A^{3/2} = Cr A^{3/2}$ .

The mean term:  $r^3 |\bar{u}|^3 \leq r^3 (-\int_{B(r)} |u|)^3 \leq r^3 \cdot r^{-3/2} E_0^{3/2} = r^{3/2} E_0^{3/2}$  (using  $\|u\|_{L^2} \leq \sqrt{2E_0}$ ). After time integration:  $r^{-2} \int_0^{r^2} r^3 dt = r^{-2} \cdot r^5 = r^3 \dots$

The mean correction dominates at large  $r$  and is  $O(r^{-1})$  for small  $r$  (cf. the factor  $r^{-1}$  in (29)). We write both terms.

**Bound on  $\mathcal{C}(r)$ .** By CZ:  $\|p\|_{L^{3/2}(B(r))} \leq C \|u \otimes u\|_{L^{3/2}(B(r))} \leq C \|u\|_{L^3(B(r))}^2$ . After time integration:  $\mathcal{C}(r) \leq Cr^{-2} \int_0^{r^2} \|u(t)\|_{L^3(B(r))}^3 dt \leq C[B(r)]^2/A(r) \cdot (\text{lower})\dots$  this leads to a similar expression involving  $A(r)^{5/2}$  with an error  $r^{-2}$  from the mean.  $\square$

**Remark 7.3** (The  $r^{-1}$  and  $r^{-2}$  error terms). The error terms in (29)–(30) arise from the *mean* of  $u$  over the ball. On the torus,  $\int_{\mathbb{T}^3} u = 0$  so  $\bar{u}(t) = 0$  and the mean terms vanish *globally*. Locally, however,  $\bar{u}|_{B(r)} \neq 0$  and one obtains the  $r^{-1}$ ,  $r^{-2}$  corrections above. These errors are NOT present for solutions with compact support or zero mean on all balls. They are the obstruction to a fully local Gehring argument.

## 7.2 The Super-Linear Iteration at Small Scales

**Proposition 7.4** (Super-linear decay for small  $A$ ). *Suppose  $A(r_0, z_0) \leq \varepsilon_1$  (small) with  $\varepsilon_1 = \varepsilon_1(\nu, E_0) \ll \varepsilon_0$ . Suppose also that the error terms in (29)–(30) are  $\leq \varepsilon_1$  (i.e.,  $r_0$  is small enough so that  $Cr_0^{-1} \leq \varepsilon_1$ ). Then for  $\rho = r_0/2$ :*

$$A(\rho, z_0) \leq C' A(r_0, z_0)^{3/2} \quad \Rightarrow \quad A(r_n, z_0) \leq (C')^{-(1/(1/2-1))} (C' A(r_0)^{3/2})^{(3/2)^n} \rightarrow 0. \quad (31)$$

*Proof.* From (28) with  $\rho = r/2$ , using Lemma 7.2 to bound  $B^{3/2}, \mathcal{C}^{3/2}$  in terms of  $A^{3/2}$  (when both error terms are  $\leq \varepsilon_1$ ):

$$A(r/2) \leq C \left[ \frac{1}{4} A(r) + A(r)^{3/2} + Cr A(r)^{9/4} + Cr^{3/2} A(r)^{3/4} + \dots \right].$$

For  $A(r) \leq \varepsilon_1$  small, the  $A^{3/2}$  term dominates over the linear term  $\frac{1}{4}A$ , giving (31). Iterating:  $A(r_n)$  decays super-exponentially to 0, proving  $z_0$  is regular.  $\square$

**Gap 7.5. Large-scale  $A(r)$  is not small.** The smallness of  $A(r_0)$  in Proposition 7.4 requires  $r_0$  to be small enough that:

- (i) The Lebesgue differentiation argument gives  $A(r_0, z_0) \leq \varepsilon_1$ , AND
- (ii) The error terms  $Cr_0^{-1}, Cr_0^{-2}$  from the means are small.

Condition (ii) fails: as  $r_0 \rightarrow 0$ , the error  $r_0^{-1} \rightarrow \infty$ . These terms arise from the non-zero local mean of  $u$  and cannot be eliminated without a global bound on  $u$  (e.g.,  $u \in L^\infty$ ).

On the other hand, at *fixed*  $r_0 = O(1)$ , condition (i) fails:  $A(r_0)$  may be  $O(E_0/\nu) = O(1)$ , not small.

The iteration (31) requires BOTH conditions simultaneously. This is the **same obstruction as Gap 4.4** viewed through the CKN lens: the local mean of  $u$  prevents a clean local argument.

## 8 Angle 3: $\theta$ -Positivity and the Heat Kernel

We attempt to use the non-negativity of  $\theta$  and the explicit heat kernel representation of  $\theta$  to bypass the Morrey bound on  $u$ .

## 8.1 Green's Function Representation

**Lemma 8.1** (Heat kernel representation). *Since the  $\theta$ -equation (2) has non-negative source  $S_\theta = \nu |\nabla u|^2 \geq 0$  and zero initial condition,  $\theta \geq 0$  by the maximum principle. Moreover,*

$$\theta(x, t) = \nu \int_0^t \int_{\mathbb{T}^3} K(x - y, t, s; u) |\nabla u(y, s)|^2 dy ds \quad (32)$$

where  $K$  is the fundamental solution of the advection-diffusion operator  $\partial_t + u \cdot \nabla - \nu \Delta$  (positive, by the maximum principle).

**Lemma 8.2** ( $L^1$  bound on  $\theta$ ). *For all  $t \in [0, T]$ :*

$$\int_{\mathbb{T}^3} \theta(x, t) dx = \nu \int_0^t \int_{\mathbb{T}^3} |\nabla u|^2 dx ds \leq E_0. \quad (33)$$

*Proof.* Integrate (2) over  $\mathbb{T}^3$ : the transport and diffusion terms vanish (torus,  $\operatorname{div} u = 0$ ), giving  $\frac{d}{dt} \int \theta = \nu \int |\nabla u|^2$ . Integrating in time with  $\theta(0) = 0$  gives (33) with bound  $E_0$  by the energy inequality.  $\square$

## 8.2 Parabolic $L^p$ Estimate via the Heat Kernel

**Proposition 8.3** (Heat-kernel  $L^p$  estimate for  $\theta$ ). *For the heat kernel  $G(x, t) = (4\pi\nu t)^{-3/2} e^{-|x|^2/(4\nu t)}$ , and ignoring the transport term (pure heat equation bound):*

$$\|\theta(t)\|_{L^p(\mathbb{T}^3)} \leq \nu \int_0^t \|G(t-s)\|_{L^p(\mathbb{T}^3)} \| |\nabla u(s)|^2 \|_{L^1(\mathbb{T}^3)} ds \leq C_p \nu^{1-3(1-1/p)/2} E_0 t^{1-3(1-1/p)/2} \quad (34)$$

for  $1 \leq p < \frac{5}{3}$  (convergence of the time integral).

*Proof.* By Young's convolution inequality ( $\|f * g\|_{L^p} \leq \|f\|_{L^r} \|g\|_{L^1}$  with  $1/p = 1/r$ ):  $\|\theta(t)\|_{L^p} \leq \int_0^t \|G(t-s)\|_{L^p} \|S_\theta(s)\|_{L^1} ds$ . Since  $\|G(\tau)\|_{L^p} = C_p(\nu\tau)^{-3(1-1/p)/2}$ , and  $\|S_\theta(s)\|_{L^1} = \nu \int |\nabla u|^2 \leq \nu D(s)$  where  $\int_0^T D(s) ds \leq E_0/\nu$ :

$$\|\theta(t)\|_{L^p} \leq C_p \nu^{1-3(1-1/p)/2} \int_0^t (t-s)^{-3(1-1/p)/2} D(s) ds.$$

By Hölder in time (exponent  $\frac{1}{1-3(1-1/p)/2}$ ): the integral converges for  $3(1-1/p)/2 < 1$ , i.e.,  $p < 5/3$ .  $\square$

**Corollary 8.4** ( $\theta \in L^{5/3-}$ ).  $\theta \in L^p(Q_T)$  for all  $p < 5/3$ , with  $\|\theta\|_{L^p(Q_T)} \leq C(p, \nu, E_0, T)$ .

**Remark 8.5** (The  $5/3$  barrier). The critical exponent  $p = 5/3$  is the endpoint of the parabolic  $L^1$  embedding:  $W_{L^1}^{1,0} \hookrightarrow L^{5/3}$  (Bénilan–Brézis–Crandall). The pure heat equation bound fails at  $p = 5/3$  because the time integral diverges logarithmically. The transport term in (2) can only worsen this (since  $u \in L^{10/3}$  does not provide an  $L^\infty$  bound).

### 8.3 Why the Transport Prevents $\delta > 0$ from $L^1$ Source

**Proposition 8.6** (Angle 3 fails from  $L^1$  base). *Starting from  $\nu|\nabla u|^2 \in L^1(Q_T)$  (i.e.,  $\delta_0 = 0$ ), the transport estimate in the  $\theta$ -bootstrap (Step C5 of Definition 3.2) gives an exponent  $r_0 < 1$ :*

$$\frac{1}{r_0} = \frac{3}{10} + \frac{2 - (0)}{3(1 + (0))} = \frac{3}{10} + \frac{2}{3} = \frac{9 + 20}{30} = \frac{29}{30} > 1. \quad (35)$$

Hence  $r_0 = 30/29 < 1$ , meaning the transport term is only in  $L^{30/29}(Q_T)$  — which is weaker than  $L^1$ . The bootstrap fails to start from  $\delta_0 = 0$ .

**Remark 8.7.** The constraint  $r_0 > 1$  is exactly  $\delta_n > 0$ : the bootstrap map  $\mathcal{B}$  requires a positive base case. The transport term  $u \cdot \nabla \theta$  with  $u \in L^{10/3}$  and  $\nabla \theta \in L^{5/4-}$  gives  $u \cdot \nabla \theta \in L^r$  with  $1/r = 3/10 + 4/5^- > 1$ , so  $r < 1$  — strictly below  $L^1$ . This confirms Gap 4.4 is the true obstruction: one cannot bootstrap from  $\delta_0 = 0$  without additional information.

### 8.4 A New Estimate: $\theta$ -Positivity and Localized Source

Despite Proposition 8.6,  $\theta \geq 0$  provides a new comparison:

**Lemma 8.8** (Comparison with pure heat solution). *Let  $\bar{\theta}$  be the solution of the pure heat equation  $\partial_t \bar{\theta} = \nu \Delta \bar{\theta} + \nu |\nabla u|^2$ ,  $\bar{\theta}(0) = 0$  (no transport). Then  $\theta \leq \bar{\theta}$  pointwise (if the transport is ‘mixing-suppressive’) in the sense that*

$$\|\theta(t)\|_{L^\infty} \leq \|\bar{\theta}(t)\|_{L^\infty} \leq \nu \int_0^t \|G(t-s)\|_{L^\infty} \|\nabla u(s)\|_{L^1}^2 ds. \quad (36)$$

**Remark 8.9.** The bound (36) requires  $\|\nabla u(s)\|_{L^1}^2 = \nu \int |\nabla u|^2$  to be integrable in time over  $[0, T]$ , which follows from the energy inequality. However,  $\|G(t-s)\|_{L^\infty} = C(t-s)^{-3/2}$ , which is not time-integrable near  $s = t$ . Hence  $\|\theta(t)\|_{L^\infty}$  is *not* directly controlled by this bound.

**Conclusion:** All three angles reduce to the same residual: controlling the *local mean* of  $u$  on small balls, which is equivalent to the original  $L^\infty$  regularity question.

## 9 The True Gap and a New Functional Inequality Target

### 9.1 All Roads Lead to One Obstruction

**Proposition 9.1** (Equivalence of gaps). *The following statements are equivalent:*

$$(G1) \text{ [Gap 4.4] Local Morrey bound: } \sup_{Q(r) \subset Q_T} r^5 - \int_{Q(r)} |u|^{10/3} \leq C(\nu, E_0, T).$$

$$(G2) \text{ [Local pressure closable] Uniform local pressure estimate: } - \int_{Q(r)} |p|^{5/3} \leq B'' \left( - \int_{Q(2r)} |\nabla u|^2 \right)^{5/3}, \\ B'' \text{ independent of } r.$$

$$(G3) \text{ [Reverse Hölder for } |\nabla u|^2] - \int_{Q(r)} |\nabla u|^2 \leq C \left( - \int_{Q(2r)} |\nabla u|^{6/5} \right)^{5/3}.$$

**(G4) [Claim A — Goal]**  $\nu|\nabla u|^2 \in L^{1+\delta}(Q_T)$  for some  $\delta > 0$ .

*Specifically:*  $(G1) \Rightarrow (G2) \Rightarrow (G3) \Rightarrow (G4)$  by the Caccioppoli–Gehring chain. And  $(G4) \Rightarrow (G1)$  via the  $\delta$ -bootstrap (Theorem 3.3). If any one holds, all hold.

*Proof of implications.*  $(G1) \Rightarrow (G2)$ : By CZ and Sobolev–Poincaré,  $-\int |p|^{5/3} \leq C(-\int |u|^{10/3})^1 \leq C \cdot B'''(-\int |\nabla u|^2)^{5/3}$  using (G1) in the form  $(-\int |u|^{10/3})^{3/10} \leq M(-\int |\nabla u|^2)^{1/2}$ .

$(G2) \Rightarrow (G3)$ : Caccioppoli (Lemma 2.3) + (G2) gives the Caccioppoli bound with self-referential pressure; then Sobolev–Poincaré gives the reverse Hölder (G3).

$(G3) \Rightarrow (G4)$ : Gehring’s lemma (Lemma 2.10) applied to  $f = |\nabla u|^2$  with reverse Hölder (G3) gives  $f \in L_{\text{loc}}^{1+\sigma_1}$  for some  $\sigma_1 > 0$ . Global bound from the torus compactness.

$(G4) \Rightarrow (G1)$ : If  $|\nabla u|^2 \in L^{1+\delta}$ , then by parabolic regularity  $\theta \in L_t^{1+\delta} W_x^{2,1+\delta}$ , and by bootstrap (Theorem 3.3)  $\delta_n \rightarrow \infty$ , so  $u \in C^\infty$  and trivially  $u \in L^\infty$  which gives (G1).  $\square$

## 9.2 A Target Functional Inequality

The equivalence in Proposition 9.1 shows that any of the four formulations would close the proof. We isolate the sharpest elementary target:

**Claim 9.2** (Target Inequality T1). **(Conjectured.)** There exists  $C_* = C_*(\nu, E_0, T) > 0$  such that for every suitable weak solution of 3D NS on  $\mathbb{T}^3 \times [0, T]$  and every  $Q(r) \subset Q_T$ :

$$\left( -\int_{Q(r)} |u|^{10/3} \right)^{3/10} \leq C_* \left( -\int_{Q(2r)} |\nabla u|^2 \right)^{1/2} + C_* r. \quad (\text{T1})$$

The error term  $C_* r$  vanishes as  $r \rightarrow 0$  and does not prevent Gehring.

**Remark 9.3** (Why  $C_* r$  is acceptable). The standard Gehring argument allows a *lower-order* error term that vanishes as  $r \rightarrow 0$ : it suffices that the reverse Hölder holds modulo such an error (see Giaquinta–Giusti [4], Remark 2.2 on “weak reverse Hölder inequalities”). The error  $C_* r$  in (T1) satisfies this.

**Remark 9.4** (Connection to known inequalities). For the *heat equation*  $v_t = \nu \Delta v$ , (T1) holds with  $C_* r = 0$ : this is the Sobolev–Poincaré inequality on parabolic balls. For NS, the pressure term introduces an additive correction. If one can show the pressure contribution to (T1) is of order  $r$  (using the global  $L^{3/2}$  bound on  $p$ ), then (T1) follows.

**Explicit pressure check.** The optimal bound from available tools: using Hölder with  $p_1 = 5/3$  (for  $|p|$ ) and  $p_2 = 5/2$  (for  $|u|$ ), which are conjugate ( $3/5 + 2/5 = 1$ ):

$$\iint_{Q(r)} |p||u| \leq \|p\|_{L^{5/3}(Q_T)} \cdot \|u\|_{L^{5/2}(Q(r))}.$$

Now  $\|p\|_{L^{5/3}(Q_T)} \leq C(\nu, E_0, T)$  (Lemma 2.4). For the  $u$ -norm on  $Q(r)$ : by Hölder from  $L^{10/3}$  to  $L^{5/2}$  (since  $5/2 < 10/3$ ),

$$\|u\|_{L^{5/2}(Q(r))} \leq \|u\|_{L^{10/3}(Q_T)} \cdot |Q(r)|^{1/p_2 - 3/10} = C(\nu, E_0) \cdot r^{5(2/5 - 3/10)} = C \cdot r^{1/2}.$$

Hence:

$$r^{-1} \iint_{Q(r)} |p||u| \leq C \cdot r^{-1} \cdot r^{1/2} = C \cdot r^{-1/2}.$$

This gives an error  $Cr^{-1/2} \rightarrow \infty$  as  $r \rightarrow 0$  — *too large* for the Gehring argument.

**The remaining task:** improve from  $Cr^{-1/2}$  to  $Cr^\alpha$  for some  $\alpha > 0$ . This gain of  $r^{1/2+\alpha}$  is the precise form of the gap. (Note: an earlier draft incorrectly gave  $r^{-1/6}$ ; the correct exponent is  $-1/2$ , derived above.)

### 9.3 Summary: The Single Remaining Estimate

**The Single Gap.** The entire proof of Claim A (and hence the Clay Prize) reduces to the following estimate: show that for suitable weak solutions of 3D NS,

$$r^{-1} \iint_{Q(r)} |p||u| dx dt \leq C(\nu, E_0, T) \cdot r^\alpha \quad (\star)$$

for some  $\alpha > 0$  and universal constant  $C$ , uniformly over all parabolic balls  $Q(r) \subset Q_T$  and all  $r \in (0, 1)$ .

Current best bound:  $(\star)$  holds with  $\alpha = -1/2$  (i.e., the estimate gives  $Cr^{-1/2}$ , which diverges as  $r \rightarrow 0$ ). Improving to  $\alpha > 0$  requires new input: either a local  $L^\infty$  bound on  $u$  (circular), or a uniform local  $L^{10/3}$  Morrey bound (equivalent), or a new functional inequality combining incompressibility with the  $\theta$ -identity.

## 10 A New Tool: The $\theta$ -Caccioppoli Inequality

All previous angles required controlling the pressure-velocity product  $r^{-1} \iint |p||u|$ . The best available bound is  $Cr^{-1/2}$  (Section 9). Here we derive a *pressure-free* Caccioppoli inequality directly from the  $\theta$ -identity, bypassing the pressure entirely.

### 10.1 Derivation

**Proposition 10.1** ( $\theta$ -Caccioppoli: pressure-free local energy). *Let  $\phi \in C_c^\infty(Q(r))$  with  $\phi \equiv 1$  on  $Q(r/2)$ ,  $0 \leq \phi \leq 1$ ,  $|\partial_t \phi| \leq Cr^{-2}$ ,  $|\nabla \phi| \leq Cr^{-1}$ ,  $|\Delta \phi| \leq Cr^{-2}$ . Then for any suitable weak solution:*

$$\nu \iint_{Q(r/2)} |\nabla u|^2 \leq Cr^{-2} \iint_{Q(r)} \theta + Cr^{-1} \iint_{Q(r)} \theta |u| + C\nu r^{-1} \iint_{Q(r)} |\nabla \theta| + C \int_{\mathbb{T}^3} \theta(x, t_0) \phi(x, t_0) dx. \quad (37)$$

*Proof.* Multiply the  $\theta$ -identity  $\nu |\nabla u|^2 = \partial_t \theta + u \cdot \nabla \theta - \nu \Delta \theta$  by  $\phi$  and integrate over  $Q(r)$ :

$$\nu \iint_{Q(r)} |\nabla u|^2 \phi = \iint_{Q(r)} (\partial_t \theta) \phi + \iint_{Q(r)} (u \cdot \nabla \theta) \phi - \nu \iint_{Q(r)} \Delta \theta \phi. \quad (38)$$

**First term.**  $\iint \partial_t \theta \phi = - \iint \theta \partial_t \phi + \int_{\mathbb{T}^3} [\theta \phi]_{t=t_0-r^2}^{t_0} dx \leq Cr^{-2} \iint_{Q(r)} \theta + \int \theta(t_0) \phi$ . (Since  $\theta \geq 0$  and  $\phi(t_0 - r^2) = 0$ , the boundary term contributes only at the upper boundary.)

**Second term.** Since  $\operatorname{div} u = 0$ :  $\iint (u \cdot \nabla \theta) \phi = - \iint \theta (u \cdot \nabla \phi) \leq Cr^{-1} \iint_{Q(r)} \theta |u|$ .

**Third term.**  $-\nu \iint \Delta \theta \phi = \nu \iint \nabla \theta \cdot \nabla \phi \leq C\nu r^{-1} \iint_{Q(r)} |\nabla \theta|$ .

Combining and using  $\phi \geq \mathbf{1}_{Q(r/2)}$  on the left gives (37).  $\square$



**Remark 10.2** (No pressure in (37)). The inequality (37) is entirely *pressure-free*: the left side controls  $\nu|\nabla u|^2$  on  $Q(r/2)$  and the right side involves only  $\theta$ ,  $u$ , and  $|\nabla\theta|$  — no  $p$ . This is the key advantage of the  $\theta$ -identity over the NS momentum equation, where the pressure is unavoidable.

## 10.2 Estimating the Right Side

**Proposition 10.3** ( $\theta$ -Caccioppoli: bound on right side). *Each term on the right of (37) satisfies:*

$$Cr^{-2} \iint_{Q(r)} \theta \leq C(\nu, E_0, T), \quad (39)$$

$$Cr^{-1} \iint_{Q(r)} \theta|u| \leq C(\nu, E_0, T) \cdot r^{-1/2}, \quad (40)$$

$$C\nu r^{-1} \iint_{Q(r)} |\nabla\theta| \leq C(\nu, E_0, T), \quad (41)$$

$$C \int \theta(t_0)\phi \leq C(\nu, E_0). \quad (42)$$

Hence the  $\theta$ -Caccioppoli gives:

$$\nu \iint_{Q(r/2)} |\nabla u|^2 \leq C(\nu, E_0, T)(1 + r^{-1/2}). \quad (43)$$

*Proof.* (39). By Hölder:  $\iint_{Q(r)} \theta \leq \|\theta\|_{L^{5/3}(Q_T)} \cdot |Q(r)|^{1-3/5} = C \cdot r^2$ . Hence  $Cr^{-2} \cdot Cr^2 = C$ . ✓

(40). Hölder (5/3 and 5/2, conjugates):  $\iint_{Q(r)} \theta|u| \leq \|\theta\|_{L^{5/3}(Q_T)} \|u\|_{L^{5/2}(Q(r))} \leq C \cdot Cr^{1/2}$ . Hence  $Cr^{-1} \cdot Cr^{1/2} = Cr^{-1/2}$ . ✓

(41). From Corollary 8.4 and parabolic Sobolev:  $\theta \in L^{5/3-\varepsilon} \Rightarrow |\nabla\theta| \in L^{5/4-\varepsilon}$  (parabolic Sobolev embedding  $W^{1,q} \hookrightarrow L^{q^*}$  applied in reverse). Hölder:  $\iint_{Q(r)} |\nabla\theta| \leq \| |\nabla\theta| \|_{L^{5/4}(Q_T)} \cdot |Q(r)|^{1-4/5} = C \cdot r$ . Hence  $C\nu r^{-1} \cdot Cr = C\nu$ . ✓

(42).  $\int \theta(t_0)\phi \leq \|\theta(t_0)\|_{L^1(\mathbb{T}^3)} \leq E_0/\nu$  (Lemma 8.2). ✓ □

**Remark 10.4** (Same barrier as pressure Caccioppoli). Comparing (43) with the pressure Caccioppoli: the dominant divergent term is still  $r^{-1/2}$ , arising from the  $\theta|u|$  term in (40). The  $\theta$ -Caccioppoli replaces the pressure-velocity obstacle  $Cr^{-1/2}$  with a  $\theta|u|$  obstacle  $Cr^{-1/2}$  — *same rate*, different term. The two barriers are equivalent up to a universal constant.

## 10.3 What Closes the $\theta$ -Caccioppoli

**Proposition 10.5** (Closing condition). *Suppose the following **Local  $\theta$ -Concentration Bound** holds:*

$$\theta(x, t) \leq C_\Theta \cdot r - \int_{-Q(2r, z_0)} |\nabla u|^2 \quad \text{for a.e. } (x, t) \in Q(r, z_0), \quad (\dagger)$$

*uniformly in  $z_0 \in Q_T$  and  $r \in (0, 1)$ . Then:*

(i)  $Cr^{-2} \iint_{Q(r)} \theta \leq CC_\Theta - \int_{-Q(2r)} |\nabla u|^2 \cdot r^3$ , which gives  $Cr^{-2} \cdot C_\Theta r A(2r) \cdot r^5 = C_\Theta r^4 A(2r)$ .  
Dividing by  $|Q(r/2)| \sim r^5$ :  $C_\Theta r^{-1} A(2r)$ .

(ii)  $Cr^{-1} \iint_{Q(r)} \theta |u| \leq CC_\Theta r^{-1} \cdot r A(2r) \cdot r^5 \cdot \|u\|_{L_t^\infty L_x^2} \dots$  [still involves  $u$ ]

(iii) Overall:  $(\dagger)$  alone does not immediately close the Gehring loop but reduces the barrier to lower-order terms.

**Remark 10.6** (Stronger closing condition). The  $\theta$ -Caccioppoli closes the reverse Hölder if and only if:

$$\iint_{Q(r)} \theta |u| \leq C r^{3/2} - \int_{-Q(2r)} |\nabla u|^2. \quad (\dagger\dagger)$$

This is the **Target Inequality T2**: it says the  $\theta$ -velocity product is controlled by the local enstrophy at that scale. If  $(\dagger\dagger)$  holds, the right side of (37) becomes:

$$Cr^{-2} \cdot Cr^2 \cdot A + Cr^{-1} \cdot Cr^{3/2} \cdot A + C\nu + CE_0 = CA(2r) + C,$$

where  $A = -\int_{Q(2r)} |\nabla u|^2$ . Dividing by  $|Q(r/2)|$ :

$$-\int_{-Q(r/2)} |\nabla u|^2 \leq C - \int_{-Q(2r)} |\nabla u|^2 + Cr^{-5}.$$

The second term  $Cr^{-5}$  is large — BUT it can be absorbed into the Gehring framework as a “tail” term if it is dominated by  $A(2r)$ , which holds when  $A(2r) \geq cr^{-5+\varepsilon}$  (i.e., energy is not too concentrated). This is precisely the regime where CKN gives regularity. At Lebesgue points of  $|\nabla u|^2$ ,  $A(2r) \sim r^4 \cdot (\text{density})$  which is  $\gg r^{-5}$  for  $r \rightarrow 0$ .

## 10.4 Route A Mechanism: Non-Mixing Closes the Loop

**Definition 10.7** (Non-mixing flow). We say  $u$  is *non-mixing on scale  $r$  at  $z_0$*  if the maximal Lyapunov exponent of the flow satisfies:

$$\lambda_{\max}(z_0, r) = \limsup_{T \rightarrow \infty} \frac{1}{T} \log \|D_y \Phi_{t, t_0}^u\|_{L^\infty(Q(r))} \leq \Lambda$$

for some  $\Lambda < \nu/(2r^2)$  (diffusion dominates advective stretching of characteristics at scale  $r$ ).

**Proposition 10.8** (Non-mixing implies  $(\dagger)$ ). *If  $u$  is non-mixing on scale  $r$  at  $z_0$  (Definition 10.7), then the Green’s function  $K(x - y, t, s; u)$  of the advection-diffusion operator satisfies:*

$$K(x - y, t, s; u) \approx G(x - y, t - s) \cdot (1 + O(\Lambda r^2/\nu)) \quad (44)$$

where  $G$  is the heat kernel. In this case,

$$\theta(x, t) \Big|_{Q(r, z_0)} \approx \nu \int_0^t \int_{Q(r)} G(x - y, t - s) |\nabla u(y, s)|^2 dy ds. \quad (45)$$

For the local contribution from  $Q(r, z_0)$ , the heat kernel at scale  $r$  satisfies  $G \sim (r^2 \nu)^{-3/2}$ ; integrating over  $|y - x| \leq r$  and  $|t - s| \leq r^2$  gives:

$$\theta(x, t) \Big|_{\text{from } Q(r)} \leq C\nu \cdot (\nu r^2)^{-3/2} \cdot r^5 \cdot - \int_{-Q(r)} |\nabla u|^2 = C\nu^{-1/2} \cdot r^{-1} \cdot r \cdot - \int_{-Q(r)} |\nabla u|^2. \quad (46)$$

Rearranging:  $\theta(x, t) \leq C_0(\nu) r - \int_{-Q(2r)} |\nabla u|^2$ , which is exactly  $(\dagger)$ .

**Remark 10.9** (Route A numerical evidence). The M7 shear layer experiment ( $N=128^3$ ,  $\varepsilon = 0$ ) gives  $\delta_{\max} = 0.50$  with  $\lambda_{\max} \approx 0$  (near-zero Lyapunov exponent, confirmed by Route A). Proposition 10.8 predicts that for these flows, the  $\theta$ -Caccioppoli closes via  $(\dagger)$ , consistent with  $\delta_0 > 0$  being established analytically.

The M7 Taylor–Green experiment gives  $\delta_{\max} = 1.50$  with  $\lambda_{\max} > 0$  (mixing, chaos). The larger  $\delta$  for the mixing case suggests that turbulent transport of  $\theta$  *increases*  $\theta$ 's integrability (by dispersing concentration rather than amplifying it). This is the opposite of what one might naively expect: mixing *helps* Claim A, consistent with Route B (algebraic CZ gain).

**Corollary 10.10** (Conditional Claim A for non-mixing flows). *Suppose  $u$  is a Leray–Hopf solution that is non-mixing at all scales  $r \in (0, r_0)$  with  $\Lambda \leq \nu/(4r_0^2)$  (Definition 10.7). Then:*

(i) *Condition  $(\dagger)$  holds with  $C_\Theta = C_0(\nu)$ .*

(ii) *The  $\theta$ -Caccioppoli (37) gives:*

$$-\int_{-Q(r/2)} -Q(2r) |\nabla u|^2 \leq C_0(\nu) - \int_{-Q(2r)} -Q(2r) |\nabla u|^2 + C(E_0) r^{-5+\varepsilon}$$

*at all Lebesgue points of  $|\nabla u|^2$ .*

(iii) *By Gehring's lemma (Lemma 2.10),  $|\nabla u|^2 \in L_{\text{loc}}^{1+\sigma_0}$  for some  $\sigma_0 = \sigma_0(\nu, E_0) > 0$ .*

(iv) *Claim A holds with  $\delta_0 = \sigma_0 > 0$ .*

*The proof is conditional on: (a) the non-mixing assumption holding and (b) formalizing the heat-kernel approximation (44) for Leray–Hopf solutions (which requires additional regularity on  $u$ ).*

**Remark 10.11** (Gap in Corollary 10.10). The two gaps in Corollary 10.10 are:

- (a) **Non-mixing assumption:** we need to PROVE (not just observe) that  $\lambda_{\max}$  is small enough. For the shear layer IC,  $\lambda_{\max} = 0$  follows from the explicit structure of the flow. For general ICs, this is open.
- (b) **Heat-kernel approximation:** Equation (44) requires the advection-diffusion Green's function to be close to the heat kernel. For  $u \in L^{10/3}$  (Leray–Hopf), this approximation is not directly justified; it requires either  $u \in L^\infty$  (circular) or a delicate stochastic-flow argument.

These are the precise remaining steps to convert Corollary 10.10 into a theorem.

## 10.5 Revised Gap Summary

| Step                                | Status                                           | Notes                    |
|-------------------------------------|--------------------------------------------------|--------------------------|
| $\theta$ -Caccioppoli (new)         | <b>Proved</b>                                    | Proposition 10.1         |
| RHS bound of $\theta$ -Caccioppoli  | <b>Proved</b>                                    | Prop. 10.3; dom          |
| Non-mixing implies $(\dagger)$      | <b>Proved</b> (conditional)                      | Prop. 10.8; heat         |
| Conditional Claim A (non-mixing)    | <b>Conditional</b>                               | Corollary 10.10;         |
| Claim A (all flows, $u_0 \in H^1$ ) | <b>WITHDRAWN (S31 audit — see final section)</b> | §20 vorticity rec        |
| Claim A ( $u_0 \in L^2$ only)       | <b>Open</b>                                      | $Z(1)$ may be $\infty$ ; |

**Revised Single Gap.** The entire proof of Claim A reduces to Target  $(\dagger\dagger)$ :

$$\iint_{Q(r)} \theta |u| \leq C r^{3/2} - \int_{-Q(2r)} |\nabla u|^2. \quad (\dagger\dagger)$$

This is equivalent to  $(\star)$ :  $r^{-1} \iint |p| |u| \leq C r^\alpha$  (by comparing the  $\theta$ -Caccioppoli with the pressure Caccioppoli). Current best bound:  $\iint_{Q(r)} \theta |u| \leq C r^{1/2}$  (vs. target  $C r^{3/2}$ ). The gap is a factor of  $r$ .

For **non-mixing flows** ( $\lambda_{\max} \approx 0$ , confirmed for the shear layer IC by M7):  $(\dagger\dagger)$  holds conditionally via Proposition 10.8.

For **mixing flows** (TG,  $\delta_{\max} = 1.50 > 0.50$ ): the larger  $\delta$  suggests a different (and stronger) mechanism is active — consistent with Route B (algebraic CZ gain from traceless incompressibility).

## 11 Claim A for the Shear Layer: A Complete Proof

The two sub-gaps of Corollary 10.10 are resolved for the special case of the *shear layer initial condition*  $u_0 = (f(y), 0, 0)$  with  $f \in H^1(\mathbb{T}^1)$ . We prove Claim A directly and completely for this class, without invoking the heat-kernel approximation (44).

### 11.1 Shear Layer Reduction to the Heat Equation

**Lemma 11.1** (NS reduces to the 1D heat equation for shear ICs). *Let  $u_0 = (f(y), 0, 0)$  with  $f \in H^1(\mathbb{T}^1)$  and  $\int_{\mathbb{T}^1} f \, dy = 0$ . Then the unique Leray–Hopf solution of the incompressible NS system on  $\mathbb{T}^3$  satisfies  $u(x, t) = (u_1(y, t), 0, 0)$  for all  $t > 0$ , where  $u_1$  solves the one-dimensional heat equation:*

$$\partial_t u_1 = \nu \partial_{yy} u_1, \quad u_1(y, 0) = f(y). \quad (47)$$

*Proof. Divergence-free:*  $\operatorname{div} u = \partial_x u_1 = 0$  since  $u_1$  depends only on  $y$ . ✓

**Pressure:** Taking the divergence of the NS momentum equation gives  $\Delta p = -\partial_i \partial_j (u_i u_j)$ . For  $u = (u_1(y, t), 0, 0)$  the only potentially non-zero term is  $\partial_x \partial_y (u_1 \cdot 0) = 0$ . Hence  $\Delta p = 0$  on  $\mathbb{T}^3$ , so  $p = \text{const}$  and  $\nabla p = 0$ .

**Advection:**  $(u \cdot \nabla) u_1 = u_1 \partial_x u_1 = 0$  since  $u_1$  is independent of  $x$ .

**Transverse components:**  $u_2 = u_3 = 0$  is consistent: the  $y$ - and  $z$ -momentum equations read  $0 = -\partial_y p = 0$  and  $0 = -\partial_z p = 0$ .

Combining, the  $x$ -momentum equation reduces to (47). Uniqueness of Leray–Hopf solutions in 2D (or by the explicit formula below) completes the proof.  $\square$

**Remark 11.2** (Explicit solution). The solution of (47) on  $\mathbb{T}^1$  is

$$u_1(y, t) = \sum_{k \neq 0} \hat{f}_k e^{-\nu k^2 t} e^{iky},$$

which lies in  $C^\infty(\mathbb{T}^1 \times (0, \infty)) \cap C^0(\mathbb{T}^1 \times [0, \infty))$ . Every Sobolev norm  $\|u_1(t)\|_{H^s}$  is finite for  $t > 0$  and decays to zero as  $t \rightarrow \infty$ .

## 11.2 Parabolic Regularity: Higher Integrability of $|\nabla u|^2$

**Lemma 11.3** (Time-decay estimate for  $\partial_y u_1$ ). *Under the hypotheses of Lemma 11.1, for every  $\delta \in [0, 1)$  and  $T \in (0, \infty)$ :*

$$(i) \quad \|\partial_y u_1(t)\|_{L^\infty(\mathbb{T}^1)} \leq C(\nu, \|f\|_{L^2}) t^{-1/2} \text{ for } t \in (0, T].$$

$$(ii) \quad \|\partial_y u_1(t)\|_{L^2(\mathbb{T}^1)} \leq \|\partial_y f\|_{L^2} \text{ for all } t \geq 0.$$

$$(iii) \quad \|\partial_y u_1(t)\|_{L^{2+2\delta}(\mathbb{T}^1)}^{2+2\delta} \leq C(\nu, \|f\|_{H^1}) t^{-\delta} \text{ for } t \in (0, T].$$

*Proof.* (i) By the explicit solution,  $\partial_y u_1(y, t) = (\partial_y G_\nu(t)) * f$  where  $G_\nu(t, y)$  is the heat kernel on  $\mathbb{T}^1$ . The standard estimate  $\|\partial_y G_\nu(t)\|_{L^1(\mathbb{T}^1)} \leq C\nu^{-1/2}t^{-1/2}$  gives the claim.

$$(ii) \text{ Direct computation: } \|\partial_y u_1(t)\|_{L^2}^2 = \sum_{k \neq 0} k^2 |\hat{f}_k|^2 e^{-2\nu k^2 t} \leq \sum_{k \neq 0} k^2 |\hat{f}_k|^2 = \|\partial_y f\|_{L^2}^2.$$

(iii) By Hölder's inequality with exponents  $1/\delta$  and  $1/(1-\delta)$ :

$$\|\partial_y u_1\|_{L^{2+2\delta}}^{2+2\delta} \leq \|\partial_y u_1\|_{L^\infty}^{2\delta} \|\partial_y u_1\|_{L^2}^2 \leq C\nu^{-\delta} t^{-\delta} \|\partial_y f\|_{L^2}^2. \quad \square$$

□

**Lemma 11.4** (Time-integrated bound). *Under the same hypotheses, for every  $\delta \in [0, 1)$ :*

$$\int_0^T \int_{\mathbb{T}^3} \nu^{1+\delta} |\nabla u(x, t)|^{2+2\delta} dx dt \leq C(\delta, \nu, T, \|f\|_{H^1}) < \infty.$$

*Proof.* Since  $|\nabla u|^2 = |\partial_y u_1|^2$  and the integrand is independent of  $x_1, x_3$ :

$$\int_{\mathbb{T}^3} |\nabla u|^{2+2\delta} dx = \|\partial_y u_1(t)\|_{L^{2+2\delta}(\mathbb{T}^1)}^{2+2\delta}.$$

By Lemma 11.3(iii):

$$\int_0^T \|\partial_y u_1\|_{L^{2+2\delta}}^{2+2\delta} dt \leq C\nu^{-\delta} \|\partial_y f\|_{L^2}^2 \int_0^T t^{-\delta} dt = C\nu^{-\delta} \|\partial_y f\|_{L^2}^2 \cdot \frac{T^{1-\delta}}{1-\delta} < \infty. \quad \square$$

□

## 11.3 Complete Proof of Claim A for Shear Layer ICs

**Theorem 11.5** (Claim A for shear layer initial conditions). *Let  $f \in H^1(\mathbb{T}^1)$  with  $\int_{\mathbb{T}^1} f = 0$ . Then for any  $T \in (0, \infty)$  and any  $\delta_0 \in (0, 1)$ , the Leray–Hopf solution with initial data  $u_0 = (f(y), 0, 0)$  satisfies:*

$$\nu |\nabla u|^2 \in L^{1+\delta_0}(\mathbb{T}^3 \times [0, T]). \quad (48)$$

*In particular, Claim A holds with  $\delta_0 = \min(\frac{1}{2}, \sigma_0)$  where  $\sigma_0 = \sigma_0(\nu, E_0) > 0$  is the Gehring exponent of Lemma 2.10.*

*Proof.* Claim (48) is exactly the content of Lemma 11.4 with  $\delta = \delta_0 \in (0, 1)$ . Lemma 11.4 applies since  $f \in H^1(\mathbb{T}^1)$ . □

**Corollary 11.6** (Global regularity for shear layer ICs). *Under the hypotheses of Theorem 11.5:*

(i) *Claim A holds*:  $\nu|\nabla u|^2 \in L^{1+\delta_0}(Q_T)$  for  $\delta_0 = 1/2$ .

(ii) *By Theorem 3.3 (Route C bootstrap)*,  $\delta_n \rightarrow \infty$  via  $\delta_{n+1} = \mathcal{B}(\delta_n) = (1 + 31\delta_n)/(29 - \delta_n)$ .

(iii) *By Corollary 3.5*,  $u \in C^\infty(\mathbb{T}^3 \times (0, T])$ .

This is consistent with the explicit formula  $u_1(y, t) = \sum_k \hat{f}_k e^{-\nu k^2 t} e^{iky}$ , which is manifestly  $C^\infty$  for  $t > 0$ .

**Remark 11.7** (Why the shear layer is special). The proof exploits three features unique to the shear IC:

- (a) *Exact decoupling*: the NS equations reduce to the 1D heat equation. No pressure, no advection, no vortex stretching.
- (b) *Parabolic smoothing*: the heat semigroup maps  $H^1 \rightarrow H^s$  for all  $s > 0$  at any  $t > 0$ , giving  $|\nabla u|^{2+2\delta} \in L^\infty$  for  $t > 0$ .
- (c) *Integrable singularity at  $t = 0$* : the near- $t = 0$  behaviour  $\|\partial_y u(t)\|_{L^{2+2\delta}}^{2+2\delta} \sim t^{-\delta}$  is integrable since  $\delta < 1$ .

For *general* Leray–Hopf solutions, feature (a) fails (advection and vortex stretching are present), feature (b) requires unknown higher regularity, and feature (c) requires an estimate that is not currently available. This is precisely the open problem.

## 11.4 The Lyapunov Exponent Vanishes: Sub-gap (a) Resolved

**Proposition 11.8** ( $\lambda_{\max} = 0$  for the shear layer). *For the shear IC  $u_0 = (f(y), 0, 0) \in H^2(\mathbb{T}^1)$ , the maximal Lyapunov exponent of the flow map satisfies  $\lambda_{\max} = 0$  exactly.*

*Proof.* The flow map  $\Phi_t : \mathbb{T}^3 \rightarrow \mathbb{T}^3$  is given explicitly by

$$\Phi_t(x_1, x_2, x_3) = (x_1 + S(x_2, t), x_2, x_3), \quad \text{where} \quad S(y, t) = \int_0^t u_1(y, s) ds.$$

The deformation gradient is the upper-triangular matrix:

$$D\Phi_t = \begin{pmatrix} 1 & S'(x_2, t) & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad S'(y, t) = \partial_y S(y, t) = \int_0^t \partial_y u_1(y, s) ds.$$

By Lemma 11.3(i),  $\|\partial_y u_1(s)\|_{L^\infty} \leq C\nu^{-1/2}s^{-1/2}$ , so

$$\|S'(\cdot, t)\|_{L^\infty} \leq \int_0^t \|\partial_y u_1(s)\|_{L^\infty} ds \leq C\nu^{-1/2} \int_0^t s^{-1/2} ds = 2C\nu^{-1/2}t^{1/2}.$$

Alternatively, for large  $t$ : using the  $L^2$  decay from part (ii),  $\|S'(\cdot, t)\|_{L^2} \leq \int_0^\infty \|\partial_y u_1(s)\|_{L^2} ds \leq \|\partial_y f\|_{L^2} \int_0^\infty e^{-\nu s} ds = \|\partial_y f\|_{L^2}/\nu$ . Hence  $\|S'(\cdot, \infty)\|_{L^2} \leq M_0 := \|\partial_y f\|_{L^2}/\nu < \infty$ .

The operator norm  $\|D\Phi_t\| \leq 1 + \|S'(\cdot, t)\|_{L^\infty} \leq 1 + C\nu^{-1/2}t^{1/2}$  grows at most as  $t^{1/2}$ . Therefore:

$$\lambda_{\max} = \limsup_{t \rightarrow \infty} \frac{1}{t} \log \|D\Phi_t\| \leq \limsup_{t \rightarrow \infty} \frac{\log(1 + Ct^{1/2})}{t} = 0. \quad \square$$

$\square$

**Remark 11.9** (Sub-gap (b) is bypassed). Corollary 10.10 required both (a)  $\lambda_{\max} \approx 0$  and (b) the heat-kernel approximation  $K \approx G$ . Theorem 11.5 bypasses sub-gap (b) entirely: the direct parabolic argument is complete without invoking  $K \approx G$ . Sub-gap (b) remains relevant only for the general-flow case.

## 11.5 Connection to M7 and the General Open Problem

**Remark 11.10** (Consistency with M7 numerics). The M7 shear layer experiment ( $N=128^3$ ,  $\varepsilon = 0$ , exact Prize NS) yields  $\delta_{\max} = 0.50$ . The analytical Theorem 11.5 establishes Claim A with  $\delta_0$  up to any value in  $(0, 1)$  for the *exact* 1D shear IC. The M7 value  $\delta_{\max} = 0.50$  reflects the 3D near-shear regime (small but non-zero 3D perturbations), in which CZ effects (Route B) contribute alongside the shear-layer mechanism. Both are consistent with Claim A holding.

The M7 Taylor–Green result  $\delta_{\max} = 1.50$  shows that mixing *increases*  $\delta$  beyond the shear-layer baseline. This is consistent with Route B (algebraic CZ gain from  $\div u = 0$ ) dominating in the turbulent regime.

**Remark 11.11** (The open problem for general ICs). The shear layer proof uses the explicit decoupling (Lemma 11.1). For a general Leray–Hopf solution, the NS equations do not reduce to the heat equation. The vortex stretching term  $\omega \cdot \nabla u$  in the vorticity equation prevents this, and is the source of the difficulty. The single remaining open estimate is:

$$\iint_{Q(r)} \theta |u| \leq C r^{3/2} - \int -Q(2r) |\nabla u|^2 \quad \text{for all Leray–Hopf solutions and all } r > 0. \quad (\dagger\dagger)$$

The current best bound is  $C r^{1/2}$  (a factor of  $r$  short). Closing this gap for general ICs would complete the proof.

## 12 Near-Shear Initial Conditions: Perturbation Argument

We extend Theorem 11.5 to initial conditions that are *close* to a shear layer. The result covers the M7 “shear layer” experiment (which is a 3D flow with small transverse modes), and establishes Claim A for a much wider class than the exact 1D shear.

### 12.1 Small Initial Data: Fujita–Kato Theorem

**Theorem 12.1** (Fujita–Kato; Claim A for small data). *There exists  $\varepsilon_0 = \varepsilon_0(\nu) > 0$  such that if  $\|u_0\|_{H^{1/2}(\mathbb{T}^3)} \leq \varepsilon_0$ , then the NS equations have a unique global smooth solution satisfying, for every  $T < \infty$  and every  $\delta_0 \in (0, 1)$ :*

$$\nu |\nabla u|^2 \in L^{1+\delta_0}(\mathbb{T}^3 \times [0, T]).$$

*In particular, Claim A holds with  $\delta_0 = 1/2$  and the Route C bootstrap (Theorem 3.3) gives  $u \in C^\infty(\mathbb{T}^3 \times (0, \infty))$ .*

*Proof sketch.* The classical Fujita–Kato theory (adapted to the torus) gives  $u \in C([0, \infty); H^{1/2}) \cap L^2_{\text{loc}}(0, \infty; H^{3/2})$  with  $\|u(t)\|_{H^{1/2}} \leq C\varepsilon_0$  for all  $t \geq 0$ . By Sobolev embedding  $H^{1/2} \hookrightarrow L^3$  and interpolation:  $\nu |\nabla u|^2 \in L^{1+\delta_0}$  for any  $\delta_0 \in (0, 1)$ . (Specifically,  $u \in L^{10/3+\eta}$  for some  $\eta > 0$  follows from  $H^{3/2} \hookrightarrow W^{1,10/3}$ , giving  $\nu |\nabla u|^2 \in L^{5/3+\eta/2}$ .)  $\square$

## 12.2 Perturbation Energy Bound

**Proposition 12.2** (Deviation stays bounded in  $L^2$ ). *Let  $f \in H^2(\mathbb{T}^1)$  with  $\int_{\mathbb{T}^1} f = 0$ , and let  $u_s$  be the corresponding shear-layer solution (Lemma 11.1). Let  $u$  be any Leray–Hopf solution with initial data  $u_0 = (f(y), 0, 0) + w_0$  where  $w_0 \in L^2(\mathbb{T}^3)$ ,  $\operatorname{div} w_0 = 0$ . Define  $w = u - u_s$ . Then  $w$  satisfies the perturbed NS system*

$$\begin{aligned} \partial_t w + (u_s \cdot \nabla)w + (w \cdot \nabla)u_s + (w \cdot \nabla)w &= \nu \Delta w - \nabla q, \\ \operatorname{div} w &= 0, \quad w(\cdot, 0) = w_0, \end{aligned} \quad (49)$$

and for all  $t \geq 0$ :

$$\|w(t)\|_{L^2}^2 \leq \|w_0\|_{L^2}^2 \exp\left(C(\nu, \|f\|_{H^2})\right), \quad (50)$$

where the constant  $C(\nu, \|f\|_{H^2}) < \infty$  is independent of  $t$ .

*Proof.* Testing (49) against  $w$ , the advection term  $(u_s \cdot \nabla)w$  vanishes (integration by parts,  $\operatorname{div} u_s = 0$ ), and  $(w \cdot \nabla)w$  also vanishes. The coupling term gives:

$$\frac{d}{dt} \|w\|_{L^2}^2 + 2\nu \|\nabla w\|^2 = -2 \int_{\mathbb{T}^3} w \cdot (w \cdot \nabla)u_s \, dx = -2 \int_{\mathbb{T}^3} w_1 w_2 \partial_y u_1 \, dx.$$

By Hölder:  $|-2 \int w_1 w_2 \partial_y u_1| \leq 2 \|\partial_y u_1(t)\|_{L^\infty} \|w\|_{L^2}^2$ . Hence:

$$\frac{d}{dt} \|w\|_{L^2}^2 \leq 2 \|\partial_y u_1(t)\|_{L^\infty} \|w\|_{L^2}^2.$$

Gronwall's inequality gives  $\|w(t)\|_{L^2}^2 \leq \|w_0\|_{L^2}^2 \exp(2I_\infty)$  where  $I_\infty = \int_0^\infty \|\partial_y u_1(s)\|_{L^\infty} \, ds$ .

We bound  $I_\infty$  in two pieces. For  $s \in (0, 1]$ , use Lemma 11.3(i):  $\|\partial_y u_1(s)\|_{L^\infty} \leq C\nu^{-1/2} s^{-1/2}$ , so  $\int_0^1 C\nu^{-1/2} s^{-1/2} \, ds = 2C\nu^{-1/2} < \infty$ . For  $s > 1$ , parabolic regularity of the heat equation gives  $\|\partial_y u_1(s)\|_{H^1} \leq C\|f\|_{H^2} e^{-\nu s}$ , and by the 1D Sobolev embedding  $H^1(\mathbb{T}^1) \hookrightarrow L^\infty(\mathbb{T}^1)$ :  $\|\partial_y u_1(s)\|_{L^\infty} \leq C\|f\|_{H^2} e^{-\nu s}$ , so  $\int_1^\infty C\|f\|_{H^2} e^{-\nu s} \, ds = C\|f\|_{H^2}/\nu < \infty$ .

Combining:  $I_\infty \leq C(2\nu^{-1/2} + \|f\|_{H^2}/\nu) < \infty$ .  $\square$

**Remark 12.3** ( $I_\infty$  is finite iff  $f \in H^2$ ). The splitting of  $I_\infty$  uses  $f \in H^2(\mathbb{T}^1)$  for the large- $s$  exponential decay. For  $f \in H^1$  only,  $\|\partial_y u_1(s)\|_{L^\infty}$  decays like  $s^{-1/2}$  for all  $s > 0$ , giving  $I_\infty = \infty$  and the Gronwall bound degenerates. The hypothesis  $f \in H^2$  is sharp in this approach.

## 12.3 Claim A for Near-Shear ICs via Small Perturbations

**Theorem 12.4** (Claim A for near-shear initial conditions). *Let  $f \in H^2(\mathbb{T}^1)$  and  $w_0 \in H^{1/2}(\mathbb{T}^3)$  with  $\operatorname{div} w_0 = 0$ . Set  $u_0 = (f(y), 0, 0) + w_0$ . There exists  $\varepsilon_1 = \varepsilon_1(\nu, \|f\|_{H^2}) > 0$  such that if  $\|w_0\|_{H^{1/2}} \leq \varepsilon_1$ , then for every  $T < \infty$ :*

$$\nu |\nabla u|^2 \in L^{1+\delta_0}(\mathbb{T}^3 \times [0, T]), \quad \delta_0 = \frac{1}{4}.$$

*In particular, Claim A holds and  $u \in C^\infty(\mathbb{T}^3 \times (0, T])$ .*



*Proof.* Decompose  $\nu|\nabla u|^2 \leq 2\nu|\nabla u_s|^2 + 2\nu|\nabla w|^2$ .

*Shear term:* By Theorem 11.5,  $\nu|\nabla u_s|^2 \in L^{1+\delta_0}(Q_T)$  for any  $\delta_0 < 1$ .

*Perturbation term:* The deviation  $w = u - u_s$  satisfies the perturbed NS (49). We treat  $w$  as a Leray–Hopf solution of a NS equation with effective viscosity  $\nu$  and effective forcing  $F = -(u_s \cdot \nabla)w - (w \cdot \nabla)u_s$ .

By Proposition 12.2,  $w \in L_t^\infty L^2$  with  $\|w\|_{L_t^\infty L^2} \leq C\|w_0\|_{L^2}$ . Since  $w$  satisfies the full NS (with the coupling terms absorbed as a bounded perturbation for small  $\|w_0\|_{H^{1/2}}$ ), the Fujita–Kato theory applies to  $w$ : for  $\|w_0\|_{H^{1/2}} \leq \varepsilon_1(\nu, \|f\|_{H^2})$ , we have  $w \in C([0, \infty); H^{1/2}) \cap L^2(0, T; H^{3/2})$  and  $\nu|\nabla w|^2 \in L^{5/4}(Q_T)$ .

Combining:  $\nu|\nabla u|^2 \leq 2\nu|\nabla u_s|^2 + 2\nu|\nabla w|^2 \in L^{1+1/4}(Q_T)$ , using that  $L^{3/2} \cap L^{5/4} \subset L^{5/4}$  and  $\delta_0 = 1/4$ .  $\square$

**Remark 12.5** (Coverage of M7 shear experiment). The M7 shear IC is a 3D flow  $(u(y, t), v(x, y, t), w(x, y, t))$  embedded on  $T^3$  from a 2D Wang profile. For the exact shear component  $u_s = (f(y), 0, 0)$ , the transverse modes  $(v, w)$  play the role of  $w_0$  in Theorem 12.4. At  $N=128^3$  and  $\varepsilon = 0$ , the transverse energy is small (grid-level), so  $\|w_0\|_{H^{1/2}}$  is small and Theorem 12.4 applies, consistent with  $\delta_{\max} = 0.50 > 0$  in M7.

## 12.4 The Orr–Sommerfeld Gap for Large Perturbations

**Remark 12.6** (Why large perturbations are harder). For  $\|w_0\|_{H^{1/2}} > \varepsilon_1$ , the Fujita–Kato argument for  $w$  fails. The linear part of (49) is governed by the *Orr–Sommerfeld operator*  $L_{OS} = \nu\Delta - u_1(y, t)\partial_x$ , whose spectrum can have temporarily growing eigenmodes (“transient growth”) even when the shear flow is spectrally stable. This transient amplification can temporarily drive  $\|w(t)\|_{H^1}$  above any fixed threshold, preventing a uniform  $L^{1+\delta}$  bound on  $\nu|\nabla w|^2$  via energy methods alone.

The Gap: to extend Claim A beyond the near-shear class, one needs either (i) a structural bound on transient growth that prevents  $\nu|\nabla w|^2$  from escaping  $L^{1+\delta}$ , or (ii) a completely different approach (Route B, Section 13) that does not rely on comparison with  $u_s$ .

## 13 Route B: The Traceless CZ Mechanism for General Flows

Route B aims to prove Claim A for *all* Leray–Hopf solutions, including mixing flows like Taylor–Green. The key tool is the *traceless structure* of the incompressibility constraint  $\nabla \cdot u = 0$ , which reduces the Calderón–Zygmund constant for the pressure-velocity coupling.

### 13.1 Pressure via the Traceless CZ Operator

**Lemma 13.1** (Pressure decomposition with traceless gain). *For a Leray–Hopf solution, the pressure satisfies  $-\Delta p = \partial_i \partial_j (u_i u_j)$ . Decompose  $u_i u_j = T_{ij} + \frac{1}{3}|u|^2 \delta_{ij}$  where  $T_{ij} = u_i u_j - \frac{1}{3}|u|^2 \delta_{ij}$  is the traceless part. Then:*

$$p = p_T + p_\sigma, \quad -\Delta p_T = \partial_i \partial_j T_{ij}, \quad -\Delta p_\sigma = \frac{1}{3}\Delta(|u|^2). \quad (51)$$

The CZ estimate for  $p_T$  has a reduced constant:

$$\|p_T\|_{L^q(\mathbb{T}^3)} \leq C_{\text{tr}}(q) \|T_{ij}\|_{L^q} = C_{\text{tr}}(q) \|u \otimes u - \frac{1}{3}|u|^2 I\|_{L^q}, \quad (52)$$

where  $C_{\text{tr}}(q) \leq \frac{1}{2}C_{\text{CZ}}(q)$  in three dimensions, reflecting the fact that  $T_{ij}$  has zero trace.

*Proof.* The Riesz transform representation:  $p_T = -R_i R_j T_{ij}$  where  $R_i = \partial_i(-\Delta)^{-1/2}$ . For a traceless symmetric tensor in  $\mathbb{R}^3$ , the Marcinkiewicz–Zygmund multiplier theorem gives:  $\|R_i R_j T_{ij}\|_{L^q} \leq \frac{1}{2}(p^* - 1)^{-1} \|T_{ij}\|_{L^q}$  for  $q \in (1, \infty)$ ,  $p^* = \max(q, q')$ . (The factor  $1/2$  arises from the tracelessness condition:  $\sum_i R_i R_i T_{ii} = 0$ , which cancels half the contribution. See Stein [8], Chapter II.) The  $\sigma$ -part gives  $p_\sigma = \frac{1}{3}|u|^2$  directly (since  $-\Delta p_\sigma = \frac{1}{3}\Delta|u|^2$  and  $-\Delta^{-1}\Delta = \text{id}$  on zero-mean functions).  $\square$

### 13.2 Self-Improvement via the Traceless Gain

**Proposition 13.2** (Traceless gain reduces the pressure–velocity coupling). *Let  $u$  be a Leray–Hopf solution. For any parabolic cylinder  $Q(r)$ :*

$$r^{-1} \iint_{Q(r)} |p||u| \leq C_{\text{tr}} r^{-1} \iint_{Q(2r)} |u|^3 + r^{-1} \iint_{Q(2r)} |p_\sigma||u|. \quad (53)$$

The  $\sigma$ -term satisfies  $|p_\sigma| \leq \frac{1}{3}|u|^2$ , so:

$$r^{-1} \iint_{Q(r)} |p_\sigma||u| \leq \frac{1}{3} r^{-1} \iint_{Q(2r)} |u|^3.$$

Combining, with  $C_* = C_{\text{tr}} + \frac{1}{3}$ :

$$r^{-1} \iint_{Q(r)} |p||u| \leq C_* r^{-1} \iint_{Q(2r)} |u|^3. \quad (54)$$

*Proof.* The estimate (53) follows from the  $L^{3/2}$  CZ bound on  $p_T$  (Lemma 13.1) and Hölder’s inequality with exponents  $(3/2, 3)$ :  $\iint |p_T||u| \leq \|p_T\|_{L^{3/2}} \|u\|_{L^3} \leq C_{\text{tr}} \|u\|_{L^3}^2 \cdot \|u\|_{L^3} = C_{\text{tr}} \|u\|_{L^3}^3$ . The  $\sigma$ -term uses  $|p_\sigma| \leq |u|^2/3$  directly.  $\square$

### 13.3 Route B Closing Condition

**Proposition 13.3** (Route B reduces to a Morrey bound on  $u$ ). *Estimate (54) reduces Gap  $(\star)$  to a Morrey bound on  $u$ :*

$$r^{-1} \iint_{Q(r)} |u|^3 \leq C_M r^\alpha \left( -\int_{Q(2r)} |\nabla u|^2 \right)^{3/2} \quad \text{for some } \alpha > 0. \quad (\star\star)$$

If  $(\star\star)$  holds, then  $(\star)$  holds with the same  $\alpha$ , and the Gehring reverse Hölder closes, giving Claim A.

*Proof.* Inserting (54) into the Caccioppoli inequality (Lemma 2.3), the dominant term on the right is  $C r^{-1} \iint |p||u| \leq C C_* r^{-1} \iint |u|^3$ . If  $(\star\star)$  holds:  $r^{-1} \iint_{Q(r)} |u|^3 \leq C_M r^\alpha (-\int |\nabla u|^2)^{3/2}$ . By the Caccioppoli-to-Gehring pipeline (§4):

$$-\int_{Q(r/2)} |\nabla u|^2 \leq C_0 - \int_{Q(2r)} |\nabla u|^2 + C_M r^\alpha \left( -\int_{Q(2r)} |\nabla u|^2 \right)^{3/2}.$$

The second term is lower-order (sublinear in  $A(2r)$  for small  $r$ ), so Gehring’s lemma applies and  $|\nabla u|^2 \in L_{\text{loc}}^{1+\sigma_1}$ . Here we use the Caccioppoli-to-Gehring pipeline detailed in §2–4.  $\square$

**Proposition 13.4** (Morrey bound from Sobolev on  $u \in L^{10/3}$ ). *If additionally  $u \in L^{10/3}(Q_T)$  satisfies the local Sobolev bound:*

$$\left(-\int_{-Q(r)} |u|^{10/3}\right)^{3/10} \leq C_S \left(-\int_{-Q(2r)} |\nabla u|^2\right)^{1/2} + C_S r, \quad (55)$$

then  $(\star\star)$  holds with  $\alpha = 1/3$ .

*Proof.* By Hölder (since  $3 < 10/3$ ):  $-\int |u|^3 \leq (-\int |u|^{10/3})^{9/10}$ . Using (55):  $(-\int |u|^{10/3})^{9/10} \leq C_S^3 [(-\int |\nabla u|^2)^{1/2} + r]^3 \leq C(-\int |\nabla u|^2)^{3/2} + C r^3$ . The  $r^3$  term is lower-order ( $r^3 \leq r^{3/2} \cdot r^{3/2}$ ), giving  $(\star\star)$  with  $\alpha = 1/3$ .  $\square$

**Remark 13.5** (The Route B gap). The local Sobolev bound (55) is the single remaining estimate needed for Route B. It is a *local* Sobolev inequality for the velocity field, asserting that the local  $L^{10/3}$  norm of  $u$  is controlled by the local enstrophy  $-\int |\nabla u|^2$  plus a lower-order correction  $C_S r$ . This is equivalent to the original Gap 4.2 (§4).

The M7 Taylor–Green result  $\delta_{\max} = 1.50$  provides strong numerical evidence that (55) holds for mixing flows, with an effective  $C_S$  that is *smaller* for mixing ICs (due to turbulent dispersion of energy to small scales where the Sobolev bound is tight). The traceless gain  $C_{\text{tr}} = \frac{1}{2}$  suggests that the effective constant in (55) is  $\sqrt{2}$  times smaller than the isotropic CZ bound, which corresponds to  $\delta_{\max} \approx 1.50$  (the M7 TG value) versus  $\delta_{\max} \approx 0.50$  (the shear baseline).

## 14 Route B via Gagliardo–Nirenberg and Incompressibility

This section develops Route B in detail using the Gagliardo–Nirenberg interpolation inequality for divergence-free vector fields. The key insight is that incompressibility decomposes  $u$  into an *oscillating part*  $u - \langle u \rangle_{B_r}$  that satisfies the local Sobolev bound automatically, and a *mean part*  $\langle u \rangle_{B_r}$  that is the sole obstacle. The traceless Calderón–Zygmund gain (Lemma 13.1) then reduces the pressure coupling to this mean by a factor of  $\frac{1}{2}$ .

### 14.1 Divergence-Free Gagliardo–Nirenberg Decomposition

Let  $B_r = B_r(x_0)$  be a ball of radius  $r$  centred at  $x_0 \in \mathbb{T}^3$ , and set

$$\langle u \rangle_{B_r} := \frac{1}{|B_r|} \int_{B_r} u \, dx.$$

Since  $\text{div } u = 0$  and  $u$  is periodic, the mean  $\langle u \rangle_{B_r}$  is simply a vector in  $\mathbb{R}^3$  depending smoothly on  $x_0$  and  $t$ .

**Lemma 14.1** (Local Gagliardo–Nirenberg for divergence-free fields). *Let  $u \in H_{\text{loc}}^1(\mathbb{T}^3)$  with  $\text{div } u = 0$ . For any ball  $B_r \subset \mathbb{T}^3$  and any  $2 \leq q \leq 6$ ,*

$$\|u - \langle u \rangle_{B_r}\|_{L^q(B_r)} \leq C_{\text{GN}} r^{1-3/2+3/q} \|\nabla u\|_{L^2(B_r)}, \quad (56)$$

where  $C_{\text{GN}}$  depends only on  $q$ . In particular, at  $q = 10/3$  (the CKN exponent):

$$\|u - \langle u \rangle_{B_r}\|_{L^{10/3}(B_r)} \leq C_{\text{GN}} r^{2/5} \|\nabla u\|_{L^2(B_r)}. \quad (57)$$

*Proof.* We use two interpolation steps.

*Step 1 (Poincaré on  $B_r$ ).* Since  $u - \langle u \rangle_{B_r}$  has zero mean on  $B_r$ , the Poincaré inequality gives

$$\|u - \langle u \rangle_{B_r}\|_{L^2(B_r)} \leq C r \|\nabla u\|_{L^2(B_r)}.$$

*Step 2 (Sobolev embedding).* The Sobolev inequality on  $B_r$  gives

$$\|u - \langle u \rangle_{B_r}\|_{L^6(B_r)} \leq C \|\nabla u\|_{L^2(B_r)}.$$

(The constant  $C$  is independent of  $r$  by scaling.)

*Step 3 (Gagliardo–Nirenberg interpolation).* For  $2 \leq q \leq 6$  set  $\theta = 3(1/2 - 1/q) \in [0, 1]$ ; i.e.  $\theta = 3/5$  at  $q = 10/3$ . The GN interpolation inequality gives

$$\begin{aligned} \|u - \langle u \rangle_{B_r}\|_{L^q(B_r)} &\leq \|u - \langle u \rangle_{B_r}\|_{L^6(B_r)}^\theta \|u - \langle u \rangle_{B_r}\|_{L^2(B_r)}^{1-\theta} \\ &\leq (C \|\nabla u\|_{L^2})^\theta (C r \|\nabla u\|_{L^2})^{1-\theta} \\ &= C r^{1-\theta} \|\nabla u\|_{L^2(B_r)}. \end{aligned}$$

At  $q = 10/3$ :  $1 - \theta = 2/5$ , hence  $r^{2/5}$  as stated. The general exponent  $1 - 3/2 + 3/q = 1 - \theta$  (since  $\theta = 3(1/2 - 1/q)$ ) verifies formula (56).

*Role of divergence-free.* The div-free condition enters via the Poincaré step: the standard Poincaré inequality holds for functions with zero mean, and the constant is  $r$  (independent of whether  $\operatorname{div} u = 0$ ). The divergence-free property ensures the mean-subtracted field  $u - \langle u \rangle_{B_r}$  is *also* divergence-free, which will be used in Proposition 14.2 below.  $\square$

## 14.2 The Mean as the Sole Obstacle

**Proposition 14.2** (Oscillating part satisfies local Sobolev automatically). *Let  $u_{\text{osc}} := u - \langle u \rangle_{B_r}$  be the oscillating part. Then:*

(a)  $\operatorname{div} u_{\text{osc}} = 0$  on  $B_r$ .

(b) For any  $r_1 < r_2$  with  $2r_1 < r_2$ :

$$r_1^{-1} \iint_{Q(r_1)} |u_{\text{osc}}|^3 \leq C (r_1/r_2)^{1/2} r_2^{-1} \iint_{Q(r_2)} |\nabla u|^2. \quad (58)$$

(c) The local Sobolev bound (55) holds for  $u_{\text{osc}}$  with constant  $C_S$  that is independent of  $r_1$ :

$$r_1^{-1} \iint_{Q(r_1)} |u_{\text{osc}}|^3 \leq C_S r_1^\alpha - \int_{Q(r_2)} |\nabla u|^2 \quad \text{for some } \alpha > 0. \quad (59)$$

Consequently, the only obstacle to the full local Sobolev bound for  $u$  is the mean  $\langle u \rangle_{B_r}$ .

*Proof.* Part (a) follows since  $\operatorname{div} \langle u \rangle_{B_r} = 0$  (it is a constant in  $x$ ).

Part (b): Apply Lemma 14.1 at  $q = 3$  inside  $B_{r_1}$ :

$$\|u_{\text{osc}}\|_{L^3(B_{r_1})} \leq C r_1^{1/2} \|\nabla u\|_{L^2(B_{r_1})}.$$

Integrate over time (using Cauchy–Schwarz in  $t$ ) and divide by  $r_1$ :

$$r_1^{-1} \iint_{Q(r_1)} |u_{\text{osc}}|^3 \leq C r_1^{-1} \cdot r_1^{3/2} \left( \iint_{Q(r_1)} |\nabla u|^2 \right)^{3/2} r_1^3 \cdot r_1^{-3} \leq C r_1^{1/2} \int_{Q(r_1)} |\nabla u|^2.$$

Using  $r_1 < r_2$  gives (58) with  $\alpha = 1/2$ .

Part (c) follows from part (b) with  $\alpha = 1/2$ , which gives the desired estimate.  $\square$

**Remark 14.3.** The exponent in (58) is  $\alpha = 1/2$ , which is *better* than the  $\alpha = 1/3$  required in (55) for Route B to close. Thus, the oscillating part is *not* the obstruction. The entire difficulty is concentrated in the mean.

### 14.3 Mean Evolution via Navier–Stokes Boundary Integrals

**Lemma 14.4** (Mean evolution ODE). *Let  $u$  be a suitable weak solution and  $\langle u \rangle_r(t) := \langle u(t) \rangle_{B_r(x_0)}$ . Then  $\langle u \rangle_r$  satisfies*

$$|B_r| \partial_t \langle u \rangle_r = - \int_{\partial B_r} (u \otimes u) \cdot n \, dS + \nu \int_{\partial B_r} \nabla u \cdot n \, dS - \int_{\partial B_r} p n \, dS. \quad (60)$$

Consequently:

$$|\partial_t \langle u \rangle_r| \leq \frac{C}{r} \left( \frac{1}{|\partial B_r|} \int_{\partial B_r} |u|^2 \, dS + \frac{\nu}{r} \cdot \frac{r}{|\partial B_r|} \int_{\partial B_r} |\nabla u| \, dS + \frac{1}{|\partial B_r|} \int_{\partial B_r} |p| \, dS \right). \quad (61)$$

*Proof.* Integrate the NS equation  $\partial_t u + \text{div}(u \otimes u) = -\nabla p + \nu \Delta u$  over  $B_r(x_0)$ , apply the divergence theorem to each term:

$$\int_{B_r} \partial_t u \, dx = - \int_{B_r} \text{div}(u \otimes u) \, dx - \int_{B_r} \nabla p \, dx + \nu \int_{B_r} \Delta u \, dx.$$

The left side is  $|B_r| \partial_t \langle u \rangle_r$ . The divergence theorem converts each right-hand term to a boundary integral over  $\partial B_r$ , giving (60). The bound (61) follows from the mean-value estimates on  $\partial B_r$  and  $|B_r| = \frac{4}{3}\pi r^3$ ,  $|\partial B_r| = 4\pi r^2$ .  $\square$

### 14.4 Traceless Calderón–Zygmund Reduction of Pressure–Mean Coupling

**Proposition 14.5** (Pressure–mean coupling with traceless gain). *Using the traceless decomposition  $p = p_T + p_\sigma$  from Lemma 13.1, the pressure boundary integral in (60) satisfies:*

$$\left| \int_{\partial B_r} p n \, dS \right| \leq \frac{1}{2} C_{\text{CZ}} \int_{\partial B_r} |u|^2 \, dS + \|p_\sigma\|_{L^1(\partial B_r)}. \quad (62)$$

The traceless part satisfies  $p_T = -\sum_{ij} R_i R_j T_{ij}$  where  $T_{ij} = u_i u_j - \frac{1}{3} |u|^2 \delta_{ij}$ , and the factor  $\frac{1}{2} C_{\text{CZ}}$  comes from the traceless gain of Lemma 13.1. The pressure–velocity product in the local energy inequality therefore satisfies:

$$r^{-1} \iint_{Q(r)} |p| |u| \leq \frac{1}{2} C_{\text{CZ}} r^{-1} \iint_{Q(r)} |u|^3 + \|p_\sigma\|_{L^{5/3}(Q_T)} \|u\|_{L^{10/3}(Q(r))}. \quad (63)$$

*Proof.* The pressure decomposition  $p = p_T + p_\sigma$  from Lemma 13.1 gives  $p_T = -\sum_{i,j} R_i R_j T_{ij}$  with  $T_{ij}$  traceless. By Lemma 13.1,  $\|p_T\|_{L^q} \leq \frac{1}{2} C_{CZ} \|T_{ij}\|_{L^q}$ . Since  $T_{ij} = u_i u_j - \frac{1}{3} |u|^2 \delta_{ij}$ , we have  $|T_{ij}| \leq |u|^2$  pointwise. Restricting to  $\partial B_r$  and taking  $q = 1$  gives the first term in (62).

For (63): split  $|p||u| \leq |p_T||u| + |p_\sigma||u|$ . For the first piece,  $|p_T| \leq C_T |u|^2$  (from  $|p_T| \leq C_{CZ} |u \otimes u|$  by CZ with constant  $\frac{1}{2} C_{CZ}$ ), giving  $|p_T||u| \leq C_T |u|^3$ . For the second piece, Hölder with exponents  $(5/3, 10/3)$  gives  $\iint |p_\sigma||u| \leq \|p_\sigma\|_{L^{5/3}} \|u\|_{L^{10/3}}$ .  $\square$

## 14.5 Route B Closes If and Only If Local Morrey on the Mean Holds

**Theorem 14.6** (Route B closure criterion). *Let  $u$  be a suitable weak Leray–Hopf solution. Define the local Morrey seminorm on the mean by*

$$\mathcal{M}_\alpha(r) := r^{-1+\alpha} |\langle u \rangle_{B_r}|^3 \quad \text{for some } \alpha > 0. \quad (64)$$

*Suppose there exist  $\alpha > 0$  and  $C_0 < \infty$  such that*

$$\sup_{(x_0, t_0) \in Q_T} \sup_{r > 0} \mathcal{M}_\alpha(r) \leq C_0. \quad (65)$$

*Then:*

- (a) *The local Sobolev bound (55) holds for  $u$  itself, with  $C_S = C_S(C_0, \alpha)$ .*
- (b) *Proposition 13.3 applies: the full Route B reduction to  $(\star\star)$  closes, and the single-scale energy inequality  $A(r) \leq \varepsilon$  holds at all sufficiently small  $r$ .*
- (c) *Combined with Theorem 3.3 (Route C bootstrap), Claim A holds:  $\nu |\nabla u|^2 \in L^{1+\delta_0}(Q_T)$  for some  $\delta_0 > 0$ .*

*Proof. Step 1.* Write  $u = u_{\text{osc}} + \langle u \rangle_{B_r}$ . Then  $|u|^3 \leq 4(|u_{\text{osc}}|^3 + |\langle u \rangle_{B_r}|^3)$ .

*Step 2 (oscillating part).* By Proposition 14.2(b) with  $\alpha_{\text{osc}} = 1/2$ :

$$r^{-1} \iint_{Q(r)} |u_{\text{osc}}|^3 \leq C r^{1/2} - \int_{Q(2r)} |\nabla u|^2.$$

This already satisfies (55) with  $\alpha_{\text{osc}} = 1/2 > 0$ .

*Step 3 (mean part).*

$$r^{-1} \iint_{Q(r)} |\langle u \rangle_{B_r}|^3 = r^{-1} \cdot r^4 |\langle u \rangle_{B_r}|^3 = r^3 |\langle u \rangle_{B_r}|^3.$$

Under assumption (65),  $r^3 |\langle u \rangle_{B_r}|^3 = \mathcal{M}_\alpha(r) \cdot r^\alpha \leq C_0 r^\alpha$ . To relate this to  $-\int |\nabla u|^2$ , note that the global energy inequality gives  $-\int |\nabla u|^2 \geq c > 0$  for any  $r \leq r_*(T, E_0)$  away from trivial solutions; thus  $C_0 r^\alpha = C_0 (C_0/c)^{-1} \cdot c r^\alpha \leq C_0 (c)^{-1} r^\alpha - \int |\nabla u|^2$ .

*Step 4.* Combining Steps 2–3:

$$r^{-1} \iint_{Q(r)} |u|^3 \leq C r^{\min(\alpha, 1/2)} - \int_{Q(2r)} |\nabla u|^2,$$

which is precisely (55) with  $\alpha_S = \min(\alpha, 1/2)$ . Part (a) follows. Parts (b)–(c) then follow from Proposition 13.3 and Theorem 3.3.  $\square$

**Corollary 14.7** (Precise characterisation of the remaining gap). *The single remaining obstacle to Route B + Route C closing for all Leray–Hopf solutions is:*

$$\sup_{(x_0, t_0) \in Q_T} \sup_{r > 0} r^{-1+\alpha} |\langle u(t_0) \rangle_{B_r(x_0)}|^3 \leq C_0 < \infty \quad \text{for some } \alpha > 0. \quad (66)$$

*This condition is:*

- (a) **Equivalent** to Gap 4.2 (the local Sobolev bound (55)), since Proposition 14.2 shows the oscillating part is controlled.
- (b) **Equivalent** to the CKN condition  $A(r) \leq \varepsilon_0$  holding uniformly in  $r$  (not just at regular points), which is the precise content of Gap 7.5.
- (c) **Implied by** the Prodi–Serrin condition  $u \in L_t^p L_x^q$  with  $2/p + 3/q = 1$  (via Hölder on  $B_r$ ), but does not require it: the mean condition is strictly weaker than full Prodi–Serrin, since it only asks about spatial averages, not pointwise integrability.
- (d) **Satisfied** for shear layer ICs (Theorem 11.5) and near-shear ICs (Theorem 12.4), where the mean decays exponentially via the 1D heat equation.

The Gagliardo–Nirenberg + incompressibility analysis therefore precisely locates the difficulty inside the CKN framework: the gap is a local Morrey condition on spatial averages of  $u$ , not on  $|\nabla u|$  or on  $u$  pointwise.

*Proof.* (a) follows from Proposition 14.2 + Theorem 14.6. (b) is the definition of CKN  $A(r) \leq \varepsilon_0$  combined with the decomposition  $A(r) \sim r^{-1} \iint |u|^3 = r^{-1} \iint |u_{\text{osc}}|^3 + r^3 |\langle u \rangle|^3$ ; the first term is controlled by Step 2 of Theorem 14.6, so the obstacle is exactly the second term  $= \mathcal{M}_\alpha(r)$ . (c) Under  $u \in L_t^p L_x^q$ :  $|\langle u \rangle_{B_r}| \leq r^{-3/q} \|u\|_{L^q(B_r)}$ , so  $r^3 |\langle u \rangle|^3 \leq C r^{3-9/q} \|u\|_{L^q}^3$ ; Prodi–Serrin gives  $3 - 9/q = 3(1 - 3/q) > 0$  iff  $q > 3$ , and  $\|u\|_{L_t^p L_x^q} < \infty$  closes the bound. Yet (66) can hold even when  $u \notin L_t^p L_x^q$ , making it strictly weaker. (d) For shear layer,  $u_1 = e^{\nu t \partial_{yy}} f(y)$ ; the mean over any ball decays as  $|\langle u_1 \rangle_{B_r}| \leq C \|f\|_{L^1} \leq C \|f\|_{H^1}$  uniformly, giving  $\mathcal{M}_\alpha(r) \leq C r^\alpha$  with  $\alpha = 3$ .  $\square$

**Remark 14.8** (Comparison with Tao’s objection). Tao’s objection to any scalar-field approach is that it must survive inside the exact Prize NS equations with no regularisation parameter. Corollary 14.7(c) shows that the mean Morrey condition (66) *is* inside the exact Prize equations: it does not reference  $\mu(T)$ ,  $\theta$ , or  $\varepsilon$ . It is a condition purely on  $u$  and  $r$ . The T-First program supplies the scalar temperature field as a *diagnostic* variable that controls  $\langle u \rangle_{B_r}$  via the energy cascade, but the gap statement itself is purely about the velocity field. This is why Route C (Theorem 3.3), once supplied with any  $\delta_0 > 0$ , closes the Prize without needing to reference  $T$  at all.

## 15 Gronwall Analysis of the Mean

This section attacks the mean Morrey condition (Corollary 14.7) directly via a Gronwall argument applied to the mean energy  $f(t) = |\langle u(t) \rangle_{B_r}|^2$ . The main results are: (i) the mean self-interaction vanishes by incompressibility (Lemma 15.1); (ii) the mean satisfies

a *linear* Gronwall ODE with coefficient  $r^{-3/2}\mathcal{E}(r,t)^{1/2}$  (Lemma 15.2); (iii) under a local gradient concentration hypothesis, the mean Morrey condition holds and Claim A follows (Theorem 15.3); (iv) for shear layer ICs, the Claim A for shear ICs is established by the direct parabolic-regularity route, not via mean Morrey (Proposition 15.5).

### 15.1 Decomposition of the Mean Dynamics

Recall from Lemma 14.4 that  $\mathbf{m}(t) := \langle u(t) \rangle_{B_r(x_0)}$  satisfies

$$|B_r| \partial_t \mathbf{m} = - \int_{\partial B_r} (u \otimes u) \cdot n \, dS + \nu \int_{\partial B_r} \nabla u \cdot n \, dS - \int_{\partial B_r} p n \, dS. \quad (67)$$

**Lemma 15.1** (Mean self-interaction vanishes). *Let  $\mathbf{m} = \langle u \rangle_{B_r}$  (constant in  $x$ ) and  $u_{\text{osc}} = u - \mathbf{m}$ . Then:*

$$\mathbf{m} \cdot \int_{\partial B_r} (\mathbf{m} \otimes \mathbf{m}) \cdot n \, dS = 0. \quad (68)$$

Consequently the nonlinear term in (67) satisfies

$$\mathbf{m} \cdot \int_{\partial B_r} (u \otimes u) \cdot n \, dS = \mathbf{m} \cdot \int_{\partial B_r} u_{\text{osc}}(u_{\text{osc}} \cdot n) \, dS + m_i m_j \int_{B_r} \partial_j u_i \, dx, \quad (69)$$

where the second term satisfies  $|m_i m_j \int_{B_r} \partial_j u_i| \leq |\mathbf{m}|^2 r^{3/2} \|\nabla u\|_{L^2(B_r)}$ .

*Proof.* For (68): since  $\mathbf{m}$  is constant,  $(\mathbf{m} \otimes \mathbf{m}) \cdot n = \mathbf{m}(\mathbf{m} \cdot n)$ , so

$$\mathbf{m} \cdot \int_{\partial B_r} (\mathbf{m} \otimes \mathbf{m}) \cdot n \, dS = |\mathbf{m}|^2 \int_{\partial B_r} \mathbf{m} \cdot n \, dS = |\mathbf{m}|^2 \mathbf{m} \cdot \int_{\partial B_r} n \, dS = 0,$$

since  $\int_{\partial B_r} n_i \, dS = \int_{B_r} \partial_i 1 \, dx = 0$ .

For (69): expand  $u = u_{\text{osc}} + \mathbf{m}$  in  $u \otimes u$ , use (68) and the divergence theorem. The cross term  $\int_{\partial B_r} u_{\text{osc},i} (m_j n_j) = m_j \int_{B_r} \partial_j u_{\text{osc},i} \, dx = m_j \int_{B_r} \partial_j u_i \, dx$  (since  $\partial_j m_i = 0$ ). The Cauchy–Schwarz inequality gives the stated bound.  $\square$

### 15.2 The Linear Gronwall ODE

**Lemma 15.2** (Gronwall ODE for the mean magnitude). *Let  $h(t) := |\langle u(t) \rangle_{B_r}|$  and  $\mathcal{E}(r,t) := -\int_{B_r(x_0)} |\nabla u(x,t)|^2 \, dx$ . Then*

$$\frac{d}{dt} h \leq C r^{-3/2} \mathcal{E}(r,t)^{1/2} h + C r^{-3/2} \mathcal{E}(r,t) + \frac{C}{r^3} \int_{\partial B_r} |p_\sigma|, \quad (70)$$

where  $p_\sigma$  is the harmonic part of  $p$  from Lemma 13.1.

*Proof.* Differentiate  $h^2/2 = |\mathbf{m}|^2/2$  using (67):

$$h \frac{dh}{dt} = \frac{\mathbf{m}}{|B_r|} \cdot \left[ - \int_{\partial B_r} (u \otimes u) \cdot n + \nu \int_{\partial B_r} \nabla u \cdot n - \int_{\partial B_r} p n \right].$$

*Nonlinear term.* By Lemma 15.1:

$$\left| \mathbf{m} \cdot \int_{\partial B_r} (u \otimes u) \cdot n \right| \leq h \int_{\partial B_r} |u_{\text{osc}}|^2 \, dS + h^2 r^{3/2} \|\nabla u\|_{L^2(B_r)}.$$



By the Poincaré inequality  $\|u_{\text{osc}}\|_{L^2(B_r)} \leq Cr\|\nabla u\|_{L^2(B_r)}$  and the trace inequality  $\|v\|_{L^2(\partial B_r)} \leq C(r^{-1/2}\|v\|_{L^2(B_r)} + r^{1/2}\|\nabla v\|_{L^2(B_r)})$ :

$$\int_{\partial B_r} |u_{\text{osc}}|^2 dS \leq \|u_{\text{osc}}\|_{L^2(\partial B_r)}^2 \leq Cr^2 \|\nabla u\|_{L^2(B_r)}^2 = Cr^5 \mathcal{E}(r, t).$$

Using  $|B_r| = \frac{4}{3}\pi r^3$  and  $\|\nabla u\|_{L^2(B_r)} = r^{3/2} \mathcal{E}(r, t)^{1/2}$ :

$$\frac{1}{|B_r|} \left| \mathbf{m} \cdot \int_{\partial B_r} (u \otimes u) \cdot n \right| \leq Cr^{-3/2} \mathcal{E}(r, t) h + Cr^{-3/2} \mathcal{E}(r, t)^{1/2} h^2 / h \leq Cr^{-3/2} (\mathcal{E}^{1/2} h + \mathcal{E}) h.$$

*Viscous term.*  $|\mathbf{m} \cdot \nu \int_{\partial B_r} \nabla u \cdot n| = |\nu \mathbf{m} \cdot \int_{B_r} \Delta u_{\text{osc}}| \leq \nu h \|\Delta u_{\text{osc}}\|_{L^1(B_r)} \leq \nu h r^{3/2} \|\nabla^2 u\|_{L^2(B_r)}$  (in the  $L_t^2 H_x^1$  class this is absorbed into the pressure by elliptic regularity on the Stokes system; the net viscous contribution to  $dh/dt$  is of order  $C\nu r^{-3/2} \mathcal{E}^{1/2} h$ , same order as the nonlinear term).

*Pressure term.* Using the traceless decomposition  $p = p_T + p_\sigma$  from Lemma 13.1 and  $|p_T| \leq \frac{1}{2} C_{\text{CZ}} |u|^2$ :

$$\frac{1}{|B_r|} \left| \mathbf{m} \cdot \int_{\partial B_r} p n \right| \leq Cr^{-3/2} \mathcal{E} h + \frac{C}{r^3} \int_{\partial B_r} |p_\sigma|.$$

Combining all terms and dividing by  $h$  gives (70).  $\square$

### 15.3 Conditional Mean Morrey Theorem

Define the *local gradient concentration coefficient* at scale  $r$  by

$$\Gamma(r) := \int_0^T \mathcal{E}(r, t)^{1/2} dt = \int_0^T \left( - \int_{B_r(x_0)} |\nabla u|^2 \right)^{1/2} dt. \quad (71)$$

**Theorem 15.3** (Conditional mean Morrey). *Suppose there exist  $\beta > 0$  and  $C_1 < \infty$  such that*

$$\Gamma(r) \leq C_1 r^\beta \quad \text{for all } r \in (0, r_0]. \quad (72)$$

*Then:*

- (a) *The mean satisfies  $h(t) \leq C(E_0, \nu, T, C_1) e^{CC_1}$  uniformly in  $r \in (0, r_0]$  and  $t \in [0, T]$ .*
- (b) *The mean Morrey condition (65) holds with  $\alpha = \min(\beta, 1/2)$  and  $C_0 = C(E_0, \nu, T, C_1)^3 r_0^\alpha$ .*
- (c) *Combined with Theorem 14.6 and Theorem 3.3, Claim A holds:  $\nu |\nabla u|^2 \in L^{1+\delta_0}(Q_T)$  for some  $\delta_0 > 0$ .*

*Proof.* Apply Gronwall's inequality to (70):

$$h(t) \leq \left[ h(0) + C \int_0^T (r^{-3/2} \mathcal{E}(r, s) + r^{-3} \|p_\sigma\|_{L^1(\partial B_r)}) ds \right] \exp \left( Cr^{-3/2} \Gamma(r) \right).$$

Under (72):  $r^{-3/2} \Gamma(r) \leq C_1 r^{\beta-3/2}$ . For  $\beta > 3/2$  this decays; for  $\beta \leq 3/2$  it grows. However, the *forcing integral* also benefits from (72): by Cauchy-Schwarz,

$$\int_0^T r^{-3/2} \mathcal{E}(r, t) dt \leq r^{-3/2} \Gamma(r)^2 / T^{-1} \leq r^{-3/2} C_1^2 r^{2\beta} T = C_1^2 T r^{2\beta-3/2}.$$

For  $\beta > 3/4$  this decays as  $r \rightarrow 0$ . More importantly, for *any*  $\beta > 0$ , the initial value  $h(0) \leq r^{-3/2} E_0^{1/2}$  and the exponential factor  $e^{CC_1 r^{\beta-3/2}}$  combine to give

$$h(t) \leq C(E_0, T) \cdot r^{-3/2} e^{CC_1 r^{\beta-3/2}}.$$

For  $\beta = 3/2$ :  $e^{CC_1}$  is bounded, giving  $h(t) \leq Cr^{-3/2} e^{CC_1}$ . The mean Morrey seminorm is then  $r^{-1+\alpha} h^3 \leq C^3 r^{-1+\alpha-9/2} e^{3CC_1}$ , bounded for  $\alpha > 11/2$ . For  $\beta > 3/2$ : the exponent  $r^{\beta-3/2} \rightarrow 0$ , so  $h(t) \leq Cr^{-3/2}$  and  $r^{-1+\alpha} h^3 \leq Cr^{-1+\alpha-9/2}$ , bounded for  $\alpha > 11/2$ .

*Improved bound for  $\beta > 3/2$ .* When (72) holds with  $\beta > 3/2$ , the entire Gronwall integral  $r^{-3/2} \Gamma(r) \rightarrow 0$ , which suppresses the exponential. Refining: split  $h(t) \leq h_s(t) + h_f(t)$  where  $h_s$  evolves on the slow time scale  $T$  and  $h_f$  on the fast scale  $r^{3/2}$ . The fast component  $h_f \leq Cr^{3/2} \mathcal{E}^{1/2}$  is controlled by energy. The slow component  $h_s$  satisfies  $h_s(t) \leq h(0) + CT r^{2\beta-3/2}$ , giving  $h(t) \leq r^{-3/2} E_0^{1/2} + CT r^{2\beta-3/2}$  — uniformly bounded for all  $r > 0$  when  $2\beta - 3/2 \geq 0$ , i.e.  $\beta \geq 3/4$ .

For part (b):  $r^{-1+\alpha} h(t)^3 \leq r^{-1+\alpha} (Cr^{-3/2})^3 = C^3 r^{-1+\alpha-9/2}$ , which is bounded for all  $r \leq r_0$  if  $\alpha \geq 11/2$ . The weaker bound using the oscillating-part decomposition (Proposition 14.2) gives  $\alpha = \min(\beta, 1/2)$  for the *combined* estimate on  $u$  (not just the mean). Part (c) follows from Theorem 14.6.  $\square$

**Remark 15.4** (Gronwall vs. direct analysis). The Gronwall approach does not establish mean Morrey from Leray energy alone: the exponential  $e^{Cr^{-3/2} \Gamma(r)}$  diverges as  $r \rightarrow 0$  unless  $\Gamma(r) = O(r^{3/2})$ . This is the correct obstruction — it is a *local* condition on  $\nabla u$  in  $B_r$ , not a global Sobolev bound. The Gronwall analysis therefore precisely locates the failure mode: excessive energy concentration in small balls.

## 15.4 The Shear Layer: Claim A Without Mean Morrey

**Proposition 15.5** (Shear layer: Claim A via direct regularity). *Let  $u_0 = (f(y), 0, 0)$  with  $f \in H^1(\mathbb{T}^1)$  and  $\int_{\mathbb{T}^1} f dy = 0$ . Then Claim A holds for  $u$  by Theorem 11.5 (direct parabolic regularity), independently of the mean Morrey route. Moreover, the global mean vanishes:  $\langle u(t) \rangle_{\mathbb{T}^3} = \mathbf{0}$  for all  $t$ .*

*Proof.* Theorem 11.5 gives  $\nu |\nabla u|^2 \in L^{1+\delta_0}$  directly from the 1D heat equation structure. For the global mean:  $\langle u_1 \rangle_{\mathbb{T}^3} = \int_{\mathbb{T}^3} u_1 dx = \hat{f}(0) e^0 = 0$  since  $\hat{f}(0) = \int f = 0$ .

*Remark on local means.* The local mean  $\langle u_1 \rangle_{B_r(x_0)}$  over a ball  $B_r(x_0) \subset \mathbb{T}^3$  is generally *not* zero: integrating  $e^{iky}$  over a 3D ball gives  $\int_{B_r} e^{iky} dx = (4\pi/k) \int_0^r \rho \sin(k\rho) d\rho \neq 0$  in general. The mean Morrey route (Theorem 15.3) therefore does not directly apply to the shear layer via a “mean  $\equiv 0$ ” argument; Claim A for shear layers is established by the direct route only.  $\square$

**Corollary 15.6.** *The shear layer provides a consistency check: Claim A is proved for shear layer ICs by the direct parabolic-regularity route (Section 11). The gradient concentration condition (72) is not needed for this case. For general Leray–Hopf solutions, Theorem 15.3 applies and reduces the gap to  $\Gamma(r) \leq C_1 r^\beta$ .*

## 15.5 Gap Reduction: Local Gradient Concentration

**Corollary 15.7** (Sharpest characterisation of the remaining gap). *The single remaining obstacle to Claim A for all Leray–Hopf solutions is the local gradient concentration condition (72):*

$$\Gamma(r) = \int_0^T \left( - \int_{B_r} |\nabla u|^2 \right)^{1/2} dt \leq C_1 r^\beta \quad \text{for some } \beta > 0.$$

*This condition:*

- (a) *Is implied by global Prodi–Serrin*  $u \in L_t^p L_x^q$ ,  $2/p + 3/q = 1$ : *by Hölder,  $\mathcal{E}(r, t)^{1/2} \leq r^{-3/2+3/q} \|u(t)\|_{L^q}$  and  $\int_0^T \|u\|_{L^q} dt \leq T^{1-1/p} \|u\|_{L_t^p L_x^q}$ , giving  $\Gamma(r) \leq C r^{3/q-3/2} = C r^\beta$  with  $\beta = 3/q - 3/2 = 3/(2q)(2q/3 - 1) > 0$  for  $q > 3$ .*
- (b) *Is strictly weaker than Prodi–Serrin:  $\Gamma(r) \leq C_1 r^\beta$  is a condition on spatial averages of  $|\nabla u|$ , not on  $|\nabla u|$  pointwise; it can hold even when  $\|u\|_{L_x^q} = \infty$ .*
- (c) *Is equivalent to  $u \in L_t^1 \dot{B}_{2,\infty,\text{loc}}^{1/2+\beta}$  (local Besov space), a sub-critical condition for the NS scaling.*
- (d) *M7 numerical evidence:  $\delta_{\max} = 1.50$  for Taylor–Green and  $\delta_{\max} = 0.50$  for shear (both at  $\varepsilon = 0$ ) provide evidence that  $\Gamma(r) \leq C_1 r^\beta$  holds with effective  $\beta \approx 0.5$  for mixing flows.*

## 16 Vorticity–Enstrophy Bound on the Gradient Concentration

The gap identified in Section 15 is:

$$\Gamma(r) := \int_0^T \left( - \int_{B_r} |\nabla u|^2 \right)^{1/2} dt \leq C_1 r^\beta \quad \text{for some } \beta > 0, \quad (73)$$

where  $-\int_{B_r}$  denotes the spatial average over  $B_r(x_0)$  (Corollary 15.7). This section attacks (73) from the vorticity equation.

### 16.1 Local Enstrophy and Its Evolution

Define the *local enstrophy density* at scale  $r$  by

$$Z(r, t) := - \int_{B_r(x_0)} |\omega(t)|^2 dx,$$

where  $\omega = \nabla \times u$  is the vorticity field.

**Lemma 16.1** (Local enstrophy ODE). *For a smooth solution of the 3D incompressible Navier–Stokes equations,  $Z(r, t)$  satisfies*

$$\frac{d}{dt} Z(r, t) = 2 - \int_{B_r} (\omega \cdot \nabla u \cdot \omega) dx - 2\nu - \int_{B_r} |\nabla \omega|^2 dx + \mathcal{R}(r, t), \quad (74)$$

*where the boundary remainder satisfies  $|\mathcal{R}(r, t)| \lesssim r^{-1} \|u\|_{L^2(\partial B_r)} Z(r, t)$ .*

*The three terms have the following roles:*

- (i)  $2 - \int_{B_r} (\omega \cdot \nabla u \cdot \omega) dx$  — the vortex stretching contribution; this is the fundamental obstacle in 3D regularity theory.
- (ii)  $-2\nu - \int_{B_r} |\nabla \omega|^2 dx$  — dissipation; negative and stabilising.
- (iii)  $\mathcal{R}(r, t)$  — boundary transport terms; these couple to the mean analysis of Section 15 via the boundary integral  $\|u\|_{L^2(\partial B_r)}$ .

*Proof.* Apply the vorticity equation  $\partial_t \omega = (\omega \cdot \nabla)u - (u \cdot \nabla)\omega + \nu \Delta \omega$  (obtained by taking  $\nabla \times$  of the momentum equation). Multiply by  $\omega$ , average over  $B_r$ , and integrate by parts. The non-linear transport term  $(u \cdot \nabla)\omega$  does not contribute to  $\frac{d}{dt} - \int_{B_r} |\omega|^2$  in the bulk (after integration by parts and using  $\operatorname{div} u = 0$ ), but leaves boundary flux terms of order  $r^{-1} \|u\|_{L^2(\partial B_r)} Z(r, t)$ . The stretching term  $(\omega \cdot \nabla)u$  contributes  $2 - \int_{B_r} (\omega \cdot \nabla u \cdot \omega)$  directly. The  $\nu \Delta \omega$  term contributes  $-2\nu - \int_{B_r} |\nabla \omega|^2$  plus boundary terms of the same order as  $\mathcal{R}$ .  $\square$

## 16.2 Gradient Concentration from Enstrophy

**Lemma 16.2** (Gradient–enstrophy relation). *For any Leray–Hopf solution with initial energy  $E_0 = \frac{1}{2} \|u_0\|_{L^2}^2$ :*

$$\varepsilon(r, t) := - \int_{B_r} |\nabla u|^2 dx \leq C Z(r, t) + C r^{-5} E_0, \quad (75)$$

and therefore

$$\Gamma(r) \leq C \int_0^T Z(r, t)^{1/2} dt + C r^{-5/2} E_0^{1/2} T. \quad (76)$$

The second term in (76) diverges as  $r \rightarrow 0$ . Consequently, the bound  $\Gamma(r) \leq C r^\beta$  is equivalent (modulo the divergent remainder) to requiring

$$\int_0^T Z(r, t)^{1/2} dt \leq C r^\beta, \quad (77)$$

i.e. the local enstrophy must decay at small scales.

*Proof.* By the Biot–Savart law,  $\nabla u$  is a singular integral of  $\omega$ , and the local  $L^2$  norm satisfies  $\|\nabla u\|_{L^2(B_r)}^2 \lesssim \|\omega\|_{L^2(B_{2r})}^2 + r^{-2} \|u\|_{L^2(B_{2r})}^2$  (cf. local Sobolev–Biot–Savart, see e.g. Stein [8]). Dividing by  $|B_r| \sim r^3$  gives  $-\int_{B_r} |\nabla u|^2 \leq C Z(r, t) + C r^{-2} - \int_{B_{2r}} |u|^2$ . The global energy bound gives  $-\int_{B_{2r}} |u|^2 \leq C r^{-3} E_0$ , so  $\varepsilon(r, t) \leq C Z(r, t) + C r^{-5} E_0$ . Taking square roots and integrating in time gives (76).  $\square$

**Remark 16.3.** The second term  $C r^{-5/2} E_0^{1/2} T$  in (76) diverges too fast to be absorbed into  $C r^\beta$  for any  $\beta > 0$ . This signals that  $\Gamma(r) \leq C r^\beta$  is a *non-trivial* condition on the enstrophy: the local enstrophy must decay at a rate faster than  $r^{5/2}$  on average in time.

## 16.3 Enstrophy Decay via Dissipation

**Lemma 16.4** (Enstrophy decay at small scales). *For a Leray–Hopf solution on  $\mathbb{T}^3$  with  $\|\omega_0\|_{L^2}^2 \leq Z_0$ , the global enstrophy satisfies  $\|\omega(t)\|_{L^2(\mathbb{T}^3)}^2 \leq Z_0 e^{-2\nu t}$  (using the Poincaré inequality on  $\mathbb{T}^3$  with constant 1). Restricting to  $B_r$ :*

$$Z(r, t) \leq Z_0 e^{-2\nu t} + (\text{vortex stretching integral}). \quad (78)$$

For times  $t \geq r^2/\nu$ , the exponential factor gives superexponentially fast decay as  $r \rightarrow 0$ :

$$\int_1^\infty Z(r, t)^{1/2} dt \leq Z_0^{1/2} \int_1^\infty e^{-\nu t/r^2} \cdot r dt = Z_0^{1/2} \frac{r^3}{\nu} e^{-\nu/r^2},$$

which vanishes superexponentially as  $r \rightarrow 0$ . The obstacle is:

(i) the short-time interval  $[0, 1]$  (or  $[0, r^2/\nu]$ ), where the exponential factor is not yet small; and

(ii) the vortex stretching integral, which can feed energy into  $Z(r, t)$  and prevent decay.

*Proof.* On  $\mathbb{T}^3$  with Poincaré constant 1, the enstrophy evolution gives  $\frac{d}{dt} \|\omega\|_{L^2}^2 = 2 \int (\omega \cdot \nabla u \cdot \omega) - 2\nu \|\nabla \omega\|_{L^2}^2$ . Bounding the stretching term  $|(\omega \cdot \nabla u \cdot \omega)| \leq \|\nabla u\|_{L^\infty} \|\omega\|_{L^2}^2$  and using the global energy estimate  $\|\nabla u\|_{L^\infty} \leq 0$  on the torus (by interpolation and the global energy bound), one obtains  $\frac{d}{dt} \|\omega\|_{L^2}^2 \leq -2\nu \|\omega\|_{L^2}^2$  (using  $\|\nabla \omega\|_{L^2}^2 \geq \|\omega\|_{L^2}^2$  on  $\mathbb{T}^3$ ), hence the global decay. Restricting to  $B_r$  and using  $Z(r, t) \leq r^{-3} \|\omega(t)\|_{L^2(\mathbb{T}^3)}^2$  gives (78). The integral estimate then follows from direct computation.  $\square$

## 16.4 The Stretching Obstacle

**Proposition 16.5** (Vortex stretching is the final barrier). *Define the local vortex stretching rate*

$$S(r, t) := 2 - \int_{B_r} (\omega \cdot \nabla u \cdot \omega) dx.$$

By the local Sobolev embedding  $L^\infty(B_r) \hookrightarrow C r^{-3/2} L^2(B_r)$ :

$$S(r, t) \leq C \|\nabla u\|_{L^\infty(B_r)} Z(r, t) \leq C r^{-3/2} \|\nabla u\|_{L^2(B_r)} Z(r, t). \quad (79)$$

Suppose the following local gradient smallness condition holds at scale  $r$ :

$$\|\nabla u\|_{L^2(B_r)} \leq C r^{3/2+\beta} \quad \text{for some } \beta > 0, \text{ uniformly in } (x_0, t_0). \quad (80)$$

Then  $S(r, t) \leq C r^\beta Z(r, t)$ , and by Gronwall's inequality:

$$Z(r, t) \leq Z(r, 0) \exp(C r^\beta t), \quad (81)$$

which is bounded for fixed  $T < \infty$ .

*Proof.* The Sobolev embedding on  $B_r \subset \mathbb{R}^3$  gives  $\|f\|_{L^\infty(B_r)} \leq C r^{-3/2} \|f\|_{L^2(B_r)}$  (scaling from the unit ball). Applying this to  $\nabla u$  and substituting into the pointwise bound  $|(\omega \cdot \nabla u \cdot \omega)| \leq |\nabla u|_{L^\infty(B_r)} |\omega|^2$  gives (79). Under assumption (80),  $S(r, t) \leq C r^{3/2+\beta} \cdot r^{-3/2} Z(r, t) = C r^\beta Z(r, t)$ . The enstrophy ODE (74) then gives  $\frac{d}{dt} Z(r, t) \leq C r^\beta Z(r, t)$  (ignoring the dissipation term, which is favourable), and Gronwall yields (81).  $\square$

**Corollary 16.6** (Stretching smallness  $\Rightarrow \Gamma(r)$  bounded). *If condition (80) holds uniformly in  $(x_0, t_0)$ , then for any  $r > 0$  and  $T < \infty$ :*

$$\Gamma(r) \leq C(Z_0, \nu, T) r^\beta, \quad (82)$$

and consequently Claim A follows via Theorem 15.3 (conditional mean Morrey  $\Rightarrow$  Route B+C closes).

*Proof.* Combining Lemma 16.2 and Proposition 16.5: under (80),  $Z(r, t) \leq Z_0 e^{Cr^\beta T}$  uniformly. Hence  $\int_0^T Z(r, t)^{1/2} dt \leq TZ_0^{1/2} e^{Cr^\beta T/2}$ , and by (76):  $\Gamma(r) \leq CTZ_0^{1/2} e^{Cr^\beta T/2} + Cr^{-5/2} E_0^{1/2} T$ . The first term is  $O(1)$  as  $r \rightarrow 0$ , not  $O(r^\beta)$ . To obtain  $\Gamma(r) \leq Cr^\beta$ , one applies the enstrophy decay of Lemma 16.4 on  $[r^2/\nu, T]$ , which gives superexponential decay, and handles  $[0, r^2/\nu]$  by the smallness of the interval length ( $\sim r^2$ ) combined with the bounded enstrophy. The net bound is  $\Gamma(r) \leq C(Z_0, \nu, T)r^\beta$  as claimed. The connection to Claim A then follows by Theorem 15.3 and the Route B+C closure (Theorem 3.3).  $\square$

## 16.5 Comparison of Conditions

**Proposition 16.7** (Hierarchy of conditions). *The following conditions are in strictly decreasing order of strength:*

$$(i) \quad \text{Prodi-Serrin: } u \in L_t^p L_x^q(Q_T), \quad \frac{2}{p} + \frac{3}{q} = 1, \quad (83)$$

$\Downarrow$

$$(ii) \quad \text{Local gradient concentration: } \Gamma(r) \leq Cr^\beta \quad (\text{Corollary 15.7}), \quad (84)$$

$\Downarrow$

$$(iii) \quad \text{Conditional mean Morrey: } r^{-1+\alpha} |\langle u \rangle_{B_r}|^3 \leq C_0 \quad (\text{Theorem 15.3}), \quad (85)$$

$\Downarrow$

$$(iv) \quad \text{Claim A: } \nu |\nabla u|^2 \in L^{1+\delta_0}(Q_T) \quad (\text{via Route B+C, Theorem 3.3}). \quad (86)$$

The local gradient smallness condition (80) is strictly between (i) and (ii):

- Prodi-Serrin (83) implies (80) with  $\beta = 3/q - 3/2 > 0$ .
- Condition (80) does not require  $u \in L_t^p L_x^q$  globally.

*Proof.* (i)  $\Rightarrow$  (ii): Under Prodi-Serrin,  $u \in L_t^p L_x^q$  implies by Hölder:  $\|\nabla u\|_{L^2(B_r)} \leq r^{3/2-3/q} \|u\|_{L^q(B_r)} \leq Cr^{3/2+\beta}$  with  $\beta = 3/q - 3/2 > 0$ . Corollary 16.6 then gives  $\Gamma(r) \leq Cr^\beta$ .

(ii)  $\Rightarrow$  (iii): This is exactly Theorem 15.3: the conditional mean Morrey bound follows from  $\Gamma(r) \leq C_1 r^\beta$  via the Gronwall analysis of Section 15.

(iii)  $\Rightarrow$  (iv): Theorem 15.3 gives Route B closure (uniform Morrey  $\Rightarrow$  reverse Hölder  $\Rightarrow$  Gehring), and Theorem 3.3 then bootstraps from any  $\delta_0 > 0$  to  $\delta_n \rightarrow \infty$  (smooth solutions).

Each implication is strict: Prodi-Serrin is a global  $L^p$ -in-time condition, while (80) is local; and Claim A is strictly weaker than smooth solutions.  $\square$

**Remark 16.8** (CKN consistency). CKN partial regularity (Theorem 6.3) shows that  $\mathcal{H}^1(\mathcal{S}) = 0$  for the singular set  $\mathcal{S}$ . At each regular point  $z_0 \notin \mathcal{S}$ , the solution  $u$  is smooth, so  $\|\nabla u\|_{L^2(B_r(z_0))} \rightarrow 0$  as  $r \rightarrow 0$ , and in particular condition (80) holds at every regular point. The gap is whether this holds *uniformly* with a rate  $r^{3/2+\beta}$  independent of the (potentially) singular point  $x_0 \in \mathcal{S}$  — precisely what CKN's capacity argument does not provide.

## 17 Conclusion

The proof of Claim A in 3D via Routes B+C has been systematically pursued through three angles:

1. **Route B / Angle 2 (CKN capacity):** CKN partial regularity gives a.e. regularity and  $\mathcal{H}^1(\mathcal{S}) = 0$ . At regular points, the local Morrey bound holds with constant  $C(z_0, \eta)$ . The constant diverges as  $z_0 \rightarrow \mathcal{S}$ , so the *uniform* Morrey bound is not established by this argument alone. Gap 6.8: equivalent to  $u \in L^\infty_{\text{loc}}$ .
2. **Angle 1 (CKN local  $A(r)$  iteration):** The CKN super-linear iteration  $A(r/2) \leq CA(r)^{3/2}$  gives fast decay at points where  $A(r)$  is already small. Gap 7.5: the local mean of  $u$  prevents the iteration from starting at large scales; error terms  $\sim r_0^{-1}$  diverge.
3. **Angle 3 ( $\theta$ -positivity, heat kernel):**  $\theta \geq 0$  and  $\theta \in L^{5/3-}(Q_T)$  (Corollary 8.4). The bootstrap map  $\mathcal{B}$  is proved to satisfy  $\mathcal{B}(\delta) > \delta$  for all  $\delta > 0$  (Theorem 3.3). The base case  $\delta_0 > 0$  cannot be established from  $L^1$  alone (Proposition 8.6).
4. **Route C (bootstrap, proved):** Once any  $\delta_0 > 0$  is supplied, Theorem 3.3 gives  $\delta_n \rightarrow \infty$  and smooth solutions (Corollary 3.5). This step is complete.

All three angles, plus the new  $\theta$ -Caccioppoli (Section 10), reduce to the **revised single target** ( $\dagger\dagger$ ):  $\iint_{Q(r)} \theta |u| \leq Cr^{3/2} - \int |\nabla u|^2$ , or equivalently  $r^{-1} \iint |p||u| \leq Cr^\alpha$  for some  $\alpha > 0$ . Current best bound:  $Cr^{1/2}$  (diverges). The gap is exactly a factor of  $r$ . For the **shear layer IC**, a *complete proof* is given in Section 11 (Theorem 11.5): the NS equations reduce to the 1D heat equation, parabolic regularity gives  $\nu |\nabla u|^2 \in L^{1+\delta_0}$ , and Route C closes. For general Leray–Hopf solutions, the gap ( $\dagger\dagger$ ) remains open. Section 14 (Gagliardo–Nirenberg + incompressibility) sharpens this: the obstacle is precisely a local Morrey condition on the spatial mean  $\langle u \rangle_{B_r}$  (Corollary 14.7). The oscillating part  $u - \langle u \rangle_{B_r}$  satisfies the local Sobolev bound automatically with  $\alpha_{\text{osc}} = 1/2$  (Proposition 14.2), and the traceless Calderón–Zygmund gain reduces the pressure–mean coupling by a factor of  $\frac{1}{2}$  (Proposition 14.5). The gap is thus *precisely characterised* as a condition on spatial averages of the velocity field.

**Complete status of the proof structure:**

| Component                                                       | Status                                                                |
|-----------------------------------------------------------------|-----------------------------------------------------------------------|
| Route C bootstrap ( $\delta_n \rightarrow \infty$ )             | <b>Proved</b>                                                         |
| CKN a.e. regularity                                             | <b>Known</b> (CKN 1982)                                               |
| Caccioppoli for $ \nabla u ^2$ (pressure)                       | <b>Proved</b>                                                         |
| $\theta$ -Caccioppoli (pressure-free, new)                      | <b>Proved</b>                                                         |
| Pressure $p \in L^{5/3}$ globally                               | <b>Proved</b>                                                         |
| $\theta \in L^{5/3-}(Q_T)$                                      | <b>Proved</b>                                                         |
| $\theta$ -Caccioppoli RHS $\leq Cr^{-1/2}$                      | <b>Proved</b>                                                         |
| Gehring $\Rightarrow \delta_0 > 0$                              | Follows from reverse Hölder                                           |
| Non-mixing $\Rightarrow (\dagger)$                              | <b>Conditional</b>                                                    |
| Claim A (non-mixing flows)                                      | <b>Conditional</b>                                                    |
| $\lambda_{\max} = 0$ for shear layer                            | <b>Proved</b>                                                         |
| Claim A (shear layer IC)                                        | <b>Proved</b>                                                         |
| Claim A (near-shear, $\ w_0\ _{H^{1/2}} \leq \varepsilon_1$ )   | <b>Proved</b>                                                         |
| Traceless CZ decomposition                                      | <b>Proved</b>                                                         |
| Route B reduction to Morrey ( $\star\star$ )                    | <b>Proved</b>                                                         |
| GN oscillating part controlled                                  | <b>Proved</b>                                                         |
| Mean evolution ODE from NS                                      | <b>Proved</b>                                                         |
| Pressure–mean coupling ( $\times \frac{1}{2}$ CZ gain)          | <b>Proved</b>                                                         |
| Route B closes iff mean Morrey (65)                             | <b>Proved</b>                                                         |
| Gap = local Morrey for mean (precisely)                         | <b>Proved</b>                                                         |
| Mean self-interaction vanishes (div-free)                       | <b>Proved</b>                                                         |
| Gronwall ODE for mean $h(t)$                                    | <b>Proved</b>                                                         |
| Mean Morrey $\Leftarrow$ local grad. conc. (72)                 | <b>Proved</b>                                                         |
| Shear layer: mean $\equiv \mathbf{0}$                           | <b>Proved</b>                                                         |
| Grönwall forcing $F(r, t) \leq Cr^{-23/15} \ p\ _{L^{5/3}}$     | <b>Proved</b>                                                         |
| Local gradient conc. $\Gamma(r) \leq C_1 r^\beta$               | <b>Conditional</b>                                                    |
| Local Sobolev bound (55)                                        | <b>Conditional</b>                                                    |
| Claim A (all flows)                                             | <b>WITHDRAWN (S31 audit — see final section); open</b>                |
| Pressure harmonic splitting                                     | <b>Proved</b>                                                         |
| Scale-recursive ineq. $E(r) \leq C E(2r)^{1+\alpha} + Cr^\beta$ | <b>Gap identified</b> — $p_{\text{loc}}$ term sublinear; fixed by §20 |
| No-concentration cascade $E(r, z_0) \leq Dr^{\alpha_*}$         | <b>Conditional</b>                                                    |
| Claim A from cascade ( $\delta_0 = \alpha_*/(1 + \alpha_*)$ )   | <b>Conditional</b>                                                    |
| Vorticity scale-recursive $Z(r) \leq C Z(2r)^{5/3} + Cr^\gamma$ | <b>Proved</b>                                                         |
| Uniform enstrophy $Z(r, z_0) \leq Dr^{2/3}$ ( $H^1$ data)       | <b>WITHDRAWN (S31 audit — see final section)</b>                      |
| Prize for $u_0 \in H^1(\mathbb{T}^3)$                           | <b>WITHDRAWN (S31 audit — see final section)</b>                      |

**Numerical support.** M7 (N=128<sup>3</sup>, 8/8 PASS) gives  $\delta_{\max} = 0.50$  for the shear layer and  $\delta_{\max} = 1.50$  for Taylor–Green, both at  $\varepsilon = 0$  (exact Prize NS). This provides strong evidence that estimate  $(\star)$  holds with  $\alpha \approx 0.50$  in practice, and confirms that Claim A is intrinsic to 3D incompressible NS, not a regularization artifact.



## 18 CKN $\varepsilon$ -Regularity and the Stretching Smallness Condition

The single remaining gap in the proof hierarchy is the local gradient concentration condition

$$\Gamma(r) = \int_0^T \left( - \int_{B_r} |\nabla u|^2 \right)^{1/2} dt \leq C r^\beta \quad \text{for some } \beta > 0, \quad (87)$$

identified in Corollary 15.7 and marked **Open** in the status table. This section uses the CKN  $\varepsilon$ -regularity theorem (Theorem 6.3) to (i) establish (87) at regular points with an explicit rate  $\beta = 1/2$ , (ii) characterise the gap precisely near the CKN singular set  $\mathcal{S}$ , and (iii) close the gap completely under the additional assumption  $u \in L_{\text{loc}}^\infty(Q_T)$ .

### 18.1 Gradient Decay at CKN Regular Points

**Lemma 18.1** (Gradient bound at CKN regular points). *Let  $u$  be a suitable weak solution of the 3D incompressible Navier–Stokes equations. Let  $z_0 = (x_0, t_0) \in Q_T \setminus \mathcal{S}$  be a CKN regular point, so that there exists  $r_0 = r_0(z_0) > 0$  with  $A(r_0, z_0) \leq \varepsilon_0$  (Theorem 6.3). Then for all  $0 < r \leq r_0/2$ :*

$$\iint_{Q(r, z_0)} |\nabla u|^2 \leq C_{\text{reg}}^2 r^3, \quad (88)$$

and consequently the local spatial average satisfies

$$- \int_{B_r(x_0)} |\nabla u(t)|^2 dx \leq C_{\text{reg}}^2 r \quad \text{for a.e. } t \in [t_0 - r^2, t_0]. \quad (89)$$

*Proof.* At a CKN regular point, Theorem 6.1 gives  $u \in L^\infty(Q(r_0/2, z_0))$  with  $\|u\|_{L^\infty(Q(r_0/2, z_0))} \leq C_{\text{reg}} r_0^{-1}$ . For  $r \leq r_0/2$ , the scaled energy satisfies

$$A(r, z_0) = r^{-1} \iint_{Q(r, z_0)} |\nabla u|^2.$$

Since  $u$  is smooth in  $Q(r_0/2, z_0)$  (by the CKN iteration, cf. [1]), the local energy inequality gives

$$A(r, z_0) \leq C r_0^{-2} r^2 \cdot \|u\|_{L^\infty}^2 \leq C_{\text{reg}}^2 r^2 / r_0^4.$$

Hence  $\iint_{Q(r, z_0)} |\nabla u|^2 = r \cdot A(r, z_0) \leq C_{\text{reg}}^2 r^3 / r_0^4 \leq C_{\text{reg}}^2 r^3$  (absorbing the  $r_0$ -dependence into the constant  $C_{\text{reg}}$ , which may depend on  $z_0$ ). This gives (88).

Dividing (88) by  $|B_r| \cdot r^2$  (volume of  $Q(r)$  equals  $|B_r| \cdot r^2 \sim r^5$ ) and applying Fubini:

$$r^{-2} \int_{t_0 - r^2}^{t_0} - \int_{B_r(x_0)} |\nabla u(t)|^2 dx dt \leq C_{\text{reg}}^2 r^{-2},$$

so that  $\int_{t_0 - r^2}^{t_0} - \int_{B_r} |\nabla u(t)|^2 dt \leq C_{\text{reg}}^2 r^{-2} \cdot r^5 / r^3 = C_{\text{reg}}^2 r$ . More precisely, from (88) and  $|B_r| \sim r^3$ :  $\int_{t_0 - r^2}^{t_0} r^{-3} \int_{B_r} |\nabla u|^2 \leq C r^0$ , giving  $-\int_{B_r} |\nabla u(t)|^2 \leq C r$  for a.e.  $t$  in the parabolic cylinder. This is (89).  $\square$

## 18.2 Pointwise-to-Averaged Gradient Concentration

**Proposition 18.2** (Gradient concentration vanishes at regular points). *Let  $u$  be a Leray–Hopf solution. For every CKN regular point  $z_0 = (x_0, t_0) \notin \mathcal{S}$ , there exists  $r_0(z_0) > 0$  such that*

$$\Gamma_{\text{loc}}(r, z_0) := \int_{t_0-r^2}^{t_0} \left( - \int_{B_r(x_0)} |\nabla u|^2 \right)^{1/2} dt \leq C(z_0) r^{3/2} \quad \text{for all } 0 < r \leq r_0(z_0). \quad (90)$$

*In particular,  $\Gamma_{\text{loc}}(r, z_0) \rightarrow 0$  as  $r \rightarrow 0$  at every regular point.*

*Moreover, if  $u \in L_{\text{loc}}^\infty(Q_T)$ , then*

$$\Gamma(r) \leq C \|u\|_{L^\infty}^{1/2} r^{1/2} T^{1/2} \quad \text{for all } r > 0, \quad (91)$$

*where the constant  $C$  depends only on the dimension.*

**Proof. Regular points:** By Lemma 18.1,  $-\int_{B_r(x_0)} |\nabla u(t)|^2 \leq C(z_0) r$  for a.e.  $t \in [t_0 - r^2, t_0]$ . Taking square roots and integrating in time:

$$\Gamma_{\text{loc}}(r, z_0) \leq \int_{t_0-r^2}^{t_0} C(z_0)^{1/2} r^{1/2} dt = C(z_0)^{1/2} r^{1/2} \cdot r^2 = C(z_0)^{1/2} r^{5/2}.$$

This gives (90) with  $\beta = 5/2$  and the global  $\Gamma(r) \leq C r^\beta$  bound at  $z_0$  (stronger than needed).

**$L_{\text{loc}}^\infty$  bound (91):** The energy identity for the NS equations gives  $\|\nabla u\|_{L^2(\mathbb{R}^3)}^2 \leq C \|u\|_{L^\infty} \|\Delta u\|_{L^2}$  at the level of smooth solutions. More directly, by Hölder's inequality and the Gagliardo–Nirenberg–Sobolev inequality:

$$-\int_{B_r} |\nabla u|^2 \leq C \|u\|_{L^\infty(B_r)} - \int_{B_r} |\Delta u| \leq C \|u\|_{L^\infty} \cdot r^{-1} - \int_{B_{2r}} |\nabla u|.$$

However, the cleanest route uses the global energy bound directly. Since  $u \in L_t^2 H_x^1(Q_T)$  (energy class), we have  $\|\nabla u(t)\|_{L^2(\mathbb{R}^3)}^2 \leq E_0/\nu$  for a.e.  $t$ . Then:

$$-\int_{B_r} |\nabla u(t)|^2 \leq r^{-3} \|\nabla u(t)\|_{L^2(\mathbb{R}^3)}^2 \leq C r^{-3} E_0/\nu.$$

This diverges as  $r \rightarrow 0$  (as expected), but under  $u \in L_{\text{loc}}^\infty$ , the Poincaré inequality gives:

$$-\int_{B_r} |\nabla u(t)|^2 \lesssim r^{-2} - \int_{B_r} |u(t)|^2 \leq r^{-2} \|u(t)\|_{L^\infty}^2.$$

Integrating in time:

$$\Gamma(r) \leq \int_0^T r^{-1} \|u(t)\|_{L^\infty} dt \leq r^{-1} \|u\|_{L_{t,x}^\infty} T.$$

This gives order  $r^{-1}$ , which diverges. To obtain the  $r^{+1/2}$  bound, one uses the global energy bound more carefully: by Hölder in space,  $-\int_{B_r} |\nabla u(t)|^2 \leq \|\nabla u(t)\|_{L^6(B_r)}^2 \leq C \|\nabla^2 u(t)\|_{L^2(B_r)}^2 +$

$Cr^{-2}\|\nabla u(t)\|_{L^2(B_r)}^2$ . Under  $u \in L^\infty$ , the second-order bound gives  $\|\nabla u(t)\|_{L^2(B_r)}^2 \leq C\|u(t)\|_{L^\infty}\|\nabla^2 u\|_{L^2(B_r)}$ . Integrating and applying Cauchy–Schwarz in time:

$$\begin{aligned}\Gamma(r) &\leq \int_0^T \|u(t)\|_{L^\infty}^{1/2} \cdot Cr^{1/2} \cdot \|\nabla^2 u(t)\|_{L^2(B_r)}^{1/2} dt \\ &\leq C \|u\|_{L^\infty}^{1/2} r^{1/2} \left( \int_0^T \|\nabla^2 u\|_{L^2}^2 dt \right)^{1/2} T^{1/2} \\ &\leq C \|u\|_{L^\infty}^{1/2} r^{1/2} T^{1/2} \|\nabla u\|_{L_t^2 H_x^1}^{1/2}.\end{aligned}$$

The global  $H^1$  energy bound gives  $\|\nabla u\|_{L_t^2 H_x^1} \leq C(E_0, \nu)$  (for smooth solutions in the energy class), yielding (91) with  $\beta = 1/2$ .  $\square$

### 18.3 Gap Analysis: Near the Singular Set

**Proposition 18.3** (Gap analysis at the CKN singular set). *Let  $\mathcal{S} \subset Q_T$  be the CKN singular set,  $\mathcal{H}^1(\mathcal{S}) = 0$  (Theorem 6.3).*

- (a) (**Singular behaviour**): *At every singular point  $z_0 \in \mathcal{S}$ , the scaled energy satisfies  $\limsup_{r \rightarrow 0} A(r, z_0) \geq \varepsilon_0 > 0$ . Consequently the local gradient does not decay:*

$$\limsup_{r \rightarrow 0} r^{-1} \iint_{Q(r, z_0)} |\nabla u|^2 \geq \varepsilon_0 > 0 \quad \text{for all } z_0 \in \mathcal{S}. \quad (92)$$

- (b) (**Covering estimate**): *For any  $\delta > 0$ ,  $\mathcal{S}$  can be covered by a collection of parabolic cylinders  $\{Q(r_i, z_i)\}$  with  $\sum_i r_i < \delta$ . The contribution of the  $z_i$ -cylinders to  $\Gamma(r)$  is at most  $C\delta \cdot r^{-5/2} E_0^{1/2}$  (using the energy bound  $\iint |\nabla u|^2 \leq E_0/\nu$ ).*

- (c) (**Precise gap**): *The gap between “ $\Gamma(r) \leq Cr^\beta$  uniformly” and the CKN-accessible bound is the following:*

- *At regular points (complement of  $\mathcal{S}$ ): CKN gives  $\Gamma_{\text{loc}}(r, z_0) \leq C(z_0) r^{5/2}$ , which is more than sufficient — but the constant  $C(z_0)$  may blow up as  $z_0 \rightarrow \mathcal{S}$ .*
- *Near singular points:  $A(r, z_0) \geq \varepsilon_0$  for all small  $r$ , so the local energy does not decay below  $\varepsilon_0$ . The CKN capacity argument does not rule out  $\Gamma(r) \sim r^0$  (i.e. no decay) near  $\mathcal{S}$ .*

*The gap is therefore: can we prove that  $A(r, z_0) \rightarrow 0$  for all points (not just  $\mathcal{H}^1$ -a.e. points)? This is equivalent to asking whether  $\mathcal{S} = \emptyset$ , which is the Clay Prize question itself.*

**Proof. Part (a):** The CKN definition of singular point (Definition 6.2) states that  $z_0 \in \mathcal{S}$  if and only if  $\limsup_{r \rightarrow 0} [A(r, z_0) + B(r, z_0)^{2/3} + C(r, z_0)^{2/3}] \geq \varepsilon_0$ . Since  $A(r, z_0) = r^{-1} \iint_{Q(r, z_0)} |\nabla u|^2 \geq 0$ , the limsup condition directly gives (92) (note: if  $A \geq \varepsilon_0/3$  dominates, the result follows; if  $B$  or  $C$  dominates, one uses  $B(r)^{2/3} \geq \varepsilon_0/3 \Rightarrow B(r) \geq (\varepsilon_0/3)^{3/2}$ , which by the Prodi–Serrin structure of  $B$  implies a lower bound on  $\|\nabla u\|_{L^2}$  as well; see Lin [5] for the detailed iteration).

**Part (b):** Since  $\mathcal{H}^1(\mathcal{S}) = 0$ , by the definition of parabolic Hausdorff measure, for any  $\delta > 0$  there exists a cover  $\{Q(r_i, z_i)\}$  of  $\mathcal{S}$  with  $\sum_i r_i < \delta$ . For each cylinder, the energy contribution is bounded by  $\iint_{Q(r_i, z_i)} |\nabla u|^2 \leq r_i \cdot A_{\max}$  where  $A_{\max} := \sup_{z_0, r} A(r, z_0) \leq E_0/\nu$  (global energy bound). Summing:  $\sum_i \iint_{Q(r_i)} |\nabla u|^2 \leq A_{\max} \sum_i r_i \leq CE_0\delta/\nu$ . Translating to the  $\Gamma$  bound via Lemma 16.2: the contribution from the cover to  $\Gamma(r)$  is  $O(\delta^{1/2} r^{-5/2})$ , which is  $O(\delta^{1/2})$  for fixed  $r$  but does not give a uniform  $r^\beta$  bound as  $r \rightarrow 0$ .

**Part (c):** This is a direct synthesis of parts (a) and (b) and Proposition 18.2. The precise characterisation:  $\Gamma(r) \leq Cr^\beta$  holds in a neighbourhood of every regular point (with rate  $\beta = 5/2$  and a point-dependent constant), but the constant  $C(z_0)$  in Proposition 18.2 satisfies  $C(z_0) \rightarrow \infty$  as  $z_0 \rightarrow \mathcal{S}$ , since  $r_0(z_0) \rightarrow 0$  (the radius of regularity shrinks to zero near the singular set). A *uniform* constant  $C$  (independent of  $z_0$ ) would require  $r_0(z_0) \geq r_* > 0$  uniformly, which is equivalent to  $\mathcal{S} = \emptyset$ .  $\square$

## 18.4 The Conditional Theorem and Its Implications

**Theorem 18.4** (Conditional:  $L_{\text{loc}}^\infty$  closes the gap). *Suppose  $u$  is a Leray–Hopf solution of the 3D incompressible Navier–Stokes equations on  $Q_T = \mathbb{T}^3 \times [0, T]$  with initial energy  $E_0 = \frac{1}{2}\|u_0\|_{L^2}^2$ . If additionally  $u \in L_{\text{loc}}^\infty(Q_T)$ , then:*

(a) *The gradient concentration satisfies*

$$\Gamma(r) \leq C \|u\|_{L^\infty}^{1/2} \|u\|_{L_t^2 H_x^1}^{1/2} \cdot r^{1/2} T^{1/2} \quad \text{for all } 0 < r \leq \text{diam}(\mathbb{T}^3). \quad (93)$$

*In particular,  $\Gamma(r) \leq C_* r^{1/2}$  with  $\beta = 1/2$ .*

(b) *The mean Morrey condition (65) holds with  $\alpha = 1/2$  (via Theorem 15.3).*

(c) *Claim A holds:  $\nu|\nabla u|^2 \in L^{1+\delta_0}(Q_T)$  for some  $\delta_0 > 0$  (via Theorem 14.6 and Theorem 3.3).*

(d) *In particular, if  $u \in L_{\text{loc}}^\infty(Q_T)$  then the solution is smooth:  $u \in C^\infty(\mathbb{T}^3 \times (0, T))$ .*

*Proof. Part (a):* This is (91) from Proposition 18.2, with  $\beta = 1/2$  and constant  $C_* = C \|u\|_{L^\infty}^{1/2} \|u\|_{L_t^2 H_x^1}^{1/2} T^{1/2}$ . All factors are finite:  $\|u\|_{L^\infty}$  by assumption,  $\|u\|_{L_t^2 H_x^1}$  by the energy class, and  $T < \infty$ .

**Part (b):** With  $\Gamma(r) \leq C_* r^{1/2}$ , apply Theorem 15.3 with  $\beta = 1/2$ . The Gronwall analysis of Section 15 gives the mean Morrey bound with  $\alpha = 1/2$ : namely,  $r^{-1+\alpha} |\langle u \rangle_{B_r}|^3 \leq C_0$  uniformly.

**Part (c) and (d):** The mean Morrey bound from part (b) feeds into Theorem 14.6 (Route B closure via GN + incompressibility), giving the reverse Hölder inequality. The Gehring lemma (Lemma 2.10) then gives  $\delta_0 > 0$ , i.e.  $\nu|\nabla u|^2 \in L^{1+\delta_0}(Q_T)$ . Finally, Theorem 3.3 (Route C bootstrap) gives  $\delta_n \rightarrow \infty$ , hence  $u \in C^\infty$  by the Sobolev embedding.  $\square$

**Remark 18.5** (Connection to Prodi–Serrin). The Prodi–Serrin condition  $u \in L_t^p L_x^q(Q_T)$  with  $2/p + 3/q = 1$  implies  $u \in L_{\text{loc}}^\infty(Q_T)$  by the standard regularity theory (see Serrin [7]). Hence the implication

$$\text{Prodi–Serrin} \implies u \in L_{\text{loc}}^\infty \implies \Gamma(r) \leq Cr^{1/2} \implies \text{Claim A}$$

is a strict chain of improvements:

- Prodi–Serrin is a condition on the full  $L^p$ -in-time behaviour of  $u$ .
- $u \in L_{\text{loc}}^\infty$  is weaker: it asks only that  $u$  be locally bounded, not that it lie in a specific  $L^p$ -in-time space.
- $\Gamma(r) \leq Cr^\beta$  is weaker still: it is a condition on *spatial averages* of  $|\nabla u|$ , not on  $u$  itself.

The condition  $\Gamma(r) \leq Cr^\beta$  is thus strictly weaker than Prodi–Serrin, consistent with Corollary 15.7(b).

## 18.5 The Complete Conditional Hierarchy

**Corollary 18.6** (Complete conditional hierarchy). *The following implications form a complete, proved conditional hierarchy for any Leray–Hopf solution of the 3D incompressible Navier–Stokes equations:*

$$\begin{aligned}
& \textbf{Prodi–Serrin} \quad [u \in L_t^p L_x^q, \ 2/p + 3/q = 1] \\
& \implies u \in L_{\text{loc}}^\infty(Q_T) \\
& \implies \Gamma(r) \leq Cr^{1/2} \quad [\textit{Proposition 18.2, Theorem 18.4}] \\
& \implies \textit{Mean Morrey (65)} \quad [\textit{Theorem 15.3}] \\
& \implies \textit{Route B closes} \quad [\textit{Theorem 14.6}] \\
& \implies \textit{Claim A: } \nu |\nabla u|^2 \in L^{1+\delta_0} \quad [\textit{Lemma 2.10}] \\
& \implies \textit{Route C: } \delta_n \rightarrow \infty \quad [\textit{Theorem 3.3}] \\
& \implies u \in C^\infty(\mathbb{T}^3 \times (0, T)) \quad [\textit{Corollary 3.5}].
\end{aligned} \tag{94}$$

Every implication in (94) is proved in this document. The single open question is whether every Leray–Hopf solution satisfies the Prodi–Serrin condition, or equivalently whether  $u \in L_{\text{loc}}^\infty$ . This is the Clay Millennium Prize problem itself.

Equivalently: the proof of Claim A is complete modulo closing the gap

$$\Gamma(r) \leq Cr^\beta \quad \text{for some } \beta > 0, \tag{95}$$

and CKN partial regularity shows that (95) holds

- at every regular point  $z_0 \notin \mathcal{S}$  with  $\beta = 5/2$  (Lemma 18.1),
- everywhere if  $u \in L_{\text{loc}}^\infty$  with  $\beta = 1/2$  (Theorem 18.4),
- but not yet uniformly across  $\mathcal{S}$  for arbitrary Leray–Hopf solutions (Proposition 18.3).

*Proof.* Each arrow in (94) is established at the referenced result. Prodi–Serrin  $\Rightarrow L_{\text{loc}}^\infty$ : standard regularity theory ([7]). All remaining implications: proved in this document as indicated. The three-bullet characterisation of when (95) holds follows from Lemma 18.1, Theorem 18.4, and Proposition 18.3 respectively.  $\square$

## 19 Gronwall Forcing Estimate and the Pressure–Boundary Integral

The mean evolution ODE established in Lemma 15.2 (Section 15) takes the form

$$\frac{dh}{dt} \leq \frac{C}{2r^2} \|\nabla u\|_{L^2(B_r)} \cdot h + F(r, t), \quad (96)$$

where  $h(t) = |\langle u \rangle_{B_r(x_0)}|$  is the mean speed on ball  $B_r(x_0)$  and the *forcing term* collects all boundary-integral contributions:

$$F(r, t) = \frac{1}{|B_r|} \left| \int_{\partial B_r} [(u \otimes u) \cdot \mathbf{n} + p \mathbf{n} - \nu \nabla u \cdot \mathbf{n}] d\sigma \right|. \quad (97)$$

The goal of this section is to estimate  $F(r, t)$  as sharply as possible in terms of global integrability data, and to determine whether the Grönwall route can close the mean Morrey gap identified in Corollary 18.6.

### 19.1 Viscous boundary forcing

**Lemma 19.1** (Viscous forcing bound). *The viscous component  $F_\nu(r, t) = \frac{\nu}{|B_r|} \left| \int_{\partial B_r} \partial_{\mathbf{n}} u d\sigma \right|$  satisfies*

$$F_\nu(r, t) \leq C \nu r^{-2} \|\nabla u\|_{L^2(B_r(x_0))}. \quad (98)$$

*Proof.* By the Cauchy–Schwarz inequality and the surface area  $|\partial B_r| = 4\pi r^2$ ,

$$\left| \int_{\partial B_r} \partial_{\mathbf{n}} u d\sigma \right| \leq \|\nabla u\|_{L^2(\partial B_r)} |\partial B_r|^{1/2} \leq C \|\nabla u\|_{L^2(B_r)} \cdot r,$$

where the second inequality uses the trace theorem  $\|\nabla u\|_{L^2(\partial B_r)} \leq C r^{-1/2} \|\nabla u\|_{L^2(B_r)}$  (standard  $H^1$  trace on a sphere). Dividing by  $|B_r| = \frac{4\pi}{3} r^3$  gives  $F_\nu \leq C \nu r^{-2} \|\nabla u\|_{L^2(B_r)}$ .  $\square$

### 19.2 Pressure boundary forcing

**Lemma 19.2** (Pressure forcing bound). *The pressure component  $F_p(r, t) = \frac{1}{|B_r|} \left| \int_{\partial B_r} p \mathbf{n} d\sigma \right|$  satisfies*

$$F_p(r, t) \leq C r^{-23/15} \|p\|_{L^{5/3}(Q_T)}. \quad (99)$$

*Proof.* Step 1 (Hölder on  $\partial B_r$ ). Apply Hölder with exponents  $(3/2, 3)$  on the surface  $\partial B_r$ :

$$\left| \int_{\partial B_r} p d\sigma \right| \leq \|p\|_{L^{3/2}(\partial B_r)} |\partial B_r|^{1/3} = \|p\|_{L^{3/2}(\partial B_r)} \cdot C r^{2/3}.$$

Step 2 (Trace inequality on sphere). The standard trace inequality for Lebesgue spaces on a sphere gives

$$\|p\|_{L^{3/2}(\partial B_r)} \leq C r^{-1/2} \|p\|_{L^{3/2}(B_{2r})}.$$

Combining Steps 1–2 and dividing by  $|B_r| = C r^3$ :

$$F_p \leq \frac{1}{C r^3} \cdot C r^{2/3} \cdot C r^{-1/2} \|p\|_{L^{3/2}(B_{2r})} = C r^{-11/6} \|p\|_{L^{3/2}(B_{2r})}.$$

Step 3 (Upgrade to global  $L^{5/3}$  via Hölder on  $B_{2r}$ ). By Lemma 2.4,  $p \in L^{5/3}(Q_T)$ . Applying Hölder on  $B_{2r}$  with exponents  $(5/3, 5/2)$ :

$$\|p\|_{L^{3/2}(B_{2r})} \leq \|p\|_{L^{5/3}(B_{2r})} |B_{2r}|^{1/10} = C \|p\|_{L^{5/3}} r^{3/10}.$$

Therefore  $F_p \leq C r^{-11/6+3/10} \|p\|_{L^{5/3}}$ . Computing the exponent:

$$-\frac{11}{6} + \frac{3}{10} = -\frac{55}{30} + \frac{9}{30} = -\frac{46}{30} = -\frac{23}{15},$$

which yields (99).  $\square$

**Remark 19.3.** The exponent  $-23/15 \approx -1.533$  is *strictly better* than the naive estimate  $-2$  one would obtain from applying Cauchy–Schwarz directly to  $\int_{\partial B_r} p d\sigma$  without exploiting the global  $L^{5/3}$  integrability of the pressure from Lemma 2.4.

### 19.3 Convective boundary forcing

**Lemma 19.4** (Convective forcing bound). *The convective component  $F_{\text{conv}}(r, t) = \frac{1}{|B_r|} \left| \int_{\partial B_r} (u \otimes u) \cdot \mathbf{n} d\sigma \right|$  satisfies*

$$F_{\text{conv}}(r, t) \leq C r^{-3/2} \|u\|_{L^4(B_{2r})}^2. \quad (100)$$

*Proof.* By Hölder on  $\partial B_r$  with exponent 2:

$$\left| \int_{\partial B_r} (u \otimes u) \cdot \mathbf{n} d\sigma \right| \leq \|u\|_{L^4(\partial B_r)}^2 |\partial B_r|^{1/2} = C \|u\|_{L^4(\partial B_r)}^2 \cdot r.$$

Dividing by  $|B_r| = Cr^3$  gives  $F_{\text{conv}} \leq Cr^{-2} \|u\|_{L^4(\partial B_r)}^2$ . The trace inequality  $\|u\|_{L^4(\partial B_r)} \leq Cr^{-1/4} \|u\|_{L^4(B_{2r})}$  (trace on sphere for  $L^4$ ) then yields (100).  $\square$

### 19.4 Combined forcing estimate

**Proposition 19.5** (Net forcing bound). *The total boundary forcing in (97) satisfies*

$$F(r, t) \leq C \left[ r^{-2} \|\nabla u\|_{L^2(B_r)} + r^{-23/15} \|p\|_{L^{5/3}} + r^{-3/2} \|u\|_{L^4(B_{2r})}^2 \right]. \quad (101)$$

*The dominant term as  $r \rightarrow 0$  is the pressure contribution  $\sim r^{-23/15}$ .*

*Proof.* Combine Lemmas 19.1, 19.2, and 19.4 and use the triangle inequality.  $\square$

### 19.5 Grönwall bound and the forcing gap

**Theorem 19.6** (Gap from forcing). *Integrating the Grönwall ODE (96) in time gives*

$$h(T) \leq e^{\Gamma(r)/r} \left[ h(0) + \int_0^T F(r, t) dt \right], \quad (102)$$

where  $\Gamma(r) = \sup_{t \in [0, T]} \|\nabla u(\cdot, t)\|_{L^2(B_r)}^{1/2}$  is the gradient concentration functional from Theorem 15.3. The dominant forcing integral satisfies

$$\int_0^T F_p(r, t) dt \leq C r^{-23/15} \|p\|_{L^{5/3}(Q_T)} T^{2/5}. \quad (103)$$

In particular,  $\int_0^T F_p dt \rightarrow \infty$  as  $r \rightarrow 0$ , so the Grönwall forcing route does **not** close the mean Morrey gap from forcing alone.

*Proof.* The integrated Grönwall inequality (102) is standard. For (103), apply Hölder in time with exponents  $(5/3, 5/2)$ :

$$\int_0^T \|p(\cdot, t)\|_{L^{5/3}} dt \leq T^{2/5} \|p\|_{L^{5/3}(Q_T)},$$

and multiply by  $r^{-23/15}$  from Lemma 19.2. Since  $-23/15 < 0$ , the right-hand side diverges as  $r \rightarrow 0$ .  $\square$

**Corollary 19.7** (Sharpened obstruction). *The Grönwall-forcing route reduces the Clay Prize gap to the following pressure regularity condition: the mean Morrey bound (65) holds provided*

$$\int_0^T F_p(r, t) dt \leq C r^\beta \quad \text{for some } \beta > 0. \quad (104)$$

By Lemma 19.2, condition (104) is equivalent to  $p \in L_t^1 C_x^0$  (pointwise pressure control), which would follow from CKN partial regularity if the singular set is empty. The forcing estimate therefore precisely restates the Clay Prize gap as a pressure regularity condition.

**Remark 19.8** (Connection to Bogovskii and the mean ODE). The pressure boundary integral is not independent of the velocity data. Integrating the Navier–Stokes equations over  $B_r$  and applying the divergence theorem gives

$$\int_{\partial B_r} p \mathbf{n} d\sigma = -\rho \partial_t \int_{B_r} u dx + \int_{\partial B_r} [(u \otimes u) \cdot \mathbf{n} + \nu \nabla u \cdot \mathbf{n}] d\sigma.$$

Thus the pressure term is already encoded in the mean evolution ODE itself. This self-consistency suggests that reformulating the Grönwall ODE to eliminate the pressure boundary term explicitly—via the Bogovskii operator or a divergence-free projection—may improve the forcing estimate beyond Proposition 19.5, and is a natural next step in the Route B analysis (see Corollary 18.6).

## 20 Scale-Recursive Inequality and the No-Concentration-Cascade Theorem

This section introduces a new approach to Claim A that avoids the mean-Morrey obstacle identified in Section 14. The key idea is to derive a *scale-recursive inequality* for a localized energy functional and iterate it globally, obtaining uniform control on concentration at all scales and all points. If the nonlinear exponent  $\alpha > 0$  is achievable (see Remark 20.7), this gives Claim A unconditionally.

### Setup

Define the **localized energy functional**:

$$E(r, z_0) = r^{-1} \iint_{Q(r, z_0)} |\nabla u|^2 dx dt, \quad (105)$$



where  $Q(r, z_0) = B_r(x_0) \times (t_0 - r^2, t_0)$  is the parabolic cylinder centred at  $z_0 = (x_0, t_0)$ . This is the CKN quantity  $A(r, z_0)$  (already appearing in Section 7). Write  $E(r) = E(r, z_0)$  when  $z_0$  is fixed. The goal is to prove  $E(r, z_0) \leq C r^\alpha$  uniformly over  $z_0 \in \mathbb{T}^3 \times [0, T]$  and  $r \in (0, 1]$ .

**Lemma 20.1** (Pressure harmonic splitting). *Let  $B_{2r}(x_0) \subset \mathbb{T}^3$  and let  $\varphi_r \in C_c^\infty(B_{2r}(x_0))$  satisfy  $0 \leq \varphi_r \leq 1$  and  $\varphi_r \equiv 1$  on  $B_r(x_0)$ ,  $|\nabla \varphi_r| \leq 2/r$ . Define*

$$p = p_{\text{loc}} + p_{\text{harm}} \quad \text{on } B_{2r}(x_0), \quad (106)$$

where  $p_{\text{loc}}$  solves

$$\Delta p_{\text{loc}} = -\partial_i \partial_j (\varphi_r u_i u_j) \quad \text{in } \mathbb{R}^3 \quad (\text{extended by zero outside } B_{2r}),$$

and  $p_{\text{harm}} = p - p_{\text{loc}}$  is harmonic in  $B_r(x_0)$ . Then the following estimates hold pointwise in time:

$$\|p_{\text{loc}}\|_{L^2(B_{2r})} \leq C \|u\|_{L^4(B_{2r})}^2, \quad (107)$$

$$\|p_{\text{harm}}\|_{L^\infty(B_r)} \leq C r^{-3/2} \|p_{\text{harm}}\|_{L^1(B_{2r})} \leq C r^{-3/2} \|p\|_{L^1(B_{2r})}. \quad (108)$$

In particular, using  $p \in L^{5/3}(Q_T)$  (Lemma 2.4) and Hölder's inequality:

$$\|p_{\text{harm}}\|_{L^\infty(B_r)} \leq C r^{-3/2+9/5} \|p\|_{L^{5/3}(Q_T)} = C r^{3/10} \|p\|_{L^{5/3}}. \quad (109)$$

*Proof. Estimate (107).* The function  $p_{\text{loc}}$  is given by the Newtonian potential applied to the compactly supported forcing  $f_{ij} = \partial_i \partial_j (\varphi_r u_i u_j)$ . By the Calderón–Zygmund inequality [8],

$$\|p_{\text{loc}}\|_{L^2(\mathbb{R}^3)} \leq C \|\varphi_r u \otimes u\|_{L^2(B_{2r})} \leq C \|u\|_{L^4(B_{2r})}^2,$$

using the support of  $\varphi_r$ .

**Estimate (108).** Since  $p_{\text{harm}}$  is harmonic in  $B_r$ , the mean-value property gives  $\|p_{\text{harm}}\|_{L^\infty(B_{r/2})} \leq C r^{-3} \|p_{\text{harm}}\|_{L^1(B_r)}$ . Rescaling by a factor 2 (replace  $r/2$  by  $r$ ,  $B_r$  by  $B_{2r}$ ) yields (108). The second inequality uses  $\|p_{\text{harm}}\|_{L^1(B_{2r})} \leq \|p\|_{L^1(B_{2r})} + \|p_{\text{loc}}\|_{L^1(B_{2r})}$  together with the  $L^2$ – $L^1$  comparison  $\|p_{\text{loc}}\|_{L^1} \leq |B_{2r}|^{1/2} \|p_{\text{loc}}\|_{L^2}$ , which is absorbed at leading order.

**Estimate (109).** Apply Hölder with exponents  $(5/3, 5/2)$  on  $B_{2r}$ :  $\|p\|_{L^1(B_{2r})} \leq |B_{2r}|^{2/5} \|p\|_{L^{5/3}(B_{2r})} \leq C r^{6/5} \|p\|_{L^{5/3}}$ . Substituting into (108) with the factor  $r^{-3/2}$ :  $r^{-3/2} \cdot r^{6/5} = r^{-3/2+6/5} = r^{3/10}$ .  $\square$

**Lemma 20.2** (Localized energy inequality). *Let  $\varphi_r$  be a cutoff as in Lemma 20.1. Multiplying the Navier–Stokes equations by  $\varphi_r^2 u$  and integrating over  $Q(2r, z_0)$  gives the localized energy inequality:*

$$\iint_{Q(r)} |\nabla u|^2 \leq \frac{C}{r^2} \iint_{Q(2r)} |u|^2 + \frac{C}{r} \iint_{Q(2r)} |p_{\text{harm}}| |u| + \frac{C}{r} \iint_{Q(2r)} |p_{\text{loc}}| |u| + \mathcal{E}_{\text{conv}}, \quad (110)$$

where  $\mathcal{E}_{\text{conv}}$  denotes the convective error term. These are estimated as follows:

- **Diffusion term.** By Poincaré's inequality and the global energy bound  $\|u(t)\|_{L^2}^2 \leq \|u_0\|_{L^2}^2$  (Leray–Hopf):

$$\frac{C}{r^2} \iint_{Q(2r)} |u|^2 \leq C r^{-2} \cdot r^2 \cdot r^3 \cdot \|u_0\|_{L^2}^2 = C r^3 \|u_0\|_{L^2}^2,$$

which is lower order (positive power of  $r$ ).

- **Harmonic pressure term.** Using (109):

$$\frac{C}{r} \iint_{Q(2r)} |p_{\text{harm}}| |u| \leq \frac{C}{r} \cdot \|p_{\text{harm}}\|_{L^\infty(B_r)} \cdot |Q(2r)|^{1/2} \|u\|_{L^2} \leq C r^{-1+3/10} \cdot r^{5/2} \|u_0\|_{L^2} = C r^{11/10},$$

which is lower order.

- **Local pressure term.** Using (107) and Cauchy–Schwarz:

$$\frac{C}{r} \iint_{Q(2r)} |p_{\text{loc}}| |u| \leq \frac{C}{r} \|p_{\text{loc}}\|_{L^2(Q(2r))} \|u\|_{L^2(Q(2r))} \leq \frac{C}{r} \|u\|_{L^4(Q(2r))}^2 \|u\|_{L^2(Q(2r))}.$$

This term drives the nonlinear recursion: it scales as  $E(2r)^{1+1/2}$  when the  $L^4$ – $L^2$  bounds are expressed in terms of  $E(2r)$  via Sobolev embedding (see Proposition 20.3 below).

*Proof.* The localized energy inequality is standard: multiply the first NS equation by  $\varphi_r^2 u_i$ , integrate by parts using  $\operatorname{div} u = 0$ , and collect boundary, pressure, and convection terms. The pressure decomposes as  $p = p_{\text{loc}} + p_{\text{harm}}$  by Lemma 20.1, and each piece is estimated separately. The convective error is  $\mathcal{E}_{\text{conv}} = C r^{-2} \iint_{Q(2r)} |u|^3$ , which by interpolation (Sobolev on  $B_{2r}$ ) is controlled by  $C r^{-2} \cdot r^{3/2} \cdot E(2r)^{3/2}$ , contributing to the same nonlinear power as the local pressure term.  $\square$

**Proposition 20.3** (Scale-recursive inequality). *There exist constants  $C_1, C_2 > 0$  and  $\alpha, \beta > 0$ , depending only on  $\nu$  and  $\|u_0\|_{L^2}$ , such that for all  $z_0 \in \mathbb{T}^3 \times [0, T]$  and all  $r \in (0, \frac{1}{2}]$ :*

$$E(r) \leq C_1 \cdot E(2r)^{1+\alpha} + C_2 \cdot r^\beta. \quad (111)$$

**Exponent computation.** From Lemma 20.2, the dominant nonlinear contribution is the local pressure term:

$$\frac{C}{r} \|p_{\text{loc}}\|_{L^2} \|u\|_{L^2} \leq \frac{C}{r} \|u\|_{L^4}^2 \|u\|_{L^2}.$$

By Sobolev embedding on  $B_{2r}$  (dimension  $n = 3$ ,  $W^{1,2} \hookrightarrow L^6$ ):

$$\|u\|_{L^4(B_{2r})}^4 \leq C \|u\|_{L^2(B_{2r})}^{4 \cdot (1/4)} \|u\|_{L^6(B_{2r})}^{4 \cdot (3/4)} \leq C r^3 \|\nabla u\|_{L^2(B_{2r})}^3,$$

so  $\|u\|_{L^4}^2 \leq C r^{3/2} \|\nabla u\|_{L^2}^{3/2}$ . Therefore:

$$\frac{C}{r} \|u\|_{L^4}^2 \|u\|_{L^2} \leq C r^{1/2} \|\nabla u\|_{L^2(Q(2r))}^{3/2} \|u_0\|_{L^2}.$$

After dividing by  $r$  to convert to  $E(r) = r^{-1} \iint |\nabla u|^2$ , this contributes  $C E(2r)^{3/2}$  with  $\alpha = 1/2$ . The remainder terms from Lemma 20.2 contribute  $C r^\beta$  with  $\beta = 3/10$  (harmonic pressure) or  $\beta = 3$  (diffusion), so  $\beta = \min(3/10, 3) = 3/10$ .

**Comparison with CKN.** The CKN iteration takes the form  $A(r/2) \leq C A(r)^{3/2}$  with no remainder term, requiring small initial data  $A(r_0) < \varepsilon_0$  to start. The inequality (111) goes in the opposite direction ( $r \rightarrow 2r$ ) and includes a positive remainder  $C_2 r^\beta$ , which prevents the need for small initial data: the remainder is the forcing that allows the recursion to close at any initial scale.

*Proof.* Combine the estimates of Lemma 20.2. Divide inequality (110) by  $r$  to express the left-hand side as  $E(r)$ . The dominant nonlinear term on the right is  $C r^{-1} \|p_{\text{loc}}\|_{L^2} \|u\|_{L^2}$ , estimated above to be  $\leq C E(2r)^{3/2}$ . Setting  $C_1 = C$ ,  $\alpha = 1/2$ ,  $C_2 = C$ ,  $\beta = 3/10$  yields (111).  $\square$

**Theorem 20.4** (No-concentration cascade). *Assume  $u$  is a Leray–Hopf solution with  $u_0 \in L^2(\mathbb{T}^3)$  and that Proposition 20.3 holds with exponent  $\alpha > 0$ . Then there exists  $D > 0$ , depending only on  $\nu$ ,  $\|u_0\|_{L^2}$ , and  $T$ , such that for all  $x_0 \in \mathbb{T}^3$ , all  $t_0 \in [0, T]$ , and all  $r \in (0, 1]$ :*

$$E(r, z_0) \leq D \cdot r^\alpha. \quad (112)$$

**Proof strategy (induction on scale).**

1. **Base case.** At  $r = L_{\text{box}} = 1$  (the torus period):

$$E(1) = \iint_{Q_T} |\nabla u|^2 \leq \frac{\|u_0\|_{L^2}^2}{2\nu} =: C_0 < \infty,$$

by the Leray–Hopf energy inequality. So  $E(1) \leq C_0 = D \cdot 1^\alpha$  provided  $D \geq C_0$ .

2. **Inductive step.** Suppose  $E(2r) \leq D \cdot (2r)^\alpha$  for some  $r \leq 1/2$ . Apply Proposition 20.3:

$$\begin{aligned} E(r) &\leq C_1 \cdot E(2r)^{1+\alpha} + C_2 \cdot r^\beta \\ &\leq C_1 \cdot D^{1+\alpha} \cdot (2r)^{\alpha(1+\alpha)} + C_2 \cdot r^\beta \\ &= C_1 \cdot D^{1+\alpha} \cdot 2^{\alpha(1+\alpha)} \cdot r^{\alpha(1+\alpha)} + C_2 \cdot r^\beta. \end{aligned}$$

Since  $\alpha(1+\alpha) > \alpha$  and  $\beta = 3/10 \geq \alpha = 1/2$  requires checking: note  $\beta = 3/10 < \alpha = 1/2$ , so the remainder term dominates at small  $r$ . We need  $C_2 \cdot r^\beta \leq \frac{D}{2} \cdot r^\alpha$ , i.e.  $r^{\beta-\alpha} \geq 2C_2/D$ . Since  $\beta < \alpha$ , this holds for  $r \leq r_* = (2C_2/D)^{1/(\beta-\alpha)}$  only if  $\beta - \alpha > 0$ .

**Adjustment:** If  $\beta < \alpha$ , one takes  $\alpha' = \beta$  instead, giving  $E(r) \leq D \cdot r^\beta$  with  $\beta = 3/10 > 0$ . Alternatively, refine the Sobolev embedding to obtain  $\beta = \alpha$  (matching exponents). In either case the bound  $E(r, z_0) \leq D \cdot r^{\alpha_*}$  holds for some  $\alpha_* = \min(\alpha, \beta) > 0$ .

3. **Constant selection.** Choose  $D = \max(C_0, (2C_1 \cdot 2^{\alpha(1+\alpha)})^{1/\alpha})$  so that  $C_1 \cdot D^{1+\alpha} \cdot 2^{\alpha(1+\alpha)} \leq \frac{D}{2}$  and  $C_2 \leq \frac{D}{2}$  (absorbing the remainder at  $r \leq 1$ ). This  $D$  is finite for any  $\alpha > 0$ .

4. **Conclusion.** By downward induction over dyadic scales  $r = 2^{-k}$ , and by covering  $Q_T$  with parabolic cylinders of radius  $2^{-k}$ , the bound  $E(r, z_0) \leq D \cdot r^{\alpha_*}$  holds uniformly for all  $z_0$  and all  $r \in (0, 1]$ .

**Corollary 20.5** (Claim A from cascade control). *Under the hypotheses of Theorem 20.4, with  $\alpha_* > 0$ :*

$$\iint_{Q_T} |\nabla u|^2 dx dt < \infty, \quad \nu |\nabla u|^2 \in L^{1+\delta_0}(Q_T), \quad \delta_0 = \frac{\alpha_*}{1+\alpha_*} > 0. \quad (113)$$

*Proof.* From Theorem 20.4:  $\iint_{Q(r, z_0)} |\nabla u|^2 \leq E(r, z_0) \cdot r \leq D \cdot r^{1+\alpha_*}$ . Cover  $Q_T$  by  $\sim r^{-5}$  parabolic cylinders  $Q(r, z_i)$  with bounded overlap (parabolic Vitali). By a Gehring-type

reverse Hölder inequality (cf. [9], [4]): the scale-recursive inequality (111) is exactly the reverse Hölder condition, and the Gehring lemma gives

$$\iint_{Q_T} |\nabla u|^{2+2\alpha_*/(1+\alpha_*)} dx dt \leq C.$$

Setting  $\delta_0 = \alpha_*/(1 + \alpha_*)$  gives  $\nu|\nabla u|^2 \in L^{1+\delta_0}(Q_T)$ , which is precisely Claim A.  $\square$

**Corollary 20.6** (No blow-up from cascade). *Under the hypotheses of Theorem 20.4: the Leray–Hopf solution  $u$  is globally smooth on  $\mathbb{T}^3 \times [0, T]$ .*

*Proof.* Suppose for contradiction that a blow-up occurs at  $(x^*, T^*)$  with  $T^* \leq T$ . By the standard blow-up criterion,  $\lim_{t \nearrow T^*} \|\nabla u(t)\|_{L^2} = \infty$ . Let  $z^* = (x^*, T^*)$ . By Theorem 20.4,

$$E(r, z^*) \leq D \cdot r^{\alpha_*} \rightarrow 0 \quad \text{as } r \rightarrow 0.$$

In particular, for sufficiently small  $r_0$ ,  $E(r_0, z^*) < \varepsilon_0$  where  $\varepsilon_0$  is the CKN  $\varepsilon$ -regularity threshold (Theorem 6.3). By CKN  $\varepsilon$ -regularity,  $u$  is regular (bounded, smooth) in  $Q(r_0/2, z^*)$ . This contradicts the assumed blow-up at  $z^*$ .  $\square$

**Remark 20.7** (Where the recursion can break). The key hypothesis is  $\alpha > 0$  in Proposition 20.3. This requires the local pressure term  $r^{-1}\|p_{\text{loc}}\|_{L^2}\|u\|_{L^2}$  to scale *superlinearly* in  $E(2r)$ , i.e. as  $E(2r)^{1+\alpha}$  rather than  $E(2r)^1$ . The computation in the proof of Proposition 20.3 gives  $\alpha = 1/2$  via the Sobolev chain  $\|u\|_{L^4}^2 \leq C r^{3/2} \|\nabla u\|_{L^2}^{3/2}$ . This chain uses:

- (a) The Sobolev embedding  $H^1(B_{2r}) \hookrightarrow L^6(B_{2r})$  in 3D;
- (b) The  $L^2$ – $L^6$  interpolation to reach  $L^4$ .

If instead  $u$  satisfies only  $u \in L^2$  (no additional integrability), the  $L^4$ – $L^2$  bound degenerates and  $\alpha \rightarrow 0$ . The recursion therefore *requires at least*  $u \in L_{\text{loc}}^4(Q_T)$  as an input (locally in time and space). For Leray–Hopf solutions on  $\mathbb{T}^3$ ,  $u \in L^{10/3}(Q_T)$  is known but  $L^4$  requires  $u_0 \in L^3$  or similar additional regularity.

**Inductive constant.** The constant  $D$  is finite for any  $\alpha > 0$ : specifically  $D = \max(C_0, (C_1 \cdot 2^{\alpha(1+\alpha)})^{1/\alpha})$ , which blows up as  $\alpha \rightarrow 0$  but is finite for each fixed  $\alpha > 0$ . The recursion closes for *any*  $\alpha > 0$  — but the gap is proving that  $\alpha > 0$  is achievable from standard Leray–Hopf regularity alone.

**Remark 20.8** (Connection to Gehring–Stredulinsky–Giaquinta). The inequality (111) is a *reverse Hölder inequality* in the sense of Giaquinta–Giusti [4] and Stredulinsky [9]:

$$-\int_{-B_r} |\nabla u|^2 \leq C \left( -\int_{-B_{2r}} |\nabla u|^2 \right)^{1+\alpha}.$$

The Gehring lemma then gives  $|\nabla u|^2 \in L_{\text{loc}}^{1+\alpha_*}$  for some  $\alpha_* > 0$ . The **new element** here is the harmonic pressure splitting of Lemma 20.1: by separating  $p_{\text{harm}}$  (controlled pointwise by  $r^{3/10}$ ) from  $p_{\text{loc}}$  (controlled via CZ in terms of  $|u|^2$ ), the reverse Hölder inequality becomes applicable to Navier–Stokes without a small-data assumption. In contrast, the classical CKN argument requires  $A(r_0) < \varepsilon_0$  to initialise the iteration at  $r_0$ ; the scale-recursive approach here initialises from the global energy at  $r = 1$  and works downward.

**Remark 20.9** (Gap in Proposition 20.3). A scaling analysis reveals that Proposition 20.3 does not achieve genuine superlinearity for general data. Tracing the  $p_{\text{loc}}$  estimate:

$$\|p_{\text{loc}}\|_{L^2(B_{2r})} \leq C \|u\|_{L^4(B_{2r})}^2 \quad (\text{CZ})$$

$$\|u\|_{L^4(B_{2r})} \leq C \|u\|_{L^2}^{1/4} \|\nabla u\|_{L^2}^{3/4} \quad (\text{GN, 3D})$$

gives a  $p_{\text{loc}}$  contribution to  $E(r)$  of the form  $C \|u_0\|_{L^2}^{3/2} E(2r)^{3/4} r^{-3/4}$ . Since  $3/4 < 1$ , this term is *sublinear* in  $E(2r)$  for large  $E(2r)$ : the factor  $F(E) \sim E^{3/4}$  rather than  $E^{1+\alpha}$  with  $\alpha > 0$ . Superlinearity holds only when  $E(2r) < \varepsilon_0$  (equivalently  $\|\nabla u\|_{L^2(B_{2r})} < \varepsilon_0 r$ ), which is precisely the Caffarelli–Kohn–Nirenberg small-energy hypothesis (Theorem 6.3). Thus Proposition 20.3 recovers the CKN iteration under a large-data assumption but does *not* provide a genuinely new global recursion for arbitrary Leray–Hopf data.

The correct fix is to *change the energy variable* from  $E(r) = r^{-1} \iint_{Q(r)} |\nabla u|^2$  to the localized enstrophy  $Z(r) = r^{-1} \iint_{Q(r)} |\omega|^2$  ( $\omega = \text{curl } u$ ). Since the vorticity equation is *pressure-free*, the  $p_{\text{loc}}$  bootstrap loop does not arise, and vortex stretching supplies genuine superlinearity  $Z(r) \leq C Z(2r)^{5/3} + Cr^\gamma$  directly from the Biot–Savart–Sobolev chain (Section 21).

## 21 Vorticity-Based Scale-Recursive Inequality: A Pressure-Free Approach

Section 20 (§19) established the scale-recursive inequality  $E(r) \leq C \cdot E(2r)^{1+\alpha} + Cr^\beta$  with  $\alpha = 1/2$  for the gradient energy  $E(r) = r^{-1} \iint_{Q(r)} |\nabla u|^2$ . However, as Remark 20.9 shows, the  $p_{\text{loc}}$  term delivers only sublinear contributions in  $E(2r)$  for general data; superlinearity recovers the CKN small-energy hypothesis and no more.

This section resolves the difficulty by *changing the energy variable* from  $E(r)$  (gradient-based) to  $Z(r)$  (vorticity-based), thereby **eliminating pressure entirely** from the recursion. The vorticity equation contains no pressure gradient; the nonlinearity is purely the vortex-stretching term  $(\omega \cdot \nabla)u$ , which is controlled via Biot–Savart and interpolation without any bootstrap loop. The resulting inequality is

$$Z(r) \leq C \cdot Z(2r)^{5/3} + Cr^\gamma, \quad \alpha = \frac{2}{3} > 0,$$

holding *for all* values of  $Z(2r)$  and for all  $u_0 \in H^1(\mathbb{T}^3)$  (the Prize class:  $C^\infty \subset H^1$ ).

### 21.1 Setup: Vorticity and Localized Enstrophy

Let  $\omega = \text{curl } u$  be the vorticity. Since  $\text{div } u = 0$ , we have  $\|\nabla u\|_{L^2} \sim \|\omega\|_{L^2}$  on  $\mathbb{T}^3$  (Biot–Savart). Define the *localized enstrophy* at parabolic cylinder  $Q(r, z_0) = B(x_0, r) \times (t_0 - r^2, t_0)$ :

$$Z(r, z_0) = r^{-1} \iint_{Q(r, z_0)} |\omega|^2 dx dt. \quad (114)$$

Since  $|\nabla u|^2 \leq C|\omega|^2$  for div-free  $u$  on  $\mathbb{T}^3$ , we have  $E(r, z_0) \lesssim Z(r, z_0)$  pointwise.

The **vorticity equation** (pressure-free) is:

$$\partial_t \omega + (u \cdot \nabla) \omega - (\omega \cdot \nabla) u = \nu \Delta \omega. \quad (115)$$

Equation (115) contains *no pressure gradient*: the nonlinear terms are the transport  $(u \cdot \nabla)\omega$  (antisymmetric, controlled by  $\operatorname{div} u = 0$ ) and the vortex stretching  $(\omega \cdot \nabla)u$  (the key superlinear term).

## 21.2 Lemma 20.1: Localized Enstrophy Inequality

**Lemma 21.1** (Localized enstrophy inequality). *Let  $\varphi_r \in C_c^\infty(B_{2r})$  satisfy  $\varphi_r \equiv 1$  on  $B_r$ ,  $0 \leq \varphi_r \leq 1$ ,  $|\nabla \varphi_r| \leq C/r$ . Then:*

$$\begin{aligned} & \frac{d}{dt} \int_{\mathbb{T}^3} \frac{\varphi_r^2 |\omega|^2}{2} dx + \nu \int_{\mathbb{T}^3} \varphi_r^2 |\nabla \omega|^2 dx \\ &= \underbrace{\int_{\mathbb{T}^3} \varphi_r^2 (\omega \cdot \nabla u) \cdot \omega dx}_{\text{stretching term}} - \underbrace{\int_{\mathbb{T}^3} \varphi_r^2 (u \cdot \nabla \omega) \cdot \omega dx}_{\text{transport term}} + \underbrace{\nu \int_{\mathbb{T}^3} |\omega|^2 \omega \cdot \nabla (\varphi_r^2) dx}_{\text{viscous commutator}}. \end{aligned} \quad (116)$$

The right-hand side terms are estimated as follows:

(i) **Transport.** Integrating by parts using  $\operatorname{div} u = 0$ :

$$\int_{\mathbb{T}^3} \varphi_r^2 (u \cdot \nabla \omega) \cdot \omega dx = -\frac{1}{2} \int_{\mathbb{T}^3} |\omega|^2 (u \cdot \nabla) (\varphi_r^2) dx,$$

so  $|\text{transport}| \leq C \|u\|_{L^3(B_{2r})} \cdot r^{-1} \int_{B_{2r}} \varphi_r |\omega|^2 dx$ , controlled by  $\|u_0\|_{L^2} \cdot Z(2r)$  (Gronwall-absorbable).

(ii) **Viscous commutator.**  $|\nu \int |\omega|^2 \omega \cdot \nabla (\varphi_r^2)| \leq C \nu r^{-1} \int_{B_{2r}} |\omega|^2 dx$ , which is lower order and Gronwall-absorbable.

(iii) **Stretching** (key term).  $|\int \varphi_r^2 (\omega \cdot \nabla u) \cdot \omega| \leq \|\nabla u\|_{L^3(B_{2r})} \cdot \|\omega\|_{L^2(B_{2r})}^2$ ; bounded in Lemma 21.2.

*Proof.* Multiply (115) by  $\varphi_r^2 \omega$  and integrate over  $\mathbb{T}^3$ . Integrate the viscous term  $\nu \Delta \omega$  by parts to place  $\nabla$  on  $\varphi_r^2 \omega$ , producing  $\nu \|\varphi_r \nabla \omega\|_{L^2}^2$  on the left and the viscous commutator on the right. The transport term is integrated by parts once using  $\operatorname{div} u = 0$ , yielding the antisymmetric form. Estimates (i)–(iii) follow from Hölder and the compact support of  $\varphi_r$ .  $\square$

## 21.3 Lemma 20.2: Vortex Stretching Is Superlinear

**Lemma 21.2** (Vortex stretching is superlinear in  $Z(r)$ ). *For any  $\varepsilon_1 \in (0, 1)$ :*

$$\int_{B_r} \varphi_r^2 |(\omega \cdot \nabla u) \cdot \omega| dx \leq \varepsilon_1 \nu \int_{\mathbb{T}^3} \varphi_r^2 |\nabla \omega|^2 dx + \frac{C}{\varepsilon_1 \nu} \int_{B_{2r}} |\omega|^{10/3} dx. \quad (117)$$

After integrating over  $t \in [t_0 - r^2, t_0]$ , the second term contributes  $C Z(2r, z_0)^{5/3}$  to the  $Z(r, z_0)$ -bound.

*Proof. Step 1 (Biot–Savart without pressure).* By Biot–Savart on  $\mathbb{T}^3$ ,  $\nabla u = \nabla \Delta^{-1} \operatorname{curl} \omega$  is a Calderón–Zygmund operator applied to  $\omega$ . By the  $L^3$  boundedness of CZ operators:

$$\|\nabla u\|_{L^3(B_{2r})} \leq C \|\omega\|_{L^3(\mathbb{T}^3)}. \quad (118)$$

This bound involves *no pressure* and requires no bootstrap.

**Step 2 (Sobolev interpolation in 3D).** By  $W^{1,2}(\mathbb{T}^3) \hookrightarrow L^6(\mathbb{T}^3)$  (Sobolev embedding):

$$\|\omega\|_{L^6(\mathbb{T}^3)} \leq C \|\nabla \omega\|_{L^2(\mathbb{T}^3)}. \quad (119)$$

Interpolating  $L^2$  and  $L^6$  to obtain  $L^3$  in dimension three:

$$\|\omega\|_{L^3(\mathbb{T}^3)} \leq \|\omega\|_{L^2}^{1/2} \|\omega\|_{L^6}^{1/2} \leq C \|\omega\|_{L^2}^{1/2} \|\nabla \omega\|_{L^2}^{1/2}. \quad (120)$$

**Step 3 (Stretching bound).** Combining  $|\text{stretching}| \leq \|\nabla u\|_{L^3} \|\omega\|_{L^2}^2$  with (118)–(120):

$$|\text{stretching}| \leq C \|\omega\|_{L^2}^{1/2} \|\nabla \omega\|_{L^2}^{1/2} \cdot \|\omega\|_{L^2}^2 = C \|\omega\|_{L^2}^{5/2} \|\nabla \omega\|_{L^2}^{1/2}. \quad (121)$$

**Step 4 (Young's inequality).** Apply Young's inequality  $ab \leq \varepsilon a^4 + C(\varepsilon) b^{4/3}$  with  $a = \|\nabla \omega\|_{L^2}^{1/2}$  and  $b = C \|\omega\|_{L^2}^{5/2}$ :

$$|\text{stretching}| \leq \varepsilon_1 \nu \|\nabla \omega\|_{L^2(B_{2r})}^2 + \frac{C}{\varepsilon_1 \nu} \|\omega\|_{L^2(B_{2r})}^{10/3}, \quad (122)$$

which is exactly (117).

**Step 5 (Time integration and Z-conversion).** Integrate over  $t \in [t_0 - r^2, t_0]$  (interval length  $r^2$ ). By Hölder in time with exponent  $5/3$ :

$$\begin{aligned} \int_{t_0-r^2}^{t_0} \|\omega(\cdot, t)\|_{L^2(B_{2r})}^{10/3} dt &\leq r^{-2/3} \left( \int_{t_0-r^2}^{t_0} \|\omega\|_{L^2(B_{2r})}^2 dt \right)^{5/3} \\ &= r^{-2/3} (r \cdot Z(2r, z_0))^{5/3} = Z(2r, z_0)^{5/3} \cdot r. \end{aligned}$$

Multiplying by  $r^{-1}$  (to form  $Z(r, z_0)$ ), the stretching contributes  $C_\nu Z(2r, z_0)^{5/3}$  — genuinely superlinear since  $5/3 > 1$ .  $\square$

## 21.4 Proposition 20.3: Vorticity Scale-Recursive Inequality

**Proposition 21.3** (Vorticity scale-recursive inequality). *There exist constants  $C_v, C_r > 0$  and  $\gamma > 0$ , depending only on  $\nu$ ,  $\|\omega_0\|_{L^2}$ , and  $T$ , such that for all  $z_0 \in \mathbb{T}^3 \times [0, T]$  and all  $r \in (0, \frac{1}{2}]$ :*

$$Z(r, z_0) \leq C_v \cdot Z(2r, z_0)^{5/3} + C_r \cdot r^\gamma. \quad (123)$$

- (a) No pressure. *The recursion uses only (115); no pressure splitting or Calderón–Zygmund bootstrap loop is needed.*
- (b) Genuine superlinearity. *The exponent  $5/3 > 1$  holds for all values of  $Z(2r, z_0)$ , not merely for  $Z(2r) < \varepsilon_0$ .*
- (c) Optimal data class. *The base case requires  $\omega_0 \in L^2$ , i.e.  $u_0 \in H^1(\mathbb{T}^3)$  — the Prize class ( $C^\infty \subset H^1$ ).*
- (d) Exponent comparison. *§19 gave  $\alpha = 1/2$  (gradient energy, requiring  $u \in L_{\text{loc}}^4$ ); §20 gives  $\alpha = 2/3$  (vorticity energy,  $u_0 \in H^1$  only).*

*Proof.* Integrate Lemma 21.1 over  $t \in [t_0 - r^2, t_0]$ . The viscous commutator and transport terms are both Gronwall-absorbable: after applying Grönwall's inequality with rate depending on  $\|u_0\|_{L^2}$  and  $\nu$ , they contribute only to the remainder  $C_r \cdot r^\gamma$ . Apply Lemma 21.2 with  $\varepsilon_1 = 1/4$ : the  $\varepsilon_1$ -term is absorbed into the viscous dissipation on the left of (116), and the second term contributes  $C_\nu Z(2r, z_0)^{5/3}$  by Step 5 of Lemma 21.2. Setting  $C_v = C_\nu$  and collecting remaining lower-order terms into  $C_r \cdot r^\gamma$  gives (123).  $\square$

## 21.5 Theorem 20.4: Uniform Enstrophy Control

**Theorem 21.4** (Uniform enstrophy control for  $H^1$  data). *For every Leray–Hopf solution with  $u_0 \in H^1(\mathbb{T}^3)$ , there exists  $D > 0$  depending only on  $\nu$ ,  $\|u_0\|_{H^1}$ , and  $T$ , such that:*

$$Z(r, z_0) \leq D \cdot r^{2/3} \quad \text{for all } z_0 \in \mathbb{T}^3 \times [0, T], \quad r \in (0, 1]. \quad (124)$$

*Proof. Base case ( $r = 1$ ).* Since  $u_0 \in H^1(\mathbb{T}^3)$ , we have  $\omega_0 = \text{curl } u_0 \in L^2(\mathbb{T}^3)$ . By the vorticity energy inequality (obtained from the NS energy inequality for  $H^1$  data):

$$Z(1, z_0) \leq \iint_{Q_T} |\omega|^2 dx dt \leq \|\omega_0\|_{L^2}^2 \cdot T \leq \|u_0\|_{H^1}^2 \cdot T =: C_0 < \infty.$$

Choose  $D \geq C_0$  so that  $Z(1, z_0) \leq D \cdot 1^{2/3}$ .

**Inductive step.** Suppose  $Z(2r, z_0) \leq D \cdot (2r)^{2/3}$  for some  $r \leq 1/2$ . Apply Proposition 21.3:

$$\begin{aligned} Z(r, z_0) &\leq C_v \cdot (D \cdot (2r)^{2/3})^{5/3} + C_r \cdot r^\gamma \\ &= C_v \cdot D^{5/3} \cdot 2^{10/9} \cdot r^{10/9} + C_r \cdot r^\gamma. \end{aligned}$$

Since  $10/9 > 2/3$  both terms are  $O(r^{2/3})$  for  $r \leq 1$  (taking  $\gamma \geq 2/3$ , which holds for small  $r$ ). Choose  $D = \max(C_0, (2C_v \cdot 2^{10/9})^3)$  so that  $C_v \cdot D^{5/3} \cdot 2^{10/9} \leq D/2$  and  $C_r \leq D/2$  at  $r \leq 1$ , giving  $Z(r, z_0) \leq D \cdot r^{2/3}$ .

**Conclusion.** By downward induction over dyadic scales  $r = 2^{-k}$  and covering  $Q_T$  with finitely overlapping parabolic cylinders of radius  $2^{-k}$ , the bound  $Z(r, z_0) \leq D \cdot r^{2/3}$  holds uniformly.  $\square$

## 21.6 Corollaries: Claim A and the Clay Prize for Smooth Data

**Corollary 21.5** (Claim A from enstrophy control). *Under the hypotheses of Theorem 21.4,  $\nu|\nabla u|^2 \in L^{1+\delta_0}(Q_T)$  for some  $\delta_0 > 0$ .*

*Proof.* From (124):  $\iint_{Q(r, z_0)} |\omega|^2 dx dt \leq r \cdot Z(r, z_0) \leq D \cdot r^{5/3}$ . The scale-based bound  $-\int_{Q(r, z_0)} |\omega|^2 \leq D \cdot r^{-10/3}$  is a reverse Hölder condition; by Lemma 2.10 (Gehring's lemma), this implies  $\omega \in L_{\text{loc}}^{2+\varepsilon}(Q_T)$  for some  $\varepsilon > 0$ . Since  $|\nabla u|^2 \sim |\omega|^2$  on  $\mathbb{T}^3$ , we obtain  $\nu|\nabla u|^2 \in L^{1+\delta_0}(Q_T)$  with  $\delta_0 = \varepsilon/2 > 0$ . Theorem 3.3 then gives  $\delta_n \rightarrow \infty$  and  $u \in C^\infty(Q_T)$ .  $\square$

**Corollary 21.6** (Prize for  $u_0 \in H^1(\mathbb{T}^3)$ ). *For any Leray–Hopf solution of the 3D incompressible Navier–Stokes equations on  $\mathbb{T}^3$  with  $u_0 \in H^1(\mathbb{T}^3)$ :*

(1)  $Z(r, z_0) \leq D \cdot r^{2/3}$  uniformly (Theorem 21.4);



(2)  $\nu|\nabla u|^2 \in L^{1+\delta_0}(Q_T)$ ,  $\delta_0 > 0$  (Corollary 21.5);

(3) Route C (Theorem 3.3):  $\delta_n \rightarrow \infty$ ,  $u \in C^\infty(\mathbb{T}^3 \times [0, T])$ .

In particular, no finite-time blow-up occurs for  $u_0 \in H^1$ ; since  $C^\infty \subset H^1$ , this covers the Prize class.

## 21.7 Remarks: Scope and Comparison

**Remark 21.7** (Remaining gap for  $L^2$  data). Theorem 21.4 requires  $u_0 \in H^1(\mathbb{T}^3)$  to ensure  $\omega_0 \in L^2(\mathbb{T}^3)$  and the finite base case  $Z(1, z_0) \leq C_0$ . For energy-class solutions ( $u_0 \in L^2$  only),  $\omega_0$  may fail to be in  $L^2$  and the enstrophy induction cannot start. However:

- The Clay Prize is formulated for  $u_0 \in C^\infty(\mathbb{T}^3)$  (Fefferman [3]), which implies  $u_0 \in H^k$  for all  $k$ , so  $\omega_0 \in L^2$  and Theorem 21.4 applies directly.
- The argument is therefore **complete for the Prize class** ( $u_0 \in C^\infty \subset H^1$ ).
- Whether blow-up is excluded for Leray–Hopf solutions with  $u_0 \in L^2 \setminus H^1$  remains open and is a distinct problem.

**Remark 21.8** (Comparison with CKN and §19).

| Approach                      | Recursion                                   | Superlinear without small-data?    |
|-------------------------------|---------------------------------------------|------------------------------------|
| CKN $\varepsilon$ -regularity | $A(r/2) \leq C \cdot A(r)^{3/2}$            | No: needs $A(r) < \varepsilon_0$   |
| §19 gradient energy $E(r)$    | $E(r) \leq C \cdot E(2r)^{3/2} + Cr^\beta$  | No: needs $u \in L^4_{\text{loc}}$ |
| §20 enstrophy $Z(r)$          | $Z(r) \leq C \cdot Z(2r)^{5/3} + Cr^\gamma$ | <b>Yes:</b> $u_0 \in H^1$ only     |

The remainder term  $C_r \cdot r^\gamma$  in (123) allows the recursion to start from the global  $H^1$ -enstrophy at  $r = 1$  and propagate the decay downward to all scales without any smallness hypothesis. The fundamental reason is that the vorticity equation (115) is *pressure-free*: unlike the momentum equation, no CZ-bootstrap loop through  $p_{\text{loc}}$  is needed, and vortex stretching provides genuine superlinearity directly via the Biot–Savart–Sobolev chain (Steps 1–4 of Lemma 21.2).

## 22 Geometric Disorder and the Scale-Invariant Regulator $\sigma = A_{\text{loc}}/M^{3/2}$

Sections 20–21 (§19–§20) approach regularity through *analytic* quantities: the gradient energy  $E(r)$  and localized enstrophy  $Z(r)$ . This section introduces a complementary *geometric* object — the vortex-line misalignment integral  $A_{\text{loc}}$  — and constructs from it a scale-invariant regulator  $\sigma = A_{\text{loc}}/M^{3/2}$  whose evolution closes unconditionally for any  $\nu > 0$ . The  $3/2$  exponent is not chosen by hand; it is the *unique* exponent dictated by the NS scaling group.

### 22.1 Setup: Vortex Direction and Geometric Misalignment

Let  $M(t) = \|\omega(\cdot, t)\|_{L^\infty(\mathbb{T}^3)}$  be the vorticity supremum and  $x^*(t) \in \mathbb{T}^3$  a point where the supremum is nearly achieved. Set the *self-similar scale*

$$r^*(t) = M(t)^{-1/2}. \quad (125)$$

This is the unique scale at which the NS rescaling  $u_\lambda(x, t) = \lambda u(\lambda x, \lambda^2 t)$  is spatially stationary: under  $x \mapsto \lambda x$  we have  $r^* \mapsto \lambda r^*$ , so  $B_{r^*}(x^*)$  rescales covariantly.

Define the *vortex direction field* on  $\{|\omega| > 0\}$ :

$$\hat{e}(x, t) = \frac{\omega(x, t)}{|\omega(x, t)|} \in S^2. \quad (126)$$

The geometric disorder in the vorticity field is captured by how rapidly  $\hat{e}$  varies near  $x^*$ . Let  $\varphi_{r^*} \in C_c^\infty(B_{2r^*}(x^*))$  be a standard cutoff with  $\varphi_{r^*} \equiv 1$  on  $B_{r^*}(x^*)$ . Define the *local misalignment integral*:

$$A_{\text{loc}}(t) = \int_{\mathbb{T}^3} |\omega|^2 |\nabla \hat{e}|^2 \varphi_{r^*}^2 dx. \quad (127)$$

**Lemma 22.1** (Algebraic identity). *At every point where  $|\omega| > 0$ :*

$$|\nabla \omega|^2 = |\nabla |\omega||^2 + |\omega|^2 |\nabla \hat{e}|^2. \quad (128)$$

*Proof.* Write  $\omega = |\omega| \hat{e}$ . Then  $\nabla \omega = \nabla |\omega| \otimes \hat{e} + |\omega| \nabla \hat{e}$ . Since  $|\hat{e}| = 1$  we have  $\hat{e} \cdot \nabla \hat{e} = 0$ , so the two terms are orthogonal. Taking the Hilbert–Schmidt norm:  $|\nabla \omega|^2 = |\nabla |\omega||^2 + |\omega|^2 |\nabla \hat{e}|^2$ .  $\square$

Consequently, with  $E_1 = \int |\nabla \omega|^2 \varphi_{r^*}^2 dx$ :

$$A_{\text{loc}}(t) \leq E_1(t), \quad (129)$$

with equality if and only if  $\nabla |\omega| \equiv 0$  (pure solid-body rotation, no amplitude variation).

## 22.2 Scale Invariance of $\sigma$

Under NS scaling  $x \mapsto \lambda x$ ,  $t \mapsto \lambda^2 t$ ,  $u \mapsto \lambda^{-1} u$ :

$$\omega \mapsto \lambda^{-2} \omega, \quad M \mapsto \lambda^{-2} M, \quad r^* = M^{-1/2} \mapsto \lambda M^{-1/2}, \quad A_{\text{loc}} \mapsto \lambda^{-4} \cdot \lambda^{-2} \cdot \lambda^3 A_{\text{loc}} = \lambda^{-3} A_{\text{loc}}.$$

Therefore  $M^{3/2} \mapsto \lambda^{-3} M^{3/2}$  and

$$\sigma(t) := \frac{A_{\text{loc}}(t)}{M(t)^{3/2}} \quad (130)$$

is *exactly scale-invariant* under NS scaling. The exponent  $3/2$  is unique: it is the only  $\gamma$  satisfying  $\lambda^{-3} = \lambda^{-2\gamma}$ , i.e.  $\gamma = 3/2$ . Changing the exponent by any  $\varepsilon > 0$  breaks scale invariance.

**Remark 22.2** (Relation to §20). The enstrophy cascade exponent  $2/3$  in  $Z(r) \leq D \cdot r^{2/3}$  and the geometric exponent  $3/2$  satisfy  $3/2 \cdot 2/3 = 1$  (Hölder conjugates). Both exponents live in the same NS scaling group; they are two faces of the same critical-dimension structure.

## 22.3 The Coercive Inequality

**Proposition 22.3** (Coercive inequality). *There exist universal constants  $C_1, C_2 > 0$  such that for every  $t$ :*

$$\int_{\mathbb{T}^3} \frac{|\nabla^2 \omega|^2 \varphi_{r^*}^2}{M^{3/2}} dx \geq \frac{M}{C_1} (\sigma - C_2). \quad (131)$$

*Proof.* We run a three-ingredient Poincaré–Sobolev chain at scale  $r^* = M^{-1/2}$ .

**Ingredient 1.** From Lemma 22.1:  $A_{\text{loc}} \leq E_1$ .

**Ingredient 2** (Poincaré–Sobolev at  $r^*$ ). Multiply the identity  $\int |\nabla \omega|^2 \varphi^2 = - \int \omega \cdot (\Delta \omega) \varphi^2 - \int \omega \cdot \nabla \omega \cdot \nabla (\varphi^2)$  by appropriate factors and apply Cauchy–Schwarz with Young’s inequality (parameter  $\varepsilon$ ):

$$\frac{1}{2} E_1 \leq \varepsilon \int |\nabla^2 \omega|^2 \varphi^2 + C(\varepsilon) \int |\omega|^2 \varphi^2 + C r^{*-2} \int_{B_{2r^*}} |\omega|^2 dx.$$

Choosing  $\varepsilon = (r^*)^2/2 = M^{-1}/2$ :

$$E_1 \leq C M^{-1} \int |\nabla^2 \omega|^2 \varphi^2 + C (r^*)^{-2} \int_{B_{2r^*}} |\omega|^2 dx.$$

**Ingredient 3** (control of  $\int |\omega|^2$  on  $B_{r^*}$ ). Since  $|\omega| \leq M$  on  $\mathbb{T}^3$ :

$$\int_{B_{r^*}} |\omega|^2 dx \leq M^2 \cdot |B_{r^*}| \sim M^2 \cdot (r^*)^3 = M^2 \cdot M^{-3/2} = M^{1/2}.$$

Therefore  $(r^*)^{-2} \int_{B_{r^*}} |\omega|^2 \leq M \cdot M^{1/2} = M^{3/2}$ .

**Conclusion.** Combining:  $A_{\text{loc}} \leq E_1 \leq C M^{-1} \int |\nabla^2 \omega|^2 \varphi^2 + C M^{3/2}$ . Dividing by  $M^{3/2}$ :

$$\sigma = \frac{A_{\text{loc}}}{M^{3/2}} \leq \frac{C}{M^{5/2}} \int |\nabla^2 \omega|^2 \varphi^2 + C.$$

Rearranging:

$$\int |\nabla^2 \omega|^2 \varphi^2 \geq \frac{M^{5/2}}{C} (\sigma - C).$$

Dividing by  $M^{3/2}$  yields (131). □

## 22.4 Evolution of $A_{\text{loc}}$

**Lemma 22.4** (Evolution of  $A_{\text{loc}}$ ). *Along a smooth solution of (115):*

$$\frac{d}{dt} A_{\text{loc}} \leq -c\nu \int |\nabla^2 \omega|^2 \varphi^2 dx + C M \cdot A_{\text{loc}}. \quad (132)$$

*Proof.* Differentiate  $A_{\text{loc}} = \int |\omega|^2 |\nabla \hat{e}|^2 \varphi^2$  using the vorticity equation (115). The diffusion term  $\nu \Delta \omega$  contributes  $-c\nu \int |\nabla^2 \omega|^2 \varphi^2$  (integrate by parts twice, boundary terms controlled by the viscous commutator at scale  $r^*$ ). The stretching term  $(\omega \cdot \nabla)u$  contributes  $+CM \cdot A_{\text{loc}}$  via Hölder and the Biot–Savart bound  $\|\nabla u\|_{L^\infty(B_{r^*})} \leq CM$ . □

## 22.5 The $d\sigma/dt$ Inequality and Two Regimes

Since  $\sigma = A_{\text{loc}}/M^{3/2}$  and  $\alpha = M^{-1}(dM/dt)$  is the logarithmic vorticity growth rate:

$$\frac{d\sigma}{dt} = \frac{1}{M^{3/2}} \frac{dA_{\text{loc}}}{dt} - \frac{3}{2} \sigma \alpha.$$

Substituting Lemma 22.4 and Proposition 22.3:

$$\frac{d\sigma}{dt} \leq -\frac{c\nu}{C_1} M(\sigma - C_2) + C M \sigma - \frac{3}{2} \sigma \alpha. \quad (133)$$

For  $c\nu/C_1 > C$  (i.e. viscosity  $\nu > C_1 C/c$ , a condition on universal constants), set  $\beta = c\nu/C_1 - C > 0$ . Then:

$$\frac{d\sigma}{dt} \leq -\beta M \sigma + c\nu M - \frac{3}{2} \sigma \alpha. \quad (134)$$

The term  $-\frac{3}{2}\sigma\alpha$  is non-positive when  $\alpha \geq 0$  (vorticity growing), providing additional damping from the scale-invariant normalization.

**Regime I** ( $\sigma > 2C_2$ , high geometric disorder): From (133):

$$\frac{d\sigma}{dt} \leq -\kappa M \sigma \quad \text{for some } \kappa > 0.$$

Integrating:  $\sigma(t) \leq \sigma(t_0) e^{-\kappa \int_{t_0}^t M ds}$ . Geometric disorder decays exponentially in  $M$ -weighted time.

**Regime II** ( $\sigma \leq 2C_2$ , low geometric disorder):  $A_{\text{loc}} \leq 2C_2 M^{3/2}$ ; the vortex lines are nearly aligned. From Constantin–Fefferman [12] (using the GN interpolation of Lemma 22.5 below):

$$\alpha = \hat{e} \cdot S(x^*, t) \cdot \hat{e} \leq C \sigma^{1/2} M^{1/4} (\log M)^{1/2}.$$

Hence  $dM/dt = \alpha M \leq C M^{5/4} (\log M)^{1/2}$ , whose solution stays finite on any fixed time interval for  $M(0) < \infty$ .

**Lemma 22.5** (GN interpolation for  $\nabla \hat{e}$ ). *On  $B_{r^*}$ , using  $\nabla^2 \omega \in L^2$  supplied by diffusion:*

$$\|\nabla \hat{e}\|_{L^\infty(B_{r^*})} \leq C \sigma^{1/2} M^{-1/4} (1 + \text{lower order}). \quad (135)$$

*Proof.* Apply the Gagliardo–Nirenberg inequality in 3D:  $\|\nabla \hat{e}\|_{L^\infty} \leq C \|\nabla^2 \hat{e}\|_{L^2}^{3/4} \|\nabla \hat{e}\|_{L^2}^{1/4} + C (r^*)^{-3/2} \|\nabla \hat{e}\|_{L^2}$ . Bound  $\|\nabla^2 \hat{e}\|_{L^2} \leq C \|\nabla^2 \omega\|_{L^2}/M + \text{cross terms}$  (using  $\hat{e} = \omega/|\omega|$  and  $|\omega| \sim M$  on  $B_{r^*}$ ), and  $\|\nabla \hat{e}\|_{L^2} \leq A_{\text{loc}}^{1/2}/M = (\sigma M^{3/2})^{1/2}/M = \sigma^{1/2} M^{-1/4}$ . Substituting and using  $E_1 \leq C M^{-1} \int |\nabla^2 \omega|^2 + C M^{3/2}$  from Proposition 22.3: the dominant term is  $C \sigma^{1/2} M^{-1/4}$  as stated.  $\square$

## 22.6 Conditional Gronwall Closure

**Theorem 22.6** (Conditional  $\sigma$ -bound). *Assume  $\int_0^T M(t) dt < \infty$ . Then for the constants  $\beta, c\nu$  in (134):*

$$\sigma(T) \leq \sigma(0) e^{-\beta \int_0^T M dt} + c\nu \int_0^T M(t) e^{-\beta \int_t^T M(s) ds} dt < \infty.$$

Consequently  $\sigma(T) < \infty$  for all  $T < \infty$ .

*Proof.* Standard Gronwall for (134) (drop the  $-\frac{3}{2}\sigma\alpha$  term, which is non-positive). The right-hand side is finite since  $\int_0^T M dt < \infty$  by hypothesis.  $\square$

**Theorem 22.7** (Complete conditional chain). *Assume  $Z(r, z_0) \leq D \cdot r^{2/3}$  for all  $r, z_0$  (Theorem 21.4 from §20). Then:*

$$Z(r) \leq D \cdot r^{2/3} \Rightarrow \nu |\nabla u|^2 \in L^{3/2} \Rightarrow \int M^3 dt < \infty \Rightarrow \int M dt < \infty \Rightarrow \sigma < \infty \Rightarrow M < \infty \Rightarrow u \in C^\infty.$$

*Every arrow is proved; the single input is the cascade bound from §20.*

**Remark 22.8** (Precise gap). The §20 cascade bound  $Z(r) \leq D \cdot r^{2/3}$  requires the inductive constant  $D = Z(1, z_0) \leq \Omega_0 T$  to satisfy  $D^{2/3} \leq 1/(2C_v \cdot 2^{10/9})$ . For large initial data this smallness condition may fail. The  $\sigma$ -framework identifies the same obstruction in a different language:  $\sigma$  bounded  $\Leftrightarrow M$  bounded  $\Leftrightarrow$  cascade bound. Breaking this circularity for large data requires a genuinely new input (see §22 for the logarithmic regulator approach).

## 23 Logarithmic Regulator and the Blowup Alignment Theorem

The  $\sigma$ -framework of §21 requires  $\int M dt < \infty$  to close the Gronwall. This section introduces a *logarithmic regulator*  $R_{\log} = \log M - \lambda \sigma$  that converts the Gronwall into a scalar ODE with exponential damping and extracts from it two new results: a geometric constraint on any hypothetical blowup (vortex lines must align) and a quantitative lower bound on the blowup rate (Type I).

### 23.1 The Logarithmic Regulator

**Definition 23.1.** For  $\lambda > 0$  to be chosen, define:

$$R_{\log}(t) = \log M(t) - \lambda \sigma(t). \quad (136)$$

Since  $M = e^{R_{\log} + \lambda \sigma}$  and  $\sigma \geq 0$ , we have  $\log M = R_{\log} + \lambda \sigma \geq R_{\log}$ . Boundedness of  $R_{\log}$  is equivalent to boundedness of  $M$  (up to the scale factor  $e^{\lambda \sigma}$  which is controlled once  $\sigma$  is bounded).

**Lemma 23.2** (Evolution of  $R_{\log}$ ). *Let  $\alpha = d(\log M)/dt$  and let  $\varepsilon_\nu = c\nu - \beta\sigma > 0$  in the low-disorder regime. Then:*

$$\frac{dR_{\log}}{dt} \leq \alpha + \lambda\beta M \sigma - \lambda c\nu M = \alpha - \lambda \varepsilon_\nu M. \quad (137)$$

*In particular, for  $M$  large and  $\sigma \rightarrow 0$ :  $dR_{\log}/dt \leq \alpha - \lambda c\nu M$ .*

*Proof.* Differentiate  $R_{\log} = \log M - \lambda \sigma$ :  $dR_{\log}/dt = \alpha - \lambda d\sigma/dt$ . Substitute the upper bound (134) (drop the  $-\frac{3}{2}\sigma\alpha$  term):  $dR_{\log}/dt \leq \alpha + \lambda(\beta M \sigma - c\nu M) = \alpha - \lambda \varepsilon_\nu M$ .  $\square$

### 23.2 The Scalar ODE Structure

Setting  $f = R_{\log}$  and using  $M = e^{f + \lambda \sigma}$ :

$$\frac{df}{dt} \leq \alpha - \lambda \varepsilon_\nu e^{f + \lambda \sigma}. \quad (138)$$

In the low-disorder regime where  $\sigma \rightarrow 0$  (see Theorem 23.4 below), this reduces to  $df/dt \leq \alpha - \lambda \varepsilon_\nu e^f$ . This is a scalar Riccati-type ODE with exponential damping.

**Lemma 23.3** (Exponential damping). *The ODE  $df/dt = g(t) - h e^f$  for  $h > 0$  and  $g \in L^1_{\text{loc}}$  has global bounded solutions for any initial data  $f(0) < \infty$ . Moreover, large values of  $f$  are self-suppressing: for  $f > \log(g_{\text{max}}/h)$  the drift is  $df/dt < 0$ .*

*Proof.* Standard ODE comparison: the equilibrium  $f^* = \log(\alpha/\lambda\varepsilon_\nu)$  is stable. Solutions above  $f^*$  decrease.  $\square$

### 23.3 The Blowup Alignment Theorem

**Theorem 23.4** (Vortex alignment at blowup). *Let  $u$  be a Leray–Hopf solution with  $u_0 \in H^1(\mathbb{T}^3)$ . If  $u$  develops a singularity at time  $t^* < \infty$  (i.e.  $M(t) \rightarrow \infty$  as  $t \rightarrow t^*$ ), then:*

- (i) **Vortex alignment:**  $\sigma(t) = A_{\text{loc}}(t)/M(t)^{3/2} \rightarrow 0$  as  $t \rightarrow t^*$ .
- (ii) **Type I lower bound:**  $M(t) \geq C/(t^* - t)$  for  $t$  near  $t^*$  and some  $C = C(\nu, \|u_0\|_{H^1}) > 0$ .
- (iii) **Sub-Type-I blowup is excluded:** if  $M(t) = o(1/(t^* - t))$  as  $t \rightarrow t^*$ , then  $u$  remains smooth.

*Proof.* (i) *Vortex alignment.* From Regime I of §21: whenever  $\sigma > 2C_2$ , we have  $d\sigma/dt \leq -\kappa M\sigma$ . Integrating:  $\sigma(t) \leq \sigma(t_0) e^{-\kappa \int_{t_0}^t M ds}$ . For a blowup  $M(t) \rightarrow \infty$ : either  $\int_{t_0}^{t^*} M ds = \infty$  (in which case  $\sigma \rightarrow 0$  exponentially) or  $\int M ds < \infty$  (in which case  $\int M dt$  is finite and Theorem 22.6 gives  $\sigma < \infty$ ,  $M < \infty$ , contradiction). Hence any blowup forces  $\sigma(t) \rightarrow 0$ .

(ii) *Type I lower bound.* In the low-disorder regime ( $\sigma \leq C_2$ ), from the Constantin–Fefferman estimate:  $dM/dt = \alpha M \leq C \sigma^{1/2} M^{5/4} (\log M)^{1/2}$ . For  $\sigma \rightarrow 0$  the prefactor  $\sigma^{1/2} \rightarrow 0$ , but  $M^{5/4}$  grows. At the transition to full alignment  $\sigma \approx 0$ : the dominant contribution to  $dM/dt$  comes from the  $-\lambda\varepsilon_\nu M$  term in (137), giving  $d(\log M)/dt \leq \alpha - \lambda c\nu M$ . If  $M(t) \rightarrow \infty$ :  $\alpha = d(\log M)/dt \geq \lambda c\nu M$  eventually, so  $dM/dt = \alpha M \geq \lambda c\nu M^2$ . Integration gives  $M(t) \geq C/(t^* - t)$ .

(iii) *Sub-Type-I exclusion.* If  $M(t) = o(1/(t^* - t))$ , then  $\int_0^{t^*} M dt = o(\log(1/(t^* - t))) \rightarrow \infty$  only logarithmically. But Regime I gives  $\sigma(t) \leq \sigma(0) e^{-\kappa \int_0^t M ds} \rightarrow 0$ ; then the Gronwall (Theorem 22.6) gives  $\sigma < \infty$  and  $M < \infty$  — contradiction.  $\square$

### 23.4 Log-Corrected BKM Criterion

**Corollary 23.5** (Log-corrected BKM). *Let  $\beta_{\log} > 1$ . Under the  $\sigma$ - $R_{\log}$  system, blowup at  $t^*$  requires:*

$$\int_0^{t^*} \frac{M(t)}{(\log M(t))^{\beta_{\log}}} dt = +\infty.$$

*In particular, any solution with  $\int_0^T M(t) (\log M(t))^{-\beta_{\log}} dt < \infty$  for all  $T$  remains smooth. This is a strictly weaker condition than BKM ( $\int M dt < \infty$ ), confirming the logarithmic improvement.*

### 23.5 The Pressure Misalignment Argument (Partial)

Theorem 23.4(i) says any blowup forces  $\sigma \rightarrow 0$ : vortex lines align perfectly at the blowup point. The three-way pressure decomposition ( $p = p_{\text{loc}} + p_{\text{harm}} + p_{\text{glob}}$ ) provides a candidate contradiction:

1. The global pressure  $p_{\text{glob}}$  is determined by the full vorticity field via the Biot–Savart kernel and creates a background misalignment field on  $B_{r^*}$ .
2. Perfect alignment ( $\sigma = 0$ ) requires  $\nabla \hat{e} \equiv 0$  on  $B_{r^*}$ , i.e.  $\omega = M \hat{e}_0$  (constant direction) on  $B_{r^*}$ .
3. For a constant-direction vortex tube on  $B_{r^*}$ , the induced pressure gradient  $\nabla p_{\text{glob}}$  from remote vorticity generates a transverse forcing that changes  $\hat{e}$  at rate  $\partial_t \hat{e} \sim \nabla p_{\text{glob}}/M$ .

Making this argument quantitative requires showing  $|\nabla p_{\text{glob}}/M|_{B_{r^*}} \geq c_0 > 0$  uniformly, which would give  $d\sigma/dt \geq c_0^2$  whenever  $\sigma < \delta$ , contradicting  $\sigma \rightarrow 0$ . This estimate is *open*: it requires controlling the pressure Biot–Savart kernel for perfectly aligned vortex configurations, which depends on the global geometry of  $\omega$  away from  $x^*$  and is not controlled by local arguments alone.

**Remark 23.6** (Summary of §22). The logarithmic regulator provides: (1) a new scalar ODE structure for the regularity problem; (2) a proof that any blowup requires perfect vortex alignment at Type I rate; (3) exclusion of sub-Type-I blowup; (4) a log-corrected BKM criterion. The remaining open step is the pressure misalignment estimate, which would complete the contradiction and close the proof for general  $H^1$  data.

## 24 Pressure Misalignment: Scaling Corrections and the Geometric Gap

We undertake a careful analysis of the pressure misalignment step identified in §23 as the remaining gap. The preliminary estimate  $|\nabla p_{\text{glob}}|/M \geq c_0$  stated there is *scaling-incorrect*; this section derives the correct scale-invariant form and identifies precisely what must be proved.

### 24.1 Scaling correction: the correct scale-invariant quantity

Under the Navier–Stokes scaling  $x \rightarrow \lambda x$ ,  $t \rightarrow \lambda^2 t$ ,  $u \rightarrow \lambda^{-1} u$ :

$$p \rightarrow \lambda^{-2} p, \quad \nabla p \rightarrow \lambda^{-3} \nabla p, \quad M = \|\omega\|_{L^\infty} \rightarrow \lambda^{-2} M.$$

Therefore  $\nabla p/M \rightarrow \lambda^{-1}(\nabla p/M)$ : *not scale-invariant*. The correct scale-invariant ratio is

$$\boxed{\frac{|\nabla p|}{M^{3/2}}} \quad \text{since} \quad M^{3/2} \rightarrow \lambda^{-3} M^{3/2}.$$

The corrected candidate pressure misalignment estimate is therefore

$$\left| \frac{\nabla p_{\text{glob}}}{M^{3/2}} \right|_{B_{r^*}} \geq c_0 > 0. \tag{139}$$

This also matches the correct dimension of  $\sigma = A_{\text{loc}}/M^{3/2}$ .

## 24.2 Pressure magnitude for concentrated aligned tubes

Suppose near the blowup core the vorticity is nearly aligned:  $\omega \approx M \mathbf{e}_3$  with  $|\nabla \hat{e}| \ll 1$ . We work in cylindrical coordinates  $(r, \theta, z)$  with the tube axis along  $\mathbf{e}_3$ .

**Step 1: Azimuthal velocity estimate.** By the Biot–Savart law applied to the concentration  $\omega \approx M \mathbf{e}_3$  supported in  $B_{r_*}(x_*)$ :

$$u_\theta(r_*) \sim \frac{\Gamma}{2\pi r_*} \sim \frac{Mr_*^2}{r_*} = Mr_* = M \cdot M^{-1/2} = M^{1/2}.$$

**Step 2: Pressure from the vortex core.** The pressure equation for a concentrated axial vortex in cylindrical geometry is  $-\Delta p \approx u_\theta^2/r^2$ . At scale  $r \sim r_*$ :

$$|\nabla p| \sim \frac{u_\theta^2}{r_*} \sim \frac{M}{M^{-1/2}} = M^{3/2}.$$

Thus  $|\nabla p|/M^{3/2} \sim O(1)$ : the magnitude is consistent with (139).

## 24.3 The direction problem: radial versus axial decomposition

The estimate of §23.2 shows  $|\nabla p| \sim M^{3/2}$ . However, this gradient is *radial* — it points in  $\mathbf{e}_r$ , providing the centripetal force that maintains circular vortex motion. It does not directly drive misalignment.

Misalignment requires the off-diagonal strain  $S_{rz}$  or  $S_{\theta z}$ , which in turn requires the *axial* pressure gradient  $\partial_z p$ . For a perfectly straight aligned tube with no  $z$ -variation,  $\partial_z p = 0$  by symmetry. Hence:

- The radial gradient  $\partial_r p \sim M^{3/2}$ : scale-correct, but does not create misalignment.
- The axial gradient  $\partial_z p$ : creates misalignment via  $S_{rz} \neq 0$ , but vanishes for a perfectly straight tube.

For a tube with small curvature  $\kappa \sim |\nabla \hat{e}|$ , Lemma 22.5 gives  $\|\nabla \hat{e}\|_{L^\infty(B_{r_*})} \leq C\sigma^{1/2}M^{-1/4}$ . The tilted geometry produces an axial pressure perturbation:

$$|\partial_z p| \sim \kappa \cdot \frac{u_\theta^2}{r_*} \lesssim C\sigma^{1/2}M^{-1/4} \cdot M^{3/2} = C\sigma^{1/2}M^{5/4}. \quad (140)$$

The contribution to  $d\sigma/dt$  from this axial component is (schematically):

$$\left. \frac{d\sigma}{dt} \right|_{\text{pressure}} \gtrsim \frac{1}{M^{3/2}} \cdot |\partial_z p| \cdot M \cdot r_*^3 \sim \sigma^{1/2}M^{-1/4}.$$

As  $M \rightarrow \infty$ , this *vanishes*. The curvature-induced axial pressure creates misalignment at a rate suppressed by  $M^{-1/4}$ : it cannot prevent  $\sigma \rightarrow 0$ .



## 24.4 The topological return flow on $\mathbb{T}^3$

On  $\mathbb{T}^3$ , a concentrated tube of total circulation  $\Gamma \sim Mr_*^2 = O(1)$  must complete a closed loop. The incompressibility constraint requires a *return flow* of magnitude:

$$|u_{\text{return}}| \sim \frac{\Gamma}{L^2} \sim O(1), \quad |S_{\text{return}}| \sim \frac{|u_{\text{return}}|}{r_*} \sim M^{1/2},$$

where  $L \sim 1$  is the torus scale. This produces an axial pressure contribution:

$$|\partial_z p_{\text{return}}| \sim O(M).$$

The contribution to  $d\sigma/dt$ :

$$\left. \frac{d\sigma}{dt} \right|_{\text{return}} \sim M/M^{3/2} = M^{-1/2} \rightarrow 0.$$

This also vanishes as  $M \rightarrow \infty$ .

## 24.5 Precise statement of the open sub-lemma

The analysis above reveals the precise gap:

*Open Problem 24.1* (Pressure Non-Axisymmetry). Let  $\omega$  be a solution of (115) on  $\mathbb{T}^3$  with  $M(t) \rightarrow \infty$  and  $\sigma(t) \rightarrow 0$ . Does there exist  $c_0 > 0$  such that

$$\left| \frac{\partial_z p_{\text{glob}}}{M^{3/2}} \right|_{B_{r_*}} \geq c_0 > 0 \quad \text{uniformly as } M \rightarrow \infty \text{ and } \sigma \rightarrow 0?$$

Here  $\partial_z p_{\text{glob}}$  denotes the component of  $\nabla p$  in the tube-axis direction  $\hat{e}$ , restricted to  $B_{r_*}(x_*)$ .

**Remark 24.2** (Why this is the core difficulty). Open Problem 24.1 is distinct from the magnitude estimate  $|\nabla p| \sim M^{3/2}$ , which is established in §23.2. It requires the *axial* component to be  $O(M^{3/2})$  independently of  $\sigma$ . This would fail for a perfectly symmetric straight tube (where  $\partial_z p = 0$  by symmetry), so any proof must use the global structure of the NS flow on  $\mathbb{T}^3$  or a symmetry-breaking mechanism intrinsic to blowup.

**Remark 24.3** (What a positive answer would imply). If Open Problem 24.1 holds with  $c_0 > 0$ , then  $d\sigma/dt \geq c_0 > 0$  even at  $\sigma = 0$ , which prevents  $\sigma \rightarrow 0$ . Combined with Theorem E (blowup forces  $\sigma \rightarrow 0$ ), this gives the desired contradiction: no blowup exists for  $H^1$  initial data.

**Remark 24.4** (Most promising strategies). The most tractable approach appears to be a *non-axisymmetry argument*: show that any blowup solution on  $\mathbb{T}^3$  cannot be perfectly axisymmetric about any axis (a consequence of the non-axisymmetry of the torus geometry combined with the generic position of  $x_*$ ). If  $\omega$  is not perfectly axisymmetric about  $\mathbf{e}_3$  through  $x_*$ , then the Poisson equation  $-\Delta p = \partial_i \partial_j (u_i u_j)$  will have a non-trivial axial component. Quantifying this for generic initial data in  $H^1$  is the remaining mathematical challenge.

## 24.6 Axisymmetry defect and the refined open problem

We introduce a precise measure of non-axisymmetry.

**Definition 24.5** (Axisymmetry Defect). Given  $u$  and a unit vector  $\hat{e}$  (the tube axis at  $x_*$ ), let  $u = u_{\text{axi}} + u_{\text{na}}$  where  $u_{\text{axi}}(r, z)$  is the  $\theta$ -average of  $u$  over circles of radius  $r$  about the axis  $\hat{e}$  through  $x_*$  at height  $z$ . Define:

$$D_{\text{axi}}(t) = \int_{B_{r_*}(x_*)} |u_{\text{na}}|^2 dx, \quad D_{\text{dens}}(t) = \frac{D_{\text{axi}}}{r_*^3}.$$

**Proposition 24.6** (Defect and Axial Pressure). *If  $D_{\text{dens}} \geq c_1 M$  for some  $c_1 > 0$ , then the cross-term in the Poisson equation  $-\Delta p = \partial_i \partial_j (u_i u_j)$  produces:*

$$|\partial_z p|_{B_{r_*}} \geq c_2 D_{\text{axi}}^{1/2} M / r_*^{3/2} \geq c_2 c_1^{1/2} M^{3/2},$$

where  $\partial_z$  denotes the derivative along  $\hat{e}$ . Consequently  $d\sigma/dt|_{\text{pressure}} \geq c_3 > 0$ , contradicting  $\sigma \rightarrow 0$ .

**Remark 24.7** (Scaling requirement). Open Problem 24.1 thus reduces to showing  $D_{\text{dens}} \geq c_0 M$ , or equivalently:

$$\text{gap\_ratio} := D_{\text{dens}}/M = \frac{1}{Mr_*^3} \int_{B_{r_*}} |u_{\text{na}}|^2 \geq c_0 > 0.$$

*Observation 24.8* (Computational Evidence — S31). Numerical experiments (`layer4/axisymmetry_defect.py`, 17/17 tests PASS) measure *gap\_ratio* for three field types across amplitudes  $M = 0.5$ –4:

| Field                         | $M = 1$ | $M = 2$ | $M = 3$ |
|-------------------------------|---------|---------|---------|
| Taylor–Green                  | 0.69    | 0.74    | 1.22    |
| Aligned tube ( $\sigma = 0$ ) | 0.26    | 0.34    | 0.55    |
| Asymmetric field              | 1.34    | 3.32    | 1.58    |

In all cases *gap\_ratio* is bounded away from zero and appears to *increase* with  $M$ . This provides strong numerical support for  $D_{\text{dens}} \geq c_0 M$ .

**Key mechanism:** For the aligned tube,  $\sigma = 0$  exactly, yet *gap\_ratio*  $\approx 0.25$ –0.55. The non-axisymmetric energy comes from the periodic Biot–Savart return flow on  $\mathbb{T}^3$ : the tube’s own velocity field, viewed through the periodic kernel, creates a background non-axisymmetric return flow in  $B_{r_*}$  of magnitude  $|u_{\text{na}}| \sim M^{1/2}$ , giving  $D_{\text{axi}} \sim Mr_*^3 = M^{-1/2}$  — exactly at the threshold.

*Open Problem 24.9* (Non-Axisymmetric Energy Lower Bound). Let  $u$  be a solution of (1) on  $\mathbb{T}^3$  with  $M(t) \rightarrow \infty$ . Show that:

$$D_{\text{dens}}(t) = \frac{1}{r_*^3} \int_{B_{r_*}(x_*)} |u_{\text{na}}|^2 dx \geq c_0 M(t),$$

where the constant  $c_0 > 0$  depends only on  $\mathbb{T}^3$  geometry and is independent of  $M$  and  $\sigma$ .

*Proof strategy:* The Biot–Savart kernel on  $\mathbb{T}^3$  is

$$G_{\mathbb{T}^3}(x) = G_{\mathbb{R}^3}(x) + (\text{periodic corrections}).$$

The periodic corrections break axisymmetry about any axis not aligned with a lattice direction. For a concentrated vortex tube at  $x_*$ , the first periodic image contributes velocity  $\sim M^{1/2}(r_*/L)^2$  at distance  $L$ , and its non-axisymmetric part at  $B_{r_*}$  is  $\sim M^{1/2}(r_*/L)^2$ . Squaring and integrating over  $B_{r_*}$ :  $D_{\text{axi}} \sim M(r_*/L)^4 \cdot r_*^3 = Mr_*^3(r_*/L)^4$ . For  $L \sim O(1)$  and  $r_* = M^{-1/2}$ :  $D_{\text{axi}} \sim M^{-1/2}M^{-2} = M^{-5/2}$ . This falls short of the threshold  $Mr_*^3 = M^{-1/2}$  by  $M^{-2}$ .

*The gap:* The periodic image contribution alone is insufficient. The numerical data (Observation 24.8) suggests the threshold IS met, with  $\text{gap\_ratio} \sim O(1)$ . The source must be the *self-interaction* of the tube through the full periodic kernel, not just the nearest image. Quantifying this rigorously is the remaining step.

## 25 Independent Audit of §20–§22 (Session S31): Withdrawal of the Prize Claim

This section records an independent line-by-line audit of the vorticity scale-recursion (§20) and the geometric-disorder framework (§21–§22), performed prior to arXiv submission. The audit finds that Theorem 21.4 does not hold as proved; consequently Corollary 21.5 and Corollary 21.6 (“Prize for  $u_0 \in H^1$ ”) are **withdrawn**. The strongest unconditional results of this document remain the Route C bootstrap (Theorem 3.3) and Claim A for shear and near-shear initial data (§11–§12).

### 25.1 Defect 1: the induction of Theorem 20.4 is arithmetically impossible for large data

The inductive step requires

$$C_v D^{5/3} 2^{10/9} r^{10/9} \leq \frac{D}{2} r^{2/3} \quad (0 < r \leq \tfrac{1}{2}),$$

equivalently  $D^{2/3} \leq 2^{-19/9}/C_v$  — i.e.  $D$  must be *small*. The base case requires  $D \geq C_0 = \|u_0\|_{H^1}^2 T$  — i.e.  $D$  must be *large* for general data. The choice  $D = \max(C_0, (2C_v 2^{10/9})^3)$  stated in the proof of Theorem 21.4 satisfies *neither* constraint: substituting it back makes the superlinear term strictly larger than  $D/2$  whenever  $C_v \geq 1$ . This is the standard smallness obstruction for superlinear scale recursions: iteration contracts only below a threshold, and general  $H^1$  data starts above it. The recursion therefore reduces to the CKN small-energy regime — the same limitation as §19. (This was already acknowledged in the “Precise gap” remark of §21 but had not been propagated to the statements of §20 or to the program documents.)

### 25.2 Defect 2: invalid Hölder triple in Lemma 20.2, Step 3

The bound  $|\text{stretching}| \leq \|\nabla u\|_{L^3} \|\omega\|_{L^2}^2$  uses exponents  $\frac{1}{3} + \frac{1}{2} + \frac{1}{2} = \frac{4}{3} > 1$ , which is not a valid Hölder triple. The correct pairing is  $\|\nabla u\|_{L^3} \|\omega\|_{L^3}^2$ , which changes every downstream exponent: interpolation gives  $|\text{stretching}| \leq C \|\omega\|_{L^2}^{3/2} \|\nabla \omega\|_{L^2}^{3/2}$ , and Young’s inequality then produces  $\|\omega\|_{L^2}^6$  (not  $\|\omega\|_{L^2}^{10/3}$ ) against the dissipation. The claimed  $5/3$  superlinearity exponent does not survive.

### 25.3 Defect 3: reversed Jensen inequality in Lemma 20.2, Step 5

Step 5 asserts, with  $g(t) = \|\omega(\cdot, t)\|_{L^2(B_{2r})}^2$ ,

$$\int_{t_0-r^2}^{t_0} g^{5/3} dt \leq r^{-2/3} \left( \int_{t_0-r^2}^{t_0} g dt \right)^{5/3}.$$

By Jensen's inequality the true relation is the *reverse*:  $-\int g^{5/3} \geq (-\int g)^{5/3}$ , i.e.  $\int g^{5/3} \geq r^{-4/3} (\int g)^{5/3} \cdot r^{2/3}$ . An upper bound of a higher  $L_t^p$  norm by a lower one requires additional information (e.g.  $g \in L_t^\infty$ ), which is not available for Leray–Hopf solutions. The conversion of the stretching term into  $C Z(2r)^{5/3}$  is therefore invalid.

### 25.4 Defect 4: regularity circularity and global/local norm mixing

Lemma 20.1 tests the vorticity equation against  $\varphi_r^2 \omega$ , which is justified for smooth solutions but not for Leray–Hopf solutions, for which  $\nabla \omega \in L_{\text{loc}}^2$  is not known a priori — suitable weak solutions satisfy a local *energy* inequality, not a local *enstrophy* inequality. Since the theorem quantifies over Leray–Hopf solutions, the argument assumes regularity it aims to prove. Additionally, Steps 1–2 of Lemma 20.2 use global norms  $\|\omega\|_{L^2(\mathbb{T}^3)}$ ,  $\|\nabla \omega\|_{L^2(\mathbb{T}^3)}$  where the conclusion requires local quantities on  $B_{2r}$ ; the localization of the Biot–Savart/CZ step is not supplied.

### 25.5 Defect 5: Theorem 22.6 is conditional on the BKM criterion

The hypothesis  $\int_0^T M(t) dt < \infty$  with  $M(t) = \|\omega(\cdot, t)\|_{L^\infty}$  is precisely the Beale–Kato–Majda criterion: under it, regularity on  $[0, T]$  already follows classically. The conditional  $\sigma$ -bound therefore cannot serve as a route to regularity; it assumes (an equivalent of) its conclusion.

### 25.6 Defect 6: the two-regime dichotomy requires a large-viscosity condition

The damping constant  $\beta = c\nu/C_1 - C > 0$  requires  $\nu > C_1 C/c$ , a relation between the fixed physical viscosity and universal constants. In the Prize formulation  $\nu$  is given; the condition cannot be arranged by scaling (NS scaling rescales data and  $M$ , not  $\nu$  relative to the constants in this argument). Regime I is therefore conditional.

### 25.7 Defect 7: the Regime II ODE conclusion is false

Regime II derives  $dM/dt \leq C M^{5/4} (\log M)^{1/2}$  and asserts the solution “stays finite on any fixed time interval.” This is incorrect: the comparison ODE  $\dot{M} = C M^{5/4}$  has solution  $M(t) = (M(0)^{-1/4} - \frac{C}{4}t)^{-4}$ , which blows up at the finite time  $t^* = 4M(0)^{-1/4}/C$ ; the logarithmic factor only accelerates this. Regime II as stated does not exclude finite-time blowup.

### 25.8 Defect 8: unproved coercivity in Lemma 22.4

The claim that diffusion contributes  $-c\nu \int |\nabla^2 \omega|^2 \varphi^2$  to  $\frac{d}{dt} A_{\text{loc}}$  is asserted without proof. Differentiating  $A_{\text{loc}} = \int |\omega|^2 |\nabla \hat{e}|^2 \varphi^2$  involves the direction field  $\hat{e} = \omega/|\omega|$ , singular on

$\{\omega = 0\}$ , and produces sign-indefinite cross terms; full  $|\nabla^2 \omega|^2$ -coercivity does not follow from the sketch. Moreover the stretching estimate invokes  $\|\nabla u\|_{L^\infty(B_{r*})} \leq CM$ , but Calderón–Zygmund operators are unbounded on  $L^\infty$ ; the correct bound carries a logarithmic factor, which matters precisely because §22 relies on log-corrected estimates.

## 25.9 What survives the audit

1. Route C bootstrap (Theorem 3.3) — unconditional. *Intact*.
2. Claim A for shear-layer initial data (§11) and near-shear data (§12). *Intact*.
3.  $\lambda_{\max} = 0$  for shear flows (Prop. 11.6). *Intact*.
4. The algebraic identity  $|\nabla \omega|^2 = |\nabla|\omega||^2 + |\omega|^2 |\nabla \hat{e}|^2$  (Lemma 22.1) and the exact scale invariance of  $\sigma = A_{\text{loc}}/M^{3/2}$ . *Intact* — these are sound foundations for a future quantitative alignment program.
5. §20 as a *conditional* small-data statement (consistent with CKN), *provided* Defects 2–4 are repaired; the superlinearity exponent must be re-derived.

## 25.10 Revised status

Global regularity for general  $u_0 \in H^1(\mathbb{T}^3)$  is **open** in this document. The Prize claim of §20 (Corollary 21.6) is withdrawn. The program’s strongest verified position: unconditional regularity for shear and near-shear initial data via Claim A and Route C, plus extensive numerical evidence (873 passing tests) consistent with — but not proving — the general case.

# 26 The Alignment Pincer: Corrected Direction-Field Framework (Session S32)

Following the S31 withdrawal, this section restarts the geometric program on rigorous footing. Every statement below is labeled *Proved*, *Conditional*, or *Open target*. Nothing here claims the Prize.

## 26.1 The exact direction-field evolution equation [Proved]

**Lemma 26.1** (Constantin’s direction equation). *Let  $u$  be a smooth solution of 3D NS and  $\hat{e} = \omega/|\omega|$  on  $\{\omega \neq 0\}$ ,  $\alpha = \hat{e} \cdot S\hat{e}$  with  $S = \frac{1}{2}(\nabla u + \nabla u^\top)$ . Then, with  $D_t = \partial_t + u \cdot \nabla$ :*

$$D_t |\omega| = \alpha |\omega| + \nu (\Delta |\omega| - |\omega| |\nabla \hat{e}|^2), \quad (141)$$

$$D_t \hat{e} = \nu \Delta \hat{e} + 2\nu (\nabla \log |\omega| \cdot \nabla) \hat{e} + \nu |\nabla \hat{e}|^2 \hat{e} + P_{\hat{e}^\perp}((\hat{e} \cdot \nabla)u), \quad (142)$$

where  $P_{\hat{e}^\perp} v = v - (v \cdot \hat{e})\hat{e}$ .

*Proof.* Write  $\omega = |\omega|\hat{e}$ . The vorticity equation gives  $D_t \omega = (\omega \cdot \nabla)u + \nu \Delta \omega = |\omega|(\hat{e} \cdot \nabla)u + \nu \Delta \omega$ . Projecting onto  $\hat{e}$ : since  $|\hat{e}| = 1$  implies  $\hat{e} \cdot \partial_k \hat{e} = 0$  for every  $k$ , we get  $\hat{e} \cdot \Delta \hat{e} = \sum_k \partial_k (\hat{e} \cdot \partial_k \hat{e}) - |\nabla \hat{e}|^2 = -|\nabla \hat{e}|^2$ , hence  $\hat{e} \cdot \Delta \omega = \Delta |\omega| - |\omega| |\nabla \hat{e}|^2$ , and  $\hat{e} \cdot (|\omega|(\hat{e} \cdot \nabla)u) = |\omega| \hat{e}_i \hat{e}_j \partial_j u_i = \alpha |\omega|$  (the antisymmetric part of  $\nabla u$  drops). This is (141). For (142), expand  $D_t(|\omega|\hat{e}) = \hat{e} D_t |\omega| + |\omega| D_t \hat{e}$

and  $\Delta(|\omega|\hat{e}) = \hat{e}\Delta|\omega| + 2(\nabla|\omega| \cdot \nabla)\hat{e} + |\omega|\Delta\hat{e}$ ; substitute (141), divide by  $|\omega|$ , and note  $(\hat{e} \cdot \nabla)u - \alpha\hat{e} = P_{\hat{e}^\perp}((\hat{e} \cdot \nabla)u)$ .  $\square$

**Remark 26.2** (No pressure). Equations (141)–(142) contain no pressure: the direction field is slaved to  $\nabla u$  through Biot–Savart only. This is the same structural advantage §20 sought, obtained without any invalid estimate.

## 26.2 A correctly scale-invariant coherence quantity [Proved]

**Definition 26.3** (Instantaneous coherence deficit). With  $M(t) = \|\omega(\cdot, t)\|_{L^\infty}$ ,  $r^* = M^{-1/2}$ , and  $x^*$  a point where  $|\omega(\cdot, t)| \geq M/2$ :

$$\sigma^*(t) = (r^*)^{-1} \int_{B_{r^*}(x^*)} |\nabla \hat{e}|^2 \mathbf{1}_{\{|\omega| \geq M/4\}} dx. \quad (143)$$

**Proposition 26.4** (Exact scale invariance). *Under  $u_\lambda(x, t) = \lambda u(\lambda x, \lambda^2 t)$  one has  $\omega_\lambda = \lambda^2 \omega(\lambda x, \lambda^2 t)$ ,  $M_\lambda = \lambda^2 M$ ,  $r_\lambda^* = r^*/\lambda$ ,  $\hat{e}_\lambda(x, t) = \hat{e}(\lambda x, \lambda^2 t)$ , so  $|\nabla \hat{e}_\lambda|^2 = \lambda^2 |\nabla \hat{e}|^2(\lambda x)$  and  $\int_{B_{r^*/\lambda}} |\nabla \hat{e}_\lambda|^2 dx = \lambda^{-1} \int_{B_{r^*}} |\nabla \hat{e}|^2 dx$ . Multiplying by  $(r_\lambda^*)^{-1} = \lambda(r^*)^{-1}$  gives  $\sigma_\lambda^* = \sigma^*$ . Moreover the indicator set is scale-covariant since  $|\omega_\lambda| \geq M_\lambda/4 \iff |\omega|(\lambda x) \geq M/4$ .*  $\square$

**Remark 26.5** (Calibration). If  $|\nabla \hat{e}| \sim 1/r^*$  throughout  $B_{r^*}$  (borderline Constantin–Fefferman coherence at the self-similar scale) then  $\sigma^* \sim 1$ . Thus  $\sigma^* \ll 1$  means *better-than-borderline* coherence, and  $\sigma^*$  is the natural small parameter for an  $\varepsilon$ -regularity statement.

**Proposition 26.6** (The S30 quantity controls  $\sigma^*$ ). *With  $\sigma = A_{\text{loc}}/M^{3/2}$  as in Definition (130) (whose scale invariance and algebraic foundation survived the S31 audit):*

$$\sigma^*(t) \leq 16\sigma(t).$$

*Proof.* On  $\{|\omega| \geq M/4\}$  we have  $|\omega|^2 \geq M^2/16$ , so  $\int_{B_{r^*}} |\nabla \hat{e}|^2 \mathbf{1} dx \leq (16/M^2) \int |\omega|^2 |\nabla \hat{e}|^2 \varphi_{r^*}^2 dx = 16 A_{\text{loc}}/M^2$  (using  $\varphi_{r^*} \equiv 1$  on  $B_{r^*}$ ). Multiply by  $(r^*)^{-1} = M^{1/2}$ :  $\sigma^* \leq 16 A_{\text{loc}}/M^{3/2} = 16\sigma$ .  $\square$

## 26.3 Criticality of the direction equation at the self-similar scale [Proved observation]

**Remark 26.7** (Marginality:  $M \cdot (r^*)^2 = 1$ ). In any parabolic estimate for (142) on the cylinder  $Q(r^*) = B_{r^*} \times [t_0 - (r^*)^2, t_0]$ , the stretching term  $P_{\hat{e}^\perp}((\hat{e} \cdot \nabla)u)$  acts as a potential of size  $\|\nabla u\|_{L^\infty} \lesssim M \log(e + \|u\|_{H^3}/M)$  (the logarithm is mandatory: Calderón–Zygmund operators are unbounded on  $L^\infty$  — this was Defect 8 of the S31 audit). Over the time span  $(r^*)^2 = M^{-1}$  its Gronwall exponent is

$$\|\nabla u\|_{L^\infty} \cdot (r^*)^2 \lesssim M \cdot M^{-1} \cdot \log(\cdot) = \log(\cdot).$$

The direction equation is therefore *exactly critical* (marginally so, up to a logarithm) at the scale  $r^* = M^{-1/2}$ : the potential cannot amplify  $\nabla \hat{e}$  by more than a power of the logarithm per self-similar time scale. This is the structural reason an  $\varepsilon$ -regularity theory for  $\hat{e}$  at scale  $r^*$  is plausible, where the corresponding statement for  $u$  itself (CKN) is supercritical.

## 26.4 The Bridge Lemma [Open target]

*Conjecture 26.8* (Bridge Lemma: averaged-to-pointwise coherence upgrade). There exist  $\varepsilon_0 > 0$  and  $C_B \geq 1$  (possibly degrading logarithmically in  $M$ ) such that: if  $u$  satisfies the Type-I bound  $\sup_{Q(r^*)} |u| \leq C_I/r^*$  and  $\sup_{t \in [t_0 - (r^*)^2, t_0]} \sigma^*(t) \leq \varepsilon_0$ , then

$$\|\nabla \hat{e}\|_{L^\infty(Q(r^*/2) \cap \{|\omega| \geq M/2\})} \leq \frac{C_B}{r^*}.$$

**Proof strategy and identified obstacles.** The natural proof is Moser iteration for  $w = |\nabla \hat{e}|^2$ , which by (142) satisfies a drift-diffusion inequality with (i) drift  $u + 2\nu \nabla \log |\omega|$  and (ii) potential  $\lesssim \|\nabla u\|_{L^\infty(B_{r^*})}$ . Obstacle (i): on  $\{|\omega| \geq M/4\}$  the singular drift satisfies  $|\nabla \log |\omega|| \leq 4|\nabla \omega|/M$ , which requires a pointwise gradient bound not currently available; a workaround is to run the iteration on level sets of  $|\omega|$  where the drift enters only through its divergence structure. Obstacle (ii): the potential is marginal by Remark 26.7, so the iteration constants degrade logarithmically — acceptable, but the bookkeeping must be done. Neither obstacle is resolved here; this is the program’s central open estimate, replacing the invalid superlinearity of §20.

## 26.5 The pincer [Conditional on two named inputs]

**Theorem 26.9** (Disorder barrier for Type-I singularities — conditional). *Assume (H1) the Bridge Lemma (Conjecture 26.8) and (H2) a localized Constantin–Fefferman criterion at scale  $r^*$ : coherence  $\|\nabla \hat{e}\|_{L^\infty} \leq C_B/r^*$  on  $Q(r^*/2) \cap \{|\omega| \geq M/2\}$  implies the depletion bound  $\alpha(x^*, t) \leq C M^{1/2+\theta}$  for some  $\theta < 1/2$  near a putative Type-I blowup time. Then no Type-I singularity can satisfy  $\liminf_{t \rightarrow T^-} \sigma^*(t) \leq \varepsilon_0$ : every Type-I singularity must maintain  $\sigma^*(t) > \varepsilon_0$  (persistent geometric disorder) for all  $t$  close to  $T$ .*

*Proof (given H1–H2).* If  $\sigma^*(t_k) \leq \varepsilon_0$  along  $t_k \rightarrow T^-$ , H1 upgrades averaged coherence to pointwise coherence on  $Q(r^*(t_k)/2)$ ; H2 then bounds the logarithmic growth rate  $\frac{d}{dt} \log M = \alpha(x^*, t) \leq C M^{1/2+\theta}$  with  $1/2+\theta < 1$ , whose integral  $\int^T M^{1/2+\theta} dt$  converges under the Type-I bound  $M \leq C_I/(T-t)$  precisely when  $1/2+\theta < 1$ . Hence  $M$  stays bounded as  $t \rightarrow T^-$ , contradicting blowup.  $\square$

**Remark 26.10** (Status of H2). H2 is a genuine second open input: the classical Constantin–Fefferman theorem [12] assumes coherence at a *fixed* scale  $\rho$ , whereas here the coherence scale  $r^*(t) \rightarrow 0$  shrinks with the putative singularity. The localized/shrinking-scale variants (Beirão da Veiga–Berselli type  $1/2$ -Hölder criteria, Grujić’s geometric depletion) are the relevant literature; adapting them to scale  $r^*$  with explicit  $\theta$  is a bounded, well-posed task.

## 26.6 Honest status table for the pincer program

| Item                                   | Status               | Where                      |
|----------------------------------------|----------------------|----------------------------|
| Direction equation (142)               | Proved               | Lemma 26.1                 |
| Scale invariance of $\sigma^*$         | Proved               | Prop. 26.4                 |
| $\sigma^* \leq 16\sigma$ (link to S30) | Proved               | Prop. 26.6                 |
| Criticality $M(r^*)^2 = 1$             | Proved observation   | Remark 26.7                |
| Bridge Lemma (H1)                      | <b>Open target</b>   | Conjecture 26.8            |
| Localized CF at scale $r^*$ (H2)       | <b>Open target</b>   | Theorem 26.9               |
| Disorder barrier                       | Conditional on H1+H2 | Theorem 26.9               |
| Numerical bridge diagnostic            | EXP-L4-BRIDGE-*      | layer4/alignment_bridge.py |

## 27 Level-Set Moser Scheme for the Bridge Lemma: The First Rung (Session S33)

This section attacks Obstacle (i) of Conjecture 26.8: the singular drift  $2\nu\nabla \log |\omega|$  in the direction equation (142), whose differentiated form produces  $\nabla^2 \log |\omega|$  — uncontrollable pointwise. The device is a *level-set formulation*: all estimates are run with a smooth cutoff in the  $|\omega|$ -variable, and every second-derivative-of-magnitude term is integrated by parts back onto first-order factors that the level set controls. The outcome (Proposition 27.8) closes the first rung of a Moser iteration with logarithmically degraded constants and *reduces* Obstacle (i) to a strictly weaker, sharply stated coupling estimate (Target 27.10).

### 27.1 Standing assumptions and non-circularity

Throughout this section  $u$  is a solution that is smooth on  $\mathbb{T}^3 \times [0, T)$  with putative first singularity at  $T$ , satisfying the Type-I bound  $M(t) \leq C_I/(T-t)$  and  $\sup |u(\cdot, t)| \leq C_I(T-t)^{-1/2}$ . All computations occur at times  $t < T$  where the solution is  $C^\infty$ .

**Remark 27.1** (No Leray–Hopf circularity). Defect 4 of the S31 audit (§25) noted that §20 applied a local enstrophy inequality to Leray–Hopf solutions for which it is not justified. The pincer route is immune to this defect by design: it is a *contradiction argument at a putative first singularity*, so every estimate is performed on the smooth interval  $[0, T)$  where all manipulations are classical. No weak-solution regularity is assumed that is not available.

### 27.2 The composite level-set cutoff [Proved]

**Definition 27.2** (Composite cutoff). Fix  $\chi \in C^\infty([0, \infty); [0, 1])$  with  $\chi \equiv 0$  on  $[0, \frac{1}{4}]$ ,  $\chi \equiv 1$  on  $[\frac{1}{2}, \infty)$ ,  $|\chi'| \leq 8$ . For scales  $\frac{r^*}{2} \leq \rho < \rho' \leq r^*$  let  $\psi \in C_c^\infty(Q(\rho'))$  be a standard space-time cutoff with  $\psi \equiv 1$  on  $Q(\rho)$ ,  $|\nabla \psi| \leq 2/(\rho' - \rho)$ ,  $|\partial_t \psi| \leq 2/(\rho' - \rho)^2$ . The composite cutoff is

$$\eta(x, t) = \psi(x, t) \chi(|\omega(x, t)|/M(t)).$$

**Lemma 27.3** (Cutoff derivative bounds). *On  $\text{supp } \eta \subseteq \{|\omega| \geq M/4\}$ :*

$$|\nabla \eta| \leq \frac{2}{\rho' - \rho} + \frac{8}{M} |\nabla |\omega||, \quad (144)$$

$$|D_t \chi(|\omega|/M)| \leq \frac{8}{M} \left( |\alpha| |\omega| + \nu |\Delta |\omega|| + \nu |\omega| |\nabla \hat{e}|^2 \right) + 8 \frac{|\dot{M}|}{M}. \quad (145)$$



*Proof.* (144): product rule and  $|\nabla\chi(|\omega|/M)| = |\chi'| |\nabla|\omega||/M$ . (145):  $D_t\chi = \chi'(D_t|\omega|/M - |\omega|\dot{M}/M^2)$ ; substitute the magnitude equation (141) and use  $|\omega|/M \leq 1$ ,  $|\chi'| \leq 8$ .  $\square$

**Remark 27.4.** The term  $\nu|\Delta|\omega||$  in (145) is second-order and is *never used pointwise*: wherever  $D_t\chi$  appears inside an integral it is integrated by parts (Lemma 27.5) so that only  $\nabla|\omega|$  survives. Under Type I,  $|\dot{M}|/M \leq \|\nabla u\|_{L^\infty} \leq CM \log(e + \|u\|_{H^3}/M)$  for a.e.  $t$  (Rademacher for the Lipschitz function  $M(t)$ ), which contributes the marginal factor  $M \log \cdot (r^*)^2 = \log$  per cylinder (Remark 26.7).

### 27.3 The three integrations by parts [Proved]

Testing the differentiated direction equation with  $\eta^2 \partial_k \hat{e}$  (summed over  $k$ ) yields, for  $w = |\nabla \hat{e}|^2$ , the weak inequality

$$\frac{1}{2} \sup_t \int w \eta^2 + \nu \iint |\nabla^2 \hat{e}|^2 \eta^2 \leq \frac{1}{2} \iint w |D_t(\eta^2)| + \nu \iint w |\Delta(\eta^2)| + \mathcal{D} + \mathcal{S}, \quad (146)$$

where  $\mathcal{D}$  collects the drift terms generated by  $2\nu(\nabla \log |\omega| \cdot \nabla) \hat{e}$  and  $\mathcal{S}$  the stretching terms generated by  $\nu w \hat{e} + P_{\hat{e}^\perp}((\hat{e} \cdot \nabla)u)$ .

**Lemma 27.5** (Three integrations by parts). *Each second-derivative-of-magnitude term in (146) can be rewritten, using only integration by parts and the support property of  $\eta$ , in forms containing at most first derivatives of  $|\omega|$ :*

(a) *the differentiated drift term*

$$2\nu \iint (\partial_k \partial_j \log |\omega|)(\partial_j \hat{e} \cdot \partial_k \hat{e}) \eta^2 = -2\nu \iint \partial_j \log |\omega| \partial_k \left[ (\partial_j \hat{e} \cdot \partial_k \hat{e}) \eta^2 \right],$$

*whose integrand is bounded by  $\frac{8\nu}{M} |\nabla|\omega|| (2|\nabla^2 \hat{e}| |\nabla \hat{e}| \eta^2 + 2w \eta |\nabla \eta|)$  on  $\text{supp } \eta$ ;*

(b) *the  $\nu \Delta|\omega|$  contribution inside  $D_t\chi$  (Lemma 27.3):*

$$\frac{\nu}{M} \iint w \eta \psi |\chi'| |\Delta|\omega|| \rightsquigarrow -\frac{\nu}{M} \iint \nabla|\omega| \cdot \nabla (w \eta \psi \chi'),$$

*with integrand bounded by  $\frac{C\nu}{M} |\nabla|\omega|| (|\nabla^2 \hat{e}| |\nabla \hat{e}| \eta + w |\nabla \eta| + \frac{w\eta}{M} |\nabla|\omega||)$ ;*

(c) *the cutoff-gradient term  $\nu \iint w |\Delta(\eta^2)|$ , whose  $\chi$ -part after one integration by parts contains only  $|\nabla|\omega||/M$  factors as in (a).*

Consequently, applying Young's inequality with parameter  $\frac{\nu}{8}$  to every product  $|\nabla^2 \hat{e}| \times (\text{first-order factor})$ , all three terms are absorbed as

$$\mathcal{D} + (b) + (c) \leq \frac{\nu}{2} \iint |\nabla^2 \hat{e}|^2 \eta^2 + \frac{C\nu}{(\rho' - \rho)^2} \iint_{Q(\rho')} w \mathbf{1}_{\{|\omega| \geq M/4\}} + C\nu K(\rho'),$$

where

$$K(\rho') = \frac{1}{M^2} \iint_{Q(\rho') \cap \{|\omega| \geq M/4\}} |\nabla|\omega||^2 (w + (\rho' - \rho)^{-2}) dx dt. \quad (147)$$

*Proof.* (a) and (b) are direct integrations by parts (no boundary terms:  $\eta$  is compactly supported in space-time, and the integrands vanish on  $\{|\omega| < M/4\}$  where  $\chi = 0$ , so the level-set boundary contributes nothing). The stated integrand bounds use  $|\nabla \log |\omega|| \leq 4|\nabla |\omega||/M$  on  $\text{supp } \eta$ ,  $|\chi'| \leq 8$ , Lemma 27.3, and  $|\nabla(w\eta\psi\chi')| \leq 2|\nabla^2 \hat{e}||\nabla \hat{e}|\eta\psi|\chi'| + w|\nabla(\eta\psi\chi')|$ . The absorption step is Young's inequality  $ab \leq \frac{\nu}{8}a^2 + \frac{2}{\nu}b^2$  applied term by term; every non-dissipative square lands either on  $w/(\rho' - \rho)^2$  or on  $(|\nabla |\omega||^2/M^2) \cdot (w + (\rho' - \rho)^{-2})$ , which is  $K(\rho')$  by definition.  $\square$

## 27.4 Localized stretching control under Type I [Proved]

**Lemma 27.6** (Local enstrophy-gradient bound, Type I). *On the smooth interval, under the Type-I bounds, the vorticity Caccioppoli inequality (classical for smooth solutions; cf. Remark 27.1) with  $\|\nabla u\|_{L^\infty} \lesssim M \log(e + \|u\|_{H^3}/M)$  gives*

$$\nu \iint_{Q(2\rho)} |\nabla \omega|^2 \lesssim M^2 \rho^3 (1 + M \rho^2 \mathcal{L}) \lesssim M^2 \rho^3 \mathcal{L}, \quad \rho \leq r^*,$$

where  $\mathcal{L} = \log(e + \|u\|_{H^3}/M)$ , using  $M \rho^2 \leq M(r^*)^2 = 1$ .

**Lemma 27.7** (Biot–Savart near/far splitting). *Write  $u = u_{\text{loc}} + u_{\text{far}}$  with  $u_{\text{loc}}$  the Biot–Savart integral of  $\omega \mathbf{1}_{B_{2r^*}}$ . Then  $\|\nabla^2 u_{\text{loc}}\|_{L^2(B_{r^*})} \leq C_{\text{CZ}} \|\nabla \omega\|_{L^2(B_{2r^*})}$  (Calderón–Zygmund,  $L^2$ ), while  $u_{\text{far}}$  is harmonic-regular on  $B_{r^*}$  with  $\|\nabla^2 u_{\text{far}}\|_{L^\infty(B_{r^*})} \leq C(r^*)^{-2} \sup |u| \leq C C_I (r^*)^{-3}$  under Type I. Hence the stretching contribution  $\mathcal{S}$  in (146) obeys*

$$\mathcal{S} \leq \frac{\nu}{4} \iint |\nabla^2 \hat{e}|^2 \eta^2 + C M \mathcal{L} \iint_{Q(\rho')} w \mathbf{1} + C \nu^{-1} \mathcal{L} M^2 (r^*)^5 + C C_I^2 r^*,$$

where the third term uses Lemma 27.6 and the fourth is the far-field contribution integrated over  $Q(r^*)$ .

## 27.5 The first rung [Proved modulo routine verifications]

**Proposition 27.8** (First-rung Caccioppoli for the angular energy). *Under the standing assumptions, for  $\frac{r^*}{2} \leq \rho < \rho' \leq r^*$ :*

$$\begin{aligned} \sup_t \int_{B_\rho} w \mathbf{1}_{\{|\omega| \geq M/2\}} dx + \nu \iint_{Q(\rho)} |\nabla^2 \hat{e}|^2 \mathbf{1}_{\{|\omega| \geq M/2\}} \\ \leq \frac{C \mathcal{L}}{(\rho' - \rho)^2} \iint_{Q(\rho')} w \mathbf{1}_{\{|\omega| \geq M/4\}} + C \nu K(\rho') + C(\nu^{-1} \mathcal{L} M^2 (r^*)^5 + C_I^2 r^*), \end{aligned}$$

with  $C = C(C_I)$  universal up to the Type-I constant and  $\mathcal{L}$  the logarithmic factor. Status: the displayed dangerous terms are handled in full by Lemmas 27.3–27.7; the routine verifications that remain for S34 are enumerated in Remark 27.9 and are bookkeeping, not new estimates.

**Remark 27.9** (Enumerated routine verifications). (i) the transport term  $\iint w u \cdot \nabla(\eta^2)$  (standard: Hölder + Type-I  $|u| \leq C_I/r^*$ ); (ii) the algebraic constant in the commutation  $[\partial_k, D_t]\hat{e} = -(\partial_k u \cdot \nabla)\hat{e}$ ; (iii) *reclassified in S34*: the  $\nu w \hat{e}$ -term yields the critical quadratic  $\nu \iint w^2 \eta^2$  ( $\partial_k \hat{e} \cdot \hat{e} = 0$  kills only the  $\partial_k w$  part); this is *not* routine — it is the standard critical nonlinearity of the iteration, handled by Gagliardo–Nirenberg interpolation plus the  $\sigma^* \leq \varepsilon_0$  smallness at the second rung (see §28); (iv) the time-slice boundary term at  $t_0 - (\rho')^2$  (vanishes by choice of  $\psi$ ). None of these involves  $\nabla^2 \log |\omega|$  or nonlocality.

## 27.6 What remains of Obstacle (i): the coupling target [Open]

*Conjecture 27.10* ( $K$ -absorption). Under the standing assumptions there exist  $\theta_K < 1$  and  $C_K$  (possibly logarithmic in  $M$ ) such that

$$\nu K(r^*) \leq \theta_K \left[ \text{LHS of Proposition 27.8} \right] + C_K (1 + \sigma^*) r^*.$$

**Remark 27.11** (Why this is strictly weaker than Obstacle (i)). Obstacle (i) demanded a *pointwise* bound on  $\nabla \log |\omega|$ . Target 27.10 demands only that the *integrated* radial-angular coupling  $M^{-2} \iint |\nabla |\omega||^2 w$  be absorbable. Two independent routes are visible: (1) interpolation —  $K$  is quartic in first derivatives and can be split by Hölder between the local enstrophy bound (Lemma 27.6) and  $\|w\|_{L^\infty}$ , making the absorption a small- $\sigma^*$  smallness argument at the second rung of the iteration; (2) the magnitude equation (141) gives its own Caccioppoli for  $|\nabla |\omega||^2$  with the *good sign* term  $-\nu |\omega| w$  appearing as a sink. Route (2) is the natural S34 starting point.

**Remark 27.12** (Numerical status). The S32–S33 diagnostics (`layer4/alignment_bridge.py`, EXP-L4-BRIDGE-\*) show the bridge ratio  $\|\nabla \hat{e}\|_{L^\infty} r^* / \sqrt{\sigma^*}$  remaining  $O(1)$  through strong vortex stretching, consistent with Conjecture 26.8 and with the absorbability of  $K$ .

## 28 The Magnitude-Equation Caccioppoli: A Priori Coherence Bound and the Refined $K$ -Target (Session S34)

### 28.1 Corrigendum to §27 [dimensional audit]

A dimensional re-audit of §27 (performed before building on it, per the S31 discipline) found three slips in the inhomogeneous terms of Lemma 27.7, Proposition 27.8 and Conjecture 27.10: in parabolic units every term of the first-rung inequality scales as  $\ell^1$ , forcing the corrected forms  $\nu^{-1} \mathcal{L} M^2 (r^*)^5$  and  $C_I^2 r^*$  (both now *vanish* as  $r^* \rightarrow 0$  — the corrected inequality is subcritically favourable, not worse). Additionally, routine item (iii) of Remark 27.9 is reclassified: it is the critical quadratic  $\nu \iint w^2 \eta^2$ , i.e. the standard critical nonlinearity of every  $\varepsilon$ -regularity iteration, handled by Gagliardo–Nirenberg plus the  $\sigma^* \leq \varepsilon_0$  hypothesis — this is precisely where the Bridge Lemma’s smallness assumption enters the scheme, as it should. The corrections are incorporated above in place.

### 28.2 The magnitude Caccioppoli and its good-sign sink [Proved]

The magnitude equation (141) carries the dissipative term  $-\nu |\omega| w$  ( $w = |\nabla \hat{e}|^2$ ): angular roughness *drains* vorticity magnitude. Testing with  $m \eta^2$  ( $m = |\omega|$ ,  $\eta$  the composite cutoff of Definition 27.2 with  $\chi$ -levels  $\frac{1}{8}, \frac{1}{4}$  so that  $\{m \geq M/4\} \subseteq \{\chi = 1\}$ ) turns this sink into a *left-hand-side* control of the coherence deficit:

$$\nu \iint m^2 w \eta^2 + \nu \iint |\nabla m|^2 \eta^2 + \frac{1}{2} \sup_t \int m^2 \eta^2 \leq \iint \alpha m^2 \eta^2 + \frac{1}{2} \iint m^2 |D_t(\eta^2)| + \nu \iint m^2 |\nabla \eta|^2. \quad (148)$$

**Lemma 28.1** (All right-hand terms are  $O(M^{1/2} \mathcal{L})$ ). *Under the standing Type-I assumptions of §27, on  $Q(r^*)$ :*

- (a)  $\iint |\alpha| m^2 \eta^2 \leq \|\nabla u\|_{L^\infty} M^2 (r^*)^5 \leq C \mathcal{L} M^{1/2}$  (using  $M(r^*)^2 = 1$ );
- (b) the  $\psi$ -part of  $D_t(\eta^2)$  contributes  $\leq C(1 + C_I) M^{1/2}$ ;
- (c) the  $\chi$ -part of  $D_t(\eta^2)$  and the  $\chi$ -part of  $|\nabla \eta|^2$  contribute  $\leq C \mathcal{L} M^{1/2}$ , with no absorption argument needed: every such term is supported on the transition annulus and is dominated pointwise via the orthogonal split (Lemma 22.1)

$$m^2 w = |\nabla \omega|^2 - |\nabla m|^2 \leq |\nabla \omega|^2, \quad |\nabla m|^2 \leq |\nabla \omega|^2,$$

after which Lemma 27.6 gives  $\nu \iint_{Q(r^*)} |\nabla \omega|^2 \leq C M^2 (r^*)^3 \mathcal{L} = C \mathcal{L} M^{1/2}$ . The  $\nu \Delta m$  term inside  $D_t \chi$  is first integrated by parts exactly as in Lemma 27.5(b), landing on the same enstrophy bound.

**Theorem 28.2** (A priori coherence bound under Type I). *There is  $C = C(C_I)$  such that for every  $t_0 < T$ , with  $r^* = M(t_0)^{-1/2}$ :*

$$\langle \sigma^* \rangle := (r^*)^{-3} \iint_{Q(r^*/2)} w \mathbf{1}_{\{|\omega| \geq M/2\}} dx dt \leq \frac{C \mathcal{L}}{\nu}.$$

*Proof.* On  $\{|\omega| \geq M/2\} \cap \{\eta = 1\}$  we have  $m^2 \geq M^2/4$ , so (148) and Lemma 28.1 give  $\nu (M^2/4) \iint_{Q(r^*/2)} w \mathbf{1} \leq C \mathcal{L} M^{1/2}$ , i.e.  $\iint w \mathbf{1} \leq (C \mathcal{L}/\nu) M^{-3/2} = (C \mathcal{L}/\nu) (r^*)^3$ .  $\square$

**Remark 28.3** (What the theorem changes). Near a Type-I singularity the coherence deficit *cannot be large*: it is a priori  $O(\mathcal{L}/\nu)$ . The Bridge Lemma requires  $\sigma^* \leq \varepsilon_0$ ; the gap between “automatic” and “needed” is now the explicit factor  $C \mathcal{L}/(\nu \varepsilon_0)$ , and closing it is a *decay* question for the  $\sigma^*$ -dynamics, not a boundedness question. This is the first unconditional (within Type I) quantitative output of the pincer program.

### 28.3 The refined $K$ -target: anti-correlation, not smallness [Open, measurable]

**Proposition 28.4** (Crude absorption fails by exactly one logarithm). *By Hölder and Lemma 27.6,*

$$\nu K_w = \frac{\nu}{M^2} \iint |\nabla m|^2 w \mathbf{1} \leq \|w\|_{L^\infty} \cdot \frac{1}{M^2} \left( \nu \iint |\nabla \omega|^2 \right) \leq C \mathcal{L} \|w\|_{L^\infty} (r^*)^3.$$

Since  $\|w\|_{L^\infty} (r^*)^3$  is comparable to the quantity the Moser scheme outputs, the coupling term re-enters at top order multiplied by  $\mathcal{L}$ : the crude route misses closing the loop by exactly one logarithm.

**Conjecture 28.5** (Refined  $K$ -absorption via equipartition). Define the coupling correlation on  $E = Q(r^*) \cap \{|\omega| \geq M/4\}$ :

$$c_K = \frac{\iint_E |\nabla m|^2 w}{\left( - \int_E |\nabla m|^2 \right) \left( - \int_E w \right) |E|} = \frac{\langle |\nabla m|^2 w \rangle_E}{\langle |\nabla m|^2 \rangle_E \langle w \rangle_E}.$$

Then  $c_K \leq C/\mathcal{L}$  near a putative Type-I singularity.

**Remark 28.6** (Why equipartition is the mechanism). The orthogonal split  $|\nabla\omega|^2 = |\nabla m|^2 + m^2 w$  makes  $|\nabla m|^2$  and  $m^2 w$  competitors for the same budget: where angular variation is maximal, radial variation is squeezed. Perfect equipartition or anti-correlation gives  $c_K \leq 1$  with room to spare; only strong *positive* correlation ( $c_K \gg 1$ ) could defeat the absorption. The S33 numerics (adversarial bridge ratio saturating  $R \approx 1$  without violation) already suggest equipartition-like behaviour;  $c_K$  is directly measurable and is added to the diagnostic suite in S34 (layer4, EXP-L4-KCORR-\*). If the measured  $c_K$  is  $O(1)$  or smaller and stable across ICs and resolutions, the conjecture’s logarithmic form is plausibly provable from the equipartition structure of (148), whose sink term penalises exactly the configurations with  $c_K$  large.

## 28.4 Updated program ledger

| Item                                                            | Status                                        | Where      |
|-----------------------------------------------------------------|-----------------------------------------------|------------|
| First-rung Caccioppoli (corrected)                              | Proved mod (i),(ii),(iv)                      | Prop. 27.8 |
| Critical quadratic (was item (iii))                             | Standard, needs $\sigma^* \leq \varepsilon_0$ | Rem. 27.9  |
| A priori bound $\langle \sigma^* \rangle \leq C\mathcal{L}/\nu$ | <b>Proved</b> (Type I)                        | Thm. 28.2  |
| Crude $K$ -absorption                                           | Fails by one log (sharp)                      | Prop. 28.4 |
| Refined $K$ -absorption ( $c_K \leq C/\mathcal{L}$ )            | <b>Open, measurable</b>                       | Conj. 28.5 |
| $H2$ (localized CF at $r^*$ )                                   | Open                                          | §26        |

## 29 The Second Rung and the $K$ -Self-Absorption Identity (Session S35)

### 29.1 The parabolic energy space and the critical quadratic [Proved given the first rung]

Let  $f = |\nabla \hat{e}| \eta$  and define the parabolic energy norm on  $Q(\rho')$ :

$$E_V = \sup_t \int f^2 dx + \nu \iint |\nabla f|^2 dx dt,$$

which is exactly what the (corrected) first rung (Proposition 27.8) controls, up to the  $K$ -term:  $E_V \leq C\mathcal{L}\sigma^* r^* + C\nu K(\rho') + o(r^*)$ . By the standard parabolic embedding  $L_t^\infty L_x^2 \cap L_t^2 H_x^1 \hookrightarrow L^{10/3}(Q)$  in three space dimensions (Ladyzhenskaya–Solonnikov–Ural’tseva),  $\iint f^{10/3} \leq C\nu^{-?} E_V^{5/3}$  with the viscosity bookkeeping carried explicitly below.

**Proposition 29.1** (Second rung: the critical quadratic yields to  $\sigma^*$ -smallness). *With  $w = |\nabla \hat{e}|^2$ , split  $w^2 = f^{10/3} \cdot w^{1/3}$  pointwise on  $\{\eta = 1\}$ . For any  $\delta \in (0, 1)$ :*

$$\nu \iint w^2 \eta^2 \leq \delta \nu (r^*)^3 \|w\|_{L^\infty(Q(\rho'))} + C \delta^{-1/2} \nu E_V^{5/2} (r^*)^{-3/2}.$$

(Dimensional check in parabolic units: every term scales as  $\ell^1$ .) Consequently, if  $\sigma^* \leq \varepsilon_0$  and the  $K$ -term is absorbed (§below), then  $E_V \leq C\mathcal{L}\varepsilon_0 r^*$  and the second term is  $C\delta^{-1/2}(\mathcal{L}\varepsilon_0)^{5/2} r^*$  — superlinear in  $\varepsilon_0$ , hence harmless; while the first term is absorbed into the Moser sup-bound with factor  $\delta$ . The critical quadratic (reclassified item (iii) of Remark 27.9) is thereby handled: the entire nonlinearity of the direction equation is subordinate to  $\sigma^*$ -smallness.

*Proof.* Hölder with exponents  $(3, \frac{3}{2})$  on  $w^2\eta^2 = w^{1/3} \cdot f^{10/3} \cdot (\eta\text{-adjust})$  gives  $\iint w^2\eta^2 \leq \|w\|_{L^\infty(Q(\rho'))}^{1/3} \iint f^{10/3}$ . Apply the parabolic embedding  $\iint f^{10/3} \leq C\nu^{-2/3} E_V^{5/3}$  (three spatial dimensions; the  $\nu^{-2/3}$  tracks the dissipation weight in  $E_V$ ), then Young's inequality  $a^{1/3}b \leq \delta a + C\delta^{-1/2}b^{3/2}$  with  $a = \nu\|w\|_\infty(r^*)^3$  and  $b = \nu^{1/3} E_V^{5/3}(r^*)^{-1}$ ; both output terms carry the dimensions  $\ell^1$  as displayed.  $\square$

## 29.2 The $K$ -self-absorption identity [Derived; two couplings pending verification]

The S34 crude route (Proposition 28.4) loses a logarithm because it decouples  $|\nabla m|^2$  from  $w$  by Hölder. The following identity keeps them coupled: *test the magnitude equation (141) with  $m w \eta^2$  instead of  $m \eta^2$ .*

**Proposition 29.2** ( $K$ -self-absorption — provisional). *Testing (141) against  $m w \eta^2$  and integrating by parts yields*

$$\nu \iint |\nabla m|^2 w \eta^2 + \nu \iint m^2 w^2 \eta^2 \leq C \nu \iint m^2 |\nabla^2 \hat{e}|^2 \eta^2 + \mathcal{R}, \quad (149)$$

where  $\mathcal{R}$  collects the transport,  $\alpha$ -, and cutoff terms. The first left-hand term is exactly  $M^2$  times the density of  $K_w$ ; the bound says the radial–angular coupling is controlled by the angular dissipation itself — with an absolute constant, no logarithm, and no correlation hypothesis. Moreover the second left-hand term is the critical quadratic with the favourable weight  $m^2 \geq M^2/16$ , reinforcing Proposition 29.1.

*Derivation, with two couplings flagged.* The viscous term gives  $\nu \iint \Delta m \cdot m w \eta^2 = -\nu \iint |\nabla m|^2 w \eta^2 - \nu \iint m \nabla m \cdot \nabla w \eta^2 - \nu \iint m w \nabla m \cdot \nabla(\eta^2)$ , and the sink term  $-\nu m w$  of (141) multiplied by  $m w \eta^2$  gives  $-\nu \iint m^2 w^2 \eta^2$  on the right, i.e. the second good term on the left of (149). For the middle viscous cross-term,  $|\nabla w| \leq 2|\nabla^2 \hat{e}| w^{1/2}$  pointwise, so by Cauchy–Schwarz

$$\nu \iint m |\nabla m| |\nabla w| \eta^2 \leq \frac{1}{2} \nu \iint |\nabla m|^2 w \eta^2 + 2 \nu \iint m^2 |\nabla^2 \hat{e}|^2 \eta^2,$$

absorbing half the  $K$ -density into the left and producing the angular dissipation on the right. **Coupling C1 (pending):** the time term  $\iint D_t m \cdot m w \eta^2 = \frac{1}{2} \iint D_t(m^2) w \eta^2$  requires substituting  $D_t w$  from the differentiated direction equation after integrating by parts in time; its dangerous part is again of the form  $\nu \iint m^2 \nabla w \cdot \nabla(\cdot)$  and is expected to close by the same Cauchy–Schwarz, but the full bookkeeping (including the  $2\nu(\nabla \log m \cdot \nabla)w$  term, which re-enters  $K$  with a constant that must come out  $< 1$ ) is deferred to S36. **Coupling C2 (pending):** the stretching term  $\iint \alpha m^2 w \eta^2 \leq \|\nabla u\|_{L^\infty} \iint m^2 w \eta^2 \leq C \mathcal{L} M^{1/2} \cdot \nu^{-1}$ -type via Theorem 28.2; marginal ( $\mathcal{L}$ -degraded), same class as all previous marginal terms, but the constant chase is part of the S36 verification. Until C1–C2 are verified, the proposition is *provisional*.  $\square$

**Remark 29.3** (S36 amendment to Proposition 29.2). The verification in §30 found that the second left-hand term  $\nu \iint m^2 w^2 \eta^2$  of (149) *cancels exactly* against the  $\nu w^2$  feedback contained in  $D_t w$  (Lemma 30.1); the displayed identity must be read without it. The  $K$ -control conclusion stands (with amended constants); the critical quadratic is handled solely by the second rung (Proposition 29.1), which suffices.

**Remark 29.4** (Consequence if C1–C2 verify). Conjecture 28.5 (the  $c_K$  decorrelation hypothesis) becomes *unnecessary*:  $K$ -absorption follows structurally from (149), and H1 (the Bridge Lemma, Conjecture 26.8) reduces to: (a)  $\sigma^* \leq \varepsilon_0$  at the working scale, plus (b) the routine items (i),(ii),(iv) of Remark 27.9. The measured  $c_K \approx 1$  (EXP-L4-KCORR-\*) is then explained rather than assumed: equipartition is enforced by the magnitude equation’s sink, which is exactly what the weighted test function exposes.

### 29.3 Honest distance to the Prize

**Remark 29.5** (What this program can and cannot yet claim). Even with H1 and H2 fully proved, the pincer yields: *Type-I singularities require persistent geometric disorder*, and — if the  $\sigma^*$ -decay gap (from the a priori  $C\mathcal{L}/\nu$  of Theorem 28.2 down to  $\varepsilon_0$ ) is closed — *no Type-I blowup*. Two further steps separate that from the Clay problem: (1) the  $\sigma^*$ -decay mechanism (open; the S31 audit removed the previous candidate), and (2) removal of the Type-I hypothesis entirely (wide open; every marginal estimate in §27–§29 uses Type-I scaling, so Type-II exclusion requires genuinely new input). Global existence of weak solutions is classical (Leray); the open question is smoothness, and this document claims no more than the ledger above.

### 29.4 Updated program ledger (post-S35)

| Item                                                               | Status                                          | Where      |
|--------------------------------------------------------------------|-------------------------------------------------|------------|
| Second rung (critical quadratic)                                   | Proved given first rung + $\varepsilon_0$       | Prop. 29.1 |
| $K$ -self-absorption identity                                      | Derived; C1–C2 pending (S36)                    | Prop. 29.2 |
| $c_K$ hypothesis                                                   | Superseded if C1–C2 verify                      | Rem. 29.4  |
| H1 (Bridge Lemma)                                                  | Reduces to $\varepsilon_0$ + routine, mod C1–C2 | Conj. 26.8 |
| $\sigma^*$ -decay ( $C\mathcal{L}/\nu \rightarrow \varepsilon_0$ ) | Open                                            | Thm. 28.2  |
| H2 (localized CF)                                                  | Open                                            | §26        |
| Type-II exclusion                                                  | Open (outside current methods)                  | —          |

## 30 Verification of C1–C2, Gram Negativity, and the Annulus Residual (Session S36)

This section carries out the verification of couplings C1–C2 flagged in Proposition 29.2, discharges the routine items (i), (ii), (iv) of Remark 27.9, and records one correction to §29.

### 30.1 C2: the $\alpha$ -weighted terms [Verified]

Every term of the form  $\iint |\alpha| m^2 w \eta^2$  or  $M\mathcal{L}$ -weighted variants obeys, by  $|\alpha| \leq \|\nabla u\|_{L^\infty} \leq CM\mathcal{L}$  and the a priori bound  $\nu \iint m^2 w \eta^2 \leq C\mathcal{L}M^{1/2}$  (the proof of Theorem 28.2):

$$\iint |\alpha| m^2 w \eta^2 \leq CM\mathcal{L} \cdot \nu^{-1} \cdot C\mathcal{L}M^{1/2} = C\mathcal{L}^2 M^{3/2} / \nu.$$

In  $K$ -units (divide by  $\nu M^2$ ) this is  $C\mathcal{L}^2 r^* / \nu^2$  — an inhomogeneous term of exactly the allowed  $C_K(1 + \sigma^*)r^*$  class (with  $\mathcal{L}^2$ -degraded constant). *C2 is verified.*

### 30.2 C1: the $D_t w$ substitution [Verified modulo the annulus residual]

Substituting the  $w$ -evolution (obtained from (142) as in §27) into  $\frac{1}{2} \iint m^2 (D_t w) \eta^2$  produces five term families. Three structural lemmas organise them.

**Lemma 30.1** (Exact cancellation of the critical quadratic). *The term  $+\nu w \hat{e}$  in (142) contributes exactly  $+\nu w^2$  to  $\frac{1}{2} D_t w$  (since  $\sum_k \partial_k (w \hat{e}) \cdot \partial_k \hat{e} = w \sum_k |\partial_k \hat{e}|^2 = w^2$ , the  $\partial_k w (\hat{e} \cdot \partial_k \hat{e})$  part vanishing). Hence  $\frac{1}{2} \iint m^2 (D_t w) \eta^2$  contains  $+\nu \iint m^2 w^2 \eta^2$ , which cancels the sink term  $-\nu \iint m^2 w^2 \eta^2$  of the weighted magnitude test identically. The second left-hand term of (149) is therefore removed; no conclusion of the program depended on it (the critical quadratic is handled by Proposition 29.1).*

**Lemma 30.2** (Gram negativity of the drift-curvature terms). *Writing  $G_{jk} = \partial_j \hat{e} \cdot \partial_k \hat{e}$  (positive semidefinite Gram matrix) and using  $\partial_j \partial_k \log m = \frac{\partial_j \partial_k m}{m} - \frac{\partial_j m \partial_k m}{m^2}$ , the drift-curvature contribution splits as*

$$2\nu \iint m^2 (\partial_j \partial_k \log m) G_{jk} \eta^2 = 2\nu \iint m (\partial_j \partial_k m) G_{jk} \eta^2 - 2\nu \iint (\partial_j m \partial_k m) G_{jk} \eta^2.$$

*The second integral is the contraction of the rank-one PSD tensor  $\nabla m \otimes \nabla m$  against  $G \succeq 0$ , hence non-negative; with its minus sign it is a good term. Integrating the first by parts,  $-2\nu \iint \partial_j m \partial_k [m G_{jk} \eta^2]$ , its leading piece is again  $-2\nu \iint (\partial_j m \partial_k m) G_{jk} \eta^2 \leq 0$ ; the remaining pieces are  $\leq \frac{\nu}{8} \iint |\nabla m|^2 w \eta^2 + C\nu M^2 \iint |\nabla^2 \hat{e}|^2 \eta^2$  (Cauchy–Schwarz via  $|\partial_k G_{jk}| \leq 2|\nabla^2 \hat{e}| |\nabla \hat{e}|$ ) plus cutoff terms in the classes below. The most dangerous couplings of C1 are good-signed, not merely bounded.*

**Lemma 30.3** (Stretching and commutator couplings). *The families  $\iint m^2 (|\nabla^2 u| |\nabla \hat{e}| + |\nabla u| w) \eta^2$  close as in Lemma 27.7: the  $|\nabla u| w$  part is  $\leq C \mathcal{L}^2 M^{3/2} / \nu$  by the a priori bound; the  $|\nabla^2 u| |\nabla \hat{e}|$  part, after the near/far Biot–Savart split and Cauchy–Schwarz, is  $\leq C \mathcal{L}^{1/2} (\sigma^*)^{1/2} \nu^{-1/2} M^{3/2}$ . Both are inhomogeneous terms of the allowed class.*

**Proposition 30.4** (Amended  $K$ -self-absorption). *For smooth Type-I solutions, with  $D_{\text{ang}} = \nu \iint |\nabla^2 \hat{e}|^2 \eta^2$ :*

$$\nu \iint |\nabla m|^2 w \eta^2 \leq C M^2 D_{\text{ang}} + C \mathcal{L}^2 M^{3/2} (\nu^{-1} + \nu^{-1/2} (\sigma^*)^{1/2}) + \mathcal{A}_{\text{ann}},$$

*where  $\mathcal{A}_{\text{ann}}$  collects the transition-annulus terms enumerated in Definition 30.5 below. Consequently the coupling  $K$  is absorbed — with an absolute constant against the angular dissipation, no logarithm and no correlation hypothesis — modulo  $\mathcal{A}_{\text{ann}}$ .*

### 30.3 The annulus residual [the single remaining technical item of H1’s estimate layer]

**Definition 30.5** (Annulus residual).  $\mathcal{A}_{\text{ann}}$  consists of the cutoff-derivative terms supported on the transition annulus  $\text{Ann} = \{M/8 \leq |\omega| \leq M/4\} \cap \text{supp } \psi$ :

$$\nu \iint_{\text{Ann}} |\nabla m|^2 w \psi |\chi'|, \quad \nu \iint_{\text{Ann}} m^2 w^2 \psi |\chi'|,$$

and lower-order variants, all weighted by  $\eta$ -incomplete cutoffs (weight  $\psi |\chi'|$ , not  $\eta^2$ ), with constants  $\leq 16$  per occurrence.



**Remark 30.6** (Level-iteration scheme — stated, to be executed).  $\mathcal{A}_{\text{ann}}$  is the classical residue of level-set truncation and is handled by the standard De Giorgi device: replace the single pair of levels  $(\frac{1}{8}, \frac{1}{4})$  by a nested sequence  $\theta_k = \frac{1}{8} + 2^{-k-3}$ ,  $k = 0, 1, 2, \dots$ , with cutoffs  $\chi_k$  at levels  $(\theta_{k+1}, \theta_k)$ ; each annulus term at level  $k$  is dominated by the full-weight quantities at level  $k+1$  with constant  $C4^k$ , and the iteration closes by choosing the level- $k$  Young parameters  $\delta_k = \delta_0 8^{-k}$  (geometric beats geometric). This is bookkeeping of the same class as Ladyzhenskaya–Uraltseva Chapter II; it involves no new estimate. Until written out it is recorded as *open-mechanical*, distinct in kind from the genuinely open items (H2,  $\sigma^*$ -decay, Type-II).

**Superseded (S41).** The scheme sketched in this remark was executed in §34 and *does not close*. The recursion it produces has coefficient  $\geq 2^8$  in front of the level- $(k+1)$  quantity for every choice of  $\delta_k$  (Proposition 34.12); the stated choice  $\delta_k = \delta_0 8^{-k}$  makes the coefficient grow like  $32^k$  rather than  $4^k$ ; and by a co-area identity no admissible level selection can reduce the residual below a level density of the underlying measure (Proposition 34.16). The classification “open-mechanical” in the last sentence above is therefore withdrawn: see Remark 34.24 and Conjecture 34.22.

### 30.4 Routine items (i), (ii), (iv) discharged [Proved]

**Lemma 30.7** (Item (i): transport).  $|\iint w u \cdot \nabla(\eta^2)| \leq \frac{C_I}{r^*} \iint w (2\eta|\nabla\psi|\chi + \frac{16}{M}\eta\psi|\chi'|)|\nabla m| \leq \frac{C C_I}{(\rho' - \rho)r^*} \iint_{Q(\rho')} w \mathbf{1} + \varepsilon \nu \frac{1}{M^2} \iint |\nabla m|^2 w \mathbf{1} + C(\varepsilon, \nu) \frac{C_I^2}{(r^*)^2} \iint w \mathbf{1}$ , by Young; the middle term joins the  $K$ -family (Proposition 30.4), the others the  $(\rho' - \rho)^{-2}$ -family of Proposition 27.8.  $\square$

**Lemma 30.8** (Item (ii): commutator).  $[\partial_k, D_t]\hat{e} = -(\partial_k u \cdot \nabla)\hat{e}$  contributes  $\leq \iint |\nabla u| w \eta^2 \leq C M \mathcal{L} \iint w \eta^2 = C \mathcal{L}(r^*)^{-2} \iint w \eta^2$  — the marginal-potential class already present in Proposition 27.8.  $\square$

**Lemma 30.9** (Item (iv): time slice). With  $\psi$  vanishing on  $\{t \leq t_0 - (\rho')^2\}$  the lower time-slice term is zero and the upper slice enters the left-hand side with positive sign.  $\square$

### 30.5 Consolidated status of H1 after S36

| Component of H1                                 | Status                                      | Where            |
|-------------------------------------------------|---------------------------------------------|------------------|
| First rung (angular Caccioppoli)                | Proved (items (i),(ii),(iv) now discharged) | Prop. 27.8       |
| Critical quadratic                              | Proved given $\sigma^* \leq \varepsilon_0$  | Prop. 29.1       |
| $K$ -absorption                                 | Proved modulo $\mathcal{A}_{\text{ann}}$    | Prop. 30.4       |
| C2 ( $\alpha$ -terms)                           | Verified                                    | this section     |
| Gram negativity                                 | Proved (good signs)                         | Lem. 30.2        |
| $w^2$ cancellation                              | Proved (corrects §29)                       | Lem. 30.1        |
| $\mathcal{A}_{\text{ann}}$ : families (G1)–(G4) | Proved good-signed, discarded (S41)         | Prop. 34.8       |
| $\mathcal{A}_{\text{ann}}$ : families (I1)–(I2) | Proved in the allowed class (S41)           | Prop. 34.8       |
| $\mathcal{A}_{\text{ann}}$ : core (S1)+(S2)     | <b>Open</b> — level iteration <i>fails</i>  | §34, Conj. 34.22 |

**Summary** (amended in S41; see §34). H1 (the Bridge Lemma) rests on: (a)  $\sigma^* \leq \varepsilon_0$  at the working scale; (b) absorption of the annulus residual  $\mathcal{A}_{\text{ann}}$ . Item (b) was classified in

Remark 30.6 as *open-mechanical*, on the expectation that the nested level iteration sketched there would dispose of it. **That expectation was wrong.** §34 executes the scheme and proves that it cannot close: the recursion coefficient is  $\geq 2^8$  for every choice of the Young parameters (Proposition 34.12), the proposed  $\delta_k = \delta_0 8^{-k}$  makes it worse, the unrolled iteration is weaker than trivial (Proposition 34.13), and by a co-area identity *no* level selection whatsoever gains anything (Proposition 34.16). What S41 does establish is that six of the eight cutoff-derivative families vanish by sign or fall in the allowed inhomogeneous class, leaving an explicit core with an unconditional bound that falls short of absorption by exactly one logarithm — the same shortfall as Proposition 28.4. Item (b) is therefore reclassified as *open*, of the same analytic type as Conjecture 28.5, and is restated as Conjecture 34.22. Everything else in H1’s estimate layer is proved. The genuinely open items of the program remain H2,  $\sigma^*$ -decay, and Type-II exclusion, as ledgered in §29.

## 31 Localized Constantin–Fefferman Depletion and the Unification of H2 with $\sigma^*$ -Decay (Session S37)

This section derives H2 (localized depletion at the self-similar scale) as a quantitative consequence of H1 (the Bridge Lemma) via the classical Constantin geometric-kernel representation, and identifies precisely when this derivation succeeds unconditionally versus when it re-invokes the  $\sigma^*$ -decay gap — the central finding of this session.

### 31.1 The geometric kernel and its exact cancellation [Classical]

*Fact 31.1* (Constantin’s stretching formula [13]). For a divergence-free field on  $\mathbb{T}^3$  (locally, treating  $y$ -integration over  $\mathbb{R}^3$  up to box-scale corrections), the stretching rate at  $x^*$  admits the representation

$$\alpha(x^*, t) = \frac{3}{4\pi} \text{p.v.} \int D(\hat{e}(x^*), \hat{e}(x^* - y), y/|y|) \frac{|\omega(x^* - y, t)|}{|y|^3} dy,$$

where  $D(e_1, e_2, \sigma) = (e_1 \cdot (\sigma \times e_2))(e_1 \cdot \sigma)$  is smooth on  $(S^2)^3$ . Since  $\sigma \times e_1 \perp e_1$  for every  $\sigma$ , one has the *pointwise algebraic identity*

$$D(e, e, \sigma) = 0 \quad \text{for every } e, \sigma \in S^2.$$

### 31.2 Near/far split and optimized cutoff [Proved, classical technique]

**Proposition 31.2** (Localized CF estimate). *Suppose  $\hat{e}$  is Lipschitz with constant  $K$  on  $B_\rho(x^*) \cap \{|\omega| \geq M/2\}$  for some  $\rho \leq r^*/2$ , i.e.  $|\hat{e}(x^* - y) - \hat{e}(x^*)| \leq K|y|$  there. Let  $\Omega = \|\omega(\cdot, t)\|_{L^2(\mathbb{T}^3)}^2$ . Then for every  $0 < \rho' \leq \rho$ :*

$$|\alpha(x^*)| \leq C K M \rho' + C \Omega^{1/2} (\rho')^{-3/2},$$

*and optimizing over  $\rho'$  (valid whenever the minimizer  $\rho'_* = C(\Omega^{1/2}/(KM))^{2/5}$  satisfies  $\rho'_* \leq \rho$ ):*

$$|\alpha(x^*)| \leq C (KM)^{3/5} \Omega^{1/5}. \tag{150}$$

*Proof. Near field* ( $|y| < \rho'$ ): by Fact 31.1 and Lipschitz continuity of the second argument of  $D$  (smooth on the compact  $(S^2)^3$ , hence Lipschitz in each slot with a universal constant),

$$|D(\hat{e}(x^*), \hat{e}(x^* - y), \sigma)| = |D(\hat{e}(x^*), \hat{e}(x^* - y), \sigma) - D(\hat{e}(x^*), \hat{e}(x^*), \sigma)| \leq C K |y| \quad (|y| < \rho' \leq \rho).$$

$$\text{Hence } |\alpha_{\text{near}}| \leq CK \int_{|y| < \rho'} |\omega(x^* - y)|/|y|^2 dy \leq CKM \int_{|y| < \rho'} |y|^{-2} dy = CKM \rho'.$$

**Far field** ( $|y| \geq \rho'$ ): by Cauchy–Schwarz,  $\int_{|y| \geq \rho'} |\omega(x^* - y)|/|y|^3 dy \leq (\int_{|y| \geq \rho'} |y|^{-6} dy)^{1/2} \|\omega\|_{L^2} = C(\rho')^{-3/2} \Omega^{1/2}$  (using  $\int_{\rho'}^{\infty} r^{-6} \cdot 4\pi r^2 dr = C(\rho')^{-3}$ ).

**Optimization.**  $f(\rho') = A\rho' + B(\rho')^{-3/2}$  with  $A = CKM$ ,  $B = C\Omega^{1/2}$  is minimized at  $\rho'_* = (3B/2A)^{2/5}$ , giving  $f(\rho'_*) \sim A^{3/5} B^{2/5} = C(KM)^{3/5} \Omega^{1/5}$ .  $\square$

### 31.3 Two regimes: the exponent depends on the growth of $\Omega$ [Central finding]

Feed the sharp Bridge Lemma output  $K \leq C_B(\sigma^*)^{1/2}/r^* = C_B(\sigma^*)^{1/2} M^{1/2}$  into (150):

$$|\alpha(x^*)| \leq C(\sigma^*)^{3/10} M^{9/10} \Omega^{1/5}.$$

**Proposition 31.3** (Regime A: subcritical enstrophy  $\Rightarrow$  unconditional depletion). *If  $\Omega(t) = O(1)$  as  $t \rightarrow T^-$  (enstrophy stays bounded), then*

$$|\alpha(x^*)| \leq C(\sigma^*)^{3/10} M^{9/10},$$

*i.e.  $\theta = 9/10 - 1/2 = 2/5 < 1/2$ : **H2 holds unconditionally given H1** (with the a priori bound  $\sigma^* \leq C\mathcal{L}/\nu$  of Theorem 28.2 supplying a finite, not necessarily small, prefactor).*

**Proposition 31.4** (Regime B: self-similar enstrophy  $\Rightarrow$  borderline). *If  $\Omega(t) \sim M(t)^{1/2}$  (the scaling of a genuine self-similar Type-I profile:  $\omega(x, t) \sim (T - t)^{-1} \Omega_0((x - x^*)/\sqrt{T - t})$  gives  $\|\omega(\cdot, t)\|_{L^2}^2 \sim (T - t)^{-1/2} \sim M(t)^{1/2}$ ), then*

$$|\alpha(x^*)| \leq C(\sigma^*)^{3/10} M^{9/10} \cdot M^{1/10} = C(\sigma^*)^{3/10} M,$$

*which is **exactly borderline** ( $\theta = 1/2$ ): the near/far optimization gains nothing beyond a constant prefactor  $(\sigma^*)^{3/10}$ . Strict depletion in Regime B requires  $\sigma^*(t) \rightarrow 0$ , not merely  $\sigma^*$  bounded.*

**Corollary 31.5** (Unification of H2 with  $\sigma^*$ -decay). *In the generic case where the local enstrophy near a putative Type-I singularity grows at least at the self-similar rate  $M^{1/2}$  (Regime B or worse), **H2's remaining content and the  $\sigma^*$ -decay gap are the same open problem**: both require  $\sigma^*(t) \rightarrow 0$  as  $t \rightarrow T^-$ , not merely the a priori bound  $\sigma^* \leq C\mathcal{L}/\nu$  (Theorem 28.2). The program's genuinely open analytical content therefore reduces from three nominally independent items (H1's annulus step, H2,  $\sigma^*$ -decay) to two: the annulus level-iteration (mechanical, §30) and a single deep gap: a mechanism forcing  $\sigma^*(t) \rightarrow 0$  along any putative Type-I blowup sequence. Regime A remains available as an unconditional fallback if a genuine (non-self-similar, subcritical) enstrophy bound can be established near the singularity — itself an interesting alternative target, since it does not require touching  $\sigma^*$  at all.*

### 31.4 Numerical status

The measured local enstrophy growth exponent relative to  $M(t)$  (whether the data sits closer to Regime A or Regime B) is the natural next diagnostic; see `layer4`, EXP-L4-ENSTROPY-EXPONENT-\* (Session S37). The existing depletion diagnostic (EXP-L4-DEPLETION-TG-001) already shows  $\theta_{\text{eff}}$  centred near  $-0.08$  (median) in the  $M > 10$  window — i.e. an *observed* exponent of  $\alpha$  near 0.42 in  $M$ , below both the naive value (1) and the Regime B borderline value (1), and close to but somewhat above the Regime A prediction (9/10 once  $\Omega$ -growth is folded in would sit between the two regimes) — consistent with the enstrophy growth in this (non-singular, decaying) numerical trajectory being sub-self-similar, as expected away from an actual singularity.

### 31.5 Updated program ledger (post-S37)

| Item                                       | Status                                     | Where      |
|--------------------------------------------|--------------------------------------------|------------|
| Localized CF estimate (near/far optimized) | Proved (classical technique)               | Prop. 31.2 |
| H2 in Regime A (bounded $\Omega$ )         | <b>Unconditional given H1</b>              | Prop. 31.3 |
| H2 in Regime B (self-similar $\Omega$ )    | Borderline; needs $\sigma^* \rightarrow 0$ | Prop. 31.4 |
| H2 $\equiv \sigma^*$ -decay (Regime B)     | <b>Unified:</b> one open problem, not two  | Cor. 31.5  |
| H1 annulus step                            | Open-mechanical                            | §30        |
| $\sigma^*$ -decay (the single deep gap)    | <b>Open</b>                                | Thm. 28.2  |
| Type-II exclusion                          | Open (outside current methods)             | §29        |

## 32 Regime A Is Impossible for Genuine Blowup; Numerical $\sigma^*$ Trend; the Closure Target (Session S38)

### 32.1 Regime A cannot occur: enstrophy blows up at any genuine singularity [Proved]

**Proposition 32.1** (Impossibility of bounded enstrophy at a genuine singularity). *Let  $u_0 \in H^1(\mathbb{T}^3)$  and let  $T < \infty$  be the maximal time of existence of the associated strong (smooth) solution, i.e.  $T$  is a genuine blowup time: the solution cannot be continued as a strong  $H^1$  solution past  $T$ . Then*

$$\limsup_{t \rightarrow T^-} \Omega(t) = \infty, \quad \Omega(t) := \|\omega(\cdot, t)\|_{L^2(\mathbb{T}^3)}^2.$$

Consequently **Regime A** of Proposition 31.3 ( $\Omega = O(1)$ ) is impossible for any genuine singularity: the only route to H2 through Proposition 31.2 is Regime B (or worse), and the unification of §31 (Corollary 31.5) is not merely the generic case — it is the only case.

*Proof.* This is the standard blow-up criterion for the  $H^1$ -Cauchy theory of 3D Navier–Stokes, included for completeness. Local well-posedness in  $H^1(\mathbb{T}^3)$  (Fujita–Kato / standard energy method) gives: for every  $R > 0$  there is  $T_0(R) > 0$ , non-increasing in  $R$ , such that any strong solution with  $\|u(t_1)\|_{H^1} \leq R$  at some time  $t_1$  extends as a strong solution to  $[t_1, t_1 + T_0(R)]$ . Suppose, for contradiction, that  $\limsup_{t \rightarrow T^-} \|u(t)\|_{H^1} = R_0 < \infty$ . Choose  $t_1 < T$  with  $T - t_1 < T_0(2R_0)$  and  $\|u(t_1)\|_{H^1} \leq 2R_0$  (possible since the limsup is finite and  $T_0$  is a fixed

positive number depending only on  $2R_0$ ). Then  $u$  extends strongly to  $[t_1, t_1 + T_0(2R_0)] \supset [t_1, T]$ , and in fact strictly past  $T$ , contradicting maximality of  $T$ . Hence  $\limsup_{t \rightarrow T^-} \|u(t)\|_{H^1} = \infty$ . Since  $\|u(t)\|_{L^2} \leq \|u_0\|_{L^2}$  (energy inequality, non-increasing), the unboundedness must come from  $\|\nabla u(t)\|_{L^2}$ , i.e.  $\limsup_{t \rightarrow T^-} \|\nabla u(t)\|_{L^2} = \infty$ . By the mean-zero Biot–Savart isomorphism on  $\mathbb{T}^3$ ,  $\|\nabla u\|_{L^2} \sim \|\omega\|_{L^2}$  with constants depending only on the domain, giving  $\limsup_{t \rightarrow T^-} \Omega(t) = \infty$ .  $\square$

**Remark 32.2.** This closes off, cleanly and unconditionally, the “local Regime A” direction previously floated as an independent alternative to  $\sigma^*$ -decay: no version of bounded-entropy escape is available at a genuine singularity, local or global, since the argument above uses only  $\|\nabla u\|_{L^2}$  on the whole torus (the domain has no smaller scale to exploit).  $\sigma^*$ -decay is not one of several possible routes to H2 — given H1, it is necessary.

### 32.2 Numerical $\sigma^*$ trend: non-monotonic, not a generic free decay

The instantaneous  $\sigma^*(t)$  was already logged (EXP-L4-BRIDGE-TG-002,  $N = 64$ ,  $\nu = 10^{-3}$ ) across the full stretching-then-decay trajectory. Read against  $M(t)$  (which rises  $2.0 \rightarrow 20.9$  over the growth phase, steps 0–190):  $\sigma^*$  is *not* monotone in  $M$ . It starts small ( $\sigma^* \approx 0.5$  at  $M \approx 2.0$ ), rises sharply through the turbulent transition to a mid-trajectory maximum ( $\sigma^* \approx 44$  at  $M \approx 2.5$ – $3.5$ ), then *partially recedes* as  $M$  climbs toward its true maximum ( $\sigma^* \approx 15$ – $27$  at  $M \approx 15$ – $21$ , steps 100–190 and 190) — markedly below the mid-trajectory peak. Past the peak of  $M$ , as the flow relaxes viscously,  $\sigma^*$  grows again into the hundreds while  $M$  falls, but this late-time regime is suspected to be a definitional artifact of a decaying, noise-dominated field rather than a physically meaningful coherence measurement: as  $M$  falls, the threshold set  $\{|\omega| \geq M/4\}$  becomes less selective and increasingly picks up turbulent debris.

**Remark 32.3** (Honest reading). This single, noisy trajectory does not show  $\sigma^*(t) \rightarrow 0$  as  $M$  grows — values in the growth phase remain  $O(10)$ – $O(50)$ , far from any plausible smallness threshold  $\varepsilon_0$ . It also does not show monotone growth: the partial recession near peak  $M$  is real in this data but should not be over-read (one run,  $N = 64$ , no ensemble averaging). The honest conclusion is that “generic” (non-singular) vortex dynamics does not naturally sit in, or approach, a small- $\sigma^*$  regime. This is consistent with — and does not contradict — the disorder-barrier picture of §26: if blowup requires  $\sigma^*$  small, and generic turbulent dynamics keeps  $\sigma^*$  large, then a genuine singularity (if one exists) would have to be a non-generic, specially aligned configuration, which is exactly the content the Constantin–Fefferman lineage of theorems is built to express. It does not, however, supply evidence that such configurations cannot occur; the numerics measures ordinary dynamics, which by construction cannot sample a genuine singularity.

### 32.3 The closure target: assembling the proved ingredients [Open, precisely stated]

**Remark 32.4** (Not yet a theorem). Propositions 27.8 (first rung), 29.1 (second rung), and 30.4 (amended  $K$ -absorption, modulo the annulus residual of §30) together bound the angular energy at scale  $\rho$  by a combination of (i) the same quantity at a larger scale  $\rho'$ , with coefficient  $C\mathcal{L}/(\rho' - \rho)^2$ , and (ii)  $\sigma^*$ -dependent inhomogeneous terms. Whether this chain, once fully

assembled and non-dimensionalized, yields a *subcritical* scale recursion (effective coefficient  $< 1$ , giving genuine self-improving decay as  $\rho \rightarrow 0$ , independent of the size of  $\sigma^*$  at the starting scale) or remains *critical/supercritical* (coefficient  $\geq 1$ , requiring a priori smallness of  $\sigma^*$  — the same obstruction that defeated §19 and §20) has **not been determined**. A preliminary scaling check on the leading terms of Proposition 30.4 (the  $M^2 D_{\text{ang}}$  term against the first-rung dissipation) shows consistent powers of  $M$ , which is necessary but not sufficient for a favourable closure; the full coefficient bookkeeping, including the annulus-iteration contribution, is deferred to a dedicated session. This is recorded as an open technical target, not a theorem, precisely to avoid repeating the error pattern of §25.

### 32.4 A pointer to Type-I compactness literature

Genuine progress on  $\sigma^*$ -decay near an *actual* singularity (as opposed to generic dynamics, which Proposition 32.1 and the numerics above show is the wrong object to study) most plausibly requires the Type-I blowup-profile compactness framework: rescaled solutions  $u_\lambda(y, s) = \lambda u(x^* + \lambda y, T + \lambda^2 s)$  converge, along a subsequence as  $\lambda \rightarrow 0$ , to a nontrivial ancient (or self-similar) solution under a Type-I bound (Nečas–Růžička–Šverák [15]; Escauriaza–Seregin–Šverák [16]). Such limiting profiles are exactly the “special, non-generic” configurations the disorder-barrier picture requires; whether every such limit profile must have  $\sigma^* = 0$  (perfect coherence) is a natural, sharply-posed question for that framework, distinct from anything the direct estimates of §27–§31 can settle alone. This is noted as the natural longer-range tool, not attempted here.

### 32.5 Updated program ledger (post-S38)

| Item                                                       | Status                                     | Where        |
|------------------------------------------------------------|--------------------------------------------|--------------|
| Regime A (bounded $\Omega$ )                               | <b>Proved impossible</b> at genuine blowup | Prop. 32.1   |
| $\sigma^*$ -decay is necessary (not one of several routes) | Established                                | Prop. 32.1   |
| Numerical $\sigma^*$ trend (generic dynamics)              | Non-monotonic; stays $O(10)$ – $O(50)$     | this section |
| Assembled scale-recursion (sub- vs supercritical)          | <b>Open, unresolved</b>                    | this section |
| Type-I profile compactness route                           | Pointed to, not attempted                  | this section |
| H1 annulus step                                            | Open-mechanical                            | §30          |
| Type-II exclusion                                          | Open (outside current methods)             | §29          |

## 33 Profile Rigidity: Converting $\sigma^*$ -Decay into a Liouville Problem (Session S39)

Corollary 31.5 and Proposition 32.1 together reduce the program’s genuinely open analytical content to a *single* deep item: forcing  $\sigma^*(t) \rightarrow 0$  along a putative Type-I blowup sequence. Every direct dynamical attempt on that item has failed in the same way (§25, Defects 1–3, and the “preliminary scaling check” remark of §32): one derives a scale recursion whose effective coefficient is critical or supercritical, so that contraction requires the very smallness one is trying to produce.

This section develops a different strategy, following the pointer left at the end of §32. Instead of asking the *dynamical* question “does  $\sigma^*$  decay?”, one uses Type-I blowup-profile

compactness to pass to a limit object — an *ancient* solution  $U$  on  $\mathbb{R}^3 \times (-\infty, 0)$  — and asks the *structural* question “what does  $\sigma^*$  do on  $U$ ?”. The value of the reframe is not that it makes the problem easy; it is that the two possible answers ( $\sigma^*(U) = 0$  and  $\sigma^*(U) > 0$ ) are attacked by two *completely different and independently developed* bodies of technique — Liouville rigidity in the first case, infinite-past accounting in the second — neither of which is a scale recursion, and neither of which can therefore fail in the S31 manner.

Every statement below is labelled. Two named inputs, **(R)** and **(N)**, are isolated in §33.2; a third, **(A-up)**, in §33.4. Nothing in this section claims the Prize, and nothing in it removes the Type-I hypothesis.

### 33.1 Conventions and dimensional normalization

**Remark 33.1** (Units). The parabolic cylinder convention  $Q(r) = B_r \times [t_0 - r^2, t_0]$  used throughout §27–§32 is dimensionally consistent only in units where  $\nu = 1$ . *We therefore fix  $\nu = 1$  for the whole of this section* and record, for the dimensional audit, the following homogeneities in physical units ( $[\cdot]$  denoting dimension,  $L$  length,  $\mathcal{T}$  time):

$$[\sigma^*] = 1, \quad [\langle \sigma^* \rangle] = \mathcal{T}L^{-2} = [\nu]^{-1}, \quad [\nu \langle \sigma^* \rangle] = 1,$$

so that Theorem 28.2 in the form  $\nu \langle \sigma^* \rangle \leq C\mathcal{L}$  is the dimensionless statement, and  $\sigma^*$  itself is already dimensionless (as it must be, by Proposition 26.4). Likewise the Type-I constant  $C_I$  in  $\sup |u| \leq C_I(T - t)^{-1/2}$  has  $[C_I] = L\mathcal{T}^{-1/2} = [\nu]^{1/2}$ , so  $c_I := C_I\nu^{-1/2}$  is the dimensionless Type-I constant; all constants below named  $C(c_I)$  are pure numbers. Every displayed inequality in this section has been checked to be homogeneous of a single degree in  $L$  (with  $\mathcal{T} \sim L^2$ ).

**Definition 33.2** (The Type-I ancient class). For  $c_I \geq 1$  let  $\mathcal{A}(c_I)$  denote the set of  $U$  such that:  $U$  is a smooth solution of the Navier–Stokes equations ( $\nu = 1$ ) on  $\mathbb{R}^3 \times (-\infty, 0)$  with  $\operatorname{div} U = 0$ , and

$$\sup_{y \in \mathbb{R}^3} |U(y, s)| \leq c_I |s|^{-1/2}, \quad M_U(s) := \sup_{y \in \mathbb{R}^3} |\omega_U(y, s)| \leq c_I |s|^{-1} \quad (s < 0). \quad (151)$$

$U$  is called *trivial* if  $\omega_U \equiv 0$ , equivalently (given (151) and  $\operatorname{div} U = 0$ ) if  $U(y, s) = b(s)$  for some bounded measurable  $b$ .

**Lemma 33.3** ( $\mathcal{A}(c_I)$  is closed under the scaling group). *If  $U \in \mathcal{A}(c_I)$  and  $\mu > 0$  then  $U_\mu(y, s) := \mu U(\mu y, \mu^2 s) \in \mathcal{A}(c_I)$ , with the same constant  $c_I$ ; moreover  $M_{U_\mu}(s) = \mu^2 M_U(\mu^2 s)$  and, by Proposition 26.4,  $\sigma^*(U_\mu)(s) = \sigma^*(U)(\mu^2 s)$ . [Proved]*

*Proof.*  $|U_\mu(y, s)| = \mu |U(\mu y, \mu^2 s)| \leq \mu c_I (\mu^2 |s|)^{-1/2} = c_I |s|^{-1/2}$ , and identically for the vorticity with  $\omega_{U_\mu}(y, s) = \mu^2 \omega_U(\mu y, \mu^2 s)$ . The scaling of  $M$  and  $\sigma^*$  is Proposition 26.4 read on  $\mathbb{R}^3$ . Smoothness and the equation are preserved because the scaling is the NS symmetry with  $\nu$  fixed.  $\square$

**Remark 33.4** (Why this closure matters). Lemma 33.3 is what makes the class  $\mathcal{A}(c_I)$  a legitimate arena for a *second* compactness argument (Proposition 33.27): one may zoom in on a profile at arbitrarily remote past times without leaving the class or degrading  $c_I$ . This is not available for the original solution  $u$ , whose Type-I bounds degenerate only at the single time  $T$ .

### 33.2 Step 1: profile extraction

Throughout,  $u$  is smooth on  $\mathbb{T}^3 \times [0, T)$  with a genuine first singularity at  $T$  and singular point  $x^*$ , satisfying the Type-I bounds  $M(t) \leq C_I/(T-t)$  and  $\sup |u(\cdot, t)| \leq C_I(T-t)^{-1/2}$  (the standing assumptions of §27). For  $\lambda \in (0, 1]$  define

$$u_\lambda(y, s) := \lambda u(x^* + \lambda y, T + \lambda^2 s), \quad (y, s) \in \Lambda_\lambda \times [-T/\lambda^2, 0), \quad (152)$$

where  $\Lambda_\lambda = \lambda^{-1}\mathbb{T}^3$  exhausts  $\mathbb{R}^3$  as  $\lambda \rightarrow 0$ . Direct computation from the Type-I bounds gives, uniformly in  $\lambda$ ,

$$|u_\lambda(y, s)| \leq c_I |s|^{-1/2}, \quad |\omega_\lambda(y, s)| \leq c_I |s|^{-1} \quad \text{on } \Lambda_\lambda \times [-T/\lambda^2, 0), \quad (153)$$

with  $\omega_\lambda = \lambda^2 \omega(x^* + \lambda \cdot, T + \lambda^2 \cdot)$ . *[Proved: this is the one place where Type-I is used exactly as designed — (153) is  $\lambda$ -independent because the Type-I rates are precisely the scaling rates.]*

The two inputs required to turn (153) into a nontrivial limit are isolated now, and neither is proved here.

- (R) *Scale-uniform interior regularity.* There are constants  $C_k(c_I)$ ,  $k \geq 0$ , such that every  $u_\lambda$  as in (152) satisfies

$$\|\nabla^k u_\lambda\|_{L^\infty(B_R \times [-S_2, -S_1])} \leq C_k(c_I) (1+R) S_1^{-(k+1)/2}$$

for all  $0 < S_1 < S_2$  and  $R \geq 1$  with  $B_R \times [-2S_2, -S_1/2]$  inside the domain of  $u_\lambda$ . *Status: Conditional. This is a scale-invariant form of Serrin’s interior regularity criterion [7] (an  $L^\infty$  velocity bound on a parabolic cylinder controls all interior spatial derivatives, quantitatively), sharpened by the local-pressure theory of Ladyzhenskaya–Seregin [19]. Two caveats are genuine and are the reason (R) is not marked Proved: (i) Serrin’s criterion as classically stated controls spatial derivatives; time regularity, and hence  $C_{\text{loc}}^\infty$  compactness, requires the local pressure decomposition and excludes the spatially-constant “parasitic” solutions only up to an additive  $b(s)$  — exactly the ambiguity that reappears in Theorem 33.17; (ii) the  $\lambda$ -uniformity of the constants must be checked against the fact that  $u_\lambda$  has no uniform global energy bound ( $\int_{\Lambda_\lambda} |u_\lambda|^2 dy = \lambda^{-1} \|u\|_{L^2(\mathbb{T}^3)}^2 \rightarrow \infty$ ), so only the local consequence  $\int_{B_R} |u_\lambda|^2 \leq CR^3 |s|^{-1}$  of (153) is available. Both caveats are standard in the Type-I literature [18] but neither is discharged in this document. **Status amended at S48: Proved.** §36 settles (R) against the literature. Theorem 36.16 proves it in a strictly stronger form (time derivatives included, no factor  $(1+R)$ ), from Serrin’s subcritical interior criterion [7, 24] together with the  $L^\infty$ -based local theory of [17, §4] and [22, Thm. 2.4]. Three corrections to the note above are recorded in Remark 36.20: caveat (ii) is void (no energy, global or local, is used anywhere in that theory); caveat (i) is real but is not a caveat on (R), which asks only for spatial derivatives, and is discharged for the objects this document forms by Theorem 36.9; and the citations to [18] and [19] in this note do not support what they are cited for.*

- (N) *Non-degeneracy at the singular point (no vorticity escape).* There exist  $\varepsilon_* > 0$ , a compact  $\mathcal{K} \subset \mathbb{R}^3 \times (-\infty, 0)$  and a sequence  $\lambda_j \downarrow 0$  with

$$\liminf_{j \rightarrow \infty} \iint_{\mathcal{K}} |\nabla u_{\lambda_j}|^2 dy ds \geq \varepsilon_*,$$



and in addition  $M_U(s) = \lim_j M_{u_{\lambda_j}}(s)$  for a.e.  $s < 0$ , where  $U$  is the limit produced by Proposition 33.6. *Status: Assumed. Discussion in Remark 33.5.*

**Remark 33.5** (What (N) really costs, and why it is not free). The Caffarelli–Kohn–Nirenberg  $\varepsilon$ -regularity theorem [1] supplies exactly the right *scale-invariant* lower bound: because  $x^*$  is a singular point, for every  $r > 0$

$$r^{-1} \iint_{Q(x^*, r)} |\nabla u|^2 dx dt \geq \varepsilon_*, \quad \text{and} \quad r^{-1} \iint_{Q(x^*, r)} |\nabla u|^2 dx dt = \iint_{Q_1(0)} |\nabla u_r|^2 dy ds,$$

the identity being the same computation as Proposition 26.4. So a uniform-in- $\lambda$  lower bound is available on the cylinder  $Q_1(0)$ . The difficulty is that  $Q_1(0)$  *touches the terminal slice*  $s = 0$ , where (153) degenerates and where the compactness of Proposition 33.6 is unavailable; the  $\varepsilon_*$  of mass could in principle live entirely in  $\{|s| < \delta\}$  for every  $\delta$ . Pushing it off the terminal slice requires a Type-I-improved  $\varepsilon$ -regularity statement (of the kind developed by Seregin [18]) and is *not* done here. The second half of (N) — convergence, not merely upper semicontinuity, of the global suprema  $M$  — excludes the vorticity maximum drifting to spatial infinity in the rescaled picture; it is the honest cost of  $\sigma^*$  having been defined with a *global* normalization  $M(t)$ . We return to this in Lemma 33.11.

**Proposition 33.6** (Extraction of an ancient profile). *Assume (R). Then there is a sequence  $\lambda_j \downarrow 0$  and a  $U \in \mathcal{A}(c_I)$  such that*

$$u_{\lambda_j} \longrightarrow U \quad \text{in } C^\infty(\mathcal{K}) \quad \text{for every compact } \mathcal{K} \subset \mathbb{R}^3 \times (-\infty, 0).$$

*If in addition (N) holds (with the same sequence), then  $U$  is nontrivial:  $\iint_{\mathcal{K}} |\nabla U|^2 \geq \varepsilon_*$ , hence  $\omega_U \not\equiv 0$  and  $M_U(s) > 0$  for every  $s < 0$ . [Proved-modulo-(R),(N)]*

*Proof. Compactness.* Fix  $R \geq 1$ ,  $0 < S_1 < S_2$ . By (R) the family  $\{u_\lambda\}$  is bounded in  $C^k(B_R \times [-S_2, -S_1])$  for every  $k$ , uniformly in  $\lambda$  small enough that the cylinder lies in the domain. In particular  $\{\partial_s u_\lambda\}$  is bounded there (the equation  $\partial_s u_\lambda = -(u_\lambda \cdot \nabla) u_\lambda + \Delta u_\lambda - \nabla p_\lambda$  with the local pressure bound supplied by (R)), so Arzelà–Ascoli and a diagonal argument over an exhausting sequence of cylinders and of  $k$  give a subsequence  $u_{\lambda_j} \rightarrow U$  in  $C_{\text{loc}}^\infty(\mathbb{R}^3 \times (-\infty, 0))$ . Locally uniform convergence of all derivatives passes to the limit in the equation, so  $U$  solves NS classically on  $\mathbb{R}^3 \times (-\infty, 0)$ , and  $\text{div } U = 0$ .

*Membership in  $\mathcal{A}(c_I)$ .* The bounds (151) follow from (153) by pointwise passage to the limit at each  $(y, s)$ , then taking suprema; note this yields the bound with the *same*  $c_I$  and in particular  $M_U(s) \leq \liminf_j M_{u_{\lambda_j}}(s)$  (upper semicontinuity of a supremum under locally uniform convergence:  $\sup_{\mathcal{K}} |\omega_U| = \lim_j \sup_{\mathcal{K}} |\omega_{\lambda_j}| \leq \liminf_j M_{u_{\lambda_j}}$  for every compact  $\mathcal{K}$ , then exhaust).

*Nontriviality.*  $C_{\text{loc}}^\infty$  convergence gives  $\iint_{\mathcal{K}} |\nabla U|^2 = \lim_j \iint_{\mathcal{K}} |\nabla u_{\lambda_j}|^2 \geq \varepsilon_*$  by (N). If  $\omega_U(\cdot, s_0) \equiv 0$  for some  $s_0$  then,  $U(\cdot, s_0)$  being bounded on  $\mathbb{R}^3$  and curl-free and divergence-free, each component is a bounded harmonic function, hence constant; the same argument at every later time via forward uniqueness for the (then linear, in fact trivial) system gives  $U(y, s) = b(s)$  on  $[s_0, 0)$  and  $\nabla U \equiv 0$  there. Since the compact  $\mathcal{K}$  may be assumed to lie in  $\{s \geq s_0\}$  after replacing  $s_0$  by  $\inf\{s : (y, s) \in \mathcal{K}\}$ , this contradicts  $\iint_{\mathcal{K}} |\nabla U|^2 \geq \varepsilon_*$ . Hence  $M_U(s) > 0$  for all  $s < 0$ .  $\square$

**Remark 33.7** (Honest statement of the convergence topology). The topology is  $C_{\text{loc}}^\infty$  on  $\mathbb{R}^3 \times (-\infty, 0)$  — *strong*, and strong at derivative level. This is far more than the weak- $*$   $L^\infty$  / weak  $L_{\text{loc}}^2$  convergence one gets from energy bounds alone, and it is bought entirely by **(R)**. It is essential to be explicit about this: *the entire derivative-level transfer problem flagged in Step 2 lives inside (R)*, not in Step 2 itself. If one weakens **(R)** to weak convergence, Lemma 33.11 degrades to Lemma 33.10 and the consequences are catalogued there.

### 33.3 Step 2: transfer audit — what survives the limit

This is the step at which an argument of this shape is most likely to be wrong, so it is treated separately and pessimistically. There are two logically distinct questions, and conflating them is the trap:

- (Q1) Does a given piece of machinery *hold for  $U$* ? Here the honest answer, for everything the program needs, is *yes and for a trivial reason*:  $U$  is a smooth solution satisfying scale-invariant Type-I bounds, so each lemma may be *re-derived on  $U$  from scratch*. Nothing is transferred as a limit of inequalities.
- (Q2) Does a *quantity* converge, i.e. is  $\sigma^*(U)$  related to  $\lim_j \sigma^*(u_{\lambda_j})$ ? Here the answer is a one-directional inequality in general, and the reframe must be designed so that it needs only the available direction.

#### (Q1): the machinery holds on $U$ — and is log-free there

**Proposition 33.8** (Re-derivation on the profile). *Let  $U \in \mathcal{A}(c_I)$  be nontrivial. Then:*

- (a) [Proved] *The magnitude and direction equations (141)–(142) hold for  $U$  pointwise on  $\{\omega_U \neq 0\}$ , and the orthogonal split Lemma 22.1 holds pointwise. Reason: both are algebraic consequences of the vorticity equation for a smooth solution, proved in Lemma 26.1 with no use of the domain, of finiteness of any global norm, or of a finite initial time.*
- (b) [Proved] *Proposition 26.4 (exact scale invariance of  $\sigma^*$ ) holds for  $U$ , and by Lemma 33.3 the induced action on the profile class is  $\sigma^*(U_\mu)(s) = \sigma^*(U)(\mu^2 s)$ .*
- (c) [Proved] *Lemma 30.2 (Gram negativity) holds for  $U$ : it is the pointwise linear-algebra fact that  $\nabla m \otimes \nabla m$  contracted against the positive semidefinite Gram tensor  $G_{jk} = \partial_j \hat{e} \cdot \partial_k \hat{e}$  is non-negative, plus one integration by parts on a compactly supported cutoff. Neither step sees the limit procedure.*
- (d) [Proved-modulo-(R)] *On the profile the logarithmic factor  $\mathcal{L}$  of §27–§31 may be replaced by an absolute constant: for every  $s < 0$ ,*

$$\|\nabla U(\cdot, s)\|_{L^\infty(\mathbb{R}^3)} \leq C_1(c_I) |s|^{-1} \quad (s < 0),$$

*and in particular  $\|\nabla U(\cdot, s)\|_{L^\infty} \leq C(c_I) M_U(s)$  at every time with  $M_U(s) \geq c_I^{-1} |s|^{-1}$  (i.e. whenever the Type-I upper bound is saturated up to a factor).*

(e) [Proved-modulo-(R),(d)] *Theorem 28.2 holds for  $U$  in the log-free form*

$$\langle \sigma_U^* \rangle(s_0) := (r^*)^{-3} \iint_{Q(r^*/2)} |\nabla \hat{e}_U|^2 \mathbf{1}_{\{|\omega_U| \geq M_U/2\}} dy ds \leq C(c_I), \quad r^* = M_U(s_0)^{-1/2}, \quad (154)$$

for every  $s_0 < 0$  with  $Q(r^*/2) \subset \mathbb{R}^3 \times (-\infty, 0)$ .

*Proof.* (a)–(c) are as stated: inspect the proofs of Lemma 26.1, Proposition 26.4 and Lemma 30.2 and observe that each uses only pointwise identities for a smooth divergence-free field and integration by parts against a compactly supported cutoff.

(d) By **(R)** applied to the limit (equivalently, by passing the **(R)** bounds through the  $C_{\text{loc}}^\infty$  convergence),  $\|\nabla U\|_{L^\infty(B_R(y_0) \times [-2|s|, -|s|/2])} \leq C_1(c_I)(1+R)|s|^{-1}$ ; the point is that the exponent  $-1$  and the constant are *independent of  $y_0$* , because (151) is uniform in  $y$ . Taking  $R = 1$  and varying  $y_0$  over  $\mathbb{R}^3$  gives the global-in-space bound. Dimensional check:  $[\nabla U] = \mathcal{T}^{-1} = L^{-2}$  in units  $\nu = 1$ , and  $[|s|^{-1}] = L^{-2}$ . *Why the logarithm disappears.* In Remark 26.7 the factor  $\mathcal{L} = \log(e + \|u\|_{H^3}/M)$  measures the ratio between the resolved scale and the smallest active scale of  $u$ ; a Calderón–Zygmund bound on  $\nabla u$  in  $L^\infty$  must pay it. On a Type-I *profile* that ratio is  $O(1)$  by scale invariance —  $U$  has, by construction, no structure below the scale  $|s|^{1/2}$  that is not already visible at scale  $|s|^{1/2}$  — so the CZ step is replaced by the interior estimate (R), which is logarithm-free. This is a genuine (if modest) gain of the reframe: every marginal  $\mathcal{L}$ -degraded estimate in §27–§30 becomes  $\mathcal{L}$ -free on the profile.

(e) Re-run the proof of Theorem 28.2 verbatim on  $U$ : test the magnitude equation (141) (valid on  $U$  by (a)) with  $m\eta^2$ ,  $\eta$  the composite cutoff of Definition 27.2 built from  $M_U(s)$ ; all cutoffs are compactly supported in  $Q(r^*)$ , so no decay of  $U$  at spatial infinity is needed. The right-hand terms are estimated by Lemma 28.1 with  $\mathcal{L}$  replaced by  $C(c_I)$  using (d) in place of the CZ-with-logarithm bound, and with Lemma 27.6 re-derived on  $U$  (its proof is the vorticity Caccioppoli inequality on a compactly supported cutoff, again local). The conclusion of Theorem 28.2 then reads  $\nu \langle \sigma_U^* \rangle \leq C(c_I)$ , i.e. (154) in units  $\nu = 1$ . Dimensional check in physical units:  $[\nu \langle \sigma^* \rangle] = 1$  by Remark 33.1. Note that (154) is invariant under  $U \mapsto U_\mu$ , consistent with (b) and Lemma 33.3.  $\square$

**Remark 33.9** (Two things that do *not* transfer, and must not be used). (i) **Proposition 31.2**

**(localized Constantin–Fefferman) does not transfer to the profile.** Its far-field step bounds  $\int_{|y| \geq \rho'} |\omega(x^* - y)| |y|^{-3} dy$  by  $C(\rho')^{-3/2} \Omega^{1/2}$  with  $\Omega = \|\omega\|_{L^2}^2$ ; for  $U \in \mathcal{A}(c_I)$  on  $\mathbb{R}^3$  there is *no* reason for  $\|\omega_U(\cdot, s)\|_{L^2(\mathbb{R}^3)}$  to be finite, and the naive substitute  $|\omega_U| \leq M_U$  makes the far-field integral logarithmically divergent at large  $|y|$ . Consequently *the entire H2/Regime-B analysis of §31 is unavailable on the profile as written*, and Step 4 below may not invoke it. This is a real obstruction discovered in this session, not a bookkeeping remark; it is recorded in the ledger.

(ii) **The Bridge Lemma may not be pulled back.** Conjecture 26.8 is an implication with hypothesis  $\sigma^* \leq \varepsilon_0$ . Even if one proved  $\sigma^*(U) = 0$ , the semicontinuity direction of Lemma 33.10 does *not* give  $\sigma^*(u_{\lambda_j}) \rightarrow 0$ , so H1 cannot be applied to the original sequence. Any argument that routes the profile conclusion back through H1 on  $u_{\lambda_j}$  is invalid. The dichotomy of §33.6 is designed to obtain its contradiction *entirely on  $U$* , for exactly this reason.

**(Q2): the  $\sigma^*$  transfer, and its direction**

**Lemma 33.10** (Lower semicontinuity — the generic case). *Suppose  $u_{\lambda_j} \rightarrow U$  only weakly in  $H_{\text{loc}}^1$ , and suppose the normalizations converge:  $M_{u_{\lambda_j}}(s) \rightarrow M_U(s)$  and the base points  $y_j^* \rightarrow y^*$ . Then*

$$\sigma^*(U)(s) \leq \liminf_{j \rightarrow \infty} \sigma^*(u_{\lambda_j})(s). \quad (155)$$

*In general no inequality holds in the opposite direction. [Proved, modulo the level-set null-set technicality noted in the proof]*

*Proof.* On the indicator set  $\{|\omega| \geq M/4\}$  the integrand is  $|\nabla \hat{e}|^2 = (|\nabla \omega|^2 - |\nabla |\omega||^2)/|\omega|^2$  (Lemma 22.1) with denominator bounded below by  $M^2/16 > 0$ , so  $|\nabla \hat{e}|^2 \mathbf{1}$  is a nonnegative quadratic form in  $\nabla \omega$  with locally uniformly continuous coefficients, and the map  $\nabla \omega \mapsto \iint |\nabla \hat{e}|^2 \mathbf{1}$  is convex and weakly lower semicontinuous on  $L_{\text{loc}}^2$ ; (155) follows. (The technicality: the indicator's boundary  $\{|\omega_U| = M_U/4\}$  must have measure zero for the coefficient convergence; this holds for a.e. choice of the threshold constant  $1/4$  by Fubini, and the definition of  $\sigma^*$  is stable under replacing  $1/4$  by any nearby constant, so this is harmless but should be stated.) Failure of the reverse: weak convergence permits oscillation to be lost in the limit, i.e.  $|\nabla \hat{e}_j|^2 \rightharpoonup |\nabla \hat{e}_U|^2 + \mu$  with  $\mu \geq 0$  a defect measure; nothing forces  $\mu = 0$ .  $\square$

**Lemma 33.11** (Equality under (R) and (N)). *Under **(R)** (so that the convergence is  $C_{\text{loc}}^\infty$ ) and the second half of **(N)** ( $M_{u_{\lambda_j}}(s) \rightarrow M_U(s)$ , with base points converging), one has for a.e.  $s < 0$*

$$\sigma^*(U)(s) = \lim_{j \rightarrow \infty} \sigma^*(u_{\lambda_j})(s),$$

*and likewise for  $\langle \sigma^* \rangle$ . [Proved-modulo-(R),(N)]*

*Proof.* Since  $M_U(s) > 0$ , the set  $\{|\omega_U| \geq M_U/4\}$  sits inside  $\{|\omega_U| > 0\}$  where  $\hat{e}_U$  is smooth; on it the integrand of  $\sigma^*$  is bounded,  $|\nabla \hat{e}|^2 \leq 16|\nabla \omega|^2/M^2$ , and  $C_{\text{loc}}^\infty$  convergence gives locally uniform convergence of  $|\nabla \hat{e}_j|^2$  there. The radius  $r_j^* = M_{u_{\lambda_j}}^{-1/2} \rightarrow M_U^{-1/2} = r_U^*$  and the threshold  $M_{u_{\lambda_j}}/4 \rightarrow M_U/4$  by hypothesis, so the domains of integration converge in measure (again up to the null level set of the previous proof). Dominated convergence concludes.  $\square$

**Remark 33.12** (The honest verdict on Step 2). (i) The derivative-level obstruction is **real but relocated**. It does not appear as an unreparable defect in Step 2; it appears as the hypothesis **(R)**. With **(R)** one has equality (Lemma 33.11); without it one has only (155).

(ii) **The available direction is the one the argument needs.** (155) says: *smallness of  $\sigma^*$  along a sequence descends to the limit*. That is precisely the implication used in Proposition 33.27 to absorb the oscillatory middle case into the aligned horn. It is *not* the implication needed to push a limit conclusion back onto the sequence, which is why Remark 33.9(ii) forbids that route.

(iii) **Which horn is hurt.** Neither horn of §33.6 needs the reverse inequality, because the dichotomy is intrinsic to  $U$ :  $\sigma^*(U)$  either vanishes identically on some past interval or it does not, and both cases are attacked on  $U$ . What semicontinuity *kills* is the apparently-attractive hybrid strategy “prove  $\sigma^*(U) = 0$ , conclude  $\sigma^*(u_{\lambda_j}) \leq \varepsilon_0$ , apply

H1+H2 to the original solution”. That strategy is unavailable, full stop. Any future draft that appears to use it should be treated as containing an error.

- (iv) **Where a defect measure would actually bite.** If one runs the argument with only weak convergence, the defect measure  $\mu$  of Lemma 33.10 is concentrated coherence deficit that the limit does not see. The disordered horn (Step 4) would then be attacking the wrong object: the disorder could be entirely in  $\mu$ . This is the precise sense in which Step 4 — not Step 3 — is the horn that depends on **(R)**.

### 33.4 Step 3: the aligned horn — $\sigma^*(U) = 0$ forces triviality

**Lemma 33.13** (Local alignment from a vanishing deficit). *Let  $U \in \mathcal{A}(c_I)$  be nontrivial and let  $s < 0$  satisfy  $\sigma^*(U)(s) = 0$ , with base point  $y^*$  (so  $|\omega_U(y^*, s)| \geq M_U(s)/2$ ) and  $r^* = M_U(s)^{-1/2}$ . Then there is  $e_0 \in S^2$  with*

$$\hat{e}_U(\cdot, s) \equiv e_0 \quad \text{on the connected component } \mathcal{O} \ni y^* \text{ of } \{|\omega_U(\cdot, s)| > M_U(s)/4\} \cap B_{r^*}(y^*).$$

[Proved]

*Proof.*  $\sigma^*(U)(s) = 0$  and  $|\nabla \hat{e}_U|^2 \geq 0$  force  $|\nabla \hat{e}_U|^2 = 0$  a.e. on  $B_{r^*}(y^*) \cap \{|\omega_U| \geq M_U/4\}$ . On the open subset  $\{|\omega_U| > M_U/4\} \cap B_{r^*}(y^*)$  the field  $\hat{e}_U$  is smooth (denominator bounded below), so  $\nabla \hat{e}_U \equiv 0$  there by continuity, and  $\hat{e}_U$  is constant on each connected component.  $y^*$  lies in this open set since  $|\omega_U(y^*, s)| \geq M_U/2 > M_U/4$ .  $\square$

**Remark 33.14** (The gap is spatial globality, and it is large). Lemma 33.13 is genuinely local: one time, one ball of radius  $r^*$ , and only the connected component of the high-vorticity set containing the base point. The Liouville mechanism of Lemma 33.16 below, by contrast, needs alignment on *all* of  $\mathbb{R}^3$ , because it runs a Liouville theorem for bounded harmonic functions on  $\mathbb{R}^3$  and no ball version of that exists. We therefore record the upgrade as a named input:

- (A-up)** *Global alignment upgrade.* If  $U \in \mathcal{A}(c_I)$  and  $\hat{e}_U(\cdot, s_1) \equiv e_0$  on a nonempty open subset of  $\{|\omega_U(\cdot, s_1)| > M_U(s_1)/4\}$  for some  $s_1 < 0$ , then  $\omega_U(y, s) \parallel e_0$  for all  $(y, s) \in \mathbb{R}^3 \times (-\infty, 0)$ . *Status: **Open**, and this is the largest single gap opened by this section. It should not be regarded as a technicality.*

Three candidate mechanisms, none executed here:  $(\alpha)$  *Unique continuation* for the differential inequality satisfied by  $w = |\nabla \hat{e}|^2$ :  $w \geq 0$  vanishes on an open set, and  $w$  satisfies a drift-diffusion inequality (§27, eq. (146)) whose drift is  $U + 2\nabla \log m$ ; strong-maximum / unique-continuation principles for such operators require the drift to lie in a suitable Kato class, which is precisely the  $K$ -family problem of §27–§30. So this route reduces (A-up) to a problem of the *same* kind the program already faces, not to a new one — a mild positive, since the  $K$ -family is now controlled modulo the annulus residual (Proposition 30.4).  $(\beta)$  *Backward uniqueness* in the sense of Escauriaza–Seregin–Šverák [16], applied to the difference between  $\omega_U$  and its projection onto  $e_0$ ; the parabolic operator is right but the required decay at spatial infinity is not available for a profile.  $(\gamma)$  *Scale exhaustion*, which is available only under a strengthened hypothesis and is the one concrete partial result we can state:

**Remark 33.15** (Partial reduction of (A-up) by scale exhaustion). Suppose, more strongly, that  $\sigma^*(U)(s) = 0$  for *every*  $s < 0$ , and that the base points are centred,  $|y^*(s)| \leq C_0|s|^{1/2}$ .

By Lemma 33.3,  $\sigma^*(U_\mu)(-1) = \sigma^*(U)(-\mu^2) = 0$  for every  $\mu > 0$ , so Lemma 33.13 applies at every scale, giving alignment on a ball of radius  $r^*(s) \asymp |s|^{1/2}$  centred within distance  $C_0|s|^{1/2}$  of the origin. As  $s \rightarrow -\infty$  these balls exhaust  $\mathbb{R}^3$ . *Residual gap*: the alignment is only on the connected component of  $\{|\omega_U| > M_U/4\}$  inside each ball, and (i) the direction  $e_0 = e_0(s)$  could a priori depend on  $s$ , (ii) the high-vorticity set need not fill, or even be connected across, the balls. So this reduces (A-up), *under the strengthened hypothesis*, to a statement about the geometry of the high-vorticity set across scales — weaker than (A-up) but still open. The dichotomy of §33.6 does *not* deliver the strengthened hypothesis (it produces vanishing at a single time), so this remark is a pointer, not a proof.

**Lemma 33.16** (Aligned profiles are two-dimensional). *Assume (R). Let  $U \in \mathcal{A}(c_I)$  satisfy  $\omega_U(y, s) \parallel e_0$  for all  $(y, s) \in \mathbb{R}^3 \times (-\infty, 0)$ , with  $e_0 \in S^2$  fixed. Choose coordinates with  $e_0 = e_3$  and write  $\omega_U = \zeta e_3$ . Then*

$$\partial_3 U \equiv 0, \quad U_3 = U_3(s), \quad U_h := (U_1, U_2) = U_h(y_1, y_2, s),$$

and  $U_h$  is a smooth solution of the two-dimensional Navier–Stokes equations on  $\mathbb{R}^2 \times (-\infty, 0)$  with  $\sup_{\mathbb{R}^2} |U_h(\cdot, s)| \leq c_I |s|^{-1/2}$ . [Proved-modulo-(R)]

*Proof. Step 1.*  $0 = \operatorname{div} \omega_U = \partial_3 \zeta$ .

*Step 2.* Put  $f := \partial_3 U$ . Then  $\nabla \times f = \partial_3 \omega_U = (\partial_3 \zeta) e_3 = 0$  and  $\nabla \cdot f = \partial_3(\operatorname{div} U) = 0$ , so  $\Delta f = \nabla(\nabla \cdot f) - \nabla \times (\nabla \times f) = 0$ : each component of  $f(\cdot, s)$  is harmonic on  $\mathbb{R}^3$ .

*Step 3.* By Proposition 33.8(d),  $\|\nabla U(\cdot, s)\|_{L^\infty(\mathbb{R}^3)} \leq C_1(c_I)|s|^{-1} < \infty$ , so  $f(\cdot, s)$  is a bounded harmonic function on  $\mathbb{R}^3$ , hence constant:  $f(\cdot, s) = f(s)$  (Liouville). Therefore  $U(y, s) = f(s) y_3 + \tilde{U}(y_1, y_2, s)$ . Since  $\sup_{y \in \mathbb{R}^3} |U(y, s)| \leq c_I |s|^{-1/2} < \infty$  is finite *uniformly* in  $y_3$ , we must have  $f(s) = 0$ . Thus  $\partial_3 U \equiv 0$ .

*Step 4.* With  $\partial_3 U = 0$ :  $\omega_1 = \partial_2 U_3 - \partial_3 U_2 = \partial_2 U_3$  and  $\omega_2 = \partial_3 U_1 - \partial_1 U_3 = -\partial_1 U_3$ ; both vanish because  $\omega_U \parallel e_3$ . Hence  $\nabla_h U_3 = 0$  and, with  $\partial_3 U_3 = 0$ ,  $U_3 = U_3(s)$ .

*Step 5.* The horizontal momentum equations read  $\partial_s U_h + (U_h \cdot \nabla_h) U_h + U_3 \partial_3 U_h = \Delta_h U_h - \nabla_h p$ , and the  $U_3 \partial_3 U_h$  term vanishes by Step 3. The third equation reads  $\partial_s U_3 = -\partial_3 p$ , so  $\partial_3 p$  is a function of  $s$  alone and  $p = p_h(y_1, y_2, s) - \partial_s U_3(s) y_3$ , whence  $\nabla_h p = \nabla_h p_h$ . Therefore  $(U_h, p_h)$  solves 2D NS. The bound is inherited from (151).  $\square$

**Theorem 33.17** (Liouville theorem for bounded ancient 2D solutions [17]). *Let  $v$  be a bounded ancient mild solution of the two-dimensional Navier–Stokes equations on  $\mathbb{R}^2 \times (-\infty, 0)$ . Then  $v(y, s) = b(s)$  for some bounded measurable  $b : (-\infty, 0) \rightarrow \mathbb{R}^2$ ; in particular  $\nabla v \equiv 0$ . [Cited. Koch–Nadirashvili–Seregin–Šverák, Acta Math. 203 (2009). The 3D analogue for general bounded ancient solutions is open; [17] also treats the 3D axisymmetric-without-swirl case. Caveat on applicability: the theorem is stated for *mild* solutions, so membership of  $U_h$  in that class must be checked — for a smooth solution with  $\|U_h\|_{L^\infty} < \infty$  and locally bounded derivatives this is the standard Duhamel representation modulo the additive  $b(s)$  ambiguity, which is exactly what the conclusion encodes; we treat this as verbatim applicable but flag it rather than assert it.*

**Theorem 33.18** (Aligned horn). *Assume (R) and (A-up). Let  $U \in \mathcal{A}(c_I)$  and suppose  $\sigma^*(U)(s_1) = 0$  for some  $s_1 < 0$  at which  $M_U(s_1) > 0$ . Then  $U$  is trivial:  $\omega_U \equiv 0$  and  $U(y, s) = b(s)$ . [Proved-modulo-(R),(A-up), and modulo the class caveat of Theorem 33.17]*

*Proof.* Lemma 33.13 gives  $\hat{e}_U(\cdot, s_1) \equiv e_0$  on a nonempty open subset of  $\{|\omega_U(\cdot, s_1)| > M_U(s_1)/4\}$ ; **(A-up)** upgrades this to  $\omega_U \parallel e_0$  on  $\mathbb{R}^3 \times (-\infty, 0)$ ; Lemma 33.16 reduces  $U$  to a bounded 2D ancient solution  $U_h$  on  $\mathbb{R}^2 \times (-\infty, -\delta]$  for each  $\delta > 0$  (bounded there by  $c_I \delta^{-1/2}$ ); Theorem 33.17 gives  $U_h(y, s) = b_h(s)$  on  $s \leq -\delta$ , hence on  $s < 0$  since  $\delta > 0$  was arbitrary. Then  $\zeta = \omega_3 = \partial_1 U_2 - \partial_2 U_1 = 0$ , so  $\omega_U \equiv 0$ , and with  $U_3 = U_3(s)$  we get  $U(y, s) = b(s)$ .  $\square$

**Remark 33.19** (Where the reframe’s leverage actually sits, in Step 3). The chain is: vanishing *scalar* coherence deficit  $\Rightarrow$  vorticity parallel to a *fixed* direction  $\Rightarrow$  the flow is two-dimensional  $\Rightarrow$  2D Liouville. The essential point is that the 2D reduction is what makes a Liouville theorem available at all: the general 3D bounded-ancient Liouville problem is open [17], so no version of Step 3 can work without producing a genuine dimensional reduction first. This is why  $\sigma^*$  — a measure of the failure of *directional* coherence — is the right quantity to have built the program around, and it is the strongest structural vindication of §26 obtained so far. Note also that Lemma 33.16 uses the Type-I *spatial* boundedness of the profile twice (Steps 3 and 4) — on the original solution, on  $\mathbb{T}^3$ , the harmonic-Liouville step is unavailable, so this argument genuinely requires the profile.

### 33.5 Step 4: the disordered horn — persistent disorder over an infinite past

The premise now is  $\sigma^*(U)(s) \geq \delta > 0$  for all  $s \leq -S_0$ . The hoped-for mechanism is that persistent geometric disorder costs persistent excess dissipation, which an infinite past cannot pay. The first task is to state precisely why the naive form of that argument *cannot* work.

**Proposition 33.20** (No summable budget exists: the naive infinite-past argument fails structurally). *Let  $U \in \mathcal{A}(c_I)$  and  $s_k = -2^k$ ,  $r_k^* = M_U(s_k)^{-1/2}$ . Then:*

- (a) [Proved] *The dyadic past shells  $(-2^{k+1}, -2^k)$  and the self-similar cylinders  $Q_k = B_{r_k^*/2}(y_k^*) \times [s_k - (r_k^*/2)^2, s_k]$  are commensurate:  $(r_k^*)^2 = M_U(s_k)^{-1} \geq |s_k|/c_I$ , so each dyadic shell carries  $O(1)$  self-similar cylinders, and the infinite past carries infinitely many.*
- (b) [Proved] *Every such cylinder satisfies the same dimensionless bound  $\langle \sigma_U^* \rangle(s_k) \leq C(c_I)$  (Proposition 33.8(e)), with no decay in  $k$ ; and if  $U$  is exactly self-similar ( $U_\mu = U$  for all  $\mu > 0$ ) then  $\langle \sigma_U^* \rangle(s_k)$  is independent of  $k$ , by Lemma 33.3.*
- (c) [Proved] *Consequently no additive scale-invariant dissipation functional can furnish a finite total budget over the infinite past:  $\sum_k \langle \sigma_U^* \rangle(s_k) = +\infty$  is consistent with (indeed forced by) the class  $\mathcal{A}(c_I)$ .*

Interpretation: the trap named in the program brief is not merely a technical difficulty — for scale-invariant quantities it is a theorem that “infinite past + finite budget” is vacuous. Any working infinite-past argument must instead use a quantity that is (i) bounded in a scale-invariant way and (ii) *monotone* in logarithmic time.

*Proof.* (a) is the identity  $r^* = M^{-1/2}$  together with (151). (b) is Proposition 33.8(e) plus Lemma 33.3 applied with  $\mu = 2^{k/2}$ , under which  $Q_k$  is the image of  $Q_0$ . (c) is immediate from (b).  $\square$

Proposition 33.20 dictates the correct framework.

**Lemma 33.21** (Logarithmic time and the profile amplitude). *For  $U \in \mathcal{A}(c_I)$  nontrivial set*

$$\tau := \log |s| \in \mathbb{R} \quad (\tau \rightarrow +\infty \iff s \rightarrow -\infty), \quad \Phi(\tau) := |s| M_U(s) \Big|_{s=-e^\tau}.$$

Then

- (a) [Proved]  $0 < \Phi(\tau) \leq c_I$  for all  $\tau$ ;
- (b) [Proved]  $\Phi$  is scale-covariant by translation only:  $\Phi_{U_\mu}(\tau) = \Phi_U(\tau + 2 \log \mu)$ . Hence  $\Phi$  is a bounded function on a line of infinite length, and the scaling group acts by shifts;
- (c) [Proved]  $M_U$  is locally Lipschitz in  $s$  and, for a.e.  $s$ ,

$$\frac{d}{d\tau} \log \Phi \geq 1 + |s| w_{\max}(s) - |s| \alpha_{\max}(s), \quad (156)$$

where  $w_{\max}(s) = |\nabla \hat{e}_U|^2(y_{\max}(s), s)$ ,  $\alpha_{\max}(s) = \alpha_U(y_{\max}(s), s)$  and  $y_{\max}(s)$  is a point at which  $|\omega_U(\cdot, s)|$  attains its supremum — provided the supremum is attained; see Remark 33.22.

*Proof.* (a) Upper bound: (151). Positivity: nontriviality via Proposition 33.6. (b) By Lemma 33.3,  $|s| M_{U_\mu}(s) = |s| \mu^2 M_U(\mu^2 s) = |\mu^2 s| M_U(\mu^2 s)$ , and  $\log |\mu^2 s| = \tau + 2 \log \mu$ . (c)  $M_U$  is locally Lipschitz because  $\nabla \omega_U$  and  $\partial_s \omega_U$  are locally bounded by **(R)**. At a point of spatial maximum of  $|\omega_U(\cdot, s)|$  one has  $\nabla |\omega_U| = 0$  and  $\Delta |\omega_U| \leq 0$ , so (141) (valid on  $U$  by Proposition 33.8(a); note  $M_U > 0$  so  $y_{\max} \in \{\omega_U \neq 0\}$ ) and Hamilton's trick give, for a.e.  $s$ ,

$$\frac{d}{ds} \log M_U \leq \alpha_{\max} - w_{\max} \quad (\nu = 1).$$

Since  $d/d\tau = -|s| d/ds$  and  $\log \Phi = \tau + \log M_U$ ,

$$\frac{d}{d\tau} \log \Phi = 1 - |s| \frac{d}{ds} \log M_U \geq 1 - |s| (\alpha_{\max} - w_{\max}),$$

which is (156). Dimensional check:  $[|s|\alpha]$  and  $[|s|w]$  are both  $\mathcal{T} \cdot L^{-2} = 1$  in units  $\nu = 1$ ;  $\tau$  and  $\log \Phi$  are dimensionless; every term of (156) is dimensionless. *Note the good sign:* the coherence deficit  $w_{\max}$  enters (156) with a **plus**, i.e. disorder *drives*  $\Phi$  *up*, and  $\Phi$  is bounded — this is the whole mechanism.  $\square$

**Remark 33.22** (Attainment of the supremum). On  $\mathbb{R}^3$  the supremum  $M_U(s)$  need not be attained. Two honest repairs: (i) include attainment in the non-degeneracy package **(N)** (it is implied by “no vorticity escape”, which (N) already asserts in the form  $M_U = \lim_j M_{u_{\lambda_j}}$  together with compactness of the maximizing points); or (ii) run the argument with almost-maximizers, replacing  $\Delta |\omega| \leq 0$  by  $\Delta |\omega| \leq \epsilon$  at an  $\epsilon$ -almost-maximum and letting  $\epsilon \rightarrow 0$  — routine but with a bookkeeping cost we do not pay here. We adopt (i) and record it.

**Proposition 33.23** (Averaged necessary condition over the infinite past). *Let  $U \in \mathcal{A}(c_I)$  be nontrivial with  $M_U$  attaining its suprema. Then for every  $\tau_0$ ,*

$$\limsup_{\tau_1 \rightarrow +\infty} \frac{1}{\tau_1 - \tau_0} \int_{\tau_0}^{\tau_1} \left[ |s| \alpha_{\max} - |s| w_{\max} \right] d\tau \geq 1. \quad (157)$$

[Proved-modulo-(R) and Remark 33.22]



*Proof.* Integrate (156) from  $\tau_0$  to  $\tau_1$ :

$$\log \Phi(\tau_1) - \log \Phi(\tau_0) \geq \int_{\tau_0}^{\tau_1} [1 + |s|w_{\max} - |s|\alpha_{\max}] d\tau.$$

By Lemma 33.21(a) the left side is at most  $\log c_I - \log \Phi(\tau_0) =: C_0 < \infty$ , a constant independent of  $\tau_1$ . Rearranging and dividing by  $\tau_1 - \tau_0$ ,

$$\frac{1}{\tau_1 - \tau_0} \int_{\tau_0}^{\tau_1} [|s|\alpha_{\max} - |s|w_{\max}] d\tau \geq 1 - \frac{C_0}{\tau_1 - \tau_0},$$

and letting  $\tau_1 \rightarrow \infty$  gives (157).  $\square$

**Remark 33.24** (What Proposition 33.23 is and is not). It *is* the correct form of the infinite-past leverage: the infinite past has infinite *logarithmic* length, and a bounded quantity cannot grow at a positive average logarithmic rate over infinite length. It *is not* yet a contradiction: it says only that, on average, the normalized stretching at the vorticity maximum must exceed the normalized coherence sink there by at least 1. Note that this is a statement about *pointwise* values at the maximum, whereas  $\sigma^*$  is a *ball average*; and note that the required margin is exactly 1, so the target below is *marginal*, not subcritical. Marginality is what defeated §19–§20, and honesty requires flagging it. The difference from §20 is nonetheless real: this is a comparison of two averages by a fixed constant, not a scale recursion requiring bootstrapped a priori smallness, so it cannot fail in the specific manner of §25, Defect 1.

*Conjecture 33.25* (Disorder–depletion target for profiles). There is a function  $c : (0, \infty) \rightarrow (0, 1]$  such that: for every  $c_I \geq 1$ , every nontrivial  $U \in \mathcal{A}(c_I)$  whose suprema are attained, and every  $\delta > 0$ , if  $\sigma^*(U)(s) \geq \delta$  for all  $s \leq -S_0$  then for some  $\tau_0$

$$\limsup_{\tau_1 \rightarrow +\infty} \frac{1}{\tau_1 - \tau_0} \int_{\tau_0}^{\tau_1} [|s|\alpha_{\max} - |s|w_{\max}] d\tau \leq 1 - c(\delta).$$

*Status: **Open target**, sharply posed. Together with Proposition 33.23 it is a contradiction, hence it closes the disordered horn.*

**Remark 33.26** (Decomposition of Conjecture 33.25 into its two hard halves). (T4a) *Average-to-point for  $w$  [Open]*. One needs the ball average  $\sigma^* \geq \delta$  to force a log-time-averaged lower bound on  $|s|w_{\max}$ , i.e. on  $w$  at the vorticity maximum. Averages do not control point values in general. The natural mechanism is a weak-Harnack inequality for the parabolic operator governing  $w$ : the  $\sigma^*$ -ball is centred at the vorticity maximum and has radius exactly  $r^*$ , so a Harnack inequality on  $Q(r^*)$  would transfer the average to the centre at a slightly later time with a universal constant. Applicability requires the drift  $U + 2\nabla \log m$  of (146) to lie in a Kato/Morrey class — which is precisely the  $K$ -family controlled, modulo the annulus residual, by Proposition 30.4. So (T4a) plugs into machinery the program already has; it is not a new front.

(T4b) *Depletion of  $|s|\alpha_{\max}$  on profiles [Open, and currently worse than on the original solution]*. One needs  $|s|\alpha_{\max} \leq 1 - c(\delta) + |s|w_{\max}$  on average — a Constantin–Fefferman depletion statement in self-similar variables. By Remark 33.9(i), Proposition 31.2 is *not available on the profile*: its far-field step needs  $\|\omega\|_{L^2(\mathbb{R}^3)} < \infty$ , which fails for  $U \in \mathcal{A}(c_I)$ , and

the crude substitute  $|\omega_U| \leq M_U$  makes the far-field kernel integral logarithmically divergent. A profile version therefore requires a genuinely different far-field control — e.g. exploiting  $\sup |U| \leq c_I |s|^{-1/2}$  through the Biot–Savart *velocity* representation rather than the vorticity one, or a local-energy Type-I bound  $\sup_s \int_{B_R} |U|^2 \leq CR^3 |s|^{-1}$ . Not attempted. *This is the harder of the two halves and is the honest reason Step 4 does not close in this session.*

### 33.6 Step 5: the reframed dichotomy, assembled

**Proposition 33.27** (Two-step reduction: the oscillatory case is aligned). *Let  $U \in \mathcal{A}(c_I)$  be nontrivial with  $\liminf_{s \rightarrow -\infty} \sigma^*(U)(s) = 0$ ; pick  $s_k \rightarrow -\infty$  with  $\sigma^*(U)(s_k) \rightarrow 0$  and set  $\mu_k = |s_k|^{1/2}$ , so that  $U_{\mu_k} \in \mathcal{A}(c_I)$  and  $\sigma^*(U_{\mu_k})(-1) \rightarrow 0$ . Assume **(R)** for the family  $\{U_{\mu_k}\}$  and*

*(N')  $\liminf_k \Phi_U(\log |s_k|) > 0$  and no vorticity escape in the second extraction (i.e. the analogue of **(N)** for  $\{U_{\mu_k}\}$ ).*

*Then there exists a nontrivial  $V \in \mathcal{A}(c_I)$  with  $\sigma^*(V)(-1) = 0$ . [Proved-modulo-(R),(N')]*

*Proof.* Lemma 33.3 places every  $U_{\mu_k}$  in  $\mathcal{A}(c_I)$  with the same constant and gives  $\sigma^*(U_{\mu_k})(-1) = \sigma^*(U)(-\mu_k^2) = \sigma^*(U)(s_k) \rightarrow 0$ . Compactness exactly as in Proposition 33.6 (the hypotheses are identical, the class being scaling-closed) yields  $V \in \mathcal{A}(c_I)$  with  $U_{\mu_k} \rightarrow V$  in  $C_{\text{loc}}^\infty$ . Nontriviality of  $V$ : by **(N')**,  $M_{U_{\mu_k}}(-1) = |s_k| M_U(s_k) = \Phi_U(\log |s_k|) \geq \phi_0 > 0$  along the sequence, and no-escape gives  $M_V(-1) \geq \phi_0 > 0$ . Finally  $\sigma^*(V)(-1) = 0$ : under **(R)** this is Lemma 33.11; even with only weak convergence it follows from the lower semicontinuity (155) of Lemma 33.10, which is the available direction.  $\square$

**Theorem 33.28** (Profile-rigidity dichotomy — conditional). *Assume **(R)**, **(N)**, **(N')**, **(A-up)**, the attainment convention of Remark 33.22, and Conjecture 33.25. Then there is no Type-I singularity at a point  $x^*$  satisfying **(N)**. [Conditional on the five named inputs. Not a theorem of this document in any stronger sense.]*

*Proof.* Suppose there were. Proposition 33.6 produces a nontrivial  $U \in \mathcal{A}(c_I)$ . Exactly one of the following holds.

*Case 1:*  $\liminf_{s \rightarrow -\infty} \sigma^*(U)(s) = 0$ . Proposition 33.27 produces a nontrivial  $V \in \mathcal{A}(c_I)$  with  $\sigma^*(V)(-1) = 0$  and  $M_V(-1) > 0$ ; Theorem 33.18 forces  $V$  trivial — contradiction.

*Case 2:*  $\liminf_{s \rightarrow -\infty} \sigma^*(U)(s) =: 2\delta > 0$ . Then  $\sigma^*(U)(s) \geq \delta$  for all  $s \leq -S_0$ , some  $S_0$ . Conjecture 33.25 and Proposition 33.23 then assert  $1 \leq 1 - c(\delta) < 1$  — contradiction.  $\square$

**Remark 33.29** (What the reframe bought, stated without inflation). The reframe does not prove  $\sigma^*$ -decay. What it does is replace one open item (“force  $\sigma^*(t) \rightarrow 0$  dynamically”, for which every candidate mechanism so far has been a critical or supercritical recursion) by a *different* set of open items, of which the two substantive ones — **(A-up)** and Conjecture 33.25 — are attacked by unrelated techniques (unique continuation / Liouville rigidity; log-time monotonicity). The concrete gains recorded here are:

- (1) *The aligned horn is genuinely closed modulo one hypothesis.* Lemmas 33.13 and 33.16 plus the cited 2D Liouville theorem give a complete argument once **(A-up)** is granted, and the 2D reduction in Lemma 33.16 is a full proof, not a sketch.

- (2) *Profiles are logarithm-free* (Proposition 33.8(d),(e)): every marginal  $\mathcal{L}$ -degraded estimate of §27–§30 improves to an absolute constant on the profile.
- (3) *The naive infinite-past argument is proved impossible* (Proposition 33.20), and the correct framework — bounded amplitude  $\Phi$  monotone in log-time — is identified, with one *proved* averaged necessary condition (157) extracted from it.
- (4) *A previously unnoticed obstruction is recorded* (Remark 33.9(i)): the §31 Constantin–Fefferman machinery does *not* survive the passage to the profile, because  $\|\omega\|_{L^2(\mathbb{R}^3)} = \infty$  there. Any future session that uses Proposition 31.2 on a profile is in error.

### 33.7 Ledger and distance to the goal (post-S39)

| Item                                            | Status                                         | Where             |
|-------------------------------------------------|------------------------------------------------|-------------------|
| Type-I ancient class scaling-closed             | Proved                                         | Lem. 33.3         |
| Rescaled bounds $\lambda$ -uniform              | Proved                                         | (153)             |
| Scale-uniform interior regularity (R)           | <b>Conditional</b> (Serrin-type)               | §33.2             |
| Non-degeneracy / no escape (N)                  | <b>Assumed</b> ; CKN gives it only on $Q_1(0)$ | Rem. 33.5         |
| Profile extraction                              | Proved mod (R),(N)                             | Prop. 33.6        |
| Eqns (141),(142), split, Gram on $U$            | <b>Proved</b> (re-derived, not transferred)    | Prop. 33.8(a)–(c) |
| Profiles are $\mathcal{L}$ -free                | Proved mod (R)                                 | Prop. 33.8(d)     |
| $\nu\langle\sigma^*\rangle \leq C(c_I)$ on $U$  | Proved mod (R)                                 | Prop. 33.8(e)     |
| $\sigma^*$ transfer: lower semicontinuity       | <b>Proved</b> (available direction)            | Lem. 33.10        |
| $\sigma^*$ transfer: equality                   | Proved mod (R),(N)                             | Lem. 33.11        |
| Pull-back of H1 to the sequence                 | <b>Forbidden</b> (no reverse inequality)       | Rem. 33.9(ii)     |
| Prop. 31.2 on profiles                          | <b>Fails</b> ( $\Omega = \infty$ )             | Rem. 33.9(i)      |
| Local alignment from $\sigma^* = 0$             | <b>Proved</b>                                  | Lem. 33.13        |
| Global alignment upgrade (A-up)                 | <b>Open</b> — largest new gap                  | Rem. 33.14        |
| Aligned $\Rightarrow$ 2D ( $\partial_3 U = 0$ ) | <b>Proved</b> mod (R)                          | Lem. 33.16        |
| 2D bounded ancient Liouville                    | Cited [17]; class caveat                       | Thm. 33.17        |
| Aligned horn                                    | Proved mod (R),(A-up)                          | Thm. 33.18        |
| Naive finite-budget argument                    | <b>Proved impossible</b>                       | Prop. 33.20       |
| Log-time amplitude ODE                          | Proved mod (R), attainment                     | Lem. 33.21        |
| Averaged necessary condition                    | <b>Proved</b> mod (R), attainment              | Prop. 33.23       |
| Disorder–depletion target                       | <b>Open target</b> , marginal                  | Conj. 33.25       |
| (T4a) average-to-point for $w$                  | Open; plugs into $K$ -family                   | Rem. 33.26        |
| (T4b) profile depletion of $\alpha$             | Open; needs new far field                      | Rem. 33.26        |
| Two-step reduction                              | Proved mod (R),(N')                            | Prop. 33.27       |
| Reframed dichotomy                              | Conditional on 5 inputs                        | Thm. 33.28        |
| Type-II exclusion                               | Open (outside all present methods)             | Rem. 33.30        |

*Amended at S48.* (R) is no longer conditional: it is proved in Theorem 36.16 from Serrin’s subcritical interior criterion [7, 24] together with the  $L^\infty$ -based local theory of [17, §4] and [22, Thm. 2.4]. Every row above reading “mod (R)” should be read with the (R) deleted — so Prop. 33.8(d) and Lem. 33.16 become **Proved**, Prop. 33.6 and Lem. 33.11 become “Proved mod (N)”, Thm. 33.18 “Proved mod (A-up)”, and Thm. 33.28 conditional on four inputs, not

five — with one exception recorded in §36.5: the row for Prop. 33.8(e) becomes “Proved mod *saturation*”, because (e) uses (d) in the form  $\|\nabla U\|_{L^\infty} \lesssim M_U$ , which (d) supplies only where the Type-I upper bound is saturated. No other row changes; in particular **(N)**, **(N')**, **(A-up)**, Conjecture 33.25 and Type-II exclusion are untouched, and Remark 33.30 stands verbatim.

**Remark 33.30** (Distance to the goal). Suppose both horns closed completely — **(A-up)** proved and Conjecture 33.25 proved, with **(R)**, **(N)**, **(N')** discharged from the Type-I literature. The conclusion would be: *no Type-I singularity*. That is strictly less than the Clay problem. What would remain:

- (1) **Type-II exclusion.** Every estimate in §27–§33 uses the Type-I rates, and the profile-extraction of §33.2 uses them *essentially*: without  $M(t) \lesssim (T-t)^{-1}$  the rescaled family (152) has no uniform bound and there is no limit object at all. A Type-II singularity would rescale to nothing, and the entire reframe would be vacuous against it. This is outside these methods, and no part of this document should be read as progress on it.
- (2) **The nontriviality input.** Even in the Type-I case, **(N)** is not a technicality: it is the assertion that the singularity’s strength does not escape the rescaling window. Discharging it needs a Type-I-improved  $\varepsilon$ -regularity theorem [18]; until then, “no Type-I singularity” should be read as “no Type-I singularity at a non-degenerate point”.
- (3) **Marginality of the surviving target.** Conjecture 33.25 must beat the constant 1 exactly. The program’s history (§25) is that marginal targets are where errors hide; the mitigating structural difference is noted after Proposition 33.23, but it is a mitigation, not a proof.

Accordingly the honest one-line summary of this section is: *the single deep gap of §32 has been replaced by two structurally unrelated gaps, one of which (the aligned horn) is now complete except for a single clearly stated propagation hypothesis, and one of which (the disordered horn) has yielded a proved averaged necessary condition and a sharply-posed marginal target*. No claim is made beyond this.

## 34 Execution of the Level Iteration for the Annulus Residual: an Exact Inventory and a Structural Obstruction (Session S41)

This section carries out the programme announced in Remark 30.6: the nested De Giorgi level iteration that was to remove the annulus residual  $\mathcal{A}_{\text{ann}}$  (Definition 30.5) from Proposition 30.4, the last item standing between H1’s estimate layer and a proof. The outcome is *mixed*, and on the decisive point *negative*; it is reported as such.

- (a) The nested levels and cutoffs are constructed and their derivative bounds computed; the constant  $C4^k$  asserted in Remark 30.6 is confirmed, and its origin identified: it is  $|\chi'_k|^2$ , produced when the second-order term  $\nu\Delta m$  inside  $D_t\chi_k$  is integrated by parts (§34.1).
- (b) An *exact* inventory of the  $\chi$ -derivative families is derived from the  $K$ -identity of Proposition 29.2 (§34.2). This is a genuine positive result: four of the eight families have a

*favourable sign* and may simply be discarded, and two more fall in the inhomogeneous class already allowed by Proposition 30.4.  $\mathcal{A}_{\text{ann}}$  shrinks to an irreducible core of two families (Proposition 34.8).

- (c) The recursion for that core is derived (§34.3). Its coefficient in front of the level- $(k+1)$  quantity is *never smaller than* 1, for *any* choice of the Young parameters  $\delta_k$ ; the proposed  $\delta_k = \delta_0 8^{-k}$  makes it strictly worse (Proposition 34.12). The unrolled iteration is weaker than the trivial inequality and carries no information (Proposition 34.13).
- (d) Neither classical iteration lemma applies: the linear (Ladyzhenskaya–Ural'tseva / Giaquinta–Giusti hole-filling) lemma needs a contraction factor  $< 1$ ; the nonlinear (Stampacchia–De Giorgi) lemma needs a superlinear gain running from coarse levels to fine ones. Here the recursion runs the *opposite* way — the level sets *grow* with  $k$  — and is homogeneous of degree one (Remark 34.14).
- (e) The  $k \rightarrow \infty$  limit does not annihilate the residual: it converges to a *flux through the level set*  $\{|\omega| = M/8\}$ , and a co-area identity shows that *no* admissible choice of levels — nested, geometric, uniform, or optimally selected — reduces the residual below the level density of the underlying measure, whose only a priori bound is a fixed multiple of the band integral (§34.4). This is the crux, and it is not bookkeeping.
- (f) The core residual does admit the unconditional bound  $\mathcal{A}_{\text{ann}}^{\text{core}} \leq C \mathcal{L} \|w\|_{L^\infty} M^{1/2}$  (Proposition 34.19) — which is *exactly* the bound of Proposition 28.4 and therefore falls short of absorption by exactly one logarithm.

Accordingly  $\mathcal{A}_{\text{ann}}$  is reclassified from *open-mechanical* to *open*, of the same analytic type as the  $c_K$ -decorrelation question (Conjecture 28.5); the consolidated table of §30 is amended accordingly. Nothing else in the document is affected: this section touches only the  $\mathcal{A}_{\text{ann}}$  row of H1's estimate layer, and in particular makes no statement about H2,  $\sigma^*$ -decay, Type-II exclusion, or the withdrawn §25 material.

### 34.1 Admissible levels, the nested family, and the derivative bounds

Throughout,  $m = |\omega|$ ,  $w = |\nabla \hat{e}|^2$ ,  $M = M(t) = \|\omega(\cdot, t)\|_{L^\infty}$ ,  $r^* = M^{-1/2}$ , and  $\psi$  is the fixed space–time cutoff of Definition 27.2 ( $\psi \equiv 1$  on  $Q(\rho)$ ,  $\text{supp } \psi \subseteq Q(\rho')$ ,  $\frac{r^*}{2} \leq \rho < \rho' \leq r^*$ ). Only the  $|\omega|$ -level of the composite cutoff is iterated below; the spatial nesting  $\rho < \rho'$  is untouched, and terms carrying  $\nabla \psi$  or  $\partial_t \psi$  remain in the  $(\rho' - \rho)^{-2}$  family of Proposition 27.8 and are not annulus terms.

**Definition 34.1** (Admissible level cutoffs). For  $\frac{1}{16} \leq \theta_- < \theta_+ \leq \frac{1}{4}$  let  $\mathcal{C}(\theta_-, \theta_+)$  be the set of nondecreasing  $\chi \in C^\infty([0, \infty); [0, 1])$  with  $\chi \equiv 0$  on  $[0, \theta_-]$  and  $\chi \equiv 1$  on  $[\theta_+, \infty)$ . For  $\chi \in \mathcal{C}$  write  $\eta_\chi = \psi \cdot \chi(m/M)$ .

**Lemma 34.2** (Uniform admissibility of every level pair). *Let  $\chi \in \mathcal{C}(\theta_-, \theta_+)$  with  $\frac{1}{16} \leq \theta_- < \theta_+ \leq \frac{1}{4}$ . Then*

- (i)  $0 \leq \chi \leq 1$  and  $\chi \equiv 1$  on  $\{m \geq M/4\}$ , hence a fortiori on  $\{m \geq M/2\}$ ;
- (ii)  $\text{supp } \chi(m/M) \subseteq \{m \geq M/16\}$ , so  $M/16 \leq m \leq M$  on  $\text{supp } \eta_\chi$ ;

(iii)  $\chi' \geq 0$ .

These, together with the derivative bound  $\|\chi'\|_\infty \leq 2/(\theta_+ - \theta_-) \leq 32$ , are the properties of the  $\chi$ -factor used in §28–§30 (see the proof). Consequently Proposition 30.4 holds verbatim with  $\eta$  replaced by  $\eta_\chi$ , with constants independent of the pair  $(\theta_-, \theta_+)$  in that range; in particular the inhomogeneous term

$$\mathcal{I} := C \mathcal{L}^2 M^{3/2} (\nu^{-1} + \nu^{-1/2}(\sigma^*)^{1/2}) \quad (158)$$

is level-independent, as is the a priori enstrophy budget

$$\mathcal{E} := \nu \iint_{Q(r^*)} |\nabla \omega|^2 = \nu \iint_{Q(r^*)} (|\nabla m|^2 + m^2 w) \leq C \mathcal{L} M^{1/2}. \quad (159)$$

*Proof.* (i)–(iii) are immediate from Definition 34.1. For the remaining assertions: Lemma 30.1 uses only  $\hat{e} \cdot \partial_k \hat{e} = 0$  and is cutoff-blind; Lemma 30.2 uses only  $\eta \geq 0$  and the Gram positivity; Lemma 30.3 and §30 (C2) use only  $\eta \leq 1$ , the bound  $|\alpha| \leq \|\nabla u\|_{L^\infty} \leq CM\mathcal{L}$  and Theorem 28.2, whose own proof needs only  $\{m \geq M/2\} \subseteq \{\chi = 1\}$ , i.e. (i). Wherever  $m$  occurs in a denominator the level enters through the lower bound of (ii), replacing the numerical factor  $m^{-1} \leq 8/M$  (levels  $\frac{1}{8}, \frac{1}{4}$ ) by  $m^{-1} \leq 16/M$ : a change of absolute constants only. Likewise Lemma 28.1(c) of §28 uses the  $\chi$ -factor only through  $\|\chi'\|_\infty$  and through the pointwise dominations  $m^2 w \leq |\nabla \omega|^2$ ,  $|\nabla m|^2 \leq |\nabla \omega|^2$ , which are level-free. (159) is Lemma 22.1 combined with Lemma 27.6 and  $M^2(r^*)^3 = M^{1/2}$ ; it involves no cutoff at all.  $\square$

**Remark 34.3** (Why the level pair is genuinely free). Lemma 34.2 is what makes any level-selection argument legitimate downstream: a bound proved for *one* admissible  $\chi \in \mathcal{C}(\theta_-, \theta_+)$ ,  $\theta_\pm$  in the stated range, feeds Proposition 27.8 and Theorem 28.2 exactly as the original  $(\frac{1}{8}, \frac{1}{4})$  pair does. This positive observation is used in §34.4; it is also the reason the failure established there is a failure of the *method* and not of a particular normalisation.

**Definition 34.4** (The nested family of Remark 30.6). Put  $\theta_k = \frac{1}{8} + 2^{-k-3}$  for  $k = 0, 1, 2, \dots$ , so  $\theta_0 = \frac{1}{4}$ ,  $\theta_k \downarrow \frac{1}{8}$ , and

$$\ell_k := \theta_k - \theta_{k+1} = 2^{-k-3} - 2^{-k-4} = 2^{-k-4}.$$

Fix once and for all a nondecreasing  $\Theta \in C^\infty(\mathbb{R}; [0, 1])$  with  $\Theta \equiv 0$  on  $(-\infty, 0]$ ,  $\Theta \equiv 1$  on  $[1, \infty)$ ,  $0 \leq \Theta' \leq 2$ ,  $|\Theta''| \leq 16$ , and set

$$\chi_k(s) = \Theta\left(\frac{s - \theta_{k+1}}{\ell_k}\right) \in \mathcal{C}(\theta_{k+1}, \theta_k), \quad \eta_k = \psi \chi_k(m/M),$$

$$\text{Ann}_k = \{\theta_{k+1}M \leq m \leq \theta_k M\} \cap \text{supp } \psi \supseteq \text{supp } [\chi'_k(m/M)] \cap \text{supp } \psi.$$

**Lemma 34.5** (Derivative bounds; the origin of  $4^k$ ). *For every  $k \geq 0$ ,*

$$0 \leq \chi'_k \leq 2/\ell_k = 2^{k+5}, \quad (160)$$

$$|\chi''_k| \leq 16/\ell_k^2 = 2^{2k+12}, \quad (161)$$

$$|\nabla[\chi_k(m/M)]| = \chi'_k |\nabla m|/M \leq 2^{k+5} |\nabla m|/M. \quad (162)$$

Hence the combinations occurring before the integration by parts of Lemma 27.5(b), namely  $\chi_k$  and  $\chi_k \chi'_k$ , obey  $\chi_k \chi'_k \leq 2^5 \cdot 2^k$ , whereas the combinations occurring after it, namely  $(\chi'_k)^2$  and  $\chi_k |\chi''_k|$ , obey  $(\chi'_k)^2 + \chi_k |\chi''_k| \leq 2^{13} \cdot 4^k$ . The constant  $C 4^k$  of Remark 30.6 is therefore correct, and it is the square of the cutoff derivative: it is created by  $\nu \Delta m$  inside  $D_t \chi_k$  when that term is integrated by parts, as it must be (Lemma 27.5(b)).

*Proof.* Chain rule:  $\chi'_k(s) = \Theta'((s - \theta_{k+1})/\ell_k)/\ell_k$  and  $\chi''_k(s) = \Theta''(\cdot)/\ell_k^2$ ; insert  $\ell_k^{-1} = 2^{k+4}$  and  $\ell_k^{-2} = 2^{2k+8}$ , and use  $\|\Theta'\|_\infty \leq 2$ ,  $\|\Theta''\|_\infty \leq 16$ ; then  $2 \cdot 2^{k+4} = 2^{k+5}$  and  $16 \cdot 2^{2k+8} = 2^{2k+12} = 2^{12} \cdot 4^k$ . (162) is  $\nabla[\chi_k(m/M)] = \chi'_k(m/M) \nabla m/M$  ( $M = M(t)$  is  $x$ -independent). Finally  $(\chi'_k)^2 \leq 2^{2k+10}$  and  $\chi_k |\chi''_k| \leq 2^{2k+12}$ , whose sum is  $\leq 2^{2k+13} = 2^{13} 4^k$ .  $\square$

**Lemma 34.6** (The level- $k$  annulus carries full level- $(k+1)$  weight). *For every  $k \geq 0$ :  $\chi_{k+1}(m/M) \equiv 1$  on  $\text{Ann}_k$ , hence  $\eta_{k+1}^2 = \psi^2$  there. Moreover  $\chi_{k+1} \geq \chi_k$  pointwise, the interiors of the  $\text{Ann}_k$  are pairwise disjoint, and  $\bigcup_{k \geq 0} \text{Ann}_k = \{M/8 < m \leq M/4\} \cap \text{supp } \psi =: \text{Ann}$ , the annulus of Definition 30.5 up to the level set  $\{m = M/8\}$ .*

*Proof.*  $\chi_{k+1} \equiv 1$  on  $[\theta_{k+1}, \infty)$  while  $\text{Ann}_k \subseteq \{m/M \geq \theta_{k+1}\}$ . For monotonicity: if  $s \geq \theta_{k+1}$  then  $\chi_{k+1}(s) = 1 \geq \chi_k(s)$ ; if  $s \leq \theta_{k+1}$  then  $\chi_k(s) = 0 \leq \chi_{k+1}(s)$ . Disjointness and the union statement follow from  $\theta_{k+1} < \theta_k$  and  $\theta_k \downarrow \frac{1}{8}$ .  $\square$

## 34.2 Exact inventory of the $\chi$ -derivative families

Definition 30.5 specifies  $\mathcal{A}_{\text{ann}}$  by the *shapes* of its terms, not by a term-by-term derivation. Since the whole question is whether these terms can be removed, we first derive them exactly. The following identity is the one behind Proposition 29.2; it is exact (no inequalities), and we record it in the form in which the cutoff derivatives are visible.

**Lemma 34.7** (The exact weighted magnitude identity). *Let  $\eta = \psi \chi(m/M)$  with  $\chi \in \mathcal{C}$ . Testing (141) against  $m w \eta^2$  and integrating by parts in space and in time (using  $\text{div } u = 0$ , so that  $\iint (D_t f) g = - \iint f D_t g$  up to time slices) gives*

$$\begin{aligned} & \nu \iint |\nabla m|^2 w \eta^2 + \nu \iint m^2 w^2 \eta^2 + [\text{time slices}] \\ &= \underbrace{\iint \alpha m^2 w \eta^2}_{R_1} - \underbrace{\nu \iint m \eta^2 \nabla m \cdot \nabla w}_{R_2} - \underbrace{\nu \iint m w \nabla m \cdot \nabla (\eta^2)}_{R_3} \\ & \quad + \underbrace{\frac{1}{2} \iint m^2 (D_t w) \eta^2}_{R_4} + \underbrace{\frac{1}{2} \iint m^2 w D_t (\eta^2)}_{R_5}. \end{aligned} \tag{163}$$

*Proof.* Multiply (141) by  $m w \eta^2$  and integrate:  $\iint (D_t m) m w \eta^2 = \iint \alpha m^2 w \eta^2 + \nu \iint (\Delta m) m w \eta^2 - \nu \iint m^2 w^2 \eta^2$ . Expand  $\nabla(m w \eta^2) = w \eta^2 \nabla m + m \eta^2 \nabla w + m w \nabla (\eta^2)$  in  $\nu \iint (\Delta m) m w \eta^2 = -\nu \iint \nabla m \cdot \nabla (m w \eta^2)$ , and  $\iint (D_t m) m w \eta^2 = \frac{1}{2} \iint D_t (m^2) w \eta^2 = -\frac{1}{2} \iint m^2 D_t (w \eta^2) + [\text{slices}]$ . Rearranging gives (163). The middle viscous cross-term  $R_2$  is the one absorbed by Cauchy-Schwarz in Proposition 29.2;  $R_1$  is C2 and  $R_4$  is C1 of §30.  $\square$

**Proposition 34.8** (Inventory: four families die by sign, two are inhomogeneous, two survive). *Write  $\chi = \chi(m/M)$ ,  $\chi' = \chi'(m/M)$ ,  $\chi'' = \chi''(m/M)$ , and isolate in (163) all terms carrying*

a  $\chi$ -derivative (all remaining terms carry either  $\eta^2$  in full, or  $\nabla\psi/\partial_t\psi$  and belong to the  $(\rho' - \rho)$ -family). Then, exactly:

$$(G1) \text{ from } R_3: -\frac{2\nu}{M} \iint m w |\nabla m|^2 \psi^2 \chi \chi' \leq 0;$$

$$(G2) \text{ from } R_5 \text{ through the sink } -\nu m w \text{ of (141): } -\frac{\nu}{M} \iint m^3 w^2 \psi^2 \chi \chi' \leq 0;$$

$$(G3) \text{ from } R_5 \text{ through } \nu \Delta m, \text{ after one integration by parts, the } (\chi')^2 \text{ piece: } -\frac{\nu}{M^2} \iint m^2 w |\nabla m|^2 \psi^2 (\chi')^2 \leq 0;$$

$$(G4) \text{ from the same integration by parts, the } \nabla(m^2) \text{ piece: } -\frac{2\nu}{M} \iint m w |\nabla m|^2 \psi^2 \chi \chi' \leq 0.$$

$$(I1) \text{ from } R_5 \text{ through } \alpha m: \frac{1}{M} \iint \alpha m^3 w \psi^2 \chi \chi', \text{ bounded in absolute value by } \|\chi \chi'\|_\infty \|\nabla u\|_{L^\infty} \nu^{-1} \mathcal{E} \leq C \|\chi \chi'\|_\infty \mathcal{L}^2 M^{3/2} \nu^{-1};$$

$$(I2) \text{ from } R_5 \text{ through } -m\dot{M}/M: \text{ the same bound, using } |\dot{M}|/M \leq \|\nabla u\|_{L^\infty} \leq C M \mathcal{L}.$$

$$(S1) \text{ the } \chi \chi'' \text{ piece of the integration by parts in (G3): } -\frac{\nu}{M^2} \iint m^2 w |\nabla m|^2 \psi^2 \chi \chi'', \text{ of indefinite sign, with } |\cdot| \leq \frac{1}{16} \|\chi \chi''\|_\infty \nu \iint_{\{\theta_- M \leq m \leq \theta_+ M\}} |\nabla m|^2 w \psi^2;$$

$$(S2) \text{ the cross family, arising both from } R_5 \text{ (through } \nu \Delta m, \text{ the } \nabla w \text{ piece) and from } R_4 \text{ (the } \nu \Delta \partial_k \hat{e} \text{ piece of the differentiated direction equation, cf. Lemma 27.5(a)): } \frac{c\nu}{M} \iint m^2 \psi^2 \chi \chi' \nabla m \cdot \nabla w, |c| \leq 2, \text{ of indefinite sign.}$$

Consequently, discarding (G1)–(G4) and absorbing (I1)–(I2) into  $\mathcal{I}$  of (158),

$$\mathcal{A}_{\text{ann}} \leq \mathcal{A}_{\text{ann}}^{\text{core}} := \left| (S1) \right| + \left| (S2) \right|. \quad (164)$$

*Proof.* Write  $\nabla(\eta^2) = \chi^2 \nabla(\psi^2) + 2\psi^2 \chi \chi' \nabla m / M$  and  $D_t(\eta^2) = \chi^2 D_t(\psi^2) + 2\psi^2 \chi \chi' (D_t m / M - m \dot{M} / M^2)$ ; the  $\nabla(\psi^2)$  and  $D_t(\psi^2)$  parts are  $(\rho' - \rho)$ -family terms by Definition 27.2, not annulus terms. Substituting the first display into  $R_3$  gives (G1), whose sign is  $\leq 0$  because  $m, w \geq 0$  and  $\chi, \chi' \geq 0$  (Lemma 34.2(iii)). Substituting the second into  $R_5$  and then  $D_t m = \alpha m + \nu \Delta m - \nu m w$  from (141) gives four contributions: the  $-\nu m w$  one is (G2),  $\leq 0$  for the same reason; the  $\alpha m$  and  $-m \dot{M} / M$  ones are (I1)–(I2), estimated by  $|\alpha| \leq \|\nabla u\|_{L^\infty}$ ,  $m/M \leq 1$ ,  $\psi^2 \leq 1$  and  $\nu \iint_{Q(\rho')} m^2 w \leq \mathcal{E}$  from (159), together with  $M \mathcal{L} \cdot \mathcal{L} M^{1/2} = \mathcal{L}^2 M^{3/2}$ ; the  $\nu \Delta m$  one is  $\frac{\nu}{M} \iint m^2 w \psi^2 \chi \chi' \Delta m = -\frac{\nu}{M} \iint \nabla m \cdot \nabla (m^2 w \psi^2 \chi \chi')$  (no boundary terms: the integrand vanishes off  $\text{supp } \psi \cap \{\chi' \neq 0\}$ ), and expanding the gradient by the product rule yields exactly the four pieces (G3), (G4), (S1), (S2) — using  $\nabla[\chi \chi' (m/M)] = ((\chi')^2 + \chi \chi'') \nabla m / M$  — plus one further piece carrying  $\nabla(\psi^2)$ , again a  $(\rho' - \rho)$ -family term. Signs: in (G3) the factor  $(\chi')^2 \geq 0$  enters with an overall minus, in (G4) the factor  $\chi \chi' \geq 0$  likewise. The stated bound in (S1) uses  $m^2 / M^2 \leq \frac{1}{16}$  on  $\text{supp } \chi'$ . For (S2), the  $R_4$  contribution has the same shape by Lemma 27.5(a) with  $\eta^2$  replaced by  $m^2 \eta^2$ , and  $|\nabla w| \leq 2w^{1/2} |\nabla^2 \hat{e}|$  is not yet used.  $\square$



**Remark 34.9** (What is and is not established by Proposition 34.8). Two honest caveats.

- (i) Families (G1)–(G4), (I1)–(I2), (S1) are derived from the *exact* identity (163) and are therefore complete for  $R_1, R_2, R_3, R_5$ . Family  $R_4$  (coupling C1) is *not* displayed term-by-term in §30; what is asserted above for it is only that its  $\chi$ -derivative terms have the shape (S2), by the structural parallel with Lemma 27.5(a). If a full audit of C1 produced  $\chi$ -derivative terms of a shape not listed here, the inventory would have to be extended — but this could only *enlarge*  $\mathcal{A}_{\text{ann}}^{\text{core}}$ , so the negative conclusion of §34.3–§34.4 is unaffected; only the positive statement (164) is conditional on that shape.
- (ii) Definition 30.5 writes the annulus weight as  $\psi|\chi'|$ . The derivation above shows the weight is in fact  $\psi^2$  times a  $\chi$ -derivative factor: the  $\chi$ -derivatives are produced by  $\nabla(\eta^2)$  and  $D_t(\eta^2)$ , both of which retain the full spatial factor  $\psi^2$ . We read Definition 30.5 accordingly. Every estimate below uses only  $\psi \leq 1$ , hence holds in either reading.

### 34.3 The recursion, and why no choice of $\delta_k$ contracts

**Definition 34.10** (Level quantities). With  $\eta_k$  as in Definition 34.4, set

$$Y_k = \nu \iint |\nabla m|^2 w \eta_k^2, \quad D_k = \nu \iint |\nabla^2 \hat{e}|^2 \eta_k^2,$$

and let  $D_b = \nu \iint_{\{m > M/8\}} |\nabla^2 \hat{e}|^2 \psi^2$ , so  $D_k \leq D_b$  for all  $k$  by Lemma 34.6. By that lemma  $\chi_{k+1} \geq \chi_k$ , hence

$$Y_0 \leq Y_1 \leq Y_2 \leq \cdots \uparrow Y_\infty = \nu \iint_{\{m > M/8\}} |\nabla m|^2 w \psi^2. \quad (165)$$

**Proposition 34.11** (The level- $k$  recursion). *For every  $k \geq 0$  and every  $\delta_k > 0$ ,*

$$Y_k \leq G_k + \Lambda_k Y_{k+1}, \quad G_k = C_*(M^2 D_b + 2^k \mathcal{I}) + \delta_k M^2 D_b, \quad \Lambda_k = 2^8 4^k + 2^8 \delta_k^{-1} 4^k. \quad (166)$$

*(The factor  $2^k$  on  $\mathcal{I}$  records that families (I1)–(I2) of Proposition 34.8 carry  $\|\chi_k \chi'_k\|_\infty \leq 2^{k+5}$ ; it only strengthens the negative conclusions below.)*

*Proof.* Proposition 30.4 at level  $k$  (legitimate by Lemma 34.2) gives  $Y_k \leq C_*(M^2 D_k + \mathcal{I}) + \mathcal{A}_k \leq C_*(M^2 D_b + 2^k \mathcal{I}) + \mathcal{A}_k^{\text{core}}$ , where  $\mathcal{A}_k^{\text{core}}$  is (164) formed with  $\chi_k$ . For (S1), by Proposition 34.8, Lemma 34.5 and Lemma 34.6 ( $\eta_{k+1}^2 = \psi^2$  on  $\text{Ann}_k$ , and  $\text{supp } \chi'_k \subseteq \text{Ann}_k$ ),

$$|(S1)_k| \leq \frac{1}{16} 2^{2k+12} \nu \iint_{\text{Ann}_k} |\nabla m|^2 w \psi^2 \leq 2^8 4^k Y_{k+1}.$$

For (S2),  $|c| \leq 2$  and  $|\nabla w| \leq 2w^{1/2} |\nabla^2 \hat{e}|$  give  $|(S2)_k| \leq \frac{4\nu}{M} \iint m^2 \psi^2 \chi_k \chi'_k |\nabla m| w^{1/2} |\nabla^2 \hat{e}|$ ; then Young's inequality  $4XZ \leq \delta_k X^2 + 4\delta_k^{-1} Z^2$  with  $X = m |\nabla^2 \hat{e}| \psi \mathbf{1}_{\text{Ann}_k}$  and  $Z = \frac{m}{M} \chi_k \chi'_k |\nabla m| w^{1/2} \psi$  gives

$$|(S2)_k| \leq \delta_k \nu \iint m^2 |\nabla^2 \hat{e}|^2 \psi^2 \mathbf{1}_{\text{Ann}_k} + 4\delta_k^{-1} \frac{\nu}{M^2} \iint m^2 (\chi_k \chi'_k)^2 |\nabla m|^2 w \psi^2 \leq \delta_k M^2 D_b + 2^8 \delta_k^{-1} 4^k Y_{k+1},$$

using  $m \leq M$ ,  $m^2/M^2 \leq \frac{1}{16}$  and  $(\chi_k \chi'_k)^2 \leq 2^{2k+10}$ , so  $4 \cdot \frac{1}{16} \cdot 2^{2k+10} = 2^8 4^k$ .  $\square$

**Proposition 34.12** (No choice of Young parameters produces a contraction). *In (166), for every sequence  $(\delta_k)$  of positive numbers and every  $k$ ,*

$$\Lambda_k \geq 2^8 4^k \geq 2^8 > 1, \quad \Lambda_k \rightarrow \infty.$$

*In particular:*

- (i) *the choice proposed in Remark 30.6,  $\delta_k = \delta_0 8^{-k}$ , gives  $\Lambda_k = 2^8 4^k + 2^8 \delta_0^{-1} 32^k$ , which grows faster than the unmodified  $4^k$ : the Young device makes the recursion strictly worse, not better;*
- (ii) *no other choice helps, because the term  $2^8 4^k$  — coming from family (S1), which is not a product of a dissipative factor with anything else — is present irrespective of  $\delta_k$ : Young’s inequality has no purchase on it;*
- (iii) *if one additionally demands, as any absorption of the extracted dissipation into  $M^2 D_b$  requires, that  $\sum_k \delta_k \leq \frac{1}{2}$ , then  $\delta_k \leq \frac{1}{2}$  and hence  $2^8 \delta_k^{-1} 4^k \geq 2^9 4^k$ : the two geometric factors act on the same side of the inequality and reinforce each other. The phrase “geometric beats geometric” has no realisation here.*

*Proof.* Immediate from (166):  $\Lambda_k$  is a sum of two nonnegative terms, the first of which is  $2^8 4^k$  and does not involve  $\delta_k$ . For (i),  $8^{-k}$  inverted is  $8^k$  and  $8^k 4^k = 32^k$ . For (iii),  $\sum_k \delta_k \leq \frac{1}{2}$  forces  $\delta_k \leq \frac{1}{2}$ , i.e.  $\delta_k^{-1} \geq 2$ .  $\square$

**Proposition 34.13** (The unrolled iteration is weaker than trivial). *With  $G_k$  and  $\Lambda_k$  as in (166), unrolling (166)  $N$  times gives*

$$Y_0 \leq \sum_{j=0}^{N-1} G_j \prod_{i < j} \Lambda_i + \left( \prod_{i < N} \Lambda_i \right) Y_N, \quad \prod_{i < N} \Lambda_i \geq 2^{8N} 4^{N(N-1)/2}.$$

*Since  $Y_N \geq Y_0$  by (165) and  $\prod_{i < N} \Lambda_i \geq 1$ , the remainder term alone already exceeds  $Y_0$ : the unrolled inequality is implied by the trivial statement  $Y_0 \leq Y_0$  and carries no information, for every  $N$ . Letting  $N \rightarrow \infty$  does not help: both the sum and the remainder coefficient diverge, and  $Y_N \uparrow Y_\infty$ , which is not known to vanish (indeed  $Y_\infty \geq Y_0$ ).*

*Proof.* Induction on  $N$  using (166) and  $\Lambda_i \geq 2^8 4^i$ , so  $\prod_{i < N} \Lambda_i \geq 2^{8N} 4^{0+1+\dots+(N-1)}$ . The monotonicity  $Y_N \geq Y_0$  is (165).  $\square$

**Remark 34.14** (Which iteration lemma would be needed, and why none applies). The task of Remark 30.6 is to decide whether the scheme is a linear telescoping (Ladyzhenskaya–Ural’tseva Ch. II, Giaquinta–Giusti [4, 26]) or needs a nonlinear iteration lemma (Stampacchia–De Giorgi type). Neither is available.

- (i) **Linear (hole-filling) lemma.** If  $f \geq 0$  is bounded on  $[\rho_0, \rho_1]$  and  $f(s) \leq \vartheta f(t) + A(t-s)^{-\gamma} + B$  for all  $\rho_0 \leq s < t \leq \rho_1$  with  $\vartheta < 1$ , then  $f(s) \leq C(\gamma, \vartheta)(A(t-s)^{-\gamma} + B)$ . The hypothesis  $\vartheta < 1$  is what makes the geometric selection of radii work. Here  $\vartheta$  is  $\Lambda_k \geq 2^8$  (Proposition 34.12), so the lemma does not apply; and by Proposition 34.12(ii) no rearrangement of the estimate reduces  $\Lambda_k$  below 1.

- (ii) **Nonlinear (Stampacchia) lemma.** If  $Y_{k+1} \leq C b^k Y_k^{1+\beta}$  with  $b > 1$ ,  $\beta > 0$  and  $Y_0 \leq C^{-1/\beta} b^{-1/\beta^2}$ , then  $Y_k \rightarrow 0$ . Three of its hypotheses fail simultaneously here: (a) the direction — our inequality bounds the *coarse* level by the *fine* one ( $Y_k$  by  $Y_{k+1}$ ), because the truncation levels  $\theta_k$  *decrease*, so the level sets  $\{m > \theta_{k+1} M\}$  *increase* with  $k$ , exactly the opposite of De Giorgi truncation, where superlevel sets shrink to measure zero; (b) homogeneity — (166) is homogeneous of degree one in  $Y$ , with no superlinear gain to beat  $b^k$ ; (c) smallness —  $Y_0$  small is the conclusion sought, not an available hypothesis.
- (iii) **The structural reason.** De Giorgi truncation gains because the sets on which the iterated quantity lives shrink. Here the truncation is in an *auxiliary* variable,  $|\omega|$ , which on the region of interest is confined to the fixed band  $[M/16, M]$ ; no set shrinks, no measure decays, and there is nothing for a nonlinear gain to feed on.

### 34.4 The $k \rightarrow \infty$ limit is a level-set flux; the co-area obstruction

The failure above is not an artefact of the geometric choice  $\theta_k = \frac{1}{8} + 2^{-k-3}$ . The following identifies the obstruction exactly and shows it is common to *all* level selections.

**Definition 34.15** (Level measure). For  $F \geq 0$  measurable with  $\iint_{Q(\rho')} F < \infty$ , let  $\mu_F$  be the push-forward of  $F dx dt$  under  $(x, t) \mapsto m(x, t)/M(t)$ , so that  $\iint F g(m/M) dx dt = \int_0^\infty g d\mu_F$  for every bounded Borel  $g$ . Write  $\mu_F = \varrho_F ds + \mu_F^{\text{sing}}$  for its Lebesgue decomposition.

**Proposition 34.16** (Co-area obstruction: no level selection gains). *Let  $\frac{1}{16} \leq \theta_- < \theta_+ \leq \frac{1}{4}$  and  $F \geq 0$  as above. Then:*

(a) *for every  $\chi \in \mathcal{C}(\theta_-, \theta_+)$ ,*

$$\iint F \chi'(m/M) dx dt = \int_{\theta_-}^{\theta_+} \chi' d\mu_F \geq \operatorname{ess\,inf}_{s \in (\theta_-, \theta_+)} \varrho_F(s),$$

*because  $\chi' \geq 0$  and  $\int_{\theta_-}^{\theta_+} \chi' ds = 1$ ;*

(b) *conversely  $\inf_{\chi \in \mathcal{C}(\theta_-, \theta_+)} \iint F \chi'(m/M) \leq \frac{\mu_F([\theta_-, \theta_+])}{\theta_+ - \theta_-}$ , attained up to  $\epsilon$  by  $\chi' \equiv (\theta_+ - \theta_-)^{-1}$  mollified;*

(c) *hence the residual produced by any admissible cutoff is a level density of  $\mu_F$ , and the only a priori information available about it — namely the total mass  $\mu_F([\theta_-, \theta_+]) \leq \iint_{Q(\rho')} F$  — bounds it from above only by  $(\theta_+ - \theta_-)^{-1} \leq 16$  times that total mass, and bounds it from below not at all;*

(d) *for the nested family of Definition 34.4,  $\chi'_k ds \rightharpoonup \delta_{1/8}$  weakly-\*, so if  $\varrho_F$  is continuous at  $\frac{1}{8}$  and  $\mu_F^{\text{sing}}$  vanishes near  $\frac{1}{8}$  then*

$$\iint F \chi'_k(m/M) \longrightarrow \varrho_F(\tfrac{1}{8}) = \int M(t) \int_{\{m=M(t)/8\}} \frac{F}{|\nabla m|} d\mathcal{H}^2 dt$$

(the last equality by the co-area formula, valid for a.e.  $t$  at which  $M(t)/8$  is a regular value of  $m(\cdot, t)$ ). The limit is a flux through the level set  $\{|\omega| = M/8\}$ ; it is zero only if  $F$  vanishes there.

*Proof.* (a) is the definition of  $\mu_F$  together with  $\int \chi' d\mu_F \geq \int \chi' \varrho_F ds \geq \text{ess inf } \varrho_F \cdot \int \chi' ds$  and  $\int_{\theta_-}^{\theta_+} \chi' = \chi(\theta_+) - \chi(\theta_-) = 1$ . (b) is the same identity for the stated  $\chi$ . (c) restates (a)–(b). (d):  $\chi'_k \geq 0$ ,  $\int \chi'_k = 1$  and  $\text{supp } \chi'_k \subseteq [\theta_{k+1}, \theta_k] \downarrow \{\frac{1}{8}\}$ , which is weak-\* convergence to  $\delta_{1/8}$ ; the co-area identity is the substitution  $s = M(t)\sigma$  in  $\iint Fg(m/M) = \int dt \int_0^\infty g(s/M) \left( \int_{\{m=s\}} F/|\nabla m| d\mathcal{H}^2 \right) ds$ .  $\square$

**Remark 34.17** (Reading of Proposition 34.16). Applied to  $F = |\nabla m|^2 w$  — the integrand of family (S1) — part (d) says that the level iteration does not annihilate the residual; it converts it into the trace  $\int M \int_{\{m=M/8\}} |\nabla m| w d\mathcal{H}^2 dt$ . Controlling that trace requires a bound for  $w$  on a level set of  $|\omega|$ , weighted by  $|\nabla m|$ : precisely the pointwise information that H1 is trying to produce. Part (c) says the same thing statically: the annulus residual is, for every admissible cutoff, a level density of  $\mu_F$ , and the only a priori bound is  $16\times$  the band mass  $\nu \iint_{\text{Ann}} |\nabla m|^2 w \psi^2$  — which is the very quantity  $Y$  restricted to the band where  $\eta^2$  degenerates. Choosing a “good” level by pigeonhole gains nothing either: with  $N$  equally spaced levels the derivative bound is  $|\chi'| \leq 16N$  while the best of  $N$  disjoint annuli carries at most  $1/N$  of the band mass, and the two factors cancel *exactly*. This is the sharp form of the obstruction: the co-area normalisation  $\int |\chi'| ds = 1$  is scale-free, so no redistribution of levels can beat it.

**Remark 34.18** (Why the same terms are harmless at the quadratic level). It is worth recording precisely why Lemma 28.1(c) could dispose of the annulus terms of the *quadratic* magnitude Caccioppoli (148) “with no absorption argument needed”, while the same device fails here. There, the annulus integrands were  $|\nabla m|^2$  and  $m^2 w$ , both dominated pointwise by  $|\nabla \omega|^2$  and hence by the unconditional budget (159). Here the integrands are  $|\nabla m|^2 w$  and  $m^2 w^2$ : each is a budgeted quadratic quantity *times*  $w$ , and  $w$  is exactly the unknown of H1. The annulus residual is thus not a defect of the cutoff, but the quartic nature of the  $K$ -identity making itself felt at the boundary of the level set.

### 34.5 What is nevertheless proved, and the exact closure target

**Proposition 34.19** (Unconditional crude bound for the core residual). *With  $\mathcal{A}_{\text{ann}}^{\text{core}}$  as in (164) formed with any  $\chi \in \mathcal{C}(\theta_-, \theta_+)$ ,  $\frac{1}{16} \leq \theta_- < \theta_+ \leq \frac{1}{4}$ , and any  $\delta > 0$ ,*

$$\mathcal{A}_{\text{ann}}^{\text{core}} \leq \delta M^2 D_b + C(1+\delta^{-1}) \|w\|_{L^\infty(Q(\rho') \cap \{m \geq M/16\})} \mathcal{E} \leq C(1+\delta^{-1}) \mathcal{L} \|w\|_{L^\infty} M^{1/2} + \delta M^2 D_b,$$

*with  $C$  depending only on  $\|\Theta'\|_\infty, \|\Theta''\|_\infty$  and  $\theta_+ - \theta_-$ . Equivalently, in the  $K$ -units of Proposition 28.4 (divide by  $M^2$  and use  $M^{-3/2} = (r^*)^3$ ):*

$$M^{-2} \mathcal{A}_{\text{ann}}^{\text{core}} \leq C(1+\delta^{-1}) \mathcal{L} \|w\|_{L^\infty} (r^*)^3 + \delta D_b.$$

*Proof.* For (S1):  $|\nabla m|^2 \leq |\nabla \omega|^2$  (Lemma 22.1),  $\psi \leq 1$ ,  $w \leq \|w\|_\infty$  on the support, and  $\|\chi \chi''\|_\infty \leq C$ ; apply (159). For (S2): Young as in the proof of Proposition 34.11 with parameter  $\delta$ , then  $|\nabla m|^2 w \leq \|w\|_\infty |\nabla \omega|^2$  in the second factor and (159) again.  $\square$

**Remark 34.20** (The shortfall is exactly one logarithm). Proposition 34.19 reproduces, term for term, the bound of Proposition 28.4. The quantity that Proposition 29.1 is able to absorb is  $\delta \nu (r^*)^3 \|w\|_{L^\infty}$ ; what Proposition 34.19 delivers is  $C\mathcal{L}\|w\|_{L^\infty}(r^*)^3$ . Absorption therefore requires  $C\mathcal{L} \leq \delta \nu$ , which fails for fixed  $\nu$  as  $\mathcal{L} = \log(e + \|u\|_{H^3}/M) \rightarrow \infty$ . So the annulus residual is short of closure by exactly the same single logarithm that Proposition 28.4 identified for  $K$  itself — and, by Remark 34.17, no level scheme recovers it.

**Remark 34.21** (The exact end target, for the record). For the “modulo  $\mathcal{A}_{\text{ann}}$ ” caveat of Proposition 30.4 to be removable, what must be proved is: *there exists one admissible  $\chi \in \mathcal{C}(\theta_-, \theta_+)$  with  $\frac{1}{16} \leq \theta_- < \theta_+ \leq \frac{1}{4}$  and constants  $C$  such that, with  $\eta = \eta_\chi$ ,*

$$\mathcal{A}_{\text{ann}} \leq \frac{1}{2} \nu \iint |\nabla m|^2 w \eta^2 + C M^2 \nu \iint |\nabla^2 \hat{e}|^2 \eta^2 + C \mathcal{L}^2 M^{3/2} (\nu^{-1} + \nu^{-1/2} (\sigma^*)^{1/2}). \quad (167)$$

Two points must be stressed. First, the absorbed first term must carry *the same*  $\eta$  as the left-hand side of Proposition 30.4: an annulus bound in terms of a *fuller* weight (e.g.  $\psi^2$  on the band, or  $\eta_{k+1}^2$ ) does not absorb, and this — not the value of any constant — is what defeats the nested scheme. Second, by Lemma 34.2 it suffices to prove (167) for a *single* admissible level pair, since all such pairs are interchangeable downstream; so the difficulty is genuinely analytic, not a matter of choosing levels cleverly.

*Conjecture 34.22* (Band absorption — the replacement for Remark 30.6). There exist  $\theta_- < \theta_+$  in  $[\frac{1}{16}, \frac{1}{4}]$ , a cutoff  $\chi \in \mathcal{C}(\theta_-, \theta_+)$  and a constant  $C$  such that, with  $\eta = \eta_\chi$  and  $\text{Band} = \{\theta_- M \leq m \leq \theta_+ M\} \cap \text{supp } \psi$ ,

$$\nu \iint_{\text{Band}} |\nabla m|^2 w \psi^2 \leq 2^{-11} \nu \iint |\nabla m|^2 w \eta^2 + C M^2 \nu \iint |\nabla^2 \hat{e}|^2 \eta^2 + C \mathcal{L}^2 M^{3/2} \nu^{-1}, \quad (168)$$

$$M^2 \nu \iint_{\text{Band}} |\nabla^2 \hat{e}|^2 \psi^2 \leq C M^2 \nu \iint |\nabla^2 \hat{e}|^2 \eta^2 + C \mathcal{L}^2 M^{3/2} \nu^{-1}. \quad (169)$$

**Remark 34.23** (Status of Conjecture 34.22). Conjecture 34.22 implies (167). Indeed, for a single level pair of width  $\theta_+ - \theta_- \geq \frac{1}{16}$  one has  $\|\chi\chi'\|_\infty \leq 32$  and  $\|\chi\chi''\|_\infty \leq 2^{12}$ , so Proposition 34.8 and the Young step of Proposition 34.11 with  $\delta = 1$  give  $|(S1)| + |(S2)| \leq 2^9 \nu \iint_{\text{Band}} |\nabla m|^2 w \psi^2 + M^2 \nu \iint_{\text{Band}} |\nabla^2 \hat{e}|^2 \psi^2$ ; inserting (168)–(169) and  $2^9 \cdot 2^{-11} = \frac{1}{4} < \frac{1}{2}$  yields (167). We do not claim the converse. The conjecture asserts that the radial-angular coupling density on the transition band is not concentrated there relative to its value on  $\{m \geq \theta_+ M\}$ . This is a statement of exactly the same analytic type as Conjecture 28.5 (a decorrelation/equipartition hypothesis), restricted to the band, and it is *open*. In particular the claim of Remark 29.4 — that verification of C1–C2 makes Conjecture 28.5 unnecessary — must be qualified: it holds on  $\{m \geq M/4\}$ , where the level cutoff is inactive, and not on the transition band, where a hypothesis of that type is still required.

**Amended (S44).** The phrase “exactly the same analytic type” above is accurate as a statement about the *form* of the two hypotheses and must not be read as an implication in either direction: §35 shows that neither conjecture implies the other, nor any nontrivial part of the other, from the a priori structure they are formulated against (Propositions 35.10 and 35.8; the exact scope of that claim is Remark 35.7). The reason is that the two are

supported on disjoint regions of the  $|\omega|$ -level decomposition (Remark 35.4). Three further amendments. (i) The description “a decorrelation/equipartition hypothesis” applies to (168) but *not* to (169), which is a bare level non-concentration statement of a different type (Remark 35.26). (ii) The absolute constants of (168)–(169) do *not* make this conjecture weaker than Conjecture 28.5: reduced to a hypothesis on the band correlation constant by the route on which the latter was calibrated, (168) requires  $c_K^{\text{band}} \leq C/\mathcal{L}$ , the same decaying form (Proposition 35.15, Remark 35.18). (iii) A single hypothesis covers the  $Y$ -halves of both targets, namely Conjecture 35.20 — the  $c_K$  hypothesis itself, asserted on  $\{m \geq M/16\}$  rather than on  $\{m \geq M/4\}$  (Proposition 35.21); the volume-normalised alternative Conjecture 35.23 also suffices but is strictly stronger in the concentrated regime (Remark 35.25).

### 34.6 Amended status of $\mathcal{A}_{\text{ann}}$

| Sub-item of $\mathcal{A}_{\text{ann}}$   | Status                                                | Where          |
|------------------------------------------|-------------------------------------------------------|----------------|
| Families (G1)–(G4)                       | <b>Proved</b> good-signed; discarded                  | Prop. 34.8     |
| Families (I1)–(I2)                       | <b>Proved</b> in the allowed class $\mathcal{I}$      | Prop. 34.8     |
| Nested cutoffs, $4^k$ constant           | <b>Proved</b> (constant confirmed)                    | Lem. 34.5      |
| Level pairs interchangeable              | <b>Proved</b>                                         | Lem. 34.2      |
| Recursion with $\delta_k$                | <b>Proved not a contraction</b>                       | Prop. 34.12    |
| $\delta_k = \delta_0 8^{-k}$ as proposed | <b>Proved counterproductive</b> ( $32^k$ )            | Prop. 34.12(i) |
| Unrolled iteration                       | <b>Proved vacuous</b>                                 | Prop. 34.13    |
| $k \rightarrow \infty$ limit             | <b>Proved</b> to be a level-set flux                  | Prop. 34.16(d) |
| Any other level selection                | <b>Proved</b> no gain (co-area)                       | Prop. 34.16(c) |
| Core residual (S1)+(S2), crude bound     | <b>Proved</b> ; one logarithm short                   | Prop. 34.19    |
| Core residual, absorption                | <b>Open</b> (was “open-mechanical”)                   | Conj. 34.22    |
| Relation to Conj. 28.5                   | <b>Proved independent</b> (S44); common hyp. isolated | §35, Conj. 35. |

**Remark 34.24** (Honest summary of S41). The level iteration of Remark 30.6 does not close, and the reason is structural rather than computational: the residue of a level-set truncation is, by the co-area normalisation  $\int |\chi'| = 1$ , a level density of the underlying measure, and no redistribution of levels — geometric, uniform, nested, or selected — makes a level density smaller than a fixed multiple of the band mass. What was gained instead is a substantial reduction: six of the eight  $\chi$ -derivative families are removed outright, four by sign and two into the already-allowed inhomogeneous class, leaving a single explicit core (164) with a sharp, unconditional, one-logarithm-short bound and a sharply stated open target (Conjecture 34.22). The correct classification of  $\mathcal{A}_{\text{ann}}$  is therefore *open*, of the same type as Conjecture 28.5, and §30 is amended accordingly. No other item of the program is affected by this section, and in particular nothing here bears on H2, on  $\sigma^*$ -decay, or on Type-II exclusion.

## 35 The Relationship Between the $c_K$ and Band-Absorption Hypotheses (Session S44)

Remark 34.24 classifies  $\mathcal{A}_{\text{ann}}$  as “open, of the same analytic type as Conjecture 28.5”, and Remark 34.23 repeats the phrase; both note that the crude bounds fail by the same single logarithm (Proposition 28.4, Proposition 34.19, Remark 34.20). This section asks whether

the resemblance is substantive, i.e. what the *logical* relationship between Conjecture 28.5 and Conjecture 34.22 actually is. The outcome is mixed, and is reported as such.

- (a) Conjecture 28.5 does *not* imply Conjecture 34.22 (Proposition 35.10); the reason is structural, not a matter of constants: the two hypotheses are supported on *disjoint* regions of the  $|\omega|$ -level decomposition (Remark 35.4). The band-transported form of the  $c_K$  hypothesis does imply the  $Y$ -half (168): through the  $\|w\|_{L^\infty}$  route with no extra ingredient (Proposition 35.15), or through the budget route with the volume hypothesis (NB) named explicitly (Proposition 35.12).
- (b) Conjecture 34.22 does not imply Conjecture 28.5, nor any upper bound on  $c_K$  whatsoever (Proposition 35.8).
- (c) There are common hypotheses from which the  $Y$ -halves of both targets follow, one for each of the two available routes (Remark 35.14). The weaker and preferred one is Conjecture 35.20: *the  $c_K$  hypothesis itself, asserted on  $\{m \geq M/16\}$  instead of on  $\{m \geq M/4\}$*  (Proposition 35.21). The alternative, Conjecture 35.23, replaces the decaying constant by an  $O(1)$  one at the cost of normalising by the ambient parabolic volume, and is a demanding hypothesis rather than a cheap one (Remark 35.25). The angular half (169) of Conjecture 34.22 is of a different type and is covered by neither (Remark 35.26).
- (d) The strength comparison suggested by the “ $C/\mathcal{L}$  versus  $O(1)$ ” discrepancy does *not* survive: (168) is  $O(1)$  in its bookkeeping only. Reduced to a hypothesis on a correlation constant by the route on which Conjecture 28.5 was calibrated, it requires  $c_K^{\text{band}} \leq C/\mathcal{L}$  — the same decaying form (Proposition 35.15). The volume-normalised alternative buys an  $O(1)$  constant only at the price of a volume hypothesis, symmetrically on both sides ((174) and (175)), and is *strictly stronger* than the corresponding correlation bound (Lemma 35.2). See Proposition 35.17, Remark 35.18 and Remark 35.19. The first draft of this section concluded the opposite from an inverted lemma; that conclusion is withdrawn and the correction is recorded in Remark 35.3.

This section changes no proved statement. It rearranges open hypotheses; its only edits elsewhere are a cross-reference appended to Remark 34.23 and one row added to the table of §34.6. Nothing here bears on H2, on  $\sigma^*$ -decay, on the hypotheses of §33, on Type-II exclusion, or on the withdrawn §25 material.

### 35.1 Normalisation; the two hypotheses side by side

Throughout, the notation is that of §34:  $m = |\omega|$ ,  $w = |\nabla \hat{e}|^2$ ,  $M = M(t)$ ,  $r^* = M^{-1/2}$ ,  $\psi$  the fixed space–time cutoff of Definition 27.2 with  $\frac{r^*}{2} \leq \rho < \rho' \leq r^*$ ,  $\mathcal{E}$  the enstrophy budget (159), and  $\mathcal{I}$  the allowed inhomogeneous class (158). We abbreviate its dominant constituent

$$\mathcal{I}_0 := \mathcal{L}^2 M^{3/2} \nu^{-1} \quad (\text{so } \mathcal{I} \geq C^{-1} \mathcal{I}_0 \text{ and } \mathcal{I}_0 \text{ is exactly the last term of (168)–(169)}).$$

Constants  $C$  depend only on the standing Type-I data  $(C_I, \nu)$  and on  $\|\Theta'\|_\infty, \|\Theta''\|_\infty$ , never on  $M$  or  $\mathcal{L}$ . We write  $|Q(\rho)| = c_5 \rho^5$  for the parabolic volume ( $c_5$  absolute) and

$$Q^\# := Q(\rho') \cap \{m \geq M/16\}, \quad E = Q(r^*) \cap \{m \geq M/4\}, \quad \text{Band} = \{\theta_- M \leq m \leq \theta_+ M\} \cap \text{supp } \psi,$$

with  $\frac{1}{16} \leq \theta_- < \theta_+ \leq \frac{1}{4}$  admissible in the sense of Definition 34.1.

**Definition 35.1** (Volume-normalised coupling excess). For a measurable weight  $\varpi$  with  $0 \leq \varpi \leq 1$  supported in  $Q(r^*)$ , and provided both factors in the denominator are positive and finite, set

$$\Xi[\varpi] := \frac{|Q(r^*)| \iint |\nabla m|^2 w \varpi}{\left( \iint |\nabla m|^2 \varpi \right) \left( \iint w \varpi \right)}.$$

Thus  $\Xi[\varpi] \leq \kappa_0$  says: *the coupling density  $|\nabla m|^2 w$  is at most  $\kappa_0$  times what it would be if  $|\nabla m|^2$  and  $w$  were uncorrelated and spread over the whole cylinder  $Q(r^*)$ .* It is the same ratio as  $c_K$  except that the normalising volume is the ambient  $|Q(r^*)|$  and not the measure of the region carrying  $\varpi$ . If either denominator vanishes then  $|\nabla m|^2 w \varpi = 0$  a.e., so  $\Xi[\varpi]$  is left undefined and every statement below in which it occurs holds trivially; we do not repeat this convention.

**Lemma 35.2** ( $\Xi$  versus  $c_K$ : the volume ratio). *With  $c_K$  as in Conjecture 28.5,*

$$\Xi[\mathbf{1}_E] = c_K \frac{|Q(r^*)|}{|E|} \geq c_K, \quad (170)$$

*and more generally, for any measurable  $R \subseteq Q(r^*)$  of positive measure and any weight  $\varpi$  supported in  $R$ ,*

$$\Xi[\varpi] = c_K[\varpi] \frac{|Q(r^*)|}{|R|}, \quad c_K[\varpi] := \frac{|R| \iint |\nabla m|^2 w \varpi}{\left( \iint |\nabla m|^2 \varpi \right) \left( \iint w \varpi \right)},$$

*$c_K[\mathbf{1}_E]$  being exactly  $c_K$  and  $c_K[\mathbf{1}_{\text{Band}\psi^2}]$  exactly the  $c_K^{\text{band}}$  of (173). Consequently:*

- (i) *a bound  $\Xi[\varpi] \leq \kappa$  is strictly stronger than the correlation bound  $c_K[\varpi] \leq \kappa$ , by exactly the volume ratio  $|Q(r^*)|/|R|$ ;*
- (ii) *Conjecture 28.5 gives only  $\Xi[\mathbf{1}_E] \leq (C/\mathcal{L}) |Q(r^*)|/|E|$ , which is  $O(1)$  only under a volume lower bound on  $E$ ; and*
- (iii)  *$\Xi[\varpi] \geq 1$  whenever  $|\nabla m|^2$  and  $w$  are supported in a common set of measure  $\leq |Q(r^*)|$  and are uncorrelated on it — for  $|\nabla m|^2 \equiv G$ ,  $w \equiv W$  on  $R$  and both vanishing off  $R$  one has  $c_K[\varpi] = 1$  but  $\Xi[\varpi] = |Q(r^*)|/|R|$ .*

*Proof.* Writing out the two averages in Conjecture 28.5 gives  $c_K = |E| \iint_E |\nabla m|^2 w / (\iint_E |\nabla m|^2 \cdot \iint_E w)$ , and Definition 35.1 with  $\varpi = \mathbf{1}_E$  gives the same expression with  $|E|$  replaced by  $|Q(r^*)|$ ; dividing yields (170). The general case is identical, and (i)–(iii) follow from  $|R| \leq |Q(r^*)|$  and direct substitution.  $\square$

**Remark 35.3** (Corrigendum: this lemma was stated inverted in the first draft of S44). In the first draft of this section Lemma 35.2 was recorded as  $\Xi[\mathbf{1}_E] = c_K |E|/|Q(r^*)| \leq c_K$  — the reciprocal of the correct volume factor, hence the wrong direction. The error was caught in the independent check of this section and is recorded here, in the style of the corrigendum of §28 and the retraction of §29, because it *reverses* the interpretation of the volume-normalised



statistic  $\Xi$  and therefore the answer to the strength question of §35.5. What the first draft concluded — that  $\Xi$ -boundedness is weaker than  $c_K$ -boundedness, that Conjecture 28.5 may be weakened to an  $O(1)$  statement without loss, and that the measured  $c_K \approx 1$  is direct evidence for  $\Xi = O(1)$  — is **withdrawn**; the corrected statements are Proposition 35.17(iii), Remark 35.18 and Remark 35.19 below. Nothing else in this section used the lemma: the two non-implications (Propositions 35.10 and 35.8), Lemma 35.5, Proposition 35.12, Proposition 35.17(i)–(ii) and Proposition 35.24 are all independent of it and were re-checked line by line. Two internal consistency checks confirm the corrected direction: Remark 35.25 computes  $\Xi = |Q(r^*)|/|S|$  under co-concentration on a set  $S$ , and Proposition 35.12 uses  $\Xi[\mathbf{1}_{\text{Band}}\psi^2] = c_K^{\text{band}}|Q(r^*)|/|\text{Band}|$ ; both agree with (170) and contradicted the first-draft lemma.

**Remark 35.4** (The two hypotheses live on disjoint regions). Conjecture 28.5 is a statement about  $E = Q(r^*) \cap \{m \geq M/4\}$ . Conjecture 34.22 is a statement about  $\text{Band} \subseteq \{m \leq M/4\}$ . The two regions meet only in the level set  $\{m = M/4\}$ , which carries no five-dimensional measure whenever  $M(t)/4$  is a regular value of  $m(\cdot, t)$  for a.e.  $t$ . Neither hypothesis constrains any integral supported in the other’s region. This already makes it implausible that either implies the other, and §35.3 confirms it. It also locates precisely what Remark 34.23 was right to flag: “the same analytic type” is a statement about the *form* of the two hypotheses, not about their content, and the qualification that remark attaches to Remark 29.4 is exactly the disjointness recorded here.

**Lemma 35.5** (Budget lemma). *Let  $0 \leq \varpi \leq 1$  be supported in  $Q^\sharp$ . Then*

$$\nu \iint |\nabla m|^2 \varpi \leq \mathcal{E}, \quad \iint w \varpi \leq \frac{256 \mathcal{E}}{\nu M^2},$$

*and consequently, whenever  $\Xi[\varpi]$  is defined,*

$$\nu \iint |\nabla m|^2 w \varpi \leq \Xi[\varpi] \frac{256 \mathcal{E}^2}{\nu M^2 |Q(r^*)|} \leq C \Xi[\varpi] \mathcal{I}_0. \quad (171)$$

*Proof.* The first bound is  $|\nabla m|^2 \leq |\nabla m|^2 + m^2 w$  together with (159) and  $\varpi \leq 1$ . The second is the same estimate after using  $m \geq M/16$  on  $\text{supp } \varpi$ , so that  $m^2 w \geq M^2 w/256$ . Multiplying the two bounds, dividing by  $|Q(r^*)|$  and using Definition 35.1 gives the first inequality of (171). For the second, insert  $|Q(r^*)| = c_5(r^*)^5 = c_5 M^{-5/2}$  and  $\mathcal{E} \leq C_\mathcal{E} \mathcal{L} M^{1/2}$ :

$$\frac{\mathcal{E}^2}{M^2 |Q(r^*)|} \leq \frac{C_\mathcal{E}^2 \mathcal{L}^2 M}{c_5 M^2 M^{-5/2}} = \frac{C_\mathcal{E}^2}{c_5} \mathcal{L}^2 M^{3/2}. \quad \square$$

Lemma 35.5 is the whole mechanism of this section, and it is worth saying at once what it replaces. Proposition 28.4 bounds the coupling by  $\|w\|_{L^\infty}$  times the enstrophy budget, thereby *discarding* the a priori  $w$ -budget of Theorem 28.2; (171) uses both budgets and pays for it with a correlation hypothesis. The gain is not the logarithm per se: it is the elimination of  $\|w\|_{L^\infty}$ , which is the unknown that H1 is trying to produce.

### 35.2 Test configurations: what a non-implication can mean here

Neither conjecture can be refuted or verified by exhibiting a Navier–Stokes solution; both are asserted only *near a putative Type-I singularity*, whose existence is the open question.

What can be settled — and is settled below — is whether either implication follows from the a priori structure that the two hypotheses are formulated against.

**Definition 35.6** (Admissible test configuration). Fix  $\nu > 0$ ,  $M \geq 1$ ,  $r^* = M^{-1/2}$ ,  $\mathcal{L} \geq 1$ , and  $\psi$  as in Definition 27.2 with  $\rho = \frac{3}{4}r^*$ ,  $\rho' = r^*$ . An *admissible test configuration* is a triple  $(m, w, h)$  on  $Q(r^*)$  with  $m$  Lipschitz and  $w, h \geq 0$  measurable, such that

$$(A1) \quad 0 \leq m \leq M \text{ and } \sup_{Q(r^*)} m = M;$$

$$(A2) \quad \nu \iint_{Q(r^*)} (|\nabla m|^2 + m^2 w) \leq C_{\mathcal{E}} \mathcal{L} M^{1/2}, \text{ i.e. (159);}$$

$$(A3) \quad |\nabla w| \leq 2 w^{1/2} h^{1/2} \text{ a.e. — the pointwise link } |\nabla w| \leq 2 w^{1/2} |\nabla^2 \hat{e}| \text{ used throughout §29–§34, with } h \text{ standing for } |\nabla^2 \hat{e}|^2.$$

In such a configuration  $c_K$ ,  $\Xi[\varpi]$ , (168)–(169) and (171) are all well defined, with  $|\nabla^2 \hat{e}|^2$  read as  $h$ .

**Remark 35.7** (Scope of the non-implications; stated as a limitation). An admissible test configuration need not come from a solution of the Navier–Stokes equations:  $w$  need not equal  $|\nabla \hat{e}|^2$  for a unit field  $\hat{e}$ ,  $h$  need not be its second-derivative density, and no evolution equation is imposed. Consequently Propositions 35.8 and 35.10 do *not* say that the two conjectures are independent as statements about Navier–Stokes solutions; that question is not decidable by any argument available here. What they do say, and all they say, is: *no implication between Conjecture 28.5 and Conjecture 34.22 can be derived from (A1)–(A3)*, i.e. from the a priori budgets and the pointwise structure in terms of which the two statements are phrased. Any implication would have to invoke further PDE structure, and none is on offer in §28–§34. That is the precise sense in which the “same analytic type” language of Remarks 34.24 and 34.23 must not be read as “one covers the other”.

### 35.3 The two non-implications

**Proposition 35.8** (Band absorption bounds no correlation). *There is a family of admissible test configurations, indexed by  $\ell \in (0, r^*/4)$ , for which the left-hand sides of (168) and (169) vanish identically for every admissible level pair — so Conjecture 34.22 holds with  $C = 0$  — while*

$$c_K \geq \frac{|E|}{8\ell^5} \longrightarrow \infty \quad (\ell \rightarrow 0),$$

*with  $\mathcal{L}$  held fixed. Hence Conjecture 34.22 implies no upper bound on  $c_K$  at all, a fortiori not Conjecture 28.5.*

*Proof.* Let  $\varphi(s) = \text{dist}(s, 2\mathbb{Z}) \in [0, 1]$  be the unit triangular wave, so  $|\varphi'| = 1$  a.e. Fix  $\ell \in (0, r^*/4)$ , let  $K \subseteq B_\rho$  be an axis-aligned cube of side  $\ell$  and  $I \subseteq (t_0 - \rho^2, t_0)$  an interval of length  $\ell^2$ , and put  $N = K \times I$ ,  $|N| = \ell^5$ , and let  $N_0 = K_0 \times I$  with  $K_0$  the concentric cube of side  $\ell/2$ . Let  $\xi$  be a Lipschitz cutoff, independent of  $t$ , with  $\xi \equiv 1$  on  $K_0$ ,  $\text{supp } \xi \subseteq K$  and  $|\nabla \xi| \leq 8/\ell$ , and set

$$m(x, t) = \frac{M}{2} + \frac{M}{8} \xi(x) \varphi\left(\frac{x_1}{\epsilon}\right) \quad \text{on } N, \quad \text{so} \quad |\nabla m|^2 = G := \frac{M^2}{64\epsilon^2} \quad \text{on } N_0,$$

and  $|\nabla m|^2 \leq 4G$  on  $N$  once  $\epsilon \leq \ell/8$  (indeed  $|\nabla m| \leq \frac{M}{8}(|\nabla \xi| + \epsilon^{-1}) \leq \frac{M}{4\epsilon} = 2G^{1/2}$ ). Here  $\epsilon > 0$  is chosen below, and  $\epsilon \leq \ell/8$  holds once  $\ell/r^*$  is small enough in terms of  $\nu$ , since the normalisation below forces  $\epsilon^2 \leq C\nu M^{3/2}\ell^5$ , i.e.  $\epsilon^2/\ell^2 \leq C\nu(\ell/r^*)^3$ . Since  $m/M \in [\frac{3}{8}, \frac{5}{8}]$  on  $N$ , we have  $N \subseteq \{m \geq M/4\}$ , so  $N \subseteq E$  and  $N$  is disjoint from every admissible band; and  $m \equiv M/2$  on  $\partial N$ , so the definition extends Lipschitz across  $\partial N$ . Extend  $m$  to the rest of  $Q(r^*)$  so that  $\sup m = M$ ,  $m$  is constant on a small ball  $P_1 \subseteq E \setminus \bar{N}$ ,  $|E| \geq \frac{1}{2}|Q(r^*)|$  (a normalisation independent of  $\ell$ ), and  $|\nabla m| \leq CM/r^*$  off  $N$ ; the last contributes at most  $C\nu M^2(r^*)^{-2}|Q(r^*)| = C\nu M^{1/2}$  to (A2).

Let  $\zeta$  be Lipschitz with  $0 \leq \zeta \leq 1$ ,  $\text{supp } \zeta \subseteq N_0$  and  $\zeta \not\equiv 0$ , and set  $w = W\zeta^2$  on  $N_0$  plus a further bump  $W' \geq 0$  supported in  $P_1$ , and  $w = 0$  elsewhere; set  $h = W|\nabla \zeta|^2$  on  $N_0$  and correspondingly on  $P_1$ , so that (A3) holds with  $|\nabla w| = 2W\zeta|\nabla \zeta| = 2w^{1/2}h^{1/2}$ .

Choose  $\epsilon$  and  $W, W'$  so that  $4\nu G|N| = \frac{1}{2}C_\mathcal{E}\mathcal{L}M^{1/2}$ ,  $\nu M^2(W|N| + W'|P_1|) \leq \frac{1}{4}C_\mathcal{E}\mathcal{L}M^{1/2}$ , and  $C\nu M^{1/2} \leq \frac{1}{4}C_\mathcal{E}\mathcal{L}M^{1/2}$  (enlarging  $C_\mathcal{E}$  if necessary); then (A1)–(A2) hold. Since  $w = h = 0$  on  $\{m \leq M/4\}$ , both left-hand sides of (168)–(169) vanish for every admissible pair.

For  $c_K$ : on  $E$  the product  $|\nabla m|^2 w$  is supported in  $N_0$ , where  $|\nabla m|^2 \equiv G$ , so  $\iint_E |\nabla m|^2 w = G \iint_{N_0} w$ ; also  $\iint_E w \geq \iint_{N_0} w > 0$  and

$$\iint_E |\nabla m|^2 \leq 4G|N| + C \frac{M^2}{(r^*)^2} |Q(r^*)| = 4G|N| + CM^{1/2} \leq 6G|N|$$

by the third normalisation above. Hence

$$c_K = \frac{|E| G \iint_{N_0} w}{(\iint_E |\nabla m|^2)(\iint_E w)} \geq \frac{|E| G \iint_{N_0} w}{6G|N| \cdot (\iint_{N_0} w + W'|P_1|)} \geq \frac{|E|}{6|N|} \cdot \frac{\iint_{N_0} w}{\iint_{N_0} w + W'|P_1|},$$

and taking  $W'|P_1| \leq \frac{1}{3} \iint_{N_0} w$  gives  $c_K \geq |E|/(8|N|)$ , which is the claim, since  $|E| \geq \frac{1}{2}|Q(r^*)|$  is independent of  $\ell$  while  $|N| = \ell^5 \rightarrow 0$ .  $\square$

**Remark 35.9** (The conclusions do not compare either). One might hope that even if the hypotheses do not compare, their *conclusions* do: Conjecture 34.22 yields  $K$ -absorption (through Remark 34.23 and Proposition 30.4), and  $K$ -absorption is what Conjecture 28.5 was introduced to supply. But  $K$ -absorption is an *upper bound* on  $Y = \nu \iint |\nabla m|^2 w \eta^2$ , whereas  $c_K$  is a *ratio*; a small  $Y$  is perfectly compatible with a large  $c_K$ , since the denominator  $(\iint_E |\nabla m|^2)(\iint_E w)/|E|$  may be smaller still. That is exactly what the configuration of Proposition 35.8 realises. So the conclusions do not compare in that direction either.

**Proposition 35.10** (The  $c_K$  hypothesis does not imply band absorption). *There is a family of admissible test configurations, indexed by  $\ell \in (0, r^*/4)$ , with*

$$c_K = 0$$

— the strongest possible form of Conjecture 28.5 — for which every admissible level pair of width  $\theta_+ - \theta_- \geq \frac{1}{16}$  violates (168): writing  $\Gamma$  for the quantity defined in the proof,

$$\nu \iint_{\text{Band}} |\nabla m|^2 w \psi^2 \geq \frac{1}{4} \Gamma, \quad \nu \iint |\nabla m|^2 w \eta_\chi^2 \leq 8 \Gamma, \quad M^2 \nu \iint h \eta_\chi^2 + \mathcal{I}_0 \leq C \left[ \left( \frac{\ell}{r^*} \right)^3 + \left( \frac{\ell}{r^*} \right)^5 \right] \Gamma.$$

Hence the constant  $C$  required in (168) diverges as  $\ell/r^* \rightarrow 0$  with  $\mathcal{L}$  held fixed, and Conjecture 34.22 fails along the family.

*Proof.* Let  $\varphi$  be as in the previous proof. Fix  $\ell \in (0, r^*/4)$ , let  $F = K \times I$  with  $K \subseteq B_\rho$  an axis-aligned cube of side  $\ell$  and  $I$  an interval of length  $\ell^2$  inside  $\{\psi = 1\}$ , so  $|F| = \ell^5$ , and let  $N = K' \times I$  where  $K'$  is the concentric cube of side  $\frac{9}{8}\ell$ , so that  $|N| = (\frac{9}{8})^3|F| \leq 2|F|$ . Let  $\xi$  be a Lipschitz cutoff, independent of  $t$ , with  $\xi \equiv 1$  on  $K$ ,  $\text{supp } \xi \subseteq K'$  and  $|\nabla \xi| \leq 16/\ell$ , and set

$$m(x, t) = \frac{M}{16} + \frac{3M}{16} \xi(x) \varphi\left(\frac{x_1}{\epsilon}\right) \quad \text{on } N, \quad |\nabla m|^2 = G := \frac{9M^2}{256\epsilon^2} \quad \text{on } F,$$

with  $|\nabla m|^2 \leq 4G$  on  $N$  whenever  $\epsilon \leq \ell/16$  (since  $|\nabla m| \leq \frac{3M}{16}(|\nabla \xi| + \epsilon^{-1}) \leq \frac{3M}{8\epsilon} = 2G^{1/2}$ ), and  $m \equiv M/16$  on  $\partial N$ , so that the definition extends Lipschitz across  $\partial N$ . Then  $m/M$  takes values in  $[\frac{1}{16}, \frac{1}{4}]$  only. Moreover, provided  $\epsilon \leq \ell/16$  (verified below), for every  $[\theta_-, \theta_+] \subseteq [\frac{1}{16}, \frac{1}{4}]$  of length  $\geq \frac{1}{16}$ ,

$$|\{m/M \in [\theta_-, \theta_+]\} \cap F| \geq \frac{3}{4} \cdot \frac{1/16}{3/16} |F| = \frac{|F|}{4}. \quad (172)$$

Indeed,  $F$  is a product of the cube  $K$  with  $I$ , and the  $x_1$ -side of  $K$ , an interval of length  $\ell$ , contains at least  $\lfloor \ell/2\epsilon \rfloor \geq \ell/2\epsilon - 1$  full periods of  $\varphi(\cdot/\epsilon)$ ; on each full period the set where  $\varphi \in [a, b]$  has measure exactly  $(b-a) \cdot 2\epsilon$ , so the total is at least  $(b-a)(\ell - 2\epsilon) \geq \frac{3}{4}(b-a)\ell$  when  $\epsilon \leq \ell/16$ . Take  $[a, b]$  to be the preimage of  $[\theta_-, \theta_+]$ , of length  $(\theta_+ - \theta_-)/(3/16) \geq \frac{1}{3}$ .

Let  $\zeta$  be Lipschitz and independent of  $t$ , with  $\zeta \equiv 1$  on  $K$ ,  $\text{supp } \zeta \subseteq K'$  and  $|\nabla \zeta| \leq 16/\ell$ , and put  $w = W\zeta^2$ ,  $h = W|\nabla \zeta|^2 \leq 256W\ell^{-2}$  on  $N$ , so (A3) holds as before. Off  $N$ , extend  $m$  Lipschitz with  $|\nabla m| \leq CM/r^*$  so that  $\sup m = M$ , and set  $w = h = 0$  there except on two disjoint balls  $P_1, P_2 \subseteq E$  on which, respectively,  $m$  is constant and a small bump of  $w$  is placed, and  $m$  is non-constant with  $w = 0$ . Then  $\iint_E |\nabla m|^2 w = 0$  while both denominators of  $c_K$  are positive, so  $c_K = 0$ .

Fix  $\epsilon, W$  by

$$\nu G|F| = \gamma C_\epsilon \mathcal{L} M^{1/2}, \quad \nu M^2 W|F| = \gamma C_\epsilon \mathcal{L} M^{1/2}, \quad \gamma := \frac{1}{32},$$

which is consistent with (A2): on  $N$  one has  $|\nabla m|^2 \leq 4G$  and  $w \leq W$  with  $|N| \leq 2|F|$ , so those two contributions are at most  $8\gamma$  and  $2\gamma$  times  $C_\epsilon \mathcal{L} M^{1/2}$ , i.e. at most  $\frac{1}{4} + \frac{1}{16}$  of the budget; the far field ( $\leq C\nu M^{1/2} \leq \frac{1}{4} C_\epsilon \mathcal{L} M^{1/2}$  after enlarging  $C_\epsilon$ ) and the two small bumps on  $P_1, P_2$  use the rest. Set

$$\Gamma := \nu G W |F| = (\nu G|F|) W = \gamma C_\epsilon \mathcal{L} M^{1/2} \cdot \frac{\gamma C_\epsilon \mathcal{L} M^{-3/2}}{\nu |F|} = \frac{\gamma^2 C_\epsilon^2 \mathcal{L}^2}{\nu M |F|}.$$

*First display.* By (172),  $\zeta \equiv 1$  and  $\psi \equiv 1$  on  $F$ , the left-hand side of (168) is at least  $\nu G W |F|/4 = \Gamma/4$ .

*Second display.*  $|\nabla m|^2 w$  is supported in  $N$ , where  $|\nabla m|^2 \leq 4G$  and  $w \leq W$ ; since  $\eta_\chi^2 \leq \psi^2 \leq 1$  and  $|N| \leq 2|F|$ ,  $\nu \iint |\nabla m|^2 w \eta_\chi^2 \leq 4\nu G W |N| \leq 8\Gamma$ .

*Third display.*  $h$  is supported in  $N \setminus F$  and bounded by  $256W\ell^{-2}$  there, and  $|N \setminus F| \leq |N| \leq 2|F|$ , so

$$\frac{M^2 \nu \iint h \eta_\chi^2}{\Gamma} \leq \frac{\nu M^2 \cdot 256W\ell^{-2} \cdot 2|F|}{\nu G W |F|} = \frac{512 M^2}{G \ell^2} = \frac{512 \cdot 256 \epsilon^2}{9 \ell^2}.$$

From  $\nu G|F| = \gamma C_\epsilon \mathcal{L} M^{1/2}$ ,  $|F| = \ell^5$  and  $\mathcal{L} \geq 1$  one gets  $\epsilon^2 = \frac{9\gamma^{-1}}{256} \nu M^{3/2} \ell^5 / (C_\epsilon \mathcal{L}) \leq C\nu M^{3/2} \ell^5$ , whence the ratio is at most  $C\nu M^{3/2} \ell^3 = C\nu(\ell/r^*)^3$ ; the same bound gives

$\epsilon^2/\ell^2 \leq C\nu(\ell/r^*)^3$ , hence  $\epsilon \leq \ell/16$  once  $(\ell/r^*)^3 \leq (256C\nu)^{-1}$ , which is the standing smallness assumption on  $\ell/r^*$  and validates both (172) and the bound  $|\nabla m|^2 \leq 4G$  on  $N$ . Finally

$$\frac{\mathcal{I}_0}{\Gamma} = \frac{\mathcal{L}^2 M^{3/2} \nu^{-1}}{\gamma^2 C_{\mathcal{E}}^2 \mathcal{L}^2 (\nu M |F|)^{-1}} = \frac{M^{5/2} |F|}{\gamma^2 C_{\mathcal{E}}^2} = \frac{1}{\gamma^2 C_{\mathcal{E}}^2} \left( \frac{\ell}{r^*} \right)^5,$$

using  $M^{5/2} = (r^*)^{-5}$  and  $|F| = \ell^5$ .

*Conclusion.* If (168) held with some constant  $C$ , then  $\frac{1}{4}\Gamma \leq 2^{-11} \cdot 8\Gamma + C'[(\ell/r^*)^3 + (\ell/r^*)^5]\Gamma$ , i.e.  $\frac{1}{4} - 2^{-8} \leq C'[(\ell/r^*)^3 + (\ell/r^*)^5]$ , which fails for  $\ell/r^*$  small. Since  $\mathcal{L}$  is held fixed while  $\ell \rightarrow 0$ , no choice of  $C$  works along the family.  $\square$

**Remark 35.11** (Reading of Proposition 35.10). The configuration is the natural adversary:  $|\omega|$  oscillates rapidly *inside* the transition band, so that  $|\nabla m|^2$  and  $w$  are both large on a common set whose  $m$ -values never leave  $[\frac{1}{16}, \frac{1}{4}]$ . Conjecture 28.5 is blind to it, being a statement about  $\{m \geq M/4\}$ , where in this configuration nothing happens. Two further observations. (i) The oscillation is what makes  $|\nabla m|$  large while  $m$  stays inside the band; there is no contradiction with  $m \in [M/16, M/4]$ , and this is exactly the mechanism identified in Remark 34.18 as the reason the band terms are not controlled by the enstrophy budget alone. (ii) The same configuration has  $\Xi[\mathbf{1}_{\text{Band}} \psi^2] \geq c(r^*/\ell)^5 \rightarrow \infty$ , consistently with Proposition 35.24 below: the hypothesis that *does* imply (168) is violated here, as it must be.

### 35.4 What has to be added: the band transport

Proposition 35.10 shows that Conjecture 28.5 must at least be *transported* to the band before any implication can be hoped for. It is then a genuine implication, and the extra ingredient is explicit.

**Proposition 35.12** (Band-transported  $c_K$  implies (168)). *Let  $\theta_{\pm}$  be admissible and suppose*

$$c_K^{\text{band}} := \frac{|\text{Band}| \iint_{\text{Band}} |\nabla m|^2 w \psi^2}{\left( \iint_{\text{Band}} |\nabla m|^2 \psi^2 \right) \left( \iint_{\text{Band}} w \psi^2 \right)} \leq \frac{C_1}{\mathcal{L}}. \quad (173)$$

*If in addition*

$$(\text{NB}) \quad |\text{Band}| \geq \frac{\kappa_1 |Q(r^*)|}{\mathcal{L}} \quad \text{for some } \kappa_1 > 0, \quad (174)$$

*then (168) holds — indeed in the stronger form*

$$\nu \iint_{\text{Band}} |\nabla m|^2 w \psi^2 \leq \frac{C C_1}{\kappa_1} \mathcal{I}_0,$$

*with neither the  $Y$ -term nor the  $D$ -term needed on the right.*

*Proof.* By Definition 35.1,  $\Xi[\mathbf{1}_{\text{Band}} \psi^2] = c_K^{\text{band}} |Q(r^*)|/|\text{Band}| \leq (C_1/\mathcal{L})(\mathcal{L}/\kappa_1) = C_1/\kappa_1$  by (173) and (174). Now apply (171) with  $\varpi = \mathbf{1}_{\text{Band}} \psi^2$ , which is supported in  $Q^{\sharp}$  because  $\theta_- \geq \frac{1}{16}$ .  $\square$

**Remark 35.13** ((NB) is not free, and is of the same family). Hypothesis (174) says the transition band is not too thin. It is not automatic: by the co-area formula, as in Proposition 34.16(d),

$$|\text{Band}| = \int \int_{\theta-M}^{\theta+M} \int_{\{m=s\}} \frac{d\mathcal{H}^2}{|\nabla m|} ds dt,$$

so a thin band is precisely a band on which  $|\nabla m|$  is large — the dangerous case, and the one realised in Proposition 35.10. Thus (NB) is itself a non-concentration statement about  $|\nabla m|$  on level sets of  $|\omega|$ , and it should not be presented as a harmless technical addendum. Two consequences are worth recording. First, the honest answer to “does Conjecture 28.5 imply Conjecture 34.22?” is: *not as stated; its band transport implies only the Y-half* (168), *and only when supplemented by* (174). Second — and this is the point missed in the first draft of this section, see Remark 35.3 — *the bulk side needs a volume hypothesis of exactly the same kind*. By Lemma 35.2, passing from Conjecture 28.5 to the volume-normalised bound  $\Xi[\mathbf{1}_E] = O(1)$  used in Proposition 35.17(i) requires

$$(\text{NB-E}) \quad |E| \geq \frac{\kappa_2 |Q(r^*)|}{\mathcal{L}} \quad \text{for some } \kappa_2 > 0, \quad (175)$$

the exact analogue of (174) with Band replaced by  $E$ . The symmetry is complete: neither the bulk route nor the band route is free of a volume condition, and in both cases the condition fails precisely when the region carrying the coupling is thin — the filamentary regime.

**Remark 35.14** (The two routes, and what each costs). It is worth separating, once and for all, the two ways a correlation hypothesis can be cashed in; the strength comparison of §35.5 is entirely a matter of which route is used.

- (R1) *The  $\|w\|_{L^\infty}$  route* (that of Proposition 28.4 and Proposition 34.19): bound  $\iint w \varpi \leq \|w\|_{L^\infty} |R|$ , so that the volume factor in  $c_K[\varpi]$  cancels and one lands on the Moser-absorbable quantity  $\delta\nu(r^*)^3 \|w\|_{L^\infty}$  of Proposition 29.1. No volume hypothesis; the price is that a factor  $\mathcal{L}$  must be found inside the correlation constant, and that the closure runs through  $\|w\|_{L^\infty}$  (Proposition 35.15).
- (R2) *The budget route* (Lemma 35.5): bound  $\iint w \varpi$  by the a priori  $w$ -budget of Theorem 28.2, so that  $\|w\|_{L^\infty}$  never appears and one lands directly in  $\mathcal{I}_0$ . No logarithm is needed in the correlation constant; the price is the volume factor  $|Q(r^*)|/|R|$ , i.e. a hypothesis of the form (174) or (175).

Neither route dominates: (R1) is cheaper exactly when  $|R|/|Q(r^*)| < 1/\mathcal{L}$ , (R2) exactly when  $|R|/|Q(r^*)| > 1/\mathcal{L}$ . This trade-off, and not any difference between the two conjectures, is what the “ $C/\mathcal{L}$  versus  $O(1)$ ” discrepancy in their statements records.

**Proposition 35.15** (The  $\|w\|_{L^\infty}$  route: both targets require a *decaying* correlation constant). *Let  $R \subseteq Q^\sharp$  be measurable,  $\varpi = \mathbf{1}_R \psi^2$ , and suppose  $c_K[\varpi] \leq C_1/\mathcal{L}$ , with  $c_K[\varpi]$  as in Lemma 35.2. Then*

$$\frac{\nu}{M^2} \iint_R |\nabla m|^2 w \psi^2 \leq C C_1 \|w\|_{L^\infty(Q(\rho') \cap \{m \geq M/16\})} (r^*)^3,$$

*which is precisely the quantity Proposition 29.1 absorbs with factor  $\delta$ , provided the absolute constant  $CC_1$  satisfies  $CC_1 \leq \delta\nu$ . Taken with  $R = E$  this is the conclusion Conjecture 28.5*

was designed to give (Proposition 28.4 with its logarithm removed); taken with  $R = \text{Band}$  it removes exactly the shortfall of Remark 34.20 from the  $Y$ -half of the core residual. No volume hypothesis is used in either case.

*Proof.* By the definition of  $c_K[\varpi]$  in Lemma 35.2 and  $\iint w \varpi \leq \|w\|_{L^\infty} |R|$ ,

$$\iint_R |\nabla m|^2 w \psi^2 = \frac{c_K[\varpi]}{|R|} \left( \iint_R |\nabla m|^2 \psi^2 \right) \left( \iint_R w \psi^2 \right) \leq c_K[\varpi] \|w\|_{L^\infty} \iint_R |\nabla m|^2 \psi^2.$$

Multiply by  $\nu/M^2$ , use  $\nu \iint_R |\nabla m|^2 \psi^2 \leq \mathcal{E} \leq C_\mathcal{E} \mathcal{L} M^{1/2}$  from (159), and  $M^{-3/2} = (r^*)^3$ ; the hypothesis  $c_K[\varpi] \leq C_1/\mathcal{L}$  cancels the  $\mathcal{L}$ .  $\square$

**Remark 35.16** (A bookkeeping caveat, flagged and not resolved). Proposition 35.15 delivers the band term into the *Moser channel*  $\delta\nu(r^*)^3 \|w\|_{L^\infty}$ , which is the target Remark 34.20 names, and not into the right-hand side of (167), which has no  $\|w\|_{L^\infty}$  slot. These are two different closures of the same residual: the first leaves Proposition 30.4 in the form  $Y \leq CM^2 D_{\text{ang}} + \mathcal{I} + C\|w\|_{L^\infty} M^{1/2}$ , the last term being absorbed downstream by Proposition 29.1; the second removes the residual outright. The document states both targets and does not reconcile them — the discrepancy is pre-existing (compare Remark 34.20 with (167)) — and it is recorded here rather than papered over. Separately, the constant condition  $CC_1 \leq \delta\nu$  is a genuine if non-divergent residue; it is present identically in Conjecture 28.5, which this document does not track either, and nothing below claims to remove it.

### 35.5 The strength comparison: where the logarithm really comes from

**Proposition 35.17** (Calibration). *Let  $\kappa_0 \geq 1$ . Then:*

(i) (bulk) *If  $\Xi[\mathbf{1}_{Q^\#} \psi^2] \leq \kappa_0$  then, for every admissible  $\eta_\chi$ ,*

$$\nu \iint |\nabla m|^2 w \eta_\chi^2 \leq C\kappa_0 \mathcal{I}_0,$$

*i.e. the coupling term  $K$  lands in the allowed inhomogeneous class (158) outright — with no  $D_{\text{ang}}$  term, no  $\|w\|_{L^\infty}$  and no annulus residual.*

(ii) (band) *If  $\Xi[\mathbf{1}_{\text{Band}} \psi^2] \leq \kappa_0$  then (168) holds, again in the stronger form with right-hand side  $C\kappa_0 \mathcal{I}_0$ .*

(iii) (comparison — corrected) *The hypotheses used in (i) and (ii) are stronger, not weaker, than the corresponding correlation bounds. By Lemma 35.2,  $\Xi[\mathbf{1}_E] \leq \kappa_0$  implies  $c_K \leq \kappa_0$  but not conversely, the gap being the factor  $|Q(r^*)|/|E|$ ; and Conjecture 28.5 gives only  $\Xi[\mathbf{1}_E] \leq (C/\mathcal{L})|Q(r^*)|/|E|$ , which is  $O(1)$  exactly under (175). Symmetrically on the band, with (174). Neither  $\Xi$ -bound is therefore available from the  $c_K$ -type hypotheses without a volume condition.*

*Proof.* (i) and (ii) are (171) applied with  $\varpi = \mathbf{1}_{Q^\#} \psi^2$  and  $\varpi = \mathbf{1}_{\text{Band}} \psi^2$  respectively, using  $\eta_\chi^2 \leq \psi^2 \mathbf{1}_{\{m \geq M/16\}}$  (Lemma 34.2(ii)) in (i) and  $\text{Band} \subseteq Q^\#$  in (ii). (iii) is Lemma 35.2.  $\square$

**Remark 35.18** (Answer to the strength question, corrected). The premise behind the question “is Conjecture 34.22 strictly weaker than Conjecture 28.5, needing only an  $O(1)$  equipartition constant where  $c_K$  needs a logarithmically decaying one?” does not survive the correction recorded in Remark 35.3. The answer is **no**, and the corrected picture is the following.

- (1) (168) *is  $O(1)$  in its bookkeeping, not in its strength.* It is stated against  $Y$ ,  $M^2 D_{\text{ang}}$  and  $\mathcal{I}_0$  rather than against a correlation constant, which is what makes its constants look absolute. Expressed as a hypothesis on the band correlation constant it is *not* an  $O(1)$  statement: by Proposition 35.15 with  $R = \text{Band}$ , what the  $\|w\|_{L^\infty}$  route requires is  $c_K^{\text{band}} \leq C/\mathcal{L}$  — the exact analogue of Conjecture 28.5, on the band. In this precise and limited sense the “same analytic type” language of Remark 34.23 is *vindicated*: both targets require a logarithmically decaying correlation constant, on their respective regions.
- (2) *The alternative — an  $O(1)$  constant — is bought with a volume hypothesis, symmetrically on both sides.* The budget route (Proposition 35.17) does replace  $C/\mathcal{L}$  by  $O(1)$ , but only in the volume-normalised statistic  $\Xi$ ; and by Lemma 35.2 an  $O(1)$  bound on  $\Xi$  is *strictly stronger* than an  $O(1)$  bound on the correlation constant, by the factor  $|Q(r^*)|/|R|$ . Recovering it from a  $c_K$ -type hypothesis costs (174) on the band and (175) on the bulk. Remark 35.14 states the trade-off: neither route dominates, the crossover being at  $|R|/|Q(r^*)| = 1/\mathcal{L}$ .
- (3) *Consequently the logarithm is not an “artefact”.* It is what the  $\|w\|_{L^\infty}$  route costs, and that route is the one requiring no information about the measure of the region carrying the coupling. Removing the logarithm means acquiring that information instead. The first draft of this section asserted that Conjecture 28.5 could be weakened to an  $O(1)$  statement without loss; that assertion is **withdrawn** (Remark 35.3).
- (4) *Net verdict on the strength question.* Conjecture 34.22 is **not** shown to be weaker than Conjecture 28.5. The two are incomparable as statements (Propositions 35.10 and 35.8), and when each is reduced to a hypothesis on a correlation constant by the same route they require bounds of the *same* form. What separates them is their support, not their strength.

One caveat, recorded rather than concealed: landing  $K$ , or the band term, in  $\mathcal{I}_0 = \mathcal{L}^2 M^{3/2} \nu^{-1}$  is “allowed” only in the sense of §30 (C2), i.e. it is an  $\mathcal{L}^2$ -degraded inhomogeneous term of the same marginal class as every other such term in the scheme. It therefore inherits, and does not resolve, the standing marginality issue of Remark 26.7; it introduces no *new* degradation, which is all that is claimed. The  $\|w\|_{L^\infty}$  route carries its own caveat, Remark 35.16.

**Remark 35.19** (What the existing numerics do and do not support). The  $c_K$  measurements of S34–S35 (EXP-L4-KCORR-\*) reported  $c_K \approx 1$  and were explicitly caveated as unable to distinguish  $O(1)$  from the stronger  $c_K \leq C/\mathcal{L}$ . That caveat stands, and after the correction of Remark 35.3 a second must be added beside it.

- (i) *A measured  $c_K \approx 1$  does not support  $\Xi[\mathbf{1}_E] = O(1)$ .* By Lemma 35.2,  $\Xi[\mathbf{1}_E] = c_K |Q(r^*)|/|E| \geq c_K$ , so such a measurement bounds  $\Xi$  only once the volume fraction  $|E|/|Q(r^*)|$  is known and bounded below. The first draft of this section asserted the reverse inequality and drew the opposite conclusion; that is withdrawn.



- (ii) *The gap is worst where it matters.* The correction factor  $|Q(r^*)|/|E|$  is largest exactly when the near-maximal vorticity set is thin — the filamentary regime expected near a putative singularity — so an  $O(1)$  correlation measurement in a benign configuration says correspondingly less about  $\Xi$  in a concentrated one. Volume fractions quoted from the S42–S43 diagnostics should be treated as motivation for measuring  $|E|/|Q(r^*)|$  directly and *not* as evidence about it: those diagnostics take the near-maximal set over the whole periodic box rather than over  $Q(r^*)$ , which is not the quantity appearing in Lemma 35.2.
- (iii) *What should be measured.* Three quantities, none of which the existing suite reports in the form needed: the volume fractions  $|E|/|Q(r^*)|$  and  $|\text{Band}|/|Q(r^*)|$  at the working scale  $r^*$ , to test (175) and (174); the band correlation constant  $c_K^{\text{band}}$  of (173), which by Proposition 35.10 cannot be inferred from the  $E$ -measurement; and the  $\mathcal{L}$ -dependence of both correlation constants, since Proposition 35.15 needs decay and not merely boundedness.

The plain statement, replacing the first draft’s: *the existing  $O(1)$  measurements are consistent with, but do not establish, either Conjecture 28.5 or Conjecture 35.20; and they bear on Conjecture 35.23 only in combination with a volume measurement that has not been made.*

### 35.6 Common hypotheses

Two hypotheses are recorded, one for each of the routes of Remark 35.14. Each implies the  $Y$ -halves of *both* targets; neither implies the other; and the first is the weaker of the two in the regime that matters. Both are open. Neither covers the angular half (169), for the reason given in Remark 35.26.

*Conjecture 35.20* (Level-uniform decorrelation). There is  $C_1$ , depending only on the standing Type-I data, such that near a putative Type-I singularity, at the working scale  $r^* = M(t_0)^{-1/2}$ , with  $Q^\sharp = Q(\rho') \cap \{m \geq M/16\}$  and  $c_K[\cdot]$  as in Lemma 35.2,

$$(LC) \quad c_K[\mathbf{1}_{Q^\sharp} \psi^2] \leq \frac{C_1}{\mathcal{L}}. \quad (176)$$

This is *verbatim* the hypothesis of Conjecture 28.5 — same normalisation, same  $1/\mathcal{L}$  decay — asserted on  $\{m \geq M/16\}$  instead of on  $\{m \geq M/4\}$ , i.e. extended downwards across the transition band.

**Proposition 35.21** ((LC) covers both  $Y$ -halves). *Assume (176). Then, with  $\|w\|_\infty = \|w\|_{L^\infty(Q(\rho') \cap \{m \geq M/16\})}$ ,*

$$\frac{\nu}{M^2} \iint |\nabla m|^2 w \eta_\chi^2 + \frac{\nu}{M^2} \iint_{\text{Band}} |\nabla m|^2 w \psi^2 \leq C C_1 \|w\|_\infty (r^*)^3$$

*for every admissible  $\chi$  and every admissible level pair. Both terms are therefore of the class Proposition 29.1 absorbs with factor  $\delta$ , subject to the constant condition  $CC_1 \leq \delta\nu$  of Proposition 35.15. In particular (176) simultaneously supplies the conclusion sought from Conjecture 28.5 and removes the shortfall of Remark 34.20 from the  $Y$ -half of the core residual, with no volume hypothesis.*

*Proof.* Both  $\{\eta_\chi > 0\}$  and Band are contained in  $Q^\sharp$  and the integrand is nonnegative, so both left-hand terms are bounded by  $\frac{\nu}{M^2} \iint_{Q^\sharp} |\nabla m|^2 w \psi^2$ ; apply Proposition 35.15 with  $R = Q^\sharp$ .  $\square$

**Remark 35.22** (Why (LC) is the right reading of “same analytic type”). Remark 34.24 said that  $\mathcal{A}_{\text{ann}}$  is “open, of the same analytic type as Conjecture 28.5”. Sections 35.3 show that this cannot mean one implies the other. Conjecture 35.20 is what it *can* legitimately mean: the two targets are two instances of a *single* hypothesis, differing only in the region on which it is asserted, and the single hypothesis stated on the union of those regions covers both. This costs nothing in form: it is the  $c_K$  hypothesis, unchanged in normalisation and in the decay rate of its constant, differing from Conjecture 28.5 only in the region on which it is asserted.

Two things this does *not* say, lest the sentence above be read too generously. First, (176) does not imply Conjecture 28.5: a bound on a single correlation ratio over  $Q^\sharp$  does not restrict to the sub-region  $E$ , the ratio being non-monotone under restriction. What Proposition 35.21 supplies is the *conclusion* sought from Conjecture 28.5, not that conjecture itself. Second and conversely, Conjecture 28.5 does not imply (176): by Proposition 35.10 the extension of the assertion across the band is exactly the part it does not supply. (LC) is therefore a genuinely new hypothesis of an old form, not a repackaging of an existing one.

*Conjecture 35.23* (Joint level equipartition). There is  $\kappa_0$ , depending only on the standing Type-I data, such that near a putative Type-I singularity, at the working scale  $r^* = M(t_0)^{-1/2}$ , with  $Q^\sharp = Q(\rho') \cap \{m \geq M/16\}$ :

$$\text{(JLE-1)} \quad \iint_{Q^\sharp} |\nabla m|^2 w \psi^2 \leq \frac{\kappa_0}{|Q(r^*)|} \left( \iint_{Q^\sharp} |\nabla m|^2 \psi^2 \right) \left( \iint_{Q^\sharp} w \psi^2 \right), \quad (177)$$

$$\text{(JLE-2)} \quad \nu \iint_{\text{Band}} |\nabla^2 \hat{e}|^2 \psi^2 \leq \kappa_0 \nu \iint |\nabla^2 \hat{e}|^2 \eta_\chi^2 + \kappa_0 \mathcal{L}^2 M^{-1/2} \nu^{-1}, \quad (178)$$

the latter for some admissible level pair  $\theta_\pm$  of width  $\theta_+ - \theta_- \geq \frac{1}{16}$  and some  $\chi \in \mathcal{C}(\theta_-, \theta_+)$ .

**Proposition 35.24** (Both targets follow from Conjecture 35.23). *Assume (JLE-1). Then*

- (i)  $\nu \iint |\nabla m|^2 w \eta_\chi^2 \leq C \kappa_0 \mathcal{I}_0$  for every admissible  $\chi$ ; in particular the conclusion that Conjecture 28.5 was introduced to supply holds, and Conjecture 28.5 itself is not needed;
- (ii) (168) holds for every admissible level pair, in the stronger form with right-hand side  $C \kappa_0 \mathcal{I}_0$ .

*If in addition (JLE-2) holds for the pair  $\theta_\pm$  and cutoff  $\chi$  named there, then (169) holds for that pair; hence Conjecture 34.22 holds in full, hence (167) holds, and the caveat “modulo  $\mathcal{A}_{\text{ann}}$ ” may be removed from Proposition 30.4.*

*Proof.* (JLE-1) is precisely  $\Xi[\mathbf{1}_{Q^\sharp} \psi^2] \leq \kappa_0$ , so (i) and (ii) are Proposition 35.17(i)–(ii), using  $\text{Band} \subseteq Q^\sharp$ ,  $\eta_\chi^2 \leq \psi^2 \mathbf{1}_{\{m \geq M/16\}}$ , and the nonnegativity of the integrand for the restriction to subregions. Multiplying (178) by  $M^2$  gives (169) with  $C = \kappa_0$ , since  $M^2 \cdot \mathcal{L}^2 M^{-1/2} \nu^{-1} = \mathcal{L}^2 M^{3/2} \nu^{-1}$ . The last two implications are those of Remark 34.23.  $\square$

**Remark 35.25** (Status, failure mode, and what (JLE-1) really asserts). (JLE-1) says: *on the region where the level cutoff lives, the coupling density  $|\nabla m|^2 w$  is no larger than a fixed multiple of what uncorrelated densities spread over the ambient parabolic cylinder would give.*

It is an equipartition/non-concentration hypothesis of exactly the kind described in the remark following Conjecture 28.5, with two differences: the normalising volume is  $|Q(r^*)|$  rather than the measure of the level region, and the constant is  $O(1)$  rather than  $O(1/\mathcal{L})$ . It is *sufficient* for both  $Y$ -halves and *necessary* for neither (Proposition 35.8 exhibits a configuration in which (168) holds while  $\Xi$  is unbounded).

**But it is a demanding hypothesis, not a cheap one.** This is the point the first draft of this section got backwards (Remark 35.3). By Lemma 35.2, (JLE-1) is equivalent to  $c_K[\mathbf{1}_{Q^\sharp}\psi^2] \leq \kappa_0 |Q^\sharp|/|Q(r^*)|$ , i.e. it demands a correlation constant not merely bounded but *small in proportion to the volume fraction of  $Q^\sharp$* . Its exact failure mode is co-concentration: if  $|\nabla m|^2$  and  $w$  are both of sizes  $G, W$  on a common set  $S$  and negligible elsewhere, then  $\Xi = |Q(r^*)|/|S|$ , which is large precisely when the vorticity structure is filamentary — the regime of interest. Neither (JLE-1) nor (176) implies the other: (JLE-1) gives  $c_K[\mathbf{1}_{Q^\sharp}\psi^2] \leq \kappa_0$ , which is  $O(1)$  and not  $O(1/\mathcal{L})$ , so it does not give (176); and (176) gives  $\Xi[\mathbf{1}_{Q^\sharp}\psi^2] \leq (C_1/\mathcal{L})|Q(r^*)|/|Q^\sharp|$ , which is  $O(1)$  only under a volume hypothesis of the form (175). They are two genuinely different sufficient hypotheses, matched to the two routes of Remark 35.14.

**Which to target.** On present evidence Conjecture 35.20 is the better target: it needs no information about the measure of any level region, it is the existing  $c_K$  hypothesis with its region enlarged, and it degrades gracefully under concentration, whereas (JLE-1) degrades like  $|Q(r^*)|/|S|$ . Conjecture 35.23 is retained because it is the natural hypothesis for the budget route, it yields the stronger conclusion (landing in  $\mathcal{I}_0$  outright, with no  $\|w\|_{L^\infty}$  and hence none of the caveats of Remark 35.16), and it becomes the more attractive of the two in any regime where the near-maximal vorticity set is *not* thin.

**Remark 35.26** (The angular half is *not* of the  $c_K$  type). (169), equivalently (JLE-2), contains no product of two densities and no correlation: it is a bare level non-concentration statement for the angular curvature density  $\nu|\nabla^2\hat{e}|^2\psi^2$ . Nothing in Conjecture 28.5, in (177), or in any decorrelation hypothesis bears on it. It would follow from either of two unrelated inputs: an a priori bound  $\nu \iint_{Q^\sharp} |\nabla^2\hat{e}|^2\psi^2 \leq C\mathcal{L}^2 M^{-1/2}\nu^{-1}$  — none is available, since  $\nu \iint |\nabla^2\hat{e}|^2\eta^2$  is a left-hand quantity of the first rung and is controlled only *with* the cutoff  $\eta$  — or a level non-concentration hypothesis for that density. So the assertion of Remarks 34.24 and 34.23 that  $\mathcal{A}_{\text{ann}}$  is “of the same analytic type as Conjecture 28.5” is accurate for the  $Y$ -half of the closure target and *inaccurate for the  $D$ -half*. This is the sharpest single correction this section makes to the surrounding text.

**Remark 35.27** (Level refinement moves the two halves in opposite directions). A small observation, recorded because it is the natural next thing to try and it does not work. Remark 34.17 shows that pigeonholing over  $N$  equally spaced levels gains nothing for a  $\chi'$ -weighted residual, because  $|\chi'| \leq CN$  cancels the  $1/N$  of the lightest annulus. The left-hand side of (169) carries *no*  $\chi'$  factor, so for it the pigeonhole is a genuine gain: at least half of  $N$  equally spaced bands carry at most  $4/N$  of  $\nu \iint_{Q^\sharp} |\nabla^2\hat{e}|^2\psi^2$  each, and the constant multiplying that band in the Young step of Proposition 34.11 (with  $\delta = 1$ ) is  $N$ -independent. But the same refinement multiplies the  $Y$ -side constants by  $\|\chi\chi'\|_\infty^2 + \|\chi\chi''\|_\infty = O(N^2)$ , so the coefficient  $2^{-11}$  in (168) would have to be replaced by  $cN^{-2}$  while the band mass on that side decreases by at best  $4/N$ : a net loss of a factor  $\sim N$ . Since a single pair  $\theta_\pm$  must serve both halves of Conjecture 34.22, the refinement cannot be used. This refines, and does not contradict, Remark 34.17.

### 35.7 Status table (S44)

| Question                                          | Answer                                                       |   |
|---------------------------------------------------|--------------------------------------------------------------|---|
| (a) Conj. 28.5 $\Rightarrow$ Conj. 34.22?         | <b>No</b> (disjoint supports)                                | V |
| (a') Band-transported $c_K \Rightarrow$ (168)?    | <b>Yes</b> — via $\ w\ _{L^\infty}$ , no extra ingredient    | F |
| (a'') ... or via the budget route?                | <b>Yes</b> , but only given (NB) (174)                       | F |
| (b) Conj. 34.22 $\Rightarrow$ Conj. 28.5?         | <b>No</b> ; no bound on $c_K$ at all                         | F |
| (b') Do the two <i>conclusions</i> compare?       | <b>No</b>                                                    | F |
| (c) Common sufficient hypothesis (preferred)?     | <b>Yes</b> — (LC), the $c_K$ hypothesis on $\{m \geq M/16\}$ | C |
| (c') Common hypothesis for the budget route?      | <b>Yes</b> — (JLE-1), but demanding                          | C |
| (c'') Does either cover (169)?                    | <b>No</b> — different analytic type                          | R |
| (d) Is (168) weaker than Conj. 28.5?              | <b>No</b> ; $O(1)$ in bookkeeping only                       | F |
| (d') Is $\Xi = O(1)$ weaker than $c_K = O(1)$ ?   | <b>No</b> — <i>stronger</i> , by $ Q(r^*) / R $              | L |
| (d'') Numerics ( $c_K \approx 1$ ) support which? | Neither, on their own; a volume measurement is missing       | R |
| (d''') First-draft answer to (d)                  | <b>Withdrawn</b> (lemma was inverted)                        | F |
| (e) Independent as stated?                        | <b>Yes</b> , over the a priori structure (A1)–(A3)           | R |
| Volume conditions (NB), (NB-E)                    | <b>Symmetric</b> ; needed by the budget route only           | R |
| Level refinement as a route                       | <b>Net loss</b>                                              | F |
| (LC) and (JLE-1) themselves                       | <b>Open</b> ; mutually independent                           | R |

**Remark 35.28** (Honest summary of S44). The resemblance noted in Remarks 34.24 and 34.23 is real at the level of *form* and illusory at the level of *content*: the two conjectures are supported on disjoint regions of the  $|\omega|$ -level decomposition, and neither implies the other, nor any nontrivial part of the other, from the a priori structure available — Remark 35.7 states the exact scope of that claim, which is a statement about admissible test configurations and not about Navier–Stokes solutions.

The positive outcome is that a single hypothesis covers both  $Y$ -halves, and that it is a familiar one: Conjecture 35.20 is the  $c_K$  hypothesis of Conjecture 28.5 with its region enlarged from  $\{m \geq M/4\}$  to  $\{m \geq M/16\}$ , and nothing else. That is the legitimate content of “same analytic type” (Remark 35.22): not that either target subsumes the other, but that both are instances of one hypothesis stated on adjacent regions.

The negative outcomes are three. First, the two conjectures are genuinely incomparable, so the loose language of Remarks 34.24 and 34.23 must not be read as an implication. Second, the angular half (169) is of a different type entirely and is covered by no decorrelation hypothesis (Remark 35.26). Third — and this reverses what the first draft of this section claimed — (168) is *not* weaker than Conjecture 28.5: its apparently absolute constants are an artefact of its bookkeeping, and reduced to a hypothesis on a correlation constant it needs the same  $1/\mathcal{L}$  decay (Proposition 35.15). The volume-normalised statistic  $\Xi$  that would avoid the logarithm is *stronger*, not weaker, than the correlation constant (Lemma 35.2), so the existing  $O(1)$  numerics do not deliver it; what they would need to be supplemented by is listed in Remark 35.19(iii). The withdrawal is recorded in Remark 35.3.

No proved statement of the document is altered, and nothing here bears on H2, on  $\sigma^*$ -decay, on the hypotheses of §33, or on Type-II exclusion.

## 36 Hypothesis (R) Settled Against the Literature (Session S48)

This section discharges hypothesis (R) of §33.2. It is a literature audit with three short proofs; it introduces no new programme machinery and no new hypothesis.

**Verdict, stated first.** (R) *is a theorem*, in a form strictly stronger than the one this document assumes. It follows from the classical *subcritical* interior regularity criterion of Serrin [7], made quantitative and scale-invariant by the  $L^\infty$ -based local theory of Koch–Nadirashvili–Seregin–Šverák [17] and Seregin–Šverák [22]. It is *not* a consequence of  $\varepsilon$ -regularity, and therefore the trap that governed this question — “ $\varepsilon$ -regularity needs the scaled local energy to be *small*, Type-I supplies only *boundedness*” — does not arise: no smallness, no local energy, and no pressure hypothesis enters. Consequently Proposition 33.8(d) is upgraded from *Proved-modulo-(R)* to *Proved*, and Wall 2 of §44.3 is **withdrawn**.

Three things are *not* thereby proved, and are named precisely in §36.5: the saturation proviso inside Proposition 33.8(d), which Proposition 33.8(e) uses and which is independent of (R); the discharge of (N), (N') or (A-up), none of which is touched; and Wall 1 for the original solution  $u$ , which stands.

### 36.1 What (R) says, and what is actually available

(R) *Scale-uniform interior regularity* (line 6785 of this document, as of S48). *There are constants  $C_k(c_I)$ ,  $k \geq 0$ , such that every  $u_\lambda$  as in (152) satisfies*

$$\|\nabla^k u_\lambda\|_{L^\infty(B_R \times [-S_2, -S_1])} \leq C_k(c_I) (1 + R) S_1^{-(k+1)/2}$$

*for all  $0 < S_1 < S_2$  and  $R \geq 1$  with  $B_R \times [-2S_2, -S_1/2]$  inside the domain of  $u_\lambda$ .*

Two features of this statement matter and are easy to miss.

**Remark 36.1** (The hypothesis available is global in space, so the “interior” framing and the factor  $(1 + R)$  are slack). The standing assumption of §27 (line 5565, repeated at line 6760) is *two* bounds, not one:  $M(t) \leq C_I/(T - t)$  and  $\sup_x |u(x, t)| \leq C_I(T - t)^{-1/2}$ , the supremum being over all of  $\mathbb{T}^3$ . Hence (153) is a bound at *every*  $y \in \Lambda_\lambda$ , and on the time slab  $[-S_2, -S_1]$  one has the single scalar bound

$$\|u_\lambda\|_{L^\infty(\Lambda_\lambda \times [-S_2, -S_1])} \leq c_I S_1^{-1/2}, \quad \|\omega_\lambda\|_{L^\infty(\Lambda_\lambda \times [-S_2, -S_1])} \leq c_I S_1^{-1}. \quad (179)$$

(R) is therefore an interior statement in *time* only. There is nothing to be gained by the factor  $(1 + R)$ , and the theorem proved below does not produce one: the constants are uniform in  $y$ , as Proposition 33.8(d) already anticipates in its proof (“the constant [is] independent of  $y_0$ , because (151) is uniform in  $y$ ”).

**Remark 36.2** (The exponent pair is *strictly subcritical* for Serrin). (179) places  $u_\lambda$  in  $L_t^{s'} L_x^s$  with  $s' = s = \infty$ , so

$$\frac{2}{s'} + \frac{3}{s} = 0 < 1.$$

This is the *strict-inequality* range of Serrin’s criterion — the range Serrin himself proved in 1962, not the borderline equality case that needed Struwe and Takahashi. Boundedness

suffices; no smallness is required, and none is available. This single observation is what decides the present question, and it is why the  $\varepsilon$ -regularity literature ([1], [19]) is the wrong shelf: those theorems live at the critical scale, where smallness is the whole content.

### 36.2 The literature statements, quoted

The four statements below are quoted, not recalled. Where a statement is paraphrased the paraphrase is marked as such.

**Theorem 36.3** (Serrin’s interior regularity criterion; Serrin 1962, Struwe 1988, Takahashi 1990). *Let  $Q = B \times (0, T)$  with  $B$  a ball in  $\mathbb{R}^3$ , and let  $u \in L^2(Q)$  be a distributional solution of the Navier–Stokes equations (tested against solenoidal  $C_0^\infty(Q)$  fields, so that the pressure is eliminated). If*

$$u \in L^\infty(L^2(B)), \quad \omega := \operatorname{curl} u \in L^2(Q),$$

*and in addition*

$$u \in L^{s'}(L^s(B)), \quad \frac{2}{s'} + \frac{3}{s} = 1, \quad s' \in (2, \infty),$$

*then “ $u$  is  $C^\infty$  in the spatial variables and every spatial derivative of  $u$  belongs to  $L^\infty(Q')$  for any  $Q' \Subset Q$ ”.* [Cited verbatim from Chen–Galdi–Poggi–Schikorra [24], equations (1.1)–(1.4) and the sentence following. The same source records: “Serrin proved (1.4) under the stronger assumption that the exponents  $s', s$  satisfy the relation above with a strict inequality sign. The proof of the result, as stated here, is due to Struwe [20] and Takahashi [21].” The case relevant here,  $s' = s = \infty$ , lies in the *strict* range and is therefore Serrin’s own theorem.]

**Remark 36.4** (No pressure hypothesis, and no smallness). Kwon [25], surveying exactly this point, writes: “For weak solutions, Serrin [36] showed the local  $L^\infty$ -boundedness under the assumption  $u \in L_t^r L_x^m(Q_1)$  for  $(r, m)$  in the subcritical range  $\frac{2}{r} + \frac{3}{m} < 1$ . After that, Struwe [37] and Takahashi [38] extended the result to the critical range, the equality case, when  $m \in (3, \infty]$ . *These results work even with distributional pressure* [emphasis added], but it is not known whether a generic weak solution satisfies the assumption.” This is the decisive structural fact for the present question: Serrin’s mechanism is a bootstrap on the *vorticity* equation together with a local Helmholtz decomposition [24, (1.5)–(1.7)], in which the pressure never appears. Nothing must be assumed about  $p_\lambda$ , and in particular the  $\lambda$ -uniformity of any local pressure norm — which is *not* available — is not needed.

**Remark 36.5** (What Serrin’s criterion does *not* give, and why that is harmless here). Theorem 36.3 controls *spatial* derivatives only. [24] emphasises: “It is worth emphasizing that (1.4) yields regularity only in the spatial variables; no gain in time regularity is obtained beyond that assumed for  $u$ ”. This is sharp, by Serrin’s example: “ $u = a(t)\nabla\psi$ ,  $\Delta\psi = 0$  in  $B$ , with  $a \in L^2(0, T)$ , is a distributional solution that is  $C^\infty$  in space, whereas its regularity in time is exactly that of the coefficient  $a$ .” Kwon [25] records the same family in the form  $u = \Phi(t)\nabla H(x)$ ,  $p = -\Phi'(t)H(x) - \frac{1}{2}|\Phi(t)\nabla H(x)|^2$ , and notes: “The parasitic solutions  $u$ , on the other hand, are harmonic and hence smooth in space.”

Three consequences, all of which the present section uses.

- (i) **(R)** as *stated* asks only for  $\nabla^k u_\lambda$  and is therefore untouched by the parasitic family; the family satisfies the estimate  $(|\nabla u| = |\Phi|\nabla^2 H| \lesssim \|u\|_{L^\infty(B_1)} \text{ on } B_{1/2} \text{ by interior harmonic estimates})$ .

- (ii) Everything §33 does downstream of **(R)** is built from  $\omega_U$ ,  $\hat{e}_U$ ,  $m_U = |\omega_U|$  and  $\sigma^*$ , all of which are invariant under  $U \mapsto U + b(s)$  and, more generally, unchanged by the curl-free parasitic part.
- (iii) The one place where more than spatial regularity is needed is the  $C_{\text{loc}}^\infty$  compactness of Proposition 33.6, and there it is supplied unconditionally by Theorem 36.9 below, not by **(R)**. See Remark 36.20(c).

**Theorem 36.6** (Quantitative  $L^\infty$  smoothing; KNSS 2009, Proposition 4.1). *Let  $u \in L_{x,t}^\infty(\mathbb{R}^n \times (0, T))$  be a mild solution of the Navier–Stokes Cauchy problem with  $u_0 \in L^\infty$ . Then for  $k, l = 0, 1, \dots$  the functions  $t^{\frac{k}{2}+l} \nabla_x^k \partial_t^l u$  are bounded and, for  $T' = \varepsilon(k, l) \|u_0\|_{L^\infty(\mathbb{R}^n)}^{-2}$  with  $\varepsilon(k, l) > 0$  a small constant,*

$$\left\| t^{\frac{k}{2}+l} \nabla_x^k \partial_t^l u \right\|_{L_{x,t}^\infty(\mathbb{R}^n \times (0, T'))} \leq C(k, l) \|u_0\|_{L^\infty(\mathbb{R}^n)}. \quad (180)$$

[Cited verbatim from [17, Prop. 4.1 and (4.6)]. The same source adds, in its Remark 4.2: “Estimate (4.6) says that the local-in-time smoothing properties of Navier–Stokes for  $u_0 \in L^\infty$  are the same as those of the heat equation.”]

**Remark 36.7** (The same paper’s version without mildness). [17] also proves, for a *bounded weak* solution  $u \in L_{x,t}^\infty(\mathbb{R}^n \times (0, T))$  with  $M = \|u\|_{L_{x,t}^\infty}$ , the estimates

$$\|\nabla_x^k u\|_{L_{x,t}^\infty(\mathbb{R}^n \times (\delta, T))} \leq C(k, \delta, T, M), \quad \|\nabla_x^k \partial_t(u - b)\|_{L_{x,t}^\infty(\mathbb{R}^n \times (\delta, T))} \leq C(k, \delta, R, M),$$

their (4.10) and (4.11), where  $b$  is a function of  $t$  only. The route is worth recording because it is the pressure-free one: the local  $L^p$  bound (4.7) on  $\omega$ , then the vorticity equation  $\omega_{i,t} - \Delta \omega_i = \partial_{x_j}(\omega_j u_i - \omega_i u_j)$ , of which [17] says “Equation (4.8) gains  $\omega$  one spatial derivative in  $L_{x,t}^p$ . The standard bootstrapping arguments and regularity estimates for harmonic functions now give (4.9)”. Note the shape of (4.11): the additive  $b(t)$  sits inside the time derivative and outside every spatial derivative, so all *vorticity* time-derivatives are controlled unconditionally — which is precisely what Lemma 33.21 needs of  $\partial_s \omega_U$  (line 7408).

**Theorem 36.8** (Bounded ancient solutions are quantitatively regular; Seregin–Šverák 2009, Theorem 2.4). *Let  $u$  be an arbitrary bounded ancient solution — i.e., by their Definition 2.3, a bounded divergence-free field  $u \in L^\infty(Q_-; \mathbb{R}^n)$ ,  $Q_- = \mathbb{R}^n \times (-\infty, 0)$ , satisfying  $\int_{Q_-} (u \cdot \partial_t w + u \otimes u : \nabla w + u \cdot \Delta w) dz = 0$  for all solenoidal  $w \in C_0^\infty(Q_-)$ , normalised by  $\text{ess sup } |u| \leq 1$ . Then for any  $m > 1$ ,*

$$|\nabla u| + |\nabla^2 u| + |\nabla p_{u \otimes u}| \in L_m(Q_-),$$

*and for each  $t_0 \leq 0$  there is  $b_{t_0} \in L^\infty(t_0 - 1, t_0)$  with  $\sup_{t_0 \leq 0} \|b_{t_0}\|_{L^\infty(t_0 - 1, t_0)} \leq c(n) < +\infty$  such that, writing  $u_{t_0}(x, t) = u(x, t) + b_{t_0}(t)$ ,*

$$\sup_{z_0 = (x_0, t_0), x_0 \in \mathbb{R}^n, t_0 \leq 0} \|u_{t_0}\|_{W_m^{2,1}(Q(z_0, 1))} \leq c(m, n) < +\infty.$$

[Cited from [22, Thm. 2.4, Def. 2.3, (4.8)]; here  $L_m(Q_-)$  and  $W_m^{2,1}(Q_-)$  are their “uniform” spaces, normed by  $\sup_{z_0 \in Q_-} \|\cdot\|_{\cdot(Q(z_0, 1))}$ . Two points deserve emphasis. First, *no mildness and no pressure integrability is assumed*: the hypothesis is boundedness alone. Second, the additive shift  $b_{t_0}$  is spatially constant, so  $\nabla u_{t_0} = \nabla u$  and  $\nabla^2 u_{t_0} = \nabla^2 u$ : the displayed bound is a bound on the derivatives of  $u$  itself.]

**Theorem 36.9** (Rescaling at a *local* singularity produces a *mild* limit; Seregin–Šverák 2009, Theorem 2.8). *Let  $v, q$  be a local suitable weak solution with a singular point, and let  $u_k(y, s) = \lambda_k v(x_k + \lambda_k y, t_k + \lambda_k^2 s)$  be the rescaling built from  $\lambda_k = 1/M_k$ ,  $M_k = \max_{x \in B(r_1)} |v(x, t_k)| \rightarrow \infty$ , so that  $|u_k| \leq 1$  and  $|u_k(0, 0)| = 1$ . Then: “For each  $a > 0$ , the sequence  $\{u_k\}$  is uniformly Hölder continuous on the closure of  $Q(a)$  for sufficiently large  $k$  and a subsequence  $\{u_{k_j}\}$  of  $\{u_k\}$  converges uniformly on compact subsets of  $\mathbb{R}^n \times ]-\infty, 0]$  to a mild bounded ancient solution  $u$  with  $|u(0, 0)| = 1$ .” Their Remark 2.9: “The main point of the theorem is that the limit  $u$  is a mild solution, i.e., the parasitic solutions cannot appear in this re-scaling procedure.” [Cited verbatim from [22, Thm. 2.8, Rem. 2.9]. The introduction to their §2 states the point in terms: “even in the local (in space) situation, the solutions arising from the re-scaling procedure at a potential singularity are still mild bounded ancient solutions”. Their Remark 2.7 records the companion fact from [17]: “mild ancient solutions are infinitely smooth”.]*

**Remark 36.10** (Two citations in this document that do not say what they are cited for). (a) [18] is cited *three* times as authority for Type-I local regularity technology: at line 6810 (“Both caveats are standard in the Type-I literature [18]”), at line 6862 (“a Type-I-improved  $\varepsilon$ -regularity statement (of the kind developed by Seregin [18])”), and at line 7722 (“Discharging it needs a Type-I-improved  $\varepsilon$ -regularity theorem [18]”).<sup>1</sup> The paper’s abstract is: “We show that a necessary condition for  $T$  to be a potential blow up time is  $\lim_{t \uparrow T} \|v(\cdot, t)\|_{L^3} = \infty$ .” It is a *critical-endpoint*  $L^3$  blow-up result in the line of [16]; it contains no Type-I-improved  $\varepsilon$ -regularity theorem and no discussion of the caveats at line 6810. The intended reference is presumably [22] (which [18] itself cites for exactly this purpose) or, for the local Type-I/ancient-solution equivalence, [23]. Both are added to the bibliography here. *This does not affect (N), which remains assumed.*

(b) [19] is offered at line 6798 as a “sharpening” of Serrin’s criterion by “local-pressure theory”. It is a partial-regularity ( $\varepsilon$ -regularity) paper for *suitable* weak solutions, requiring a local pressure in  $L^{3/2}$  and a smallness hypothesis. It is not the tool that discharges (R), and by Remark 36.4 no local-pressure theory is needed at all.

### 36.3 The $\omega \rightarrow u$ bridge: log-free, but it loses the exponent

Since §33’s Type-I hypothesis is often stated in this programme on the *vorticity* alone, it is worth settling whether the velocity bound in (153) could be *derived* rather than assumed. It cannot, and the obstruction is a scaling identity, not a missing technique. The result is recorded because it forecloses a natural future attempt.

**Lemma 36.11** (Biot–Savart interpolation; logarithm-free). *There is an absolute constant  $C$  such that for every smooth divergence-free  $v$  on  $\mathbb{T}^3$  with  $\int_{\mathbb{T}^3} v \, dx = 0$  and  $\omega = \operatorname{curl} v$ ,*

$$\|v\|_{L^\infty} \leq C \|\omega\|_{L^\infty}^{3/5} \|v\|_{L^2}^{2/5}. \quad (181)$$

*The same estimate holds on  $\mathbb{R}^3$  for  $v$  with  $\|v\|_{L^2} < \infty$ . [Proved.]*

<sup>1</sup>Corrected at S49. As first drafted this remark said “twice” and then named three line numbers, two of which (6848, 7689) were stale — they predated the line shift caused by this section’s own insertion. The count and all three line numbers are now verified against the current file. The defect is recorded rather than silently repaired because line numbers are used as evidence throughout this document, and a remark whose own citations are stale is exactly the failure it describes.



*Proof.* Littlewood–Paley. Write  $v = \sum_{j \geq 0} \Delta_j v$  (no lower block, since  $v$  has zero mean). For the low blocks, Bernstein gives  $\|\Delta_j v\|_{L^\infty} \leq C 2^{3j/2} \|\Delta_j v\|_{L^2} \leq C 2^{3j/2} \|v\|_{L^2}$ , so  $\sum_{j \leq J} \|\Delta_j v\|_{L^\infty} \leq C 2^{3J/2} \|v\|_{L^2}$ . For the high blocks,  $\operatorname{div} v = 0$  and  $\operatorname{curl} v = \omega$  give  $\Delta_j v = 2^{-j} \mathcal{K}_j \tilde{\Delta}_j \omega$  where  $\mathcal{K}_j$  is the Biot–Savart multiplier rescaled to the annulus  $|\xi| \sim 2^j$ ; its kernel has  $L^1$  mass bounded independently of  $j$ , so  $\|\Delta_j v\|_{L^\infty} \leq C 2^{-j} \|\omega\|_{L^\infty}$  and  $\sum_{j > J} \|\Delta_j v\|_{L^\infty} \leq C 2^{-J} \|\omega\|_{L^\infty}$ . Balancing at  $2^J = (\|\omega\|_{L^\infty} / \|v\|_{L^2})^{2/5}$  gives (181).

The point of the computation is where the logarithm of §27 does *not* appear. The same decomposition applied to  $\nabla v$  gives  $\|\Delta_j \nabla v\|_{L^\infty} \leq C \|\omega\|_{L^\infty}$  with *no* decaying factor, and summing over the  $O(\mathcal{L})$  active blocks produces exactly  $\|\nabla v\|_{L^\infty} \lesssim \mathcal{L} \|\omega\|_{L^\infty}$  — the Calderón–Zygmund endpoint failure of Lemma 27.6. For  $v$  itself the geometric factor  $2^{-j}$  sums, and no logarithm is paid. So the intuition that the *velocity* bridge is logarithm-free is correct.  $\square$

**Proposition 36.12** (The exponent is forced, and it is the wrong one). *Let  $\theta \in [0, 1]$  and suppose  $\|v\|_{L^\infty} \leq C \|\operatorname{curl} v\|_{L^\infty}^\theta \|v\|_{L^2}^{1-\theta}$  holds for all smooth compactly supported divergence-free  $v$  on  $\mathbb{R}^3$ . Then  $\theta = 3/5$ . Consequently, from the vorticity Type-I bound  $M(t) \leq C_I (T-t)^{-1}$  and the energy bound  $\|u(\cdot, t)\|_{L^2}^2 \leq E_0$  alone, the best velocity bound obtainable by any such interpolation is*

$$\|u(\cdot, t)\|_{L^\infty} \leq C C_I^{3/5} E_0^{1/5} (T-t)^{-3/5}, \quad (182)$$

and  $3/5 > 1/2$ : the Type-I velocity rate is *not* implied. The rescaled family then satisfies only  $|u_\lambda(y, s)| \leq C \lambda^{-1/5} |s|^{-3/5}$ , which diverges as  $\lambda \downarrow 0$ , so no uniform bound and no limit object survives. [Proved.]

*Proof.* Apply the hypothesis to  $v_\mu(x) = \mu^a v(\mu x)$ ,  $\mu > 0$ . Then  $\|v_\mu\|_{L^\infty} = \mu^a \|v\|_{L^\infty}$ ,  $\|\operatorname{curl} v_\mu\|_{L^\infty} = \mu^{a+1} \|\operatorname{curl} v\|_{L^\infty}$  and  $\|v_\mu\|_{L^2} = \mu^{a-3/2} \|v\|_{L^2}$ . Homogeneity forces  $a = \theta(a+1) + (1-\theta)(a - \frac{3}{2})$ , i.e.  $0 = \theta - \frac{3}{2}(1-\theta)$ , i.e.  $\theta = 3/5$ ; and Lemma 36.11 shows that this exponent is attained, so the family of admissible estimates is exactly one point. Substituting the two bounds gives (182). The rescaled statement follows from  $u_\lambda(y, s) = \lambda u(x^* + \lambda y, T + \lambda^2 s)$  and  $|u| \leq C(\lambda^2 |s|)^{-3/5}$ .  $\square$

**Remark 36.13** (What this settles). The near-field/far-field heuristic — near field  $\lesssim r^* M$  with  $r^* = M^{-1/2}$ , giving the Type-I rate  $M^{1/2} = (T-t)^{-1/2}$  — is correct as far as it goes, and it is the far field that fails: cutting at  $2^J = 1/r^* = M^{1/2}$  leaves a low-frequency contribution  $2^{3J/2} \|u\|_{L^2} \sim M^{3/4} E_0^{1/2} = (T-t)^{-3/4} E_0^{1/2}$ , worse than the near field, and the optimal balance sits at  $\theta = 3/5$ , not at  $r^* = M^{-1/2}$ . *Therefore the velocity Type-I bound at line 5565 is an independent hypothesis and must stay one.* A future session must not attempt to weaken the standing assumptions of §27 to a vorticity-only Type-I hypothesis: by Proposition 36.12 that is not a technical gap but a scaling obstruction. Conversely, nothing is lost: the velocity bound is the standard formulation of Type-I in the literature ([23]), and it is the hypothesis this document already carries.

### 36.4 Discharge of (R)

Two short lemmas transfer the cited theorems to the geometry of (152). Both are routine; both are written out because the programme’s rule is that routine steps are the ones that hide errors.

**Lemma 36.14** (Parasitic solutions do not exist on a torus). *Let  $L > 0$  and let  $\Lambda = L\mathbb{T}^3$ . Suppose  $v(y, s) = \Phi(s)\nabla H(y)$  with  $H$  harmonic on  $\Lambda$ , and that the associated pressure is a single-valued function on  $\Lambda$ . Then  $H$  is constant and  $v \equiv 0$ . In particular the family  $u(y, s) = b(s)$  with  $b$  non-constant is not a solution on  $\Lambda$ . [Proved.]*

*Proof.*  $H$  is harmonic on the closed manifold  $\Lambda$ , hence constant (maximum principle), so  $\nabla H \equiv 0$ . For the special case  $H(y) = b(s) \cdot y$  the function  $H$  is not single-valued on  $\Lambda$  unless  $b = 0$ ; equally, the pressure  $q = -b'(s) \cdot y$  of  $u = b(s)$  is not single-valued on  $\Lambda$  unless  $b' = 0$ .  $\square$

**Lemma 36.15** (Torus form of Theorem 36.6, uniform in the period). *For every  $k, l \geq 0$  there are  $\varepsilon(k, l) > 0$  and  $C(k, l)$ , independent of  $L$ , with the following property. Let  $L > 0$  and let  $w$  be a smooth solution of the Navier–Stokes equations on  $(L\mathbb{T}^3) \times [0, T]$  with single-valued pressure, and put  $A = \|w(\cdot, 0)\|_{L^\infty}$ . Then for  $0 < t \leq \min(T, \varepsilon(k, l)A^{-2})$ ,*

$$\|\nabla^k \partial_t^l w(\cdot, t)\|_{L^\infty(L\mathbb{T}^3)} \leq C(k, l) A t^{-\frac{k}{2}-l}.$$

[Proved, by the argument of [17, §4] with the Oseen kernel periodised; the one estimate used is  $L$ -uniform.]

*Proof.* The proof of Theorem 36.6 in [17] is a perturbation series around the heat extension: with  $B$  the bilinear form built from the Oseen kernel  $K_{ijk}$ , the equation is  $w = W + B(w, w)$  and the entire argument rests on [17, (4.5)],  $\|B(u, v)\|_{L^\infty} \leq C\sqrt{T}\|u\|_{L^\infty}\|v\|_{L^\infty}$ , which in turn rests only on the kernel bound  $\int |K(\cdot, t)| \leq Ct^{-1/2}$ . On  $L\mathbb{T}^3$  the corresponding kernel is the periodisation  $K^{(L)}(z, t) = \sum_{n \in \mathbb{Z}^3} K(z + Ln, t)$ , and

$$\int_{L\mathbb{T}^3} |K^{(L)}(z, t)| dz \leq \sum_{n \in \mathbb{Z}^3} \int_{L\mathbb{T}^3} |K(z + Ln, t)| dz = \int_{\mathbb{R}^3} |K(z, t)| dz \leq Ct^{-1/2},$$

uniformly in  $L$ . The Leray projector on  $L\mathbb{T}^3$  is bounded on every  $L^p$ ,  $1 < p < \infty$ , with constants independent of  $L$  (it is a Fourier multiplier with the same symbol as on  $\mathbb{R}^3$ ), and the pressure is fixed by the mean-zero normalisation, which is legitimate by Lemma 36.14. A smooth solution with single-valued pressure satisfies the corresponding Duhamel identity by variation of constants, and the  $L^\infty$  mild solution with datum  $w(\cdot, 0)$  is unique [17, §3]; hence  $w$  is the fixed point, and (180) applies verbatim with the periodised kernel.  $\square$

**Theorem 36.16** ((**R**) is a theorem, in a stronger form). *Assume the standing Type-I bounds of §27 (line 5565), i.e. both  $M(t) \leq C_I/(T-t)$  and  $\sup_{\mathbb{T}^3} |u(\cdot, t)| \leq C_I(T-t)^{-1/2}$ , and let  $u_\lambda$  be as in (152). Then for every  $k, l \geq 0$  there is  $C_{k,l}(c_I)$  such that*

$$\|\nabla^k \partial_s^l u_\lambda(\cdot, s)\|_{L^\infty(\Lambda_\lambda)} \leq C_{k,l}(c_I) |s|^{-\frac{k+1}{2}-l} \quad (183)$$

for every  $\lambda \in (0, 1]$  and every  $s < 0$  with  $\frac{3}{2}|s| \leq T/\lambda^2$ . In particular (**R**) holds, with constants  $C_k(c_I) = C_{k,0}(c_I)$ , without the factor  $(1+R)$  and without any dependence on  $S_2$ . [Proved.]

*Proof.* Fix  $s_0 < 0$  with  $\frac{3}{2}|s_0| \leq T/\lambda^2$  and write  $S = |s_0|$ . By (153), for every  $s$  with  $|s| \in [S, \frac{3}{2}S]$  we have  $\|u_\lambda(\cdot, s)\|_{L^\infty(\Lambda_\lambda)} \leq c_I |s|^{-1/2} \leq c_I S^{-1/2} =: A$ , and the whole time interval lies inside the domain of  $u_\lambda$  by the standing restriction on  $s_0$ .

Set  $\tau = \min(\varepsilon(k, l)c_I^{-2}, \frac{1}{2})S$ , so that  $\tau \leq \frac{1}{2}S$  and  $\tau \leq \varepsilon(k, l)A^{-2}$ . Apply Lemma 36.15 on  $L\mathbb{T}^3$  with  $L = \lambda^{-1}$  to  $w(\cdot, t) = u_\lambda(\cdot, s_0 - \tau + t)$ ,  $t \in [0, \tau]$ , whose initial datum has  $\|w(\cdot, 0)\|_{L^\infty} \leq A$ . At  $t = \tau$ ,

$$\|\nabla^k \partial_s^l u_\lambda(\cdot, s_0)\|_{L^\infty} \leq C(k, l) A \tau^{-\frac{k}{2}-l} = C(k, l) c_I S^{-\frac{1}{2}} \left( \min(\varepsilon(k, l)c_I^{-2}, \frac{1}{2})S \right)^{-\frac{k}{2}-l} = C_{k,l}(c_I) S^{-\frac{k+1}{2}-l},$$

which is (183). The bound is uniform in  $y \in \Lambda_\lambda$  because (179) is, so no factor  $(1+R)$  appears; and it depends on  $s_0$  only, so no  $S_2$  appears. Since  $S \geq S_1$  on  $[-S_2, -S_1]$  and the right-hand side is decreasing in  $S$ , (R) follows.

*Dimensional check.* In units  $\nu = 1$  (Remark 33.1),  $[\nabla^k \partial_s^l u] = L^{-1-k-2l}$  and  $[|s|^{-(k+1)/2-l}] = L^{-(k+1)-2l}$ ; the two agree, and  $c_I$  is dimensionless.

*Independent confirmation without Lemma 36.15.* The qualitative content — that every spatial derivative of  $u_\lambda$  is locally bounded — is Theorem 36.3 applied on any coordinate ball, with  $s' = s = \infty$  in the strict range (Remark 36.2), needing neither periodicity nor pressure nor mildness. Lemma 36.15 is used only to make the constants scale-invariant and to supply  $l \geq 1$ .  $\square$

**Corollary 36.17** (Proposition 33.8(d) is Proved). *Let  $U \in \mathcal{A}(c_I)$ . Then  $\|\nabla U(\cdot, s)\|_{L^\infty(\mathbb{R}^3)} \leq C_1(c_I)|s|^{-1}$  for every  $s < 0$ ; i.e. Proposition 33.8(d) holds unconditionally, and its label changes from Proved-modulo-(R) to Proved. [Proved.]*

*Proof.* Two independent arguments; the second needs nothing beyond Definition 33.2.

(1) *Via the family.* If  $U$  arises from Proposition 33.6, pass (183) with  $k = 1$ ,  $l = 0$  through the locally uniform convergence.

(2) *Via Theorem 36.8, for an arbitrary member of the class.* Fix  $s_0 < 0$ , set  $r = |s_0|^{1/2}$ , and define

$$V(y, s) := c_I^{-1} r U(c_I^{-1} r y, r^2(c_I^{-2} s - 1)).$$

This is a composition of three Navier–Stokes symmetries — the parabolic scaling of Lemma 33.3 applied twice (with factors  $r$  and  $c_I^{-1}$ ) and one translation in time — so  $V$  is a smooth divergence-free solution, hence a distributional one in the sense of [22, Def. 2.1]. Its time argument is  $r^2(c_I^{-2} s - 1) < 0$  for every  $s \leq 0$ , so  $V$  is defined on all of  $\mathbb{R}^3 \times (-\infty, 0)$ , and

$$|V(y, s)| \leq c_I^{-1} r \cdot c_I |r^2(c_I^{-2} s - 1)|^{-1/2} = |c_I^{-2} s - 1|^{-1/2} \leq 1 \quad (s \leq 0),$$

since  $c_I^{-2} s \leq 0$  forces  $|c_I^{-2} s - 1| \geq 1$ . So  $V$  is a bounded ancient solution normalised as in [22, (4.8)], with no mildness and no pressure integrability assumed. Theorem 36.8 gives, for any  $m > 1$ ,  $\sup_{z_0} \|V_{t_0}\|_{W_m^{2,1}(Q(z_0, 1))} \leq c(m, 3)$ ; taking  $m > 5$  and using the parabolic embedding  $W_m^{2,1}(Q(z_0, 1)) \hookrightarrow C^{1+\alpha, (1+\alpha)/2}$  with  $\alpha = 1 - 5/m$  yields  $\|\nabla V\|_{L^\infty(Q(z_0, 1))} \leq c$  uniformly in  $z_0 = (x_0, t_0)$  with  $t_0 \leq 0$ , the shift  $b_{t_0}$  being spatially constant so that  $\nabla V_{t_0} = \nabla V$ . Now  $\nabla V(y, s) = c_I^{-2} r^2 (\nabla U)(c_I^{-1} r y, r^2(c_I^{-2} s - 1))$ , and at  $s = 0$  the inner time argument is exactly  $-r^2 = s_0$ ; taking  $t_0 = 0$  and the supremum over  $y$ ,

$$c_I^{-2} r^2 \|\nabla U(\cdot, s_0)\|_{L^\infty(\mathbb{R}^3)} \leq c, \quad \text{i.e.} \quad \|\nabla U(\cdot, s_0)\|_{L^\infty(\mathbb{R}^3)} \leq c c_I^2 |s_0|^{-1},$$

which is the claim with  $C_1(c_I) = c c_I^2$ .

Neither argument uses the pressure, a local energy, or a smallness hypothesis.  $\square$

**Remark 36.18** (The mechanism, corrected). The proof of Proposition 33.8(d) explains the disappearance of  $\mathcal{L}$  by: “ $U$  has, by construction, no structure below the scale  $|s|^{1/2}$  that is not already visible at scale  $|s|^{1/2}$ ”. That is a heuristic, it is not proved anywhere in this document, and it is not the reason. The reason is Remark 36.2: the available hypothesis is an  $L_t^\infty L_x^\infty$  velocity bound, which is *strictly subcritical* for Serrin’s criterion, so  $\nabla U$  is obtained by parabolic smoothing from  $U$  — not by a Calderón–Zygmund bound on  $\omega_U$  — and parabolic smoothing has no endpoint failure. The correct one-line statement is: *the logarithm is not removed on the profile; it is never incurred, because the profile route does not pass through the Calderón–Zygmund operator at all*. The conclusion of Proposition 33.8(d) is unaffected; only its stated reason is.

### 36.5 Verdict, residue, and what does *not* follow

**Verdict: (i). (R)** is a theorem. Precisely: under the standing hypotheses of §27 — *both* Type-I bounds of line 5565, and smoothness of  $u$  on  $\mathbb{T}^3 \times [0, T)$  — the estimate displayed at line 6785 holds, with constants depending only on  $c_I$ , by Theorem 36.16; and the strengthened form (183), which additionally controls  $l$  time derivatives, holds by the same proof. The hypotheses under which the cited theorems apply are checked in Theorem 36.16 and Corollary 36.17. **(R)** is removed from the hypothesis list of §33.

The three traps named for this question are disposed of as follows. *Smallness vs. boundedness*: does not arise, because the route is not  $\varepsilon$ -regularity (Remark 36.2, Remark 36.4). *Circularity*: none of Theorems 36.3, 36.6, 36.8, 36.9 assumes any conclusion of §33; they assume boundedness of the velocity and nothing about alignment,  $\sigma^*$ , or nontriviality. *Interior vs. global*: **(R)** is interior in time and global in space (Remark 36.1); the theorem proved respects that, and asserts nothing at  $s = 0$ , where (153) degenerates.

**Residue.** Four items, none of which is **(R)**.

- (1) **The saturation proviso, and it is load-bearing.** Proposition 33.8(d) states two things: the bound  $\|\nabla U\|_{L^\infty} \leq C_1(c_I)|s|^{-1}$ , now proved, *and* the consequence “ $\|\nabla U(\cdot, s)\|_{L^\infty} \leq C(c_I)M_U(s)$  at every time with  $M_U(s) \geq c_I^{-1}|s|^{-1}$  (i.e. whenever the Type-I upper bound is saturated up to a factor)”. Part (e) uses the *second* form — it replaces  $\mathcal{L}$  inside Lemma 28.1, where the gradient is measured against  $M_U$  and  $r^* = M_U(s_0)^{-1/2}$ . The two forms differ by the factor  $(|s|M_U(s))^{-1}$ , which is  $O(1)$  only on the saturated set. *Whether  $M_U(s) \gtrsim |s|^{-1}$  holds for all  $s < 0$ , or only on a set, is not settled by (R) and is not settled here.* Under **(N)**,  $M_U(s) > 0$  for every  $s < 0$  (Proposition 33.6), but positivity is weaker than saturation. This is the sharpest surviving gap on the profile branch and it is a good target: it is a statement about one scalar function of one variable on an ancient solution. *Amended at S50.* It is a sharply posed target and, on the evidence of §37, a poor one. The strength (e) actually needs is saturation at the single reference time  $s_0$  (Lemma 37.4), not at every  $s$ ; (e)’s only mathematical consumer is Proposition 33.20(b), which itself has none (Observation 37.6) and can be discharged without saturation at the Type-I scale (Corollary 37.26); and on the global branch logarithm-free saturation is equivalent to Wall 1 (Theorem 37.21). The status of (e) itself is unchanged.
- (2) **(R) as used exceeded (R) as stated, and the excess is now supplied elsewhere.** See Remark 36.20(c). No circularity results, because the replacement input

(Theorem 36.9, or (183) with  $l \geq 1$ ) is independent of §33.

- (3) **Nothing else in the S39 ledger moves.** (N), (N') and (A-up) are untouched; Conjecture 33.25 is untouched; Remark 33.30 stands verbatim, including its ceiling (“no Type-I singularity ... strictly less than the Clay problem”) and its Type-II reservation. Wall 3 is unaffected. In the ledger of §33 the rows reading “mod (R)” become *Proved*; the rows reading “mod (R),(N)”, “mod (R),(N')” and “mod (R),(A-up)” lose only the (R).
- (4) **The  $\omega$ -only route is closed, permanently.** Proposition 36.12: the velocity Type-I bound cannot be derived from the vorticity Type-I bound and the energy. This is not a residue of the present argument but a constraint on future ones.

*Observation 36.19* (The same estimate applies to  $u$  itself — recorded, not claimed). Lemma 36.15 applies verbatim to  $u$  on  $\mathbb{T}^3$  (take  $L = 1$ , restart at  $t - \tau$  with  $\tau = \min(\varepsilon(1, 0)c_I^{-2}, \frac{1}{2})(T - t)$ ), giving, under the standing hypotheses of §27,

$$\|\nabla u(\cdot, t)\|_{L^\infty(\mathbb{T}^3)} \leq C(c_I)(T - t)^{-1}, \quad \text{logarithm-free.}$$

This does *not* remove Wall 1. Every estimate of §27–§31 needs  $\nabla u$  measured against  $M(t)$ , not against  $(T - t)^{-1}$ , and the two agree only where the Type-I upper bound is saturated — the same proviso as residue (1) above, one level down. The observation is recorded here because it is cheap, because it identifies *saturation*, not the logarithm, as the quantity both branches actually need, and because a future session should test it before attacking either wall separately. It is not a revision of §44.2 and is not used anywhere in this document. *Amended at S50*. The test was carried out in §37 and the diagnosis is half right: saturation *is* what both branches need, and it is *not* a separate, cheaper question. By Theorem 37.21 the bound displayed above, combined with logarithm-free saturation, is equivalent to removing Wall 1 in the sense of §44.2(a), losslessly in the constants. The observation is used in that proof — in the direction saturation  $\Rightarrow$  Wall 1 only — and nowhere else.

**Remark 36.20** (Three places where this document’s own text needs correcting). (a) The status note of (R) (line 6785ff) gives caveat (ii) as: the  $\lambda$ -uniformity “must be checked against the fact that  $u_\lambda$  has *no* uniform global energy bound”. The premise is true and the inference is not: the  $L^\infty$ -based local theory ([17, §4], [22, Thm. 2.4]) uses no energy at any point, global or local. Caveat (ii) is void.

(b) The same note gives caveat (i) as the time-regularity gap. It is correctly identified — Remark 36.5 confirms it is sharp for general distributional solutions — but it is not a caveat on (R), which asks only for spatial derivatives, and it is discharged for the objects this document actually forms by Theorem 36.9.

(c) The proof of Proposition 33.6 bounds  $\partial_s u_\lambda$  using “the local pressure bound supplied by (R)”. (R) supplies no pressure bound: its statement (line 6785) bounds  $\nabla^k u_\lambda$  only. This is a genuine gap between the hypothesis as stated and the hypothesis as used, and it is exactly the parasitic-solution issue in disguise. The repair does not need a pressure bound at all: (183) with  $l = 1$  bounds  $\partial_s u_\lambda$  directly, and the mildness of the limit — the property that rules out  $U \mapsto U + b(s)$  — is supplied by Theorem 36.9. The sentence in the proof should be replaced by a reference to (183).

| Item                                              | Status before S48                | Status after S48             |
|---------------------------------------------------|----------------------------------|------------------------------|
| <b>(R)</b> scale-uniform interior regularity      | <b>Conditional</b> (Serrin-type) | <b>Proved</b> (Thm. 36.16)   |
| Prop. 33.8(d), profiles $\mathcal{L}$ -free       | Proved mod (R)                   | <b>Proved</b> (Cor. 36.17)   |
| Prop. 33.8(e), $\nu\langle\sigma^*\rangle \leq C$ | Proved mod (R)                   | Proved mod <i>saturation</i> |
| Prop. 33.6, profile extraction                    | Proved mod (R),(N)               | Proved mod (N)               |
| Lem. 33.11, $\sigma^*$ equality                   | Proved mod (R),(N)               | Proved mod (N)               |
| Lem. 33.16, aligned $\Rightarrow$ 2D              | Proved mod (R)                   | <b>Proved</b>                |
| Thm. 33.18, aligned horn                          | Proved mod (R),(A-up)            | Proved mod (A-up)            |
| Wall 2 (§44.3)                                    | OPEN; relocated into (R)         | WITHDRAWN                    |
| Wall 1 (§44.2)                                    | OPEN, structural                 | OPEN, structural (unchanged) |
| Wall 3 (§44.4)                                    | OPEN, outside methods            | unchanged                    |
| Type-II exclusion                                 | Open                             | unchanged                    |

**Remark 36.21** (Scope discipline). This section changes the status of one hypothesis and one proposition. It does not close either horn of the pincer, does not bear on  $\sigma^*$ -decay, does not touch Type-II, and does not bring the programme nearer to the Clay problem than Remark 33.30 already states. What it does is remove a hypothesis that had been carried for nine sessions and was, on inspection, classical — and, in doing so, replace the question “is the profile route logarithm-free?” with the sharper question of residue (1): *is the Type-I upper bound saturated?*

## 37 Saturation of the Type-I Bound (Session S50)

Session S48 removed hypothesis **(R)** and, in doing so, left two statements standing on a single common proviso: the second clause of Proposition 33.8(d), which Part (e) uses, and Observation 36.19, which records a logarithm-free bound on  $\|\nabla u\|_{L^\infty}$  that §27–§31 cannot consume. Remark 36.21 named the resulting question — *is the Type-I upper bound saturated?* — and proposed it as a target cheaper and better defined than either wall. This section settles what can be settled and reports, as the principal finding, that the proposal was wrong: log-free saturation is not a cheaper target than Wall 1, because *it is the same target* (Theorem 37.21).

Three things are proved here. First, an audit of who actually needs saturation and in what strength (§37.2), which finds the requirement both weaker and less consequential than S48 recorded. Second, a theorem: under the standing hypotheses the Type-I vorticity bound *is* saturated up to a single logarithm, at every time and not merely along a sequence (Theorem 37.19); and a corollary that this is worth nothing, being a restatement of an estimate the document already carried (Corollary 37.22). Third, the equivalence with Wall 1, in both directions, on both branches (Theorem 37.21, Proposition 37.23).

One route is opened rather than closed: Proposition 37.25 discharges the only mathematical consumer of Part (e) without saturation, by running the estimate at the Type-I scale  $|s_0|^{1/2}$  instead of the amplitude scale  $M_U(s_0)^{-1/2}$ .

**Standing hypotheses.** Throughout this section  $u$  is smooth on  $\mathbb{T}^3 \times [0, T]$  with putative first singularity at  $T$  and satisfies the standing Type-I bounds of §27 (line 5565): *both*  $M(t) = \|\omega(\cdot, t)\|_{L^\infty} \leq C_I/(T - t)$  and  $\sup_{\mathbb{T}^3} |u(\cdot, t)| \leq C_I(T - t)^{-1/2}$ . We write  $c_I = C_I$  for

consistency with §36, normalise  $|\mathbb{T}^3| = 1$  and  $\nu = 1$  (as in Remark 33.1), and set

$$\Lambda(t) := \log \frac{1}{T-t}, \quad T-t < 1.$$

By Proposition 36.12 the velocity bound is an *independent* hypothesis and is used as such; nothing below attempts to derive it from the vorticity bound.

### 37.1 What saturation is, and what is free

**Definition 37.1** (Saturation). Define the *Type-I amplitude*

$$\Phi(t) := (T-t) M(t) \in (0, C_I],$$

the torus analogue of the profile amplitude  $\Phi$  of Lemma 33.21. The *saturated set* at level  $c > 0$  is  $\mathcal{S}_c := \{t < T : \Phi(t) \geq c\}$ . We distinguish:

**(Sat-1) Log-free saturation.** There are  $c = c(c_I) > 0$  and  $t^* < T$  with  $\Phi(t) \geq c$  for *all*  $t \in (t^*, T)$ ; equivalently  $M(t) \geq c(T-t)^{-1}$  there.

**(Sat-2) Saturation up to a logarithm.** There are  $c > 0$ ,  $t^* < T$  with  $\Phi(t) \geq c/(1 + \Lambda(t))$  for all  $t \in (t^*, T)$ .

**(Sat-3) Sequential saturation.**  $\limsup_{t \uparrow T} \Phi(t) \geq c$ .

**(Sat-4) Saturation on a set.**  $\mathcal{S}_c$  accumulates at  $T$ , or meets infinitely many dyadic shells  $(T - 2^{-k-1}, T - 2^{-k})$ .

The profile forms are the same with  $U \in \mathcal{A}(c_I)$ ,  $\Phi_U(s) = |s|M_U(s)$  and  $|s|$  in place of  $T-t$ , asserted as  $s \rightarrow -\infty$  (the regime Proposition 33.20 inhabits).

**Remark 37.2** (Velocity saturation is free; it is the vorticity that is open). The name “Type I” is the assertion that the velocity bound is saturated. Cheskidov, Dai and Palasek state it thus [34, §2.2]:

*A basic consequence of Leray’s local theory is that  $\|u(t)\|_{L^\infty}$  diverges at least as fast as  $(T_* - t)^{-\frac{1}{2}}$  before a finite-time blow-up at  $t = T_*$ . A blow-up that saturates this critical lower bound is referred to as Type I.*

So the standing hypothesis  $\sup |u| \leq C_I(T-t)^{-1/2}$  together with the classical lower bound gives  $\|u(\cdot, t)\|_{L^\infty} \asymp (T-t)^{-1/2}$  exactly, with no gap. The question of this section is *not* of that kind. It asks for the matching lower bound on  $\|\omega\|_{L^\infty}$ , whose Type-I upper bound is one Biot–Savart degree away; and the distance between the two is precisely the distance Wall 1 measures. That observation is not a proof, but it already indicates where the difficulty must live, and Theorem 37.21 confirms it.

## 37.2 Audit: who needs saturation, and how much

The S48 residue (§36.5, item (1)) records the proviso as “load-bearing”. That is correct about the step (d) → (e) and incorrect about (e)’s own weight. The following audit re-derives every line number against the present file.

**Proposition 37.3** (Audit of the three users). *In the present document saturation is required at exactly three places, in the following strengths.*

- (a) **Proposition 33.8(d), second clause** (lines 6983–6985). *The clause  $\|\nabla U(\cdot, s)\|_{L^\infty} \leq C(c_I)M_U(s)$  is asserted exactly on  $\{s : M_U(s) \geq c_I^{-1}|s|^{-1}\}$ , i.e. on the saturated set. It is not a hypothesis of the proposition but a restriction on its conclusion, and it is equivalent to saturation at  $s$ : given the first clause (Corollary 36.17) the second holds at  $s$  iff  $\Phi_U(s) \gtrsim 1$ .*
- (b) **Proposition 33.8(e)** (lines 6986–6995), through Lemma 28.1(a) (line 5852), where  $\|\nabla u\|_{L^\infty} M^2(r^*)^5$  is converted using  $M(r^*)^2 = 1$  with  $r^* = M_U(s_0)^{-1/2}$ . *The requirement is saturation at the single reference time  $s_0$ , not throughout  $Q(r^*)$ : by Lemma 37.4 saturation at  $s_0$  propagates over the whole time extent of the cube.*
- (c) **Observation 36.19** (line 10408). *Its bound  $\|\nabla u\|_{L^\infty} \leq C(c_I)(T - t)^{-1}$  is usable by §27–§31 at time  $t$  iff  $t \in \mathcal{S}_c$ . Every estimate there measures  $\nabla u$  against  $M(t)$ .*

[Proved — (a) and (c) by inspection of the cited lines; (b) is Lemma 37.4.]

**Lemma 37.4** (Saturation at one time propagates across the cube). *Let  $U \in \mathcal{A}(c_I)$ , let  $s_0 < 0$  satisfy  $\Phi_U(s_0) \geq c$ , and put  $r^* = M_U(s_0)^{-1/2}$ ,  $h = (r^*/2)^2 = \frac{1}{4}M_U(s_0)^{-1}$ . Then for all  $s \in [s_0 - h, s_0]$ ,*

$$M_U(s) \geq \left(1 + \frac{1}{4c}\right)^{-C_1(c_I)} M_U(s_0),$$

where  $C_1(c_I)$  is the constant of Corollary 36.17. The identical statement holds on  $\mathbb{T}^3$  with  $(T - t)$  for  $|s|$  and Observation 36.19 for Corollary 36.17. [Proved.]

*Proof.* Saturation at  $s_0$  gives  $h = \frac{1}{4}M_U(s_0)^{-1} \leq |s_0|/(4c)$ , so  $|s| \leq |s_0|(1 + \frac{1}{4c})$  on the interval. By Lemma 33.21(c) (Hamilton’s trick applied to (141); the supremum is attained here because  $M_U(s) > 0$  and the estimate is local) and  $w \geq 0$ ,

$$\frac{d}{ds} \log M_U \leq \alpha_{\max} \leq \|\nabla U(\cdot, s)\|_{L^\infty} \leq C_1(c_I)|s|^{-1},$$

the last step by Corollary 36.17. Integrating from  $s$  to  $s_0$  gives  $\log M_U(s_0) - \log M_U(s) \leq C_1 \log(|s|/|s_0|) \leq C_1 \log(1 + \frac{1}{4c})$ .  $\square$

**Remark 37.5** (The requirement is weaker than S48 recorded — and this is the correct reading of the proviso). Proposition 37.3(b) with Lemma 37.4 says that (e) holds *at every  $s_0$  of the saturated set*. So the saturated set is not merely where (d)’s second clause is stated: it is exactly the set of times at which (e)’s conclusion (154) is available. Consequently (**Sat-4**) — not (**Sat-1**) — is the form (e) needs, provided its consumers only sample times.



*Observation 37.6* (Part (e) is a leaf). In the present document, (154) and Proposition 33.8(e) are invoked at exactly one mathematical location: Proposition 33.20(b) (lines 7364–7370), whose proof cites “Proposition 33.8(e) plus Lemma 33.3” (line 7384). Every other occurrence of (e) is a ledger row or a summary sentence (lines 7629, 7661, 7696, 9775, 10466, 10726). Proposition 33.20 is itself cited only three times (lines 7389, 7634, 7671), all three rhetorical: a transitional sentence, a summary bullet and a ledger row. It has no mathematical consumer.

*Consequence.* (e) is a leaf of the dependency graph, and the statement it supports is an *impossibility* result — “the naive infinite-past argument fails” — so if saturation were false, the effect on the programme would be to *weaken an obstruction*, not to lose a tool. The live profile machinery (Lemma 33.21, Proposition 33.23, Conjecture 33.25) is saturation-free: the proof of Proposition 33.23 uses only  $\Phi_U \leq c_I$  and  $\Phi_U(\tau_0) > 0$ , i.e. the Type-I *upper* bound and nontriviality. [*Proved, by exhaustive search of the cross-reference graph.*]

**Remark 37.7** (Two inaccuracies in the document’s own handling of saturation). (a) The proof of Theorem 28.2 (line 5883) writes “on  $\{|\omega| \geq M/2\} \cap \{\eta = 1\}$  we have  $m^2 \geq M^2/4$ ” and then removes  $M^2$  from a space-time integral, although the cutoff  $\chi(|\omega(x, t)|/M(t))$  of Definition 27.2 makes  $M$  time-dependent while  $r^*$  is built from  $M(t_0)$ . The step therefore carries an unstated requirement,  $M(t) \gtrsim M(t_0)$  on  $Q(r^*/2)$ , which Type I does not supply. It is not an additional gap: Lemma 37.4 shows it follows from saturation at  $t_0$  together with the logarithm-free gradient bound. But it was not stated, it is inherited verbatim by Proposition 33.8(e), and it is a second, independent reason (e) needs saturation.

(b) Lemma 27.6 (line 5701) uses  $\|\nabla u\|_{L^\infty} \lesssim M \log(e + \|u\|_{H^3}/M)$  with no citation, and Wall 1 (§44.2) rests on it. The inequality is Beale–Kato–Majda [28, (15)], refined by Kozono–Taniuchi [14] — the latter *is* in the bibliography but is never cited for this. BKM’s literal form is  $\|\nabla u\|_{L^\infty} \leq C\{(1 + \log^+ \|u\|_{H^3})\|\omega\|_{L^\infty} + \|\omega\|_{L^2}\}$ ; the document’s  $\mathcal{L}$  is the scale-invariant variant. Under the standing hypotheses the two agree up to constants (Lemma 37.18), so nothing downstream changes; the citation is added here.

### 37.3 What the literature actually proves

Statements are quoted, not recalled. Where a display failed to extract from a scanned source this is said explicitly rather than reconstructed.

*Fact 37.8* (Leray; Giga). The classical lower bounds, as recalled verbatim in [32, §1]: “ $\|u(t)\|_{L^p} \geq c(T - t)^{-(1-3/p)/2}$ ” for  $p \in [3, \infty)$  and “ $\|u(t)\|_{L^\infty} \geq c(T - t)^{-1/2}$ ”. Their provenance, quoted from the same source: “These blow-up estimates were first stated for the  $L^p$  spaces,  $p > 3$ , without proof by Leray in his remarkable paper [29], and proved by Giga in [30] via semigroup theory.” (In this quotation the source’s reference numerals [9] and [7] have been replaced by the corresponding keys of this bibliography; nothing else is altered.) These hold for *all*  $t$  in a left neighbourhood of  $T$ ; they are not limsup statements. The mechanism is local well-posedness run backwards, and the exponent is the scaling deficit of the norm halved.

*Fact 37.9* (Cortissoz–Montero, optimal Sobolev rate on the torus). [32, Thm. 2.1], verbatim: “Let  $u(x, t)$  be a solution Navier–Stokes equations whose maximum interval of existence is  $(0, T)$ ,  $0 < T < \infty$ , and such that  $u \in C((0, T), \dot{H}^s(\mathbb{T}^3) \cap \dot{H}^{s+1}(\mathbb{T}^3))$ , with  $\frac{1}{2} < s < \frac{5}{2}$ . Then the following estimate holds  $C_s t^{-\frac{1}{2}(s-\frac{1}{2})} \leq \|u(T - s)\|_{\dot{H}^s(\mathbb{T}^3)}$ .” The paper uses  $s$  for both the Sobolev index and the elapsed time in this display; read unambiguously it is

$$\|u(\cdot, t)\|_{\dot{H}^\sigma(\mathbb{T}^3)} \geq C_\sigma (T - t)^{-\frac{2\sigma-1}{4}}, \quad \frac{1}{2} < \sigma < \frac{5}{2},$$

for *all*  $t$  in the interval, matching the exponent  $-(2s-1)/4$  the same paper attributes to Robinson–Sadowski–Silva [31]. Taking  $\sigma = 1$  and  $\|\nabla u\|_{L^2} = \|\omega\|_{L^2}$  for divergence-free fields on  $\mathbb{T}^3$ :

$$M(t) \geq \|\omega(\cdot, t)\|_{L^2(\mathbb{T}^3)} \geq C(T-t)^{-1/4} \quad \text{for all } t < T. \quad (184)$$

(With  $|\mathbb{T}^3| = 1$ ; otherwise a factor  $|\mathbb{T}^3|^{-1/2}$  appears.)

*Fact 37.10* (Beale–Kato–Majda). [28, Theorem], verbatim: “Let  $u$  be a solution of Euler’s equations as described above, and suppose there is a time  $T_*$  such that the solution cannot be continued in the class (2) to  $T = T_*$ . Assume that  $T_*$  is the first such time. Then  $\int_0^{T_*} |\omega(t)|_{L^\infty} dt = \infty$ , and in particular [...]” The paper adds: “Similar statements hold for the Navier–Stokes equations; the arguments below apply equally well in this case.” The displayed “in particular” consequence did not extract from the scanned original and is therefore not quoted from it; it is quoted instead from [33, (1.4)], which renders BKM as  $\int_0^{T_*} \|\omega(t)\|_{L^\infty} = \infty$  “and, consequently,”  $\limsup_{t \rightarrow T_*^-} \|\omega(t)\|_{L^\infty} = \infty$ . **BKM is an integral-divergence statement whose pointwise shadow is a lim sup.**

**Lemma 37.11** (BKM does not deliver saturation on any set). *Non-integrability of  $M$  implies none of (Sat-1)–(Sat-4), even given the Type-I upper bound.* [Proved.]

*Proof.* Pass to logarithmic time  $x = \log \frac{1}{T-t}$  and set  $g(x) = \Phi(t) = (T-t)M(t)$ , so that  $\int_{t_0}^T M dt = \int_{x_0}^\infty g dx$  and the Type-I upper bound reads  $0 < g \leq C_I$ . The pair of conditions “ $\int_{t_0}^\infty g = \infty$  and  $g \leq C_I$ ” is satisfied by  $g(x) = 1/x$ , for which  $g \rightarrow 0$ : (Sat-3) fails, hence so do (Sat-1) and (Sat-2); and for every  $c > 0$  the set  $\{g \geq c\}$  is the bounded interval  $[x_0, 1/c]$ , so (Sat-4) fails as well. Thus BKM constrains  $\Phi$  only through a divergent integral in logarithmic time, which is compatible with  $\Phi \rightarrow 0$ .  $\square$

*Fact 37.12* (Ingimarsen–Kukavica: the best known vorticity rate, and it is a limsup with a logarithm). [33, §1], on the state of the art for *Euler*: “However, not much is known regarding the rate at which either of the two quantities (1.3) or (1.4) blow up.” Their Theorem 2.1, verbatim: under the hypothesis that  $T_*$  is the first time the solution leaves the class (1.2),

$$\int_0^t \|\omega(s)\|_{L^\infty} ds \geq \frac{1}{C} \log \log \frac{1}{T_* - t}, \quad T_* - t < \frac{1}{C},$$

“and, moreover,”

$$\limsup_{t \rightarrow T_*^-} (T_* - t) \log \left( \frac{1}{T_* - t} \right) \|\omega(t)\|_{L^\infty} \geq \frac{1}{C}.$$

Two features are decisive for us. (i) The bound is a lim sup, and the authors are deliberate about this: in the same paper Theorems 2.5–2.7 state lower bounds on  $\|D^k \omega\|_{L^\infty}$  “for all  $t \in [0, T_*)$ ”, so the lim sup in Theorem 2.1 is not a stylistic default. (ii) It is weaker than exponent 1 by exactly one logarithm: it asserts  $\|\omega\|_{L^\infty} \geq c / ((T_* - t) \log \frac{1}{T_* - t})$  along a sequence, not  $c / (T_* - t)$ .

Their mechanism is the vorticity ODE. Verbatim from their proof: “Following Lagrangian trajectories, we obtain  $\frac{d}{dt} \|\omega\|_{L^\infty} \lesssim \|Du\|_{L^\infty} \|\omega\|_{L^\infty}$ ”, combined with “ $\|Du\|_{L^\infty} \leq \tilde{C}(1 + \|\omega\|_{L^\infty}(1 + \log^+ \|u\|_{H^3}))$ , see [14]”. Lacking a Type-I hypothesis they must close  $\|u\|_{H^3}$  by Grönwall, obtaining  $1 + \log^+ \|u(t)\|_{H^3} \lesssim \exp(\tilde{C} \int_0^t \|\omega\|_{L^\infty})$ , which is what degrades the conclusion to log log and to a lim sup. §37.4 runs the same ODE *with* the Type-I hypothesis and does better on both counts.

*Fact 37.13* (Cheskidov–Dai–Palasek: a constructed Type-I solution whose vorticity lower bound is sequential). [34, Thm. 1.1] constructs, on  $\mathbb{T}^d$  for any smooth divergence-free datum, a weak solution blowing up instantaneously at  $T_*$  with, verbatim, “for some  $c > 0$ ,  $\|u(t_n)\|_{L^\infty} \geq \frac{c}{\sqrt{t_n - T_*}}$ ,  $\|\nabla \wedge u(t_n)\|_{L^\infty} \geq \frac{c}{t_n - T_*}$  along a sequence of times  $t_n \rightarrow T_+$ ” (Property 2), while “the blow-up is of Type I:  $\|u(t)\|_{L^\infty} \leq \frac{C}{\sqrt{t - T_*}}$ ,  $\|\nabla u(t)\|_{L^\infty} \leq \frac{C}{t - T_*}$  for all  $t \in (T_*, T]$ ” (Property 4).

*What this is and is not.* It is *not* a counterexample to saturation: the construction is a forward (instantaneous) blow-up, the bounds run for  $t > T_*$ , and the Navier–Stokes equations are not time-reversible, so the maximum-principle argument of Lemma 37.16 — which runs *toward* the singularity — has no image in that setting. It is evidence of a different kind: in a bespoke example engineered to be Type I in both  $u$  and  $\nabla u$ , the authors assert the *upper* bounds at all times and the *vorticity lower* bound only along a sequence. If a general pointwise vorticity lower bound were available from Type I, it would have been free to them.

**Remark 37.14** (The limsup/liminf finding, stated plainly). Collecting Facts 37.8–37.13: every known *pointwise* (all-times) lower bound is a *velocity* or Sobolev bound whose exponent is the halved scaling deficit —  $1/2$  for  $\|u\|_{L^\infty}$ ,  $(2\sigma - 1)/4$  for  $\dot{H}^\sigma$ , hence  $1/4$  for  $\|\omega\|_{L^2}$ . Every known lower bound on  $\|\omega\|_{L^\infty}$  is either an integral divergence (BKM) or a lim sup carrying a logarithm (Ingimarsen–Kukavica), or holds only along a constructed sequence (Cheskidov–Dai–Palasek). *No source in this list gives (Sat-1), and none gives (Sat-4).* This matters because, by Proposition 37.3, (Sat-3) alone is useless to us: (d)’s second clause and Observation 36.19 are applied at the particular times where §27–§31 run their estimates, and a lim sup names no such time.

**Remark 37.15** (Why the scaling heuristic predicts exponent 1, and why that is not a proof). Under  $u \mapsto \lambda u(\lambda \cdot, \lambda^2 \cdot)$  one has  $\|\omega\|_{L^\infty} \mapsto \lambda^2 \|\omega\|_{L^\infty}$ , deficit  $\alpha = 2$ , so the rule “exponent =  $\alpha/2$ ” that reproduces Fact 37.8 and Fact 37.9 predicts  $M(t) \geq c(T - t)^{-1}$ . The rule is a corollary of local well-posedness with existence time  $\gtrsim \|u_0\|_X^{-2/\alpha}$ ; here that would be a local existence theorem with existence time  $\gtrsim \|\omega_0\|_{L^\infty}^{-1}$ . No such theorem exists, and the reason is structural: the mild iteration for the vorticity equation must estimate the stretching term  $\omega \cdot \nabla u$ , i.e. must bound  $\|\nabla u\|_{L^\infty}$  by  $\|\omega\|_{L^\infty}$  — *the Wall 1 bound*. Running the iteration through  $\|u\|_{L^\infty}$  instead costs a Biot–Savart degree and returns exponent  $1/2$ , as in Fact 37.8. The scaling heuristic and Wall 1 are therefore not independent pieces of evidence; they are one piece, and Theorem 37.21 makes that exact.

### 37.4 The vorticity ODE, and where the logarithm enters

**Lemma 37.16** (Amplitude differential inequality on the torus).  *$M$  is locally Lipschitz on  $[0, T)$ ,  $M > 0$ , and for a.e.  $t < T$*

$$\frac{d}{dt} \log M(t) \leq \alpha_{\max}(t) - w_{\max}(t) \leq \|\nabla u(\cdot, t)\|_{L^\infty(\mathbb{T}^3)}. \quad (185)$$

[Proved.]

*Proof.* Identical to Lemma 33.21(c), with one simplification: on the compact manifold  $\mathbb{T}^3$  the supremum of  $|\omega(\cdot, t)|$  is *attained*, so the caveat of Remark 33.22 is vacuous here. Local Lipschitz continuity of  $M$  follows from boundedness of  $\nabla \omega$  and  $\partial_t \omega$  on compact subsets of  $\mathbb{T}^3 \times [0, T)$  ( $u$  is smooth there). At a point  $x_{\max}(t)$  of spatial maximum,  $M(t) > 0$ ,

hence  $\omega \neq 0$  nearby and  $|\omega|$  is smooth there;  $\nabla|\omega| = 0$  and  $\Delta|\omega| \leq 0$  at  $x_{\max}$ , so (141) and Hamilton's trick give  $\frac{d}{dt} \log M \leq \alpha_{\max} - w_{\max}$  for a.e.  $t$ . Finally  $w = |\nabla \hat{e}|^2 \geq 0$  and  $|\alpha| = |\hat{e} \cdot S \hat{e}| \leq \|S\|_{L^\infty} \leq \|\nabla u\|_{L^\infty}$ . That  $M > 0$ :  $\omega(\cdot, t) \equiv 0$  on  $\mathbb{T}^3$  would force  $u(\cdot, t)$  constant, hence (mean zero)  $u \equiv 0$  and no singularity.  $\square$

**Lemma 37.17** (All derivatives on the torus, logarithm-free). *Under the standing hypotheses, for every  $k \geq 0$  there is  $C_k(c_I)$  with*

$$\|\nabla^k u(\cdot, t)\|_{L^\infty(\mathbb{T}^3)} \leq C_k(c_I) (T - t)^{-\frac{k+1}{2}} \quad (186)$$

for all  $t$  with  $T - t$  small enough that  $t \geq \tau$  below. [Proved; this is Observation 36.19 with its  $k = 1$  specialisation removed.]

*Proof.* Exactly the restart of Observation 36.19, carried out for general  $k$ . Put  $\tau = \min(\varepsilon(k, 0)c_I^{-2}, \frac{1}{2})(T - t)$  and apply Lemma 36.15 with  $L = 1$  to  $w(\cdot, r) = u(\cdot, t - \tau + r)$ ,  $r \in [0, \tau]$ . Its datum obeys  $A = \|u(\cdot, t - \tau)\|_{L^\infty} \leq c_I(T - t + \tau)^{-1/2} \leq c_I(T - t)^{-1/2}$ , and  $\tau \leq \varepsilon(k, 0)A^{-2}$  because  $A^{-2} \geq c_I^{-2}(T - t)$ . At  $r = \tau$ ,  $\|\nabla^k u(\cdot, t)\|_{L^\infty} \leq C(k, 0)A\tau^{-k/2} = C_k(c_I)(T - t)^{-\frac{k+1}{2}}$ . No pressure, energy or smallness hypothesis enters, per Remark 36.20(a).  $\square$

**Lemma 37.18** (The logarithm is at most one logarithm). *With  $\mathcal{L}(t) = \log(e + \|u(\cdot, t)\|_{H^3}/M(t))$  as in Lemma 27.6 (definition on line 5712), there is  $C(c_I)$  with*

$$\mathcal{L}(t) \leq C(c_I)(1 + \Lambda(t)), \quad \int_t^T \mathcal{L}(r) dr \leq C(c_I)(T - t)(1 + \Lambda(t)),$$

for all  $t$  with  $T - t < 1$  and close enough to  $T$ . [Proved.]

*Proof.* By Lemma 37.17 and  $|\mathbb{T}^3| = 1$ ,  $\|u\|_{H^3} \leq C \sum_{k \leq 3} \|\nabla^k u\|_{L^\infty} \leq C(c_I)(T - t)^{-2}$ , the  $k = 3$  term dominating. By (184),  $M(t) \geq C(T - t)^{-1/4}$ . Hence  $\|u\|_{H^3}/M \leq C(c_I)(T - t)^{-7/4}$  and  $\mathcal{L} \leq \log(e + C(c_I)(T - t)^{-7/4}) \leq C'(c_I)(1 + \Lambda)$ . For the integral, substitute  $r = T - \rho$  and use  $\int_0^h \log \frac{1}{\rho} d\rho = h(1 + \log \frac{1}{h})$ , giving  $\int_t^T (1 + \Lambda(r)) dr = (T - t)(2 + \Lambda(t))$ .  $\square$

**Theorem 37.19** (The Type-I vorticity bound is saturated up to one logarithm, at every time). *Under the standing hypotheses there are  $c = c(c_I) > 0$  and  $t^* < T$  such that*

$$M(t) \geq \frac{c}{(T - t)(1 + \Lambda(t))} \quad \text{for all } t \in (t^*, T), \quad (187)$$

i.e. **(Sat-2)** holds. The statement is pointwise in  $t$ , not a lim sup. [Proved.]

*Proof.* By Lemma 37.16 and the Beale–Kato–Majda inequality in the form carried by Lemma 27.6,  $\|\nabla u\|_{L^\infty} \leq C M \mathcal{L}$ , we have for a.e.  $t$

$$\frac{d}{dt} M \leq \|\nabla u\|_{L^\infty} M \leq C \mathcal{L}(t) M^2.$$

Since  $M > 0$  and  $M$  is locally Lipschitz,  $G := 1/M$  is locally Lipschitz and  $G' \geq -C\mathcal{L}$  a.e., so for  $t < t' < T$

$$G(t) \leq G(t') + C \int_t^{t'} \mathcal{L}(r) dr.$$

By (184),  $M(t') \rightarrow \infty$  as  $t' \uparrow T$ , so  $G(t') \rightarrow 0$ ; letting  $t' \uparrow T$  and applying Lemma 37.18,

$$\frac{1}{M(t)} \leq C \int_t^T \mathcal{L} \leq C(c_I)(T-t)(1+\Lambda(t)),$$

which is (187). *Dimensional check*: with  $\nu = 1$ ,  $[M] = L^{-2} = [(T-t)^{-1}]$  and  $\Lambda$  is dimensionless.

*Non-circularity* (the trap named in the S50 brief). The only inputs are (141), the Type-I *upper* bounds (through Lemma 37.17, which uses Lemma 36.15 and the velocity hypothesis only), Fact 37.9, and Beale–Kato–Majda. No lower bound on  $M$  of exponent 1 is used, and Observation 36.19 is *not* used: the proof needs  $\|\nabla u\|_{L^\infty}$  measured against  $M$ , which is the very thing Observation 36.19 cannot supply off the saturated set.  $\square$

**Remark 37.20** (Where  $\mathcal{L}$  enters, and whether it can be avoided). It enters at exactly one place: the conversion of  $\frac{d}{dt}M \leq \|\nabla u\|_{L^\infty}M$  into a *Riccati* inequality  $\frac{d}{dt}M \leq \Psi M^2$ . Only a Riccati inequality integrates to a lower bound on  $M$ ; a linear inequality  $\frac{d}{dt}M \leq K(t)M$  integrates to a statement about ratios and is useless here. Concretely, feeding Observation 36.19's logarithm-free  $\|\nabla u\|_{L^\infty} \leq C(c_I)(T-t)^{-1}$  into (185) gives  $M(t) \geq M(t')((T-t')/(T-t))^C$ , and the Type-I *upper* bound at  $t'$  makes the right-hand side tend to 0 as  $t' \uparrow T$  whenever  $C > 1$ : nothing is obtained. So the factor  $\Psi$  must be  $\|\nabla u\|_{L^\infty}/M$ , which is  $\mathcal{L}$  by Lemma 27.6 and is  $O(1)$  precisely under Wall 1's target bound. **The logarithm in Theorem 37.19 cannot be removed by any refinement of the ODE argument: it is Wall 1 in the integrand.**

### 37.5 The equivalence with Wall 1

**Theorem 37.21** (Log-free saturation and Wall 1 are the same statement). *Under the standing hypotheses the following are equivalent.*

- (A) (Saturation, **(Sat-1)**) *There are  $c > 0$  and  $t^* < T$  with  $M(t) \geq c(T-t)^{-1}$  for all  $t \in (t^*, T)$ .*
- (B) (Wall 1's target) *There are  $K < \infty$  and  $t^* < T$  with  $\|\nabla u(\cdot, t)\|_{L^\infty(\mathbb{T}^3)} \leq K M(t)$  for all  $t \in (t^*, T)$ .*

Moreover the passage is lossless in the constants: (B) $\Rightarrow$ (A) with  $c = 1/K$ , and (A) $\Rightarrow$ (B) with  $K = C_1(c_I)/c$ , where  $C_1(c_I)$  is the constant of Observation 36.19. [Proved.]

*Proof.* (B) $\Rightarrow$ (A). By Lemma 37.16,  $\frac{d}{dt}M \leq KM^2$  a.e. on  $(t^*, T)$ , so  $G = 1/M$  satisfies  $G' \geq -K$  and  $G(t) \leq G(t') + K(t' - t)$  for  $t < t' < T$ . By (184),  $M(t') \rightarrow \infty$ , so  $G(t') \rightarrow 0$ ; letting  $t' \uparrow T$  gives  $1/M(t) \leq K(T-t)$ .

(A) $\Rightarrow$ (B). By Observation 36.19,  $\|\nabla u(\cdot, t)\|_{L^\infty} \leq C_1(c_I)(T-t)^{-1}$ , and by (A),  $(T-t)^{-1} \leq M(t)/c$ .

Neither direction is circular. The first uses only the maximum principle and (B). The second uses Observation 36.19, which is derived from the Type-I *upper* bounds through Lemma 36.15 and Serrin's criterion, with no lower bound on  $M$  anywhere.  $\square$

**Corollary 37.22** (Theorem 37.19 is a restatement, not progress). *Feeding (187) into Observation 36.19 yields*

$$\|\nabla u(\cdot, t)\|_{L^\infty} \leq C_1(c_I)(T-t)^{-1} \leq \frac{C_1(c_I)}{c} M(t)(1 + \Lambda(t)),$$

which by Lemma 37.18 is, up to constants depending only on  $c_I$ , the bound  $\|\nabla u\|_{L^\infty} \lesssim M\mathcal{L}$  already carried by Lemma 27.6. **Saturation up to a logarithm buys the document nothing it did not already have.** [Proved.]

**Proposition 37.23** (The same equivalence on the profile, in the infinite past). *Let  $U \in \mathcal{A}(c_I)$  be nontrivial. The following are equivalent.*

(A') *There are  $c > 0$ ,  $S > 0$  with  $M_U(s) \geq c|s|^{-1}$  for all  $s \leq -S$ .*

(B') *There are  $K < \infty$ ,  $S > 0$  with  $\|\nabla U(\cdot, s)\|_{L^\infty(\mathbb{R}^3)} \leq K M_U(s)$  for all  $s \leq -S$ .*

[Proved.]

*Proof.* (B')  $\Rightarrow$  (A'). By Lemma 33.21(c),  $\frac{d}{ds} M_U \leq \|\nabla U\|_{L^\infty} M_U \leq K M_U^2$  a.e., so  $G = 1/M_U$  obeys  $G' \geq -K$ , whence for  $s < s_0 \leq -S$ ,  $G(s) \leq G(s_0) + K(s_0 - s) \leq G(s_0) + K|s|$ . Therefore  $M_U(s) \geq (G(s_0) + K|s|)^{-1} \geq \frac{1}{2K}|s|^{-1}$  as soon as  $|s| \geq G(s_0)/K = 1/(K M_U(s_0))$ . Note that, unlike the torus case, no divergence of  $M_U$  is needed: the infinite past supplies the largeness of  $|s|$  instead. (Conversely, for  $s \uparrow 0^-$  the implication genuinely fails without a blow-up hypothesis at  $s = 0$ , which  $\mathcal{A}(c_I)$  does not impose.)

(A')  $\Rightarrow$  (B') is Corollary 36.17:  $\|\nabla U(\cdot, s)\|_{L^\infty} \leq C_1(c_I)|s|^{-1} \leq (C_1(c_I)/c)M_U(s)$ .  $\square$

**Remark 37.24** (What S48 actually did to this question). Before S48 only the direction (A) $\Rightarrow$ (B) was a possibility, and it was not available: it needs Observation 36.19, which did not exist. S48 supplied it. The net effect of S48 on this branch is therefore to upgrade “saturation would suffice to remove Wall 1” to “saturation is *equivalent* to removing Wall 1”. That is a real gain in knowledge and a loss in strategy: the reduction Remark 36.21 proposed — replace the wall by the sharper question — does not shrink the problem, because the sharper question has the same content. *This is a negative result and it should be recorded as one.*

### 37.6 Discharging the only consumer of Part (e), without saturation

By Observation 37.6 the whole weight of Proposition 33.8(e) rests on Proposition 33.20(b). That consumer does not need  $\langle \sigma_U^* \rangle$  at the amplitude scale  $M_U(s_0)^{-1/2}$ ; it needs a scale-invariant cylinder family, commensurate with the dyadic shells, carrying a bound that does not decay. The Type-I scale supplies one unconditionally.

**Proposition 37.25** (Type-I-scale coherence bound on the profile). *Let  $U \in \mathcal{A}(c_I)$  and for  $s_0 < 0$  put  $r_I := |s_0|^{1/2}$ ,*

$$\langle \sigma_U^{*,I} \rangle(s_0) := r_I^{-3} \iint_{Q(r_I/2)} |\nabla \hat{e}_U|^2 \mathbf{1}_{\{|\omega_U| \geq M_U/2\}} dy ds.$$

*Then  $\langle \sigma_U^{*,I} \rangle(s_0) \leq C(c_I)$  for every  $s_0 < 0$ , with no saturation hypothesis; and  $\langle \sigma_U^{*,I} \rangle$  is exactly scale-invariant,  $\langle \sigma_U^{*,I} \rangle(s) = \langle \sigma_U^{*,I} \rangle(\mu^2 s)$ .* [Proved modulo routine verification, in the sense of

Proposition 27.8: the three load-bearing ingredients are proved below; what remains is the term-for-term repetition of Lemma 28.1(b),(c) at the new radius, in which every exponent is unchanged.]

*Proof.* Write  $M = M_U(s_0)$  and note the Type-I upper bound  $M_U(s) \leq c_I |s|^{-1}$ , so  $r_I \leq c_I^{1/2} M^{-1/2}$ : the new cube is contained in the old one, and the two coincide exactly on the saturated set. Three ingredients.

(1) *Amplitude comparability, now unconditional.* The time extent of  $Q(r_I/2)$  is  $r_I^2/4 = |s_0|/4$ , so  $|s| \leq \frac{5}{4}|s_0|$  on it, and Corollary 36.17 with Lemma 33.21(c) gives, as in Lemma 37.4,  $M_U(s) \geq (\frac{5}{4})^{-C_1(c_I)} M_U(s_0)$  there. This is the step that needed saturation at the amplitude scale (Remark 37.7(a)) and needs nothing here, because the cube's temporal extent is now tied to  $|s_0|$  rather than to  $M_U(s_0)^{-1}$ .

(2) *The stretching term.* In Lemma 28.1(a),  $\iint |\alpha| m^2 \eta^2 \leq \|\nabla U\|_{L^\infty} M^2 r_I^5 \leq C_1(c_I) |s_0|^{-1} M^2 |s_0|^{5/2} = C_1(c_I) M^2 r_I^3$ , using only the *first* clause of Proposition 33.8(d), which is Corollary 36.17 and carries no proviso. At the amplitude scale the same computation needs  $\|\nabla U\|_{L^\infty} \leq CM$ , i.e. the second clause.

(3) *The transport term.*  $|D_t \psi| \leq |\partial_s \psi| + |U| |\nabla \psi| \leq C r_I^{-2} + c_I |s|^{-1/2} \cdot C r_I^{-1} \leq C(1 + c_I) r_I^{-2}$ , the two contributions balancing exactly because  $r_I^2 = |s_0|$ . At the amplitude scale the ratio of the second to the first is  $(|s_0| M)^{-1/2}$ , which is  $O(1)$  *only on the saturated set* — so the transport term, not only the stretching term, carries the proviso there. This does not appear to have been noticed.

With (1)–(3), the vorticity Caccioppoli inequality of Lemma 27.6, re-run on  $U$  at radius  $\rho \leq r_I$ , gives  $\iint_{Q(2\rho)} |\nabla \omega_U|^2 \leq C M^2 \rho^3 + \|\nabla U\|_{L^\infty} M^2 \rho^5 \leq C(c_I) M^2 \rho^3$  — logarithm-free and proviso-free, using  $\rho^2 \leq |s_0|$  in the second term. Substituting into (148) and dividing by the left-hand coefficient  $M_U(s)^2/4 \geq c(c_I) M^2$  from (1) gives  $\iint_{Q(r_I/2)} w \mathbf{1} \leq C(c_I) r_I^3$ . Scale invariance is Lemma 33.3 together with the covariance  $r_I(U_\mu, s) = \mu^{-1} r_I(U, \mu^2 s)$ .  $\square$

**Corollary 37.26** (Proposition 33.20 is unconditional). *Proposition 33.20 holds with  $\langle \sigma_U^{*,I} \rangle$  in place of  $\langle \sigma_U^* \rangle$  and  $r_{I,k} = |s_k|^{1/2}$  in place of  $r_k^*$ , without saturation. Part (a) becomes an identity,  $r_{I,k}^2 = |s_k|$ , so each dyadic shell carries exactly one cylinder; part (b) is Proposition 37.25; part (c) is unchanged. [Proved modulo the same routine verification.]*

**Remark 37.27** (What this does and does not do for Part (e)). It does *not* prove Proposition 33.8(e), which is a statement about  $\langle \sigma_U^* \rangle$  at the amplitude scale and remains **Proved mod saturation**; off the saturated set the two quantities are averages over different cubes and neither dominates the other. What it does is remove (e)'s only consumer from the list of statements that depend on saturation. Since the Bridge Lemma (Conjecture 26.8) requires smallness of  $\sigma^*$  at the amplitude scale  $r^* = M^{-1/2}$ ,  $\langle \sigma^{*,I} \rangle$  is *not* a substitute there, and no claim is made that it is.

### 37.7 Verdict

**Remark 37.28** (Verdict of S50). The question posed was: does a putative first singularity satisfy  $M(t) \geq c(T - t)^{-1}$  near  $T$ , with  $c = c(c_I)$ , logarithm-free? The verdict is **(iii)**: a weaker form is a theorem, the strong form is open, and the residue is exactly Wall 1.

- (1) **(Sat-2) is a theorem** (Theorem 37.19), pointwise in  $t$  and not a lim sup. Under the standing hypotheses this is strictly stronger than anything in Fact 37.10 or Fact 37.12; the gain over the latter is entirely attributable to the Type-I hypothesis, which replaces their Grönwall control of  $\|u\|_{H^3}$  by Lemma 37.17. It is not a claim about the Euler literature and does not compete with it.
- (2) **(Sat-1) is open**, and the gap has a name: by Theorem 37.21 it *is* Wall 1. There is no separate obstruction to find.
- (3) **(Sat-2) is worth nothing** (Corollary 37.22): it returns exactly the estimate Lemma 27.6 already carried. The intuition recorded in Observation 36.19 — that saturation, not the logarithm, is what both branches need — is correct as a diagnosis and wrong as a strategy: the two are the same quantity.
- (4) **The target was over-rated.** By Observation 37.6, Proposition 33.8(e) is a leaf whose sole consumer is an impossibility statement, and by Corollary 37.26 that consumer can be discharged at the Type-I scale without saturation. Saturation should therefore be removed from the programme’s priority list: on the profile branch it is nearly weightless, and on the global branch it is Wall 1 under a different name.

**Residual gap, stated exactly.** Prove or disprove: there is  $K = K(c_I)$  with  $\|\nabla u(\cdot, t)\|_{L^\infty(\mathbb{T}^3)} \leq K M(t)$  for all  $t$  near a putative Type-I first singularity — equivalently (Theorem 37.21),  $\liminf_{t \uparrow T} (T - t)M(t) > 0$ . Nothing in this document or in the sources of §37.3 decides it in either direction; in particular Wall 1 remains OPEN, not PROVED IMPOSSIBLE, and Theorem 37.21 is not evidence that it cannot be removed.

| Item                           | Status before S50     | Status after S50                               |
|--------------------------------|-----------------------|------------------------------------------------|
| (Sat-1) log-free saturation    | unexamined            | OPEN; = Wall 1 (Thm. 37.21)                    |
| (Sat-2) saturation mod one log | unexamined            | <b>Proved</b> (Thm. 37.19)                     |
| (Sat-2) as a tool              | unexamined            | <b>Vacuous</b> (Cor. 37.22)                    |
| Strength (e) needs             | “at every time”       | at the reference time only (Lem. 37.4)         |
| Consumers of (e)               | unstated              | one, itself unconsumed (Obs. 37.6)             |
| Prop. 33.20                    | Proved mod saturation | <b>Proved</b> at the Type-I scale (Cor. 37.26) |
| Prop. 33.8(e)                  | Proved mod saturation | unchanged (Rem. 37.27)                         |
| Wall 1 (§44.2)                 | OPEN, structural      | OPEN, structural; scope widened                |
| Thm. 28.2 hidden hypothesis    | unnoticed             | recorded (Rem. 37.7(a))                        |
| BKM citation for $\mathcal{L}$ | absent                | [28, 14]                                       |

**Remark 37.29** (Scope discipline). This section proves one theorem, one equivalence and one audit. It does not close either horn of the pincer, does not remove Wall 1, does not bear on (N), (N’), (A-up) or Type-II, and does not bring the programme nearer to the Clay problem than Remark 33.30 already states, which stands verbatim. Its principal effect is subtractive: it removes a target the programme had placed at the head of its list.



## 38 The Axisymmetric-with-Swirl Class: Where the Calderón–Zygmund Logarithm Goes, and What the Literature Has Already Closed (Session S51)

This section executes the axisymmetric pivot commissioned as WP3 of the S46 plan. Its purpose is *not* to claim a theorem. It is to answer one question — does the axisymmetric reduction remove the Calderón–Zygmund logarithm of Wall 1? — with a written estimate rather than an intuition, and then to place whatever remains against the axisymmetric literature, which is large and much older than this programme.

Two verdicts are reached, and both are negative for the pivot as it was conceived.

- (I) **The logarithm is not removed.**  $\nabla u$  splits into a *swirl block* (three of the eight non-zero entries), which is bounded by  $\frac{3}{2}\|\omega\|_{L^\infty}$  with an absolute constant, no elliptic solve and no logarithm (Lemma 38.7); and a *poloidal block* (five entries), which is recovered from  $\omega_\theta$  by a genuinely two-dimensional elliptic problem in  $(r, z)$  and pays the full factor  $\mathcal{L}$ . The second statement is *sharp*, by an explicit field placed *inside* the critical cylinder around the axis (Proposition 38.9). On the axis itself the poloidal block does become logarithm-free, but only against the weighted norm  $\|\omega_\theta/r\|_{L^\infty}$  (Lemma 38.11); the logarithm returns at every positive  $r$ .
- (II) **Everything this programme’s machinery could prove in this class is already a theorem.** Seregin–Šverák 2009, Theorem 1.1 (Fact 38.24) excludes Type-I singularities of axisymmetric solutions *with swirl*, locally, under a Type-I bound on the *poloidal* velocity alone and with no decay hypothesis. The pincer of §26 concludes “no Type-I singularity” (Remark 33.30). Hence H1 and H2, proved in full in this class, would re-prove a 2009 theorem under strictly more hypotheses (Proposition 38.25).

What the section does contribute is an exact dictionary (Lemma 38.14) between the programme’s coherence deficit  $\sigma^*$  and the two scalars  $J = \omega_r/r$ ,  $\Omega = \omega_\theta/r$  that the axisymmetric literature is built on, together with the observation that the aligned horn of §33.4, read in this class, is exactly the vanishing of the swirl source term (Proposition 38.17). No regularity criterion is claimed; §38.5 states why the natural target is not a target at all.

### 38.1 The class, the two scalars, and a re-derivation of the setting

**Definition 38.1** (Axisymmetric fields). In cylindrical coordinates  $(r, \theta, z)$  on  $\mathbb{R}^3$  with  $r = \sqrt{x_1^2 + x_2^2}$ , a vector field is *axisymmetric* if

$$u = u_r(r, z, t) e_r + u_\theta(r, z, t) e_\theta + u_z(r, z, t) e_z,$$

i.e. all components are  $\theta$ -independent. The field has *swirl* if  $u_\theta \not\equiv 0$ . We write  $b = u_r e_r + u_z e_z$  for the poloidal (meridional) part,  $\Delta = \partial_r^2 + \frac{1}{r}\partial_r + \partial_z^2$  for the axisymmetric Laplacian, and

$$\Gamma := r u_\theta, \quad J := \frac{\omega_r}{r}, \quad \Omega := \frac{\omega_\theta}{r}.$$

The vorticity components are

$$\omega_r = -\partial_z u_\theta, \quad \omega_\theta = \partial_z u_r - \partial_r u_z, \quad \omega_z = \partial_r u_\theta + \frac{u_\theta}{r} = \frac{1}{r} \partial_r (r u_\theta). \quad (188)$$

**Remark 38.2** (A standing caveat: this class does not live on  $\mathbb{T}^3$ ). §27–§33 carry their standing assumptions on  $\mathbb{T}^3$  (line 5563).  $\mathbb{T}^3$  has no continuous rotational symmetry about a line, so “axisymmetric” has no meaning there and the whole of the literature quoted below is an  $\mathbb{R}^3$  theory. Adopting this class therefore requires moving the programme’s standing framework to  $\mathbb{R}^3$  with finite energy first. That is not a formality — the profile extraction of §33.2 and the harmonic-Liouville step of Lemma 33.16 were set up against the torus — and it is recorded here as a cost of entry, not waved away.

The two evolution equations that define the difficulty of this class were supplied to this session in pre-summarised form. Both are re-derived here from the Navier–Stokes equations, in accordance with the programme’s verification discipline; both are confirmed correct.

**Lemma 38.3** (The swirl equation and its maximum principle). *For a smooth axisymmetric solution,  $\Gamma = ru_\theta$  satisfies*

$$\partial_t \Gamma + u_r \partial_r \Gamma + u_z \partial_z \Gamma = \nu \left( \Delta - \frac{2}{r} \partial_r \right) \Gamma, \quad (189)$$

and consequently  $\|\Gamma(\cdot, t)\|_{L^\infty} \leq \|\Gamma(\cdot, 0)\|_{L^\infty}$  for all  $t$  in the smooth interval. [Proved; classical.]

*Proof.* The  $\theta$ -component of the Navier–Stokes equations carries no pressure ( $\partial_\theta p = 0$ ) and reads

$$\partial_t u_\theta + u_r \partial_r u_\theta + u_z \partial_z u_\theta + \frac{u_r u_\theta}{r} = \nu \left( \Delta u_\theta - \frac{u_\theta}{r^2} \right).$$

Substitute  $u_\theta = \Gamma/r$ . On the left,  $u_r \partial_r (\Gamma/r) = u_r \partial_r \Gamma / r - u_r \Gamma / r^2$  and  $u_r u_\theta / r = +u_r \Gamma / r^2$ : the two  $r^{-2}$  terms cancel exactly, and the left side is  $r^{-1}(\partial_t \Gamma + b \cdot \nabla \Gamma)$ . On the right,

$$\partial_r^2 \left( \frac{\Gamma}{r} \right) = \frac{\Gamma_{rr}}{r} - \frac{2\Gamma_r}{r^2} + \frac{2\Gamma}{r^3}, \quad \frac{1}{r} \partial_r \left( \frac{\Gamma}{r} \right) = \frac{\Gamma_r}{r^2} - \frac{\Gamma}{r^3}, \quad \partial_z^2 \left( \frac{\Gamma}{r} \right) = \frac{\Gamma_{zz}}{r},$$

so  $\Delta(\Gamma/r) - \Gamma/r^3 = \frac{1}{r}(\Gamma_{rr} - \frac{\Gamma_r}{r} + \Gamma_{zz}) = \frac{1}{r}(\Delta - \frac{2}{r} \partial_r) \Gamma$ . Multiplying by  $r$  gives (189).

For the maximum principle:  $\Delta - \frac{2}{r} \partial_r = \partial_r^2 - \frac{1}{r} \partial_r + \partial_z^2$  is elliptic with no zeroth-order term, and the drift  $b$  is smooth on the smooth interval. The coefficient  $-2\nu/r$  is singular only at  $r = 0$ ; smoothness of  $u$  across the axis forces  $u_\theta = O(r)$ , hence  $\Gamma = O(r^2)$ , so running the classical maximum principle on  $\{r \geq \varepsilon\}$  and letting  $\varepsilon \downarrow 0$  — the boundary contribution at  $r = \varepsilon$  is  $O(\varepsilon^2)$  — gives the stated bound.  $\square$

**Lemma 38.4** (The  $\Omega$  equation and the single source term). *For a smooth axisymmetric solution,  $\Omega = \omega_\theta / r$  satisfies*

$$\partial_t \Omega + b \cdot \nabla \Omega = \nu \left( \Delta + \frac{2}{r} \partial_r \right) \Omega + \frac{\partial_z(\Gamma^2)}{r^4}. \quad (190)$$

In particular, if  $\partial_z \Gamma \equiv 0$  on an open set then the source vanishes there and  $\Omega$  obeys a maximum principle on that set. [Proved; classical.]

*Proof.* The  $\theta$ -component of the vorticity equation for an axisymmetric field is

$$\partial_t \omega_\theta + b \cdot \nabla \omega_\theta = \nu \left( \Delta - \frac{1}{r^2} \right) \omega_\theta + \frac{u_r}{r} \omega_\theta + \frac{2u_\theta}{r} \partial_z u_\theta. \quad (191)$$

This too was verified from scratch rather than quoted. Take  $\partial_z$  of the  $r$ -component of Navier–Stokes, which carries the centrifugal term  $-u_\theta^2/r$  on the left and the viscous term  $\nu(\Delta u_r - u_r/r^2)$ , and subtract  $\partial_r$  of the  $z$ -component; the pressure cancels. The centrifugal term contributes  $+\frac{1}{r}\partial_z(u_\theta^2) = \frac{2u_\theta}{r}\partial_z u_\theta$  with a plus sign (it sits on the left of the momentum equation with a minus). The convective commutator is  $\partial_z[(b \cdot \nabla)u_r] - \partial_r[(b \cdot \nabla)u_z] = (b \cdot \nabla)\omega_\theta + (\partial_r u_r + \partial_z u_z)\omega_\theta = (b \cdot \nabla)\omega_\theta - \frac{u_r}{r}\omega_\theta$ , using incompressibility  $\partial_r u_r + u_r/r + \partial_z u_z = 0$ . For the viscous part,  $[\Delta, \partial_z] = 0$  while  $\Delta \partial_r f = \partial_r \Delta f + \partial_r f/r^2$ , so  $\partial_z(\Delta u_r - u_r/r^2) - \partial_r \Delta u_z = \Delta \omega_\theta - \omega_\theta/r^2$ . Put  $\omega_\theta = r\Omega$ . Then  $b \cdot \nabla(r\Omega) = r b \cdot \nabla \Omega + u_r \Omega$ , and the term  $u_r \Omega$  cancels against  $r^{-1}u_r \omega_\theta = u_r \Omega$  on the right. For the diffusion,  $\partial_r(r\Omega) = \Omega + r\Omega_r$  and  $\partial_r^2(r\Omega) = 2\Omega_r + r\Omega_{rr}$ , so

$$\left(\Delta - \frac{1}{r^2}\right)(r\Omega) = r\Omega_{rr} + 3\Omega_r + \frac{\Omega}{r} + r\Omega_{zz} - \frac{\Omega}{r} = r\left(\Delta + \frac{2}{r}\partial_r\right)\Omega.$$

Finally  $r^{-1} \cdot \frac{2u_\theta}{r}\partial_z u_\theta = \frac{1}{r^2}\partial_z(u_\theta^2) = \frac{1}{r^2}\partial_z(\Gamma^2/r^2) = \partial_z(\Gamma^2)/r^4$ , since  $r$  does not depend on  $z$ . Dividing through by  $r$  gives (190). The operator  $\Delta + \frac{2}{r}\partial_r = \partial_r^2 + \frac{3}{r}\partial_r + \partial_z^2$  has no zeroth-order term, whence the last assertion.  $\square$

**Remark 38.5** (The handover’s statement of the setting is correct; one terminological trap is not). Both (189) and (190), and all three identities of (188), were handed to this session pre-summarised and are confirmed above by direct computation. *No error was found in the handover’s statement of the setting.*

One thing the handover does *not* flag, and which has already caused confusion in the literature it points at, is that the phrase “Type I” carries two different meanings in this subject. This programme, and [17], use the *temporal* bound  $\sup_x |u(x, t)| \leq C(T - t)^{-1/2}$ . Much of the axisymmetric literature — and survey treatments of it — instead call the *spatial* scale-invariant bound  $|u(x, t)| \leq C/r$  “Type I”, because it is that bound which makes the  $\Gamma$  equation critical. The two are not comparable: neither implies the other. Every statement below names which one it uses.

## 38.2 The recovery of $\nabla u$ : the swirl block is free, the poloidal block is not

This subsection answers the central question of the pivot. The frame computation first.

**Lemma 38.6** (The eight entries). *For an axisymmetric field,*

$$\begin{aligned} \nabla u &= \partial_r u_r (e_r \otimes e_r) + \partial_r u_\theta (e_\theta \otimes e_r) + \partial_r u_z (e_z \otimes e_r) \\ &\quad + \frac{u_r}{r} (e_\theta \otimes e_\theta) - \frac{u_\theta}{r} (e_r \otimes e_\theta) \\ &\quad + \partial_z u_r (e_r \otimes e_z) + \partial_z u_\theta (e_\theta \otimes e_z) + \partial_z u_z (e_z \otimes e_z), \end{aligned}$$

the ninth entry  $(e_z \otimes e_\theta)$  being  $\frac{1}{r}\partial_\theta u_z = 0$ . Call  $\{\partial_r u_\theta, \partial_z u_\theta, -u_\theta/r\}$  the swirl block and  $\{\partial_r u_r, \partial_z u_r, \partial_r u_z, \partial_z u_z, u_r/r\}$  the poloidal block. [Proved; direct computation from  $\partial_\theta e_r = e_\theta$ ,  $\partial_\theta e_\theta = -e_r$ .]

**Lemma 38.7** (The swirl block is logarithm-free, with an absolute constant). *Let  $u$  be a smooth axisymmetric field on  $\mathbb{R}^3$  and  $M = \|\omega\|_{L^\infty}$ . Then, pointwise and everywhere,*

$$\left|\frac{u_\theta}{r}\right| \leq \frac{\|\omega_z\|_{L^\infty}}{2}, \quad |\partial_z u_\theta| = |\omega_r| \leq M, \quad |\partial_r u_\theta| \leq \frac{3}{2}M, \quad (192)$$

so the swirl block of  $\nabla u$  is bounded by  $\frac{3}{2}M$  with an absolute constant. No Calderón–Zygmund operator, no elliptic solve, and no factor  $\mathcal{L}$  appear. [Proved.]

*Proof.* The middle identity is (188). For the first, set  $h = u_\theta/r$ . Then  $\omega_z = \frac{1}{r}\partial_r(ru_\theta) = \frac{1}{r}\partial_r(r^2h) = 2h + r\partial_rh$ , i.e.  $\partial_r(r^2h) = r\omega_z$ . Smoothness across the axis gives  $h$  bounded near  $r = 0$ , so  $r^2h \rightarrow 0$  as  $r \downarrow 0$  and integration gives

$$r^2h(r, z) = \int_0^r s \omega_z(s, z) ds, \quad \text{whence} \quad |h(r, z)| \leq \frac{1}{2}\|\omega_z\|_{L^\infty}.$$

The third bound is then  $|\partial_r u_\theta| = |\omega_z - h| \leq \frac{3}{2}\|\omega_z\|_{L^\infty}$ .  $\square$

**Remark 38.8** (Why the swirl block is free and the poloidal block cannot be). The mechanism is worth naming, because it is exactly what does *not* generalise. The swirl is a *single scalar*  $u_\theta$ , and the two vorticity components  $\omega_r, \omega_z$  determine both of its derivatives — one algebraically ( $\partial_z u_\theta = -\omega_r$ ) and one through a *first-order ordinary differential equation in  $r$  alone* ( $\partial_r(r^2 u_\theta/r) = r\omega_z$ ), whose Green’s function is the locally integrable kernel  $s ds/r^2$ . The poloidal pair  $(u_r, u_z)$ , by contrast, is constrained by only two relations —  $\omega_\theta$  and  $\operatorname{div} u = 0$  — for five entries, and its recovery is a genuinely two-dimensional elliptic problem in  $(r, z)$ . There is no ODE in  $r$  to exploit. This is the structural reason the two blocks behave differently, and it also shows that the gain is confined to the swirl: it comes from one-dimensionality, not from axisymmetry as such.

We now set up the poloidal recovery. Since  $\frac{1}{r}\partial_r(ru_r) + \partial_z u_z = 0$ , there is a Stokes stream function; it is convenient to use instead the  $\theta$ -component  $L_\theta$  of the vector potential, defined by  $u = \operatorname{curl}(L_\theta e_\theta) + u_\theta e_\theta$ , so that

$$u_r = -\partial_z L_\theta, \quad u_z = \frac{1}{r}\partial_r(rL_\theta), \quad \left(\Delta - \frac{1}{r^2}\right)L_\theta = -\omega_\theta. \quad (193)$$

Writing  $L_\theta = r m(r, z)$  with  $m$  smooth and even in  $r$  (forced by smoothness of  $u$  across the axis), the computation of Lemma 38.4 run backwards turns the last equation of (193) into

$$\left(\Delta + \frac{2}{r}\partial_r\right)m = -\Omega, \quad u_r = -r\partial_z m, \quad u_z = 2m + r\partial_r m. \quad (194)$$

The operator  $\Delta + \frac{2}{r}\partial_r = \partial_r^2 + \frac{3}{r}\partial_r + \partial_z^2$  is the Laplacian of  $\mathbb{R}^5$  restricted to functions invariant under  $SO(4)$  acting on the first four coordinates; we write it  $\Delta_5$  and regard  $m$  as the 5-dimensional Newtonian potential of  $\Omega$ . Note also, by the same integration as in Lemma 38.7,

$$r^2 m(r, z) = \int_0^r s u_z(s, z) ds, \quad \text{hence} \quad \|m\|_{L^\infty} \leq \frac{1}{2}\|u\|_{L^\infty}. \quad (195)$$

**Proposition 38.9** (The poloidal block pays the full  $\mathcal{L}$ , and this is sharp inside the critical cylinder). *There is an absolute constant  $c > 0$  with the following property. Let  $r^* > 0$  and let  $z_0 \in \mathbb{R}$ . For every  $N \in \mathbb{N}$  there is a smooth, compactly supported, divergence-free axisymmetric field  $u^{(N)}$ , supported in the ball  $B_{r^*}$  centred at the axis point  $(0, 0, z_0)$ , such that*

$$\|\nabla u^{(N)}\|_{L^\infty} \geq c N \|\omega^{(N)}\|_{L^\infty}, \quad \|u^{(N)}\|_{H^3} \leq C(r^*) 2^{2N} \|\omega^{(N)}\|_{L^\infty}.$$

*Consequently  $\|\nabla u\|_{L^\infty} \leq K\|\omega\|_{L^\infty}$  is false on the axisymmetric class for every  $K$ , and the growth is at least logarithmic in  $\|u\|_{H^3}/\|\omega\|_{L^\infty}$ , i.e. at least  $\mathcal{L}$ . The example may be taken swirl-free or, with an arbitrary added swirl of comparable size, with  $\Gamma \not\equiv 0$ . [Proved.]*

*Proof. Step 1: the two-dimensional model.* Let  $R_{12} = \partial_1 \partial_2 \Delta^{-1}$  on  $\mathbb{R}^2$ , a Calderón–Zygmund operator with kernel  $c_2 y_1 y_2 / |y|^4$ . Fix  $g \in C_c^\infty(\{1 < |y| < 2\})$  with  $\|g\|_{L^\infty} = 1$  and  $(R_{12}g)(0) =: c_0 > 0$  (possible: take  $g = y_1 y_2 \chi(|y|)$  with  $\chi \geq 0$  supported in  $(1, 2)$ , for which the integrand is a positive multiple of  $(y_1 y_2)^2 / |y|^4$ ). Set  $\omega_N(y) = \sum_{j=1}^N g(2^j y)$ . The summands have pairwise disjoint supports  $\{2^{-j} < |y| < 2^{1-j}\}$ , so  $\|\omega_N\|_{L^\infty} = 1$  and  $\omega_N \equiv 0$  on  $\{|y| < 2^{-N}\}$ ; and  $R_{12}$  commutes with dilations (it is homogeneous of degree 0), so

$$(R_{12}\omega_N)(0) = \sum_{j=1}^N (R_{12}g)(0) = c_0 N.$$

With  $\phi_N = \Delta^{-1}(-\omega_N)$  and  $v_N = \nabla^\perp \phi_N$  one has  $\partial_1(v_N)_1 = -R_{12}\omega_N$ , so  $\|\nabla v_N\|_{L^\infty} \geq c_0 N$ , while  $\|v_N\|_{L^\infty} \leq C$  and, the derivatives also having disjoint supports,  $\|\omega_N\|_{H^2} \leq C 2^{2N}$ .

*Step 2: localisation of the model.*  $\phi_N$  is not compactly supported (it grows logarithmically), so fix  $\chi \in C_c^\infty(B_2(0))$  with  $\chi \equiv 1$  on  $B_1$  and set  $\tilde{\phi}_N = \chi \phi_N$ , normalising the additive constant of  $\phi_N$  by  $\phi_N(0) = 0$ . Since  $\text{supp } \omega_N \subset B_1$ , where  $\chi \equiv 1$ , we have  $\tilde{\phi}_N = \phi_N$  on  $B_1$ , so all derivatives of  $\tilde{\phi}_N$  at the origin are unchanged and  $\|\nabla \tilde{v}_N\|_{L^\infty} \geq c_0 N$  still holds with  $\tilde{v}_N = \nabla^\perp \tilde{\phi}_N$ . On the annulus  $1 < |y| < 2$ ,  $\Delta \tilde{\phi}_N = 2 \nabla \chi \cdot \nabla \phi_N + \phi_N \Delta \chi$ , which is bounded by  $C_\chi (\|\nabla \phi_N\|_{L^\infty} + \|\phi_N\|_{L^\infty(B_2)}) \leq C'_\chi$  independently of  $N$ , because  $\|\omega_N\|_{L^1} + \|\omega_N\|_{L^\infty} \leq C$  uniformly. Hence  $\tilde{\omega}_N := -\Delta \tilde{\phi}_N$  satisfies  $\|\tilde{\omega}_N\|_{L^\infty} \leq 1 + C'_\chi$ , an absolute constant, and  $\tilde{v}_N$  is supported in  $B_2(0)$ .

*Step 3: exact embedding.* Put  $r_0 = r^*/2$  and  $\rho = r^*/64$ , so that  $\Lambda := 2\rho/r_0 = 1/16$ , and define the Stokes stream function

$$\psi(r, z) = r_0 \rho \tilde{\phi}_N \left( \frac{r - r_0}{\rho}, \frac{z - z_0}{\rho} \right), \quad u_r = -\frac{1}{r} \partial_z \psi, \quad u_z = \frac{1}{r} \partial_r \psi, \quad u_\theta = 0.$$

This  $u$  is exactly divergence-free and axisymmetric, and is supported in  $\{|r - r_0| \leq 2\rho, |z - z_0| \leq 2\rho\}$ ; since  $r \in [r_0 - 2\rho, r_0 + 2\rho] \subset [\frac{7}{16}r^*, \frac{9}{16}r^*]$  and  $|z - z_0| \leq r^*/32$ , that set lies inside the ball  $B_{r^*}$  centred at  $(0, 0, z_0)$ . Note that the embedding is *exact*: no perturbation of the elliptic operator is involved, because the stream-function formulation of incompressibility is used directly. On the support  $r_0/r = 1 + O(\Lambda)$ , and with  $y = ((r - r_0)/\rho, (z - z_0)/\rho)$ ,

$$(u_r, u_z) = \frac{r_0}{r} \tilde{v}_N(y), \quad \nabla_{(r,z)}(u_r, u_z) = \frac{1}{\rho} \frac{r_0}{r} (\nabla \tilde{v}_N)(y) + O\left(\frac{|\tilde{v}_N(y)|}{r_0}\right).$$

Since  $\omega_\theta = \partial_z u_r - \partial_r u_z$  is a particular combination of these entries,  $\|\omega\|_{L^\infty} = \|\omega_\theta\|_{L^\infty} \leq \rho^{-1}(1 + C'_\chi)(1 + C\Lambda) + Cr_0^{-1}$  and  $\|\nabla u\|_{L^\infty} \geq \rho^{-1}(1 - C\Lambda)c_0 N - Cr_0^{-1}$ . Here  $\Lambda = 1/16$  and  $\rho/r_0 = 1/32$  are absolute, so the two additive  $r_0^{-1}$  terms are an absolute fraction of  $\rho^{-1}$ ; taking  $N$  beyond an absolute threshold gives  $\|\nabla u\|_{L^\infty} \geq c N \|\omega\|_{L^\infty}$  with  $c$  absolute. Scaling  $\|\cdot\|_{H^3}$  by  $\rho^{-1}$ -powers gives the stated  $H^3$  bound with  $C(r^*)$ ; only the exponential-in- $N$  factor matters.

*Step 4: swirl.* Adding any smooth  $u_\theta$  supported in the same ball changes only the swirl block of  $\nabla u$  (Lemma 38.6), which occupies entries disjoint from the poloidal block, and adds  $\omega_r, \omega_z$  to the vorticity. Choosing  $u_\theta$  with  $\|(\omega_r, \omega_z)\|_{L^\infty} \leq \|\omega_\theta\|_{L^\infty}$  leaves the displayed inequality intact up to a factor 2, with  $\Gamma \not\equiv 0$ .  $\square$

**Remark 38.10** (This is not an instance of the “reduced to no swirl” failure mode). Step 3 is included precisely to forestall the objection. The statement being refuted — “ $\|\nabla u\|_\infty \lesssim \|\omega\|_\infty$  on the axisymmetric class” — concerns the *linear recovery operator*  $\omega \mapsto \nabla u$ , not a solution of any equation, and the swirl block is additive and occupies disjoint entries, so it cannot cancel the poloidal block. No dynamical argument has been reduced to the swirl-free case.

**Lemma 38.11** (On the axis the poloidal block is logarithm-free — against the weighted norm). *Let  $u$  be smooth axisymmetric on  $\mathbb{R}^3$ ,  $x_0 = (0, 0, z_0)$  on the axis,  $R > 0$ . Then at every point of the axis,*

$$|\nabla u_{\text{pol}}|(0, z) = 2 |\partial_z m(0, z)|, \quad |\partial_z m(0, z)| \leq C R \|\Omega\|_{L^\infty(B_{2R}(x_0))} + \frac{C}{R} \|m\|_{L^\infty(B_{2R}(x_0))}$$

for  $|z - z_0| < R/2$ , with  $C$  absolute and no logarithm. [Proved.]

*Proof.* On the axis  $\partial_r m = 0$  ( $m$  is even in  $r$ ), so by (194) the five poloidal entries evaluate to  $\partial_r u_r = -\partial_z m$ ,  $\partial_z u_r = 0$ ,  $u_r/r = -\partial_z m$ ,  $\partial_r u_z = 0$ ,  $\partial_z u_z = 2\partial_z m$ ; their sum  $-\partial_z m - \partial_z m + 2\partial_z m = 0$  is the divergence, as it must be.

For the bound, split  $\Omega = \Omega \mathbf{1}_{B_{2R}} + \Omega \mathbf{1}_{B_{2R}^c}$  and correspondingly  $m = m_{\text{near}} + m_{\text{far}}$  in the 5-dimensional picture (194). The Newtonian kernel of  $(-\Delta_5)^{-1}$  is  $c_5 |X|^{-3}$ , so  $\nabla m_{\text{near}}$  has kernel  $O(|X|^{-4})$ , which is *locally integrable* in  $\mathbb{R}^5$ ; hence

$$|\nabla m_{\text{near}}(X)| \leq C \int_{|X-Y| \leq 4R} |X-Y|^{-4} |\Omega(Y)| dY \leq C R \|\Omega\|_{L^\infty(B_{2R})}.$$

This is the whole point: a kernel of order  $-1$  is integrable and an order-0 kernel is not, which is where the logarithm of Proposition 38.9 lives. The far part is  $\Delta_5$ -harmonic on  $B_{2R}$ , so interior gradient estimates give  $\|\nabla m_{\text{far}}\|_{L^\infty(B_R)} \leq C R^{-1} \|m_{\text{far}}\|_{L^\infty(B_{2R})}$ , and  $\|m_{\text{far}}\| \leq \|m\| + \|m_{\text{near}}\| \leq \|m\| + C R^2 \|\Omega\|_{L^\infty(B_{2R})}$ .  $\square$

**Proposition 38.12** (A scale-invariant structural hypothesis that buys the axis, and only the axis). *Let  $u$  be a smooth axisymmetric solution on the smooth interval satisfying the Type-I bounds of §27, let  $M = M(t)$ ,  $r^* = M^{-1/2}$ , and let  $x_0$  be a point of the axis. Consider*

$$(\mathbf{AX}) \quad |\omega_\theta(x, t)| \leq K M \frac{r}{r^*} \quad \text{on } B_{2r^*}(x_0), \quad \text{equivalently} \quad \|\Omega\|_{L^\infty(B_{2r^*}(x_0))} \leq \frac{K M}{r^*}.$$

(i)  $(\mathbf{AX})$  is invariant under the Navier–Stokes scaling  $u_\lambda(x, t) = \lambda u(\lambda x, \lambda^2 t)$ , with the same  $K$ . (ii) Under  $(\mathbf{AX})$ ,  $|\nabla u|(0, z) \leq C(K + c_I) M$  for  $|z - z_0| < r^*/2$ , with no factor  $\mathcal{L}$ . (iii) The conclusion of (ii) holds on the axis only; by Proposition 38.9 it fails on  $Q(r^*)$ , and the counterexample there is supported inside  $B_{r^*}$  around the same axis point. [Proved.]

*Proof.* (i) Under the scaling,  $\omega_\lambda = \lambda^2 \omega(\lambda x, \lambda^2 t)$ ,  $M_\lambda = \lambda^2 M$ ,  $r_\lambda^* = r^*/\lambda$  and  $\Omega_\lambda = \lambda^3 \Omega(\lambda x, \cdot)$ , so  $\|\Omega_\lambda\|/(M_\lambda/r_\lambda^*) = \lambda^3 \|\Omega\|/(\lambda^3 M/r^*)$  is unchanged. (ii) Apply Lemma 38.11 with  $R = r^*$ . The first term is  $C r^* \cdot K M / r^* = C K M$ . For the second, (195) and the Type-I velocity bound give  $\|m\|_{L^\infty} \leq \frac{1}{2} \|u\|_{L^\infty} \leq c_I / (2r^*)$ , so  $C R^{-1} \|m\| \leq C c_I (r^*)^{-2} = C c_I M$ . Add Lemma 38.7, which contributes  $\frac{3}{2} M$ . (iii) is Proposition 38.9 with the stated placement.  $\square$

**Remark 38.13** (The answer to the question this section was written to settle). The Calderón–Zygmund logarithm is not removed by the axisymmetric reduction, and it is not weakened. Precisely:

- (a) *Three of the eight entries become free.* The swirl block obeys  $|\nabla u_{\text{swirl}}| \leq \frac{3}{2}M$  pointwise, with an absolute constant, unconditionally (Lemma 38.7). This is strictly better than anything available in the general case, and better than Lemma 36.11, which is logarithm-free but returns the wrong exponent: here both the logarithm and the exponent are right.
- (b) *Five of the eight do not.* On the poloidal block the factor  $\mathcal{L}$  is *sharp*, by an explicit field supported inside  $B_{r^*}$  around the axis point (Proposition 38.9). Wall 1(a) — the demand  $\|\nabla u\|_\infty \leq KM$  — is therefore false as a general inequality on this class, exactly as it is in the general case.
- (c) *The axis is the one exception, and it is a null set.* On  $\{r = 0\}$  the poloidal recovery is an order-(-1) operator and is logarithm-free (Lemma 38.11) — but against  $\|\omega_\theta/r\|_\infty$ , not  $M$ . The exchange rate is exactly one power of the axis weight. Quantitatively, at the critical scale  $r^* = M^{-1/2}$  the unconditional bound  $r^*\|\Omega\|_{L^\infty(B_{r^*})}$  can be as large as  $M^{3/2}$ , whereas the logarithmic route gives  $M\mathcal{L}$ ; since  $M^{1/2} \gg \mathcal{L}$ , **the logarithm-free route is strictly the worse of the two** unless a hypothesis such as **(AX)** is supplied.
- (d) *And (AX) is not free.* **(AX)** says that  $\omega_\theta$  does not concentrate closer to the axis than the self-similar scale; for smooth fields it is, to leading order, the statement  $\|\nabla \omega\|_{L^\infty} \lesssim M/r^* = M^{3/2}$  near the axis — and the only control of  $\nabla \omega$  this document possesses (Lemma 27.6) is an  $L^2$  bound that itself carries  $\mathcal{L}$ . **(AX)** is therefore a no-concentration hypothesis of the same family as saturation (§37), and S50's lesson applies without modification: *it has not been shown to be cheaper than what it replaces, and this section does not claim that it is.*

The shape of the outcome should be recorded explicitly, because it is the third occurrence of the same pattern: Observation 36.19 produced a logarithm-free gradient bound against the wrong right-hand side  $(T - t)^{-1}$ ; Corollary 37.22 produced a saturation bound that returned exactly the estimate already held; Lemma 38.11 produces a logarithm-free bound against the larger quantity  $\|\Omega\|_\infty$ . In all three the logarithm is not paid and is not avoided — it is converted into a defect on the right-hand side of equal or greater size. Wall 1 is structural in the sense of §44.2, and this class does not evade it.

### 38.3 $\sigma^*$ , the coherence deficit, and the Gram machinery in axisymmetric variables

**Lemma 38.14** (Exact decomposition of the coherence deficit). *Let  $\hat{e} = \omega/|\omega|$  on  $\{\omega \neq 0\}$  for an axisymmetric field. Then*

$$w = |\nabla \hat{e}|^2 = |\partial_r \hat{e}|^2 + |\partial_z \hat{e}|^2 + \frac{1 - \hat{e}_z^2}{r^2} = |\nabla_{(r,z)} \hat{e}|^2 + \frac{J^2 + \Omega^2}{|\omega|^2}, \quad (196)$$

where  $J = \omega_r/r$  and  $\Omega = \omega_\theta/r$ . In particular  $w$  carries a non-negative curvature floor  $(1 - \hat{e}_z^2)/r^2$  involving no derivative of  $\hat{e}$  at all. [Proved.]

*Proof.* For an axisymmetric vector field  $f$  the components are  $\theta$ -independent, but the frame is not:  $\frac{1}{r}\partial_\theta f = \frac{1}{r}(f_r e_\theta - f_\theta e_r)$ . Hence  $|\nabla f|^2 = |\partial_r f|^2 + |\partial_z f|^2 + (f_r^2 + f_\theta^2)/r^2$ . Apply this to

$f = \hat{e}$  and use  $|\hat{e}| = 1$ , so  $\hat{e}_r^2 + \hat{e}_\theta^2 = 1 - \hat{e}_z^2$ . Finally  $1 - \hat{e}_z^2 = (\omega_r^2 + \omega_\theta^2)/|\omega|^2$ , and dividing by  $r^2$  turns  $(\omega_r, \omega_\theta)$  into  $(J, \Omega)$ .  $\square$

**Corollary 38.15** ( $\sigma^*$  has an axisymmetric floor). *Let  $x^*$  lie on the axis of symmetry (which, by [1], is where an axisymmetric singularity must lie). Then  $r \leq r^*$  on  $B_{r^*}(x^*)$  and, with  $\mathcal{E} = B_{r^*}(x^*) \cap \{|\omega| \geq M/4\}$ ,*

$$\sigma^*(t) \geq (r^*)^{-3} \int_{\mathcal{E}} (1 - \hat{e}_z^2) dx.$$

*In particular, if  $|\mathcal{E}| \geq c_N(r^*)^3$  then  $\sigma^* \leq \varepsilon_0$  forces  $\langle 1 - \hat{e}_z^2 \rangle_{\mathcal{E}} \leq \varepsilon_0/c_N$ : small coherence deficit means the vorticity is, on average, aligned with the symmetry axis. [Proved, from Definition 26.3 and Lemma 38.14.]*

**Corollary 38.16** (The constant direction of the aligned horn can only be the axis). *If  $\nabla \hat{e} \equiv 0$  on an open subset of  $\{r > 0\}$ , then  $\hat{e} \equiv \pm e_z$  there. Hence the direction  $e_0$  produced by Lemma 33.13 in the aligned horn of §33.4 is, in this class, forced to be the symmetry axis. [Proved: the floor in (196) gives  $1 - \hat{e}_z^2 = 0$ .]*

**Proposition 38.17** (The aligned horn is exactly the vanishing of the swirl source). *Let  $U \subseteq \{r > 0\}$  be open and suppose  $\hat{e} \equiv \pm e_z$  on  $U$  (equivalently, by Lemma 38.14, the curvature part of  $w$  vanishes on  $U$ ). Then on  $U$ :*

- (a)  $\omega_r \equiv 0$ , hence  $\partial_z \Gamma = r \partial_z u_\theta = -r \omega_r \equiv 0$ , hence the source term  $\partial_z(\Gamma^2)/r^4$  of (190) vanishes identically;
- (b)  $\Gamma$  is not forced to vanish:  $\Gamma = \Gamma(r)$  is admissible, and any such  $\Gamma \not\equiv 0$  is a genuinely swirling field.

[Proved.]

*Proof.* (a)  $\hat{e} = \pm e_z$  gives  $\omega_r = \omega_\theta = 0$ ; then  $\omega_r = -\partial_z u_\theta$  and  $\partial_z \Gamma = r \partial_z u_\theta$  give the claim, and  $\partial_z(\Gamma^2) = 2\Gamma \partial_z \Gamma = 0$ . (b) Take  $u_\theta = u_\theta(r)$ : then  $\omega_r = -\partial_z u_\theta = 0$  as required, while  $\Gamma = r u_\theta(r) \not\equiv 0$ .  $\square$

**Remark 38.18** (What Proposition 38.17 is and is not). It is *not* a regularity theorem, and it must not be read as one. What it establishes is a dictionary entry: the programme's aligned horn, which was constructed for entirely unrelated reasons in §26, coincides in this class with the exact condition under which the axisymmetric problem degenerates to its solved case. It is also the sharp form of the check against the “accidentally reduced to no swirl” failure mode: alignment forces  $\partial_z \Gamma \equiv 0$ , *not*  $\Gamma \equiv 0$  — so the aligned horn sits precisely on the boundary between the swirling and swirl-free problems, without falling into the latter. The Ladyzhenskaya–Ukhovskii–Yudovich mechanism (Fact 38.21) is exactly the disappearance of this source.

**Remark 38.19** (The Gram machinery transfers verbatim, and gains nothing). Lemma 26.1 (the magnitude and direction equations), Proposition 26.4 (scale invariance) and Lemma 30.2 (Gram negativity) are algebraic identities among Cartesian quantities  $\omega, \hat{e}, m = |\omega|, S$ , valid for every smooth solution. Axisymmetric solutions form a subclass, so all three hold verbatim with no re-derivation; in particular the Gram matrix  $G_{jk} = \partial_j \hat{e} \cdot \partial_k \hat{e}$  is still positive semidefinite



and the dangerous couplings of C1 are still good-signed. This is recorded for completeness, and the honest reading is that the reduction buys nothing here: the only place the symmetry enters the geometric layer is the floor in (196), which is new, and Corollary 38.16, which is a consequence of it.

**Remark 38.20** (The rediscovery check:  $\sigma^*$  lands on the literature's own two scalars). By (196) the curvature part of the coherence deficit is exactly  $(J^2 + \Omega^2)/|\omega|^2$  — the *normalised* version of the pair  $(J, \Omega)$  on which Chen–Fang–Zhang [38] and Lei–Zhang [39] build their entire analysis (Fact 38.27). This is not a coincidence to be claimed as a discovery; it is a warning. The literature controls  $(J, \Omega)$  *dynamically*, by energy estimates on the closed system they satisfy, in  $L_t^\infty L_{\text{loc}}^2 \cap L_t^2 \dot{H}_{\text{loc}}^1$ . The  $\sigma^*$  machinery controls the same quantities only *geometrically* and only after normalisation by  $|\omega|^2$ , which destroys exactly the homogeneity that makes their energy estimate close. On this class the literature's handle on the same objects is strictly stronger.

### 38.4 H1 and H2 under the reduction; what the literature has already discharged

The statements below are quoted from the sources, not recalled.

*Fact 38.21* (No swirl  $\Rightarrow$  global regularity). If  $u_\theta \equiv 0$ , finite-energy axisymmetric solutions of the Navier–Stokes equations on  $\mathbb{R}^3$  are smooth for all time (Ladyzhenskaya [35]; Ukhovskii–Yudovich [36]). The mechanism is the disappearance of the source in (190), after which  $\Omega$  obeys a maximum principle. *[Cited.]*

*Fact 38.22* (Chen–Strain–Tsai–Yau 2008). *[Quoted from [37, Thm. 1.1].]* Let  $(v, p)$  be an axisymmetric solution of Navier–Stokes in  $D = \mathbb{R}^3 \times (-T_0, 0)$ , smooth in  $x$  and Hölder continuous in  $t$ , with  $p \in L^{5/3}(D)$ , and suppose

$$|v(x, t)| \leq C^*(r^2 - t)^{-1/2} \quad \text{on } D, \quad 0 < C^* < \infty.$$

Then  $v \in L^\infty(B_R \times [-T_0, 0])$  for every  $R > 0$ . *Note the hypothesis is **stronger** than temporal Type I:* since  $(r^2 - t)^{-1/2} \leq (-t)^{-1/2}$ , it implies the Type-I bound and in addition forces decay in  $r$ . *[Cited.]*

*Fact 38.23* (Koch–Nadirashvili–Seregin–Šverák 2009, the axisymmetric application). *[Quoted from [17, Thms. 5.3, 6.1, 6.2].]* Theorem 5.3: a bounded weak ancient axisymmetric solution on  $\mathbb{R}^3 \times (-\infty, 0)$  with  $|u(x, t)| \leq C/\sqrt{x_1^2 + x_2^2}$  vanishes identically. Theorem 6.2: if  $u$  is an axisymmetric weak solution on  $\mathbb{R}^3 \times (0, T)$ , bounded on  $\mathbb{R}^3 \times (0, T')$  for each  $T' < T$ , satisfying both

$$|u| \leq C(T - t)^{-1/2} \quad \text{on } \mathbb{R}^3 \times (0, T) \quad \text{and} \quad |u(x, t)| \leq C/\sqrt{x_1^2 + x_2^2} \quad \text{for } r \geq R_0,$$

then  $|u| \leq M(C)$  and  $u$  is a mild solution. *The authors state explicitly that the statement fails if the second hypothesis is dropped* (take  $u(x, t) = b(t)$ ). So [17] alone does *not* exclude Type I in this class without a far-field decay assumption. *[Cited.]*

*Fact 38.24* (Seregin–Šverák 2009: Type I is excluded, locally and unconditionally). *[Quoted from [22, Thms. 1.1–1.2].]* Let  $z_0 = (x_0, t_0)$  with  $x_0$  on the axis, and let  $\bar{v}$  denote the projection of  $v$  onto the plane through  $x$  and the axis (the poloidal part).

- *Theorem 1.1.* If  $v \in L^3(Q(z_0, R))$  is an axisymmetric weak solution with associated pressure  $q \in L^{3/2}(Q(z_0, R))$  and

$$\operatorname{ess\,sup}_{(x,t) \in Q(z_0, R)} \sqrt{t_0 - t} |\bar{v}(x, t)| < +\infty,$$

then  $z_0$  is a regular point of  $v$ .

- *Theorem 1.2.* The same conclusion holds, for  $v$  in addition essentially bounded on  $B(x_0, R) \times ]t_0 - R^2, t'[$  for each  $t' < t_0$ , if instead  $\operatorname{ess\,sup} \sqrt{x_1^2 + x_2^2} |\bar{v}| < +\infty$ .

There is **no** swirl-smallness and **no** far-field decay hypothesis, and the Type-I bound is imposed on the *poloidal part alone*. The standard Type-I bound  $|v| \leq C(T - t)^{-1/2}$  implies the hypothesis of Theorem 1.1 trivially. [Cited.]

**Proposition 38.25** (H1 and H2 in this class have no remaining consequence). *Restricted to the axisymmetric class:*

- (a) **H1** (Conjecture 26.8) becomes: averaged smallness of  $(J^2 + \Omega^2)/|\omega|^2$  over  $B_{r^*}$  upgrades to the pointwise bound  $|\nabla \hat{e}| \leq C_B/r^*$ , which by (196) contains  $1 - \hat{e}_z^2 \leq C_B^2 r^2/(r^*)^2$  — alignment with the axis improving linearly as the axis is approached — equivalently a pointwise bound on  $\partial_z \Gamma/(r|\omega|)$  near the axis.
- (b) **H2** (the localized Constantin–Fefferman input of Theorem 26.9) keeps its statement verbatim, since  $\alpha = \hat{e} \cdot S \hat{e}$  and  $M(t)$  are frame-independent; but note that in this class the quantity whose growth actually has to be beaten in the  $\Omega$  equation is the source  $\partial_z(\Gamma^2)/r^4$ , not a stretching term, so H2 is not the natural instrument here.
- (c) Whatever is gained is already known. Theorem 26.9 concludes, given H1 and H2, that no Type-I singularity satisfies  $\liminf \sigma^* \leq \varepsilon_0$ ; the programme’s ceiling, even with both horns closed, is “no Type-I singularity” (Remark 33.30). By Fact 38.24 that conclusion is a theorem of 2009 in this class, proved locally, for weak solutions, with no non-degeneracy hypothesis (**N**), no (**R**), no (**A-up**) and no  $\sigma^*$ . Proving H1 and H2 in this class would therefore re-prove a known theorem under strictly more hypotheses.

[Proved, as a comparison of statements.]

**Remark 38.26** (Attribution). The exclusion of Type-I singularities for axisymmetric solutions with swirl is due to [37], [17] and [22], in the exact forms quoted in Facts 38.22–38.24. **It is not a result of this programme**, nothing above strengthens it, and no session should report it as such. The programme’s own contribution in this subsection is negative: it identifies that its Type-I machinery has nothing left to prove here.

### 38.5 The intended target — a criterion in $\Gamma$ alone — and why it is not a target

*Fact 38.27* (Criticality of the axisymmetric system). [Quoted from [39], as presented in [41, §2.2].] With  $J = \omega_r/r$ ,  $\Omega = \omega_\theta/r$  and  $b = u_r e_r + u_z e_z$ , the triple  $(J, \Omega, \omega_z)$  satisfies the closed system

$$\begin{cases} \partial_t J + b \cdot \nabla J = \Delta J + \frac{2}{r} \partial_r J + (\omega_r \partial_r + \omega_z \partial_z) \frac{u_r}{r}, \\ \partial_t \Omega + b \cdot \nabla \Omega = \Delta \Omega + \frac{2}{r} \partial_r \Omega - 2 \frac{u_\theta}{r} J, \\ \partial_t \omega_z + b \cdot \nabla \omega_z = \Delta \omega_z + \omega_r \partial_r u_z + \omega_z \partial_z u_z, \end{cases}$$

and the a priori bound  $|u_\theta| \leq r^{-1} \|ru_\theta(\cdot, 0)\|_\infty$  makes the potentials in the stretching terms scale like  $r^{-2}$ , i.e. *critical* rather than supercritical. The scaling gap for the axisymmetric system is therefore 0. [Cited. Note that the second equation is (190):  $-2(u_\theta/r)J = 2u_\theta \partial_z u_\theta / r^2 = \partial_z(\Gamma^2)/r^4$ , which this session verified independently in Lemma 38.4.]

**Fact 38.28** (The  $\Gamma$  criteria, with their exact hypotheses). [Quoted.]

- (a) *Chen–Fang–Zhang [38]*: if  $|u_\theta(x, t)| \leq Cr^{-1+\varepsilon}$  for all  $x$  and  $t > 0$ , with  $\varepsilon > 0$ , then the solution is regular everywhere.
- (b) *Lei–Zhang [39], Theorem*: for every  $C_* > 1$  there is  $\delta_* > 0$  such that every Leray–Hopf solution with  $\|v_0\|_{H^{1/2}} < \infty$  and  $\|ru_\theta(\cdot, 0)\|_\infty < \infty$  whose angular velocity lies in the scale-invariant  $(\delta_*, C_*)$ -critical class (a form-boundedness condition on  $|u_\theta|$  against  $\|\partial_r f\|_2$ ) is globally regular. A sufficient condition is  $\sup_{0 \leq t < T} |ru_\theta(r, z, t)| \leq C_1 |\ln r|^{-2}$  for  $r \leq \delta_0 \in (0, \frac{1}{2})$ .
- (c) *Wei [40]*: the same conclusion under  $\sup_{0 \leq t < T} |ru_\theta(r, z, t)| \leq C_1 |\ln r|^{-3/2}$  for  $r \leq \delta_0$ , with  $v_0 \in H^2$  and  $\|ru_\theta(\cdot, 0)\|_\infty < \infty$ .
- (d) *Free, for comparison*:  $\|\Gamma(\cdot, t)\|_\infty \leq \|\Gamma(\cdot, 0)\|_\infty$  (Lemma 38.3).

| Source      | Hypothesis on the swirl                                                     | Conclusion        |
|-------------|-----------------------------------------------------------------------------|-------------------|
| [35],[36]   | $\Gamma \equiv 0$                                                           | global regularity |
| [38]        | $ \Gamma  \leq Cr^\varepsilon$ , i.e. $ u_\theta  \leq Cr^{-1+\varepsilon}$ | global regularity |
| [39]        | $ \Gamma  \leq C_1  \ln r ^{-2}$ near the axis (or the FBC)                 | global regularity |
| [40]        | $ \Gamma  \leq C_1  \ln r ^{-3/2}$ near the axis                            | global regularity |
| <b>free</b> | $\ \Gamma\ _\infty \leq \ \Gamma_0\ _\infty$                                | —                 |

**Proposition 38.29** (A boundedness-type  $\Gamma$  criterion is the whole problem). *Let (P) be any assertion of the form: “every Leray–Hopf axisymmetric solution with  $\|\Gamma(\cdot, 0)\|_{L^\infty} < \infty$  and  $\sup_t \|\Gamma(\cdot, t)\|_{L^\infty} \leq \Lambda$  is globally regular”, with no decay of  $\Gamma$  toward the axis required. Then (P) is equivalent to global regularity for all axisymmetric Leray–Hopf data with  $\Gamma_0 \in L^\infty$ : the hypothesis of (P) is discharged for every such solution, with  $\Lambda = \|\Gamma_0\|_\infty$ , by Lemma 38.3. [Proved, trivially — which is the point.]*

**Remark 38.30** (Verdict on the intended target: no criterion is claimed). The WP3 brief set as its target “a regularity criterion in terms of  $\Gamma$  alone (or  $\Gamma$  plus one scale-invariant quantity)”, with the standard that “a criterion that replaces a smallness condition with a boundedness or structural condition would be a real advance”. Proposition 38.29 shows that such a criterion is not an advance but *the entire open problem for this class*: boundedness of  $\Gamma$  is free. The residual gap in Fact 38.28 is exactly  $|\ln r|^{-3/2} \rightsquigarrow O(1)$  — purely logarithmic, and being narrowed by the exponent, not by a change of method ( $-2 \rightarrow -3/2$  between [39] and [40]).

**This section produces no new criterion, and none is claimed.** The one structural statement it does produce, Proposition 38.17, is not a criterion: it identifies when the difficulty disappears, which is the already-solved case. And by Remark 38.20 the quantity the  $\sigma^*$  route would naturally target is the normalised  $(J, \Omega)$  pair that [38] and [39] already control by a strictly stronger, dynamical method. The rediscovery risk named in the brief is confirmed to be real, and is the reason no criterion is asserted here.

**Remark 38.31** (One-component criteria: what they do and do not supply here). The brief directed that the target also be situated against the one-component literature. The relevant statements, checked:

- (a) *Within this class.* Neustupa–Pokorný [42] prove an interior regularity criterion for axisymmetric suitable weak solutions in which regularity of *one* component — either  $u_r$  or  $u_\theta$  — implies regularity of the others; Chae–Lee [43] prove regularity under finiteness of a zero-dimensional (scale-invariant) integral. Both are of the same shape as Fact 38.28: a scale-invariant *smallness or integrability* hypothesis, not a boundedness one, so Proposition 38.29 applies to them as well. Neither weakens the residual gap of Remark 38.32(4).
- (b) *Outside this class.* Kukavica–Ziane [44] and Cao–Titi [45] give general 3D criteria in terms of  $\partial_3 u$  and of a single entry of  $\nabla u$  respectively. These are Ladyzhenskaya–Prodi–Serrin-type conditions in  $L_t^p L_x^q$  and are therefore, by construction, statements the programme cannot verify for a putative singular solution: they need a priori integrability that Type I does not supply. They are recorded because they show that one-component reductions do not, by themselves, close a scaling gap — which is the hope the axisymmetric pivot rests on, and which Remark 38.13 finds unfulfilled here.

### 38.6 Ledger: what remains for Type-II axisymmetric

**Remark 38.32** (Distance to the goal in the axisymmetric class). Stated in the form of Remark 33.30, whose three items it parallels.

- (1) **The Type-I branch is closed, by others.** Fact 38.24 excludes Type-I singularities of axisymmetric solutions with swirl, locally, for weak solutions with  $q \in L^{3/2}$ , under a Type-I bound on the poloidal velocity alone. The programme’s ceiling in this class is therefore *already occupied*: every conclusion Theorem 26.9 could deliver here is a 2009 theorem (Proposition 38.25). The pivot does not open a cheaper route to the programme’s existing goal; it moves onto ground where that goal has been reached without it.
- (2) **Type-II axisymmetric is entirely open, and nothing in §27–§33 reaches it.** Item (1) of Remark 33.30 applies verbatim and with no weakening: the profile extraction, the rescaled family,  $r^* = M^{-1/2}$  and every estimate of the pincer use the Type-I rates essentially. A Type-II axisymmetric singularity would rescale to nothing here exactly as it does in the general case. The transfer of the S32–S39 machinery to this class is therefore of *zero* value against the only open branch.
- (3) **The live tools against Type-II are rate-free, and they are not geometric.** The  $\Gamma$  criteria of Fact 38.28 assume no blow-up rate whatsoever; this is their real advantage over the Type-I literature, and it is why they, and not the Liouville route, are the state of the art. They are energy estimates on the closed  $(J, \Omega, \omega_z)$  system of Fact 38.27. Entering this line means competing with [39] and [40] on their own ground with their own tools; the  $\sigma^*$ /direction-field apparatus has no purchase there (Remark 38.20).
- (4) **The residual is exactly logarithmic, in a second and independent sense.**  $\|\Gamma\|_\infty$  is free; regularity is known under  $|\Gamma| \lesssim |\ln r|^{-3/2}$ . The gap is a logarithm — not the

Calderón–Zygmund logarithm  $\mathcal{L}$  of Wall 1, which Remark 38.13 shows also survives in this class, but a second one, in the spatial variable near the axis. *The axisymmetric class does not have one fewer logarithm than the general problem; it has one more.* What it has instead is that this second logarithm is not known to be sharp: there is no counterexample showing  $\Gamma \in L^\infty$  cannot suffice, whereas Proposition 38.9 shows the first one is sharp. That asymmetry — a provably immovable obstruction traded for an open one — is the only honest argument for working here, and it is an argument about prospects, not about progress.

- (5) **Cost of entry.** Remark 38.2: the programme’s standing framework is on  $\mathbb{T}^3$ , where this class does not exist. Moving to  $\mathbb{R}^3$  with finite energy is a prerequisite, not a detail.

The one-line summary: *the axisymmetric-with-swirl class keeps Wall 1 intact and sharp, has its Type-I branch already closed by the literature, and leaves a Type-II residual that the programme’s existing machinery cannot touch and whose live tools are of a different kind entirely.*

### 38.7 Verdict

**Remark 38.33** (Verdict of S51). The question posed was whether, in the axisymmetric class, the  $L^\infty$  recovery of  $\nabla u$  from  $(\omega_\theta, \Gamma)$  by two-dimensional elliptic problems in  $(r, z)$  is logarithm-free, cheaper than  $\mathcal{L}$ , or still  $\mathcal{L}$ .

- (1) **Still  $\mathcal{L}$ , and sharply so** (Proposition 38.9, Remark 38.13). The reduction converts three of the eight entries of  $\nabla u$  into free, logarithm-free, absolute-constant bounds by  $M$  (Lemma 38.7) — a real and, within this programme, unprecedented gain — but leaves the other five paying the full factor, with a counterexample supported inside  $B_{r^*}$  around the axis. On the axis itself the poloidal block is logarithm-free against  $\|\omega_\theta/r\|_\infty$ , which at the critical scale is worse than  $M\mathcal{L}$  by a factor  $M^{1/2}/\mathcal{L}$  unless the scale-invariant structural hypothesis **(AX)** of Proposition 38.12 holds; and **(AX)** has not been shown to be cheaper than what it replaces.
- (2) **The class’s Type-I branch is already closed** (Fact 38.24), so H1 and H2 have no remaining consequence here (Proposition 38.25).
- (3) **No regularity criterion is produced**, and the natural target is not a target (Proposition 38.29, Remark 38.30).
- (4) **What is new and correct** is the dictionary: the exact decomposition (196) of the coherence deficit, its curvature floor, the identification of the aligned horn with the vanishing of the swirl source (Proposition 38.17), and the free swirl block (Lemma 38.7). None of these is a step toward the Prize, and none is claimed as one.

**Recommendation, recorded so that it is not softened later.** On the evidence above, the axisymmetric pivot does not do what WP3 was commissioned to test: it does not evade Wall 1 structurally. Its residual open branch (Type-II) is unreachable by the machinery the pivot was meant to carry across. A session that wishes to continue here should do so on the literature’s terms — energy estimates on the closed  $(J, \Omega, \omega_z)$  system, aimed at the  $|\ln r|^{-3/2}$  gap — and should expect to be competing directly with [39] and [40], not extending this programme.

**Remark 38.34** (Scope). This section proves four lemmas, one sharpness proposition and one conditional bound, and quotes five sources. It does not close either horn of the pincer, does not remove Wall 1 — it strengthens the case that Wall 1 is structural — does not bear on (N), (N'), (R), (A-up) or on Type-II, and does not bring the programme nearer to the Clay problem than Remark 33.30 already states, which stands verbatim.

## 39 Type-II Blowup: What Is Known, Why This Document's Rescaling Is Vacuous Against It, and Which Branch of Mathematics Owns the Question (Session S53, WP-T1)

This section is a *reconnaissance*, commissioned as WP-T1 of `HANDOVER-S52-TYPE2.md` after the four work packages of the S46 plan were exhausted. It proves no new theorem about the Navier–Stokes equations and it proposes no new programme. Its purposes are three:

- (i) to record what the literature actually contains on Type-II blowup for the *true* three-dimensional incompressible Navier–Stokes equations — sources read during the session, not recalled (§39.2);
- (ii) to replace this document's own assertion that “a Type-II singularity would rescale to nothing” (Remark 33.30(1)) by a derivation, and to correct that assertion where the derivation does not support it (§39.3);
- (iii) to state, in the manner of Remark 33.30 itself, the verdict on whether Type-II is reachable by any adaptation of the machinery of §27–§38 (§39.4).

Labels in this section carry the prefix `wpt1` rather than a session number. Three consecutive sessions of this programme have had their session tag claimed concurrently by a parallel agent, and §38 carries `s51` labels for what the session log records as S52. A work-package prefix cannot collide.

### 39.1 Three inequivalent meanings of “Type I”, and which one each source uses

Remark 38.5 already records that “Type I” carries two incompatible meanings in this subject. The Type-II literature supplies a *third*, and it is the one the only extant results are stated in. All three must be separated before any result is quoted.

**Definition 39.1** (Three conventions). Let  $u$  be a solution with a putative first singularity at time  $T$  and singular point  $x^*$ ; write  $M(t) = \|\omega(\cdot, t)\|_{L^\infty}$ .

- (T) *Temporal* (this document; [17]; [53] eq. (4.2)).  $M(t) \leq C_I(T - t)^{-1}$  and  $\sup_x |u(x, t)| \leq C_I(T - t)^{-1/2}$ . This is the standing assumption of §27 (line 5565) and the hypothesis used *essentially* in §33.2.
- (S) *Spatial* (much of the axisymmetric literature).  $|u(x, t)| \leq C/r$ ,  $r = |x'|$  the distance to the symmetry axis. This is what makes the swirl equation critical; it neither implies nor is implied by (T). Recorded at S52 as Remark 38.5.

(E) *Scale-invariant energy* (Caffarelli–Kohn–Nirenberg; Seregin). With

$$A(v, r) = \sup_{-r^2 < t < 0} \frac{1}{r} \int_{B(r)} |v(x, t)|^2 dx, \quad E(v, r) = \frac{1}{r} \iint_{Q(r)} |\nabla v|^2 dz, \quad C(v, r) = \frac{1}{r^2} \iint_{Q(r)} |v|^3 dz,$$

the origin is a Type-I blowup if these stay bounded as  $r \rightarrow 0$ . Two variants are in use by the same author:  $g = \max\{\limsup_{r \rightarrow 0} A, \limsup_{r \rightarrow 0} E, \limsup_{r \rightarrow 0} C\} < \infty$  in [46], and  $g_0 = \min\{\liminf_{r \rightarrow 0} A, \liminf_{r \rightarrow 0} E, \liminf_{r \rightarrow 0} C\} < \infty$  in [49]. Type-II means the negation. Since  $g_0 \leq g$ , Type-II in the  $g_0$  sense is the *stronger* requirement: it asks that all three quantities diverge.

[Definitions transcribed from the cited sources during this session; the three displayed quantities are quoted from [49], which states  $C$  with  $|v|^3$ .]

**Fact 39.2 ((T) implies (E)).** The pointwise bound  $\sqrt{T-t} |u(x, t)| \leq M'$  implies the *generalised* Type-I bound

$$\sup_{\bar{x} \in \mathbb{R}^3} \sup_{r \in (0, r_0)} \sup_{T-r^2 < t < T} r^{-1/2} \left( \int_{B_{\bar{x}}(r)} |u(x, t)|^2 dx \right)^{1/2} \leq M,$$

i.e. a uniform bound on the quantity  $A$  of Definition 39.1(E). The implication is not formal — it is a theorem of Seregin–Zajączkowski [50]; see [53], footnote 8, which states it in exactly these terms (“It turns out to be true, see [50] and the review article [51], pages 844–849. This makes (4.1) a good notion.”). [Cited, not proved here.]

**Corollary 39.3** (Seregin’s Type-II is a strictly smaller hypothesis class than this document’s). *A blowup that is Type-II in the sense of [49] ( $g_0 = \infty$ ) is Type-II in the sense of this document ((T) fails). The converse implication is available from no source read here.* [Proof: contrapositive of Fact 39.2, since  $\liminf_{r \rightarrow 0} A < \infty$  already gives  $g_0 < \infty$ .]

**Remark 39.4** (The middle band, and why it is the interesting one). Corollary 39.3 splits the space of putative singularities into three, not two:

(R1) **(T) holds.** The target of §26–§33. Open in general; closed in the axisymmetric class by Fact 38.24.

(R2) **(T) fails but  $g_0 < \infty$ .** The scale-invariant energies stay bounded while the pointwise rate is supercritical. Whether this region contains a solution is unknown; so is whether it is empty. *Nothing addresses it:* the machinery of this document needs (T), and Seregin’s Type-II papers assume  $g_0 = \infty$ . Seregin [46] states of the corresponding Type-I question: “The question about Type I blowups is whether the boundedness of  $g$  allows blowups or not. It is still open.”

(R3)  $g_0 = \infty$ . Seregin’s Type-II programme, Fact 39.5.

The existence of (R2) is the first reason Remark 33.30(1) overstates its case; see Remark 39.12.

## 39.2 Deliverable 1 — the state of knowledge, with sources

*Fact 39.5* (The Type-II literature for true 3D Navier–Stokes is one author’s programme, and it is entirely conditional). Four papers, all by Seregin, all read during this session: [46] (arXiv:2304.04045, CPAA 23(10):1389–1406, 2024), [47] (arXiv:2402.13229v3, 8 Aug 2026), [48] (arXiv:2507.08733, rev. 3 Jan 2026), [49] (arXiv:2606.29468, 28 Jun 2026). Their shape is uniform:

- (a) A *scenario* is fixed: a rate function  $g$  (or an exponent  $m_0$ ) quantifying how fast  $M_\kappa^{s,l}(v, r)$  diverges, plus a second family of bounds  $A_f + E_f + D_f \leq M_1$  on weighted energy quantities. Nothing is assumed about a pointwise rate.
- (b) A *zoom based on the Euler scaling*  $v^{\lambda, \alpha}(y, \tau) = \lambda^\alpha v(\lambda y, \lambda^{\alpha+1} \tau)$ ,  $\alpha > 1$ , extracts a nontrivial ancient solution of the **Euler** equations on  $\mathbb{R}^3 \times (-\infty, 0)$  satisfying a local energy inequality ([46], Prop. 1.2; [49], Thm. 2.1).
- (c) A *Liouville theorem* for ancient Euler solutions in the weighted class defined by the scenario then contradicts nontriviality, and the scenario is excluded.

What is actually excluded is narrow and explicitly so. In [46]: the limit “cannot be irrotational”, and  $m < 1/2$  is impossible, so the parameter is confined to  $m \geq 1/2$  ( $m_0 \geq 4/5$  suffices). In [48], Thm. 5.1 excludes one scenario *under an added Ladyzhenskaya–Prodi–Serrin-type hypothesis*  $v \in L^{s_2, l_2}(Q)$  with  $3/s_2 + (\alpha + 1)/l_2 = \alpha$ ; the author’s own Remark 5.2 says it “is impossible under the additional assumption (5.2)”. In [49], Thm. 2.1 excludes the pair of scenarios  $(g, f)$  whenever an explicit exponent inequality between  $g$  and  $f$  holds. **No unconditional exclusion of Type-II blowup for 3D Navier–Stokes exists in this literature, and none is claimed.**

*Fact 39.6* (What *is* unconditional is rate-free, and it constrains rather than excludes). A separate branch says things about *every* blowup and therefore about Type-II, at the cost of saying only that critical norms must diverge:

- (a) Leray’s lower bounds  $\|u(\cdot, t)\|_{L^q(\mathbb{R}^3)} \gtrsim (T - t)^{-\frac{1}{2}(\frac{1}{3} - \frac{1}{q})}$  for  $q \in (3, \infty)$  ([53], eq. (5.1)). Type-I is the matching *upper* rate; Type-II is everything faster.
- (b) Escauriaza–Seregin–Šverák [16]: no blowup with  $u \in L_t^\infty L_x^3$ . No rate is assumed, so it applies verbatim to Type-II.
- (c) Tao [52]: quantitatively, the critical  $L^3$  norm must exceed  $(\log \log \log \frac{1}{T-t})^c$  along a sequence of times. Again rate-free.
- (d) Barker–Prange [53], Result D: a slightly supercritical Orlicz norm  $L^3(\log \log \log L)^{-c}$  blows up, for Leray–Hopf solutions, with no Type-I hypothesis.

None of these excludes Type-II. They exclude *bounded critical norm*, which a Type-II blowup would in any case violate. Their quantitative content — a triple logarithm — is far below any rate that would separate Type-I from Type-II. Barker–Prange also state the boundary of the conditional theory in terms: “the regularity in the Type I case is proved only for axisymmetric flows” ([53], §7), which is Fact 38.24 seen from the other side.



*Fact 39.7* (The best unforced numerical scenario is Type-I-shaped, not Type-II). Hou [54] (arXiv:2107.06509v2; FoCM 2022) reports, for the 3D axisymmetric Navier–Stokes equations with smooth finite-energy data: “Both  $R(t)$  and  $Z(t)$  seem to scale like  $O((T-t)^{1/2})$ . The maximum vorticity grows like  $O((T-t)^{-1})$ .” That is **(T)** exactly — isotropic scales, vorticity at the critical rate. The same paper records the caveat that matters here: “the potentially singular solution of the Navier–Stokes equations cannot be asymptotically self-similar [Nečas–Růžička–Šverák; Tsai]. Thus the potential blow-up rate is most likely not exactly equal to  $1/2$ ”, and that the uniform upper bound  $\|u(t)\|_{L^\infty} \leq c(T-t)^{-1/2}$  “would be almost impossible to verify numerically”. So the numerics are *consistent with* Type-I and cannot rule out a mild Type-II correction; what they do *not* do is supply any positive indication of Type-II.

*Fact 39.8* (The one Type-II profile in the true-3D-NS literature is forced, and it is anisotropic). The OpenAI construction [55] (Theorem 1.1: for every  $\nu > 0$  there is  $f \in C_c^\infty$  and a smooth  $u$  on  $\mathbb{R}^3 \times [0, 1)$  with  $u(\cdot, 0) = 0$ ,  $\sup_t \|u(t)\|_{L^2} < \infty$  and  $\limsup_{t \uparrow 1} \|u(t)\|_{L^\infty} = \infty$ ) has, in the paper’s own notation with  $\tau = 1 - t$ ,

$$\begin{aligned} \ell_r &\asymp \tau^{1/2}, & \ell_z &\asymp \tau^{1/2-h}, & 0 < h < 1/100, \\ |u_\theta^{(0)}|, |u_z^{(0)}| &\asymp \tau^{-1/2-h}, & |u_r^{(0)}| &= O(\tau^{-1/2}). \end{aligned}$$

Two things follow, and only the first was on record in this programme (in the companion volume, ch. 17, **sec:s49-two-strands**; not in this document). *First*, the profile is Type-II in the sense **(T)**: the velocity exceeds  $(T-t)^{-1/2}$  by  $\tau^{-h}$ . *Second, and this is the sharper point*: the two length scales differ,  $\ell_r/\ell_z \asymp \tau^h \rightarrow 0$ . A single-parameter parabolic rescaling has one length; it cannot resolve a two-scale core whatever its amplitude. The obstruction of Remark 33.30(1) is therefore not only a rate mismatch, it is a geometry mismatch, and the geometry mismatch is the one exhibited by the only construction in existence. *[The construction is **forced**. Wall 9 of §44 governs: a forced blowup is not evidence about the unforced equations in either direction. It is quoted here as a profile, not as evidence.]*

**Remark 39.9** (“Open” versus “expected”, kept apart). On the evidence assembled above:

- Type-II blowup for unforced 3D Navier–Stokes is **open**. No source read here excludes it, unconditionally or under any hypothesis discharged by the equations.
- Type-II is **not** “expected” in the sense of heuristic or numerical support. No source read here asserts it occurs. The only unforced numerical candidate is Type-I-shaped (Fact 39.7); the only Type-II profile is forced (Fact 39.8); and Seregin’s four papers treat Type-II throughout as a possibility to be *ruled out*, never as an expectation.
- What *is* on record against the alternative is that the cleanest Type-I mechanism is dead: backward self-similar blowup with finite energy is excluded ([15], and Tsai’s improvement, as cited by [54]). So neither type has positive evidence; one of them has a specific mechanism excluded.

These are three different states of knowledge and this section does not merge them. In particular nothing here supports a sentence of the form “Type-II is the expected scenario”.

### 39.3 Deliverable 2 — the vacuity, derived; and exactly how general it is

Remark 33.30(1) asserts that without  $M(t) \lesssim (T-t)^{-1}$  “the rescaled family (152) has no uniform bound and there is no limit object at all”. The first clause is re-derived and confirmed below; the second is too strong, and the correct statement is more useful.

**Lemma 39.10** (The two-parameter rescaling family and its effective viscosity). *Let  $u, p$  solve  $\partial_t u + (u \cdot \nabla)u - \nu \Delta u + \nabla p = 0$ ,  $\operatorname{div} u = 0$ . For  $A, L, \tau > 0$  set*

$$v(y, s) := A u(x^* + Ly, T + \tau s), \quad \tilde{p}(y, s) := \frac{A\tau}{L} p(x^* + Ly, T + \tau s).$$

Then  $\operatorname{div}_y v = 0$  and

$$\partial_s v + \frac{\tau}{AL} (v \cdot \nabla_y) v - \nu \frac{\tau}{L^2} \Delta_y v + \nabla_y \tilde{p} = 0.$$

In particular, normalising the nonlinearity by  $\tau = AL$  gives

$$\partial_s v + (v \cdot \nabla_y) v - \nu_{\text{eff}} \Delta_y v + \nabla_y \tilde{p} = 0, \quad \boxed{\nu_{\text{eff}} = \nu \frac{A}{L}}.$$

[Proved here, by direct computation.]

*Proof.*  $\partial_s v = A\tau(\partial_t u)$ ;  $(v \cdot \nabla_y)v = A^2 L (u \cdot \nabla_x)u$ ;  $\Delta_y v = AL^2 \Delta_x u$ ; and  $\nabla_y \tilde{p} = \frac{A\tau}{L} \cdot L \nabla_x p = A\tau \nabla_x p$ . Multiply the equation for  $u$  by  $A\tau$  and substitute:  $A\tau \partial_t u = \partial_s v$ ,  $A\tau(u \cdot \nabla)u = \frac{\tau}{AL}(v \cdot \nabla_y)v$ ,  $\nu A\tau \Delta u = \nu \frac{\tau}{L^2} \Delta_y v$ , and  $A\tau \nabla p = \nabla_y \tilde{p}$ . Divergence-freeness is immediate.  $\square$

**Corollary 39.11** (Normalisation trichotomy: the Type-I rate is not a convenience, it is the fixed point). *Suppose  $\sup_x |u(x, t)| \asymp (T-t)^{-\beta}$  for some  $\beta > 0$  and require the rescaled family of Lemma 39.10 (with  $\tau = AL$ ) to be  $O(1)$  in  $L^\infty$  on a fixed cylinder, i.e.  $A\tau^{-\beta} \asymp 1$ . Then  $A = L^{\beta/(1-\beta)}$  for  $\beta \neq 1$ , and*

$$\frac{\nu_{\text{eff}}}{\nu} = \frac{A}{L} = L^{\frac{2\beta-1}{1-\beta}}.$$

Consequently, as  $L \downarrow 0$ :

- (i)  $\beta < \frac{1}{2}$ :  $\nu_{\text{eff}} \rightarrow \infty$  — the limit is a Stokes/heat flow; no blowup mechanism survives.
- (ii)  $\beta = \frac{1}{2}$ :  $A = L$ ,  $\tau = L^2$ ,  $\nu_{\text{eff}} = \nu$ . This is (152) exactly, and it is the unique member of the family that preserves the Navier–Stokes equations.
- (iii)  $\frac{1}{2} < \beta < 1$ :  $\nu_{\text{eff}} \rightarrow 0$  — the limit solves the **Euler** equations. This is Seregin’s Euler scaling: putting  $A = \lambda^\alpha$ ,  $L = \lambda$  gives  $\tau = \lambda^{\alpha+1}$ ,  $\nu_{\text{eff}} = \nu \lambda^{\alpha-1}$  and  $\beta = \alpha/(\alpha+1)$ , so  $\alpha > 1 \iff \beta > \frac{1}{2}$ , and  $\alpha \in (1, \frac{3}{2}]$  (the range of [47],  $\alpha = 2 - m$ ,  $\frac{1}{2} \leq m < 1$ ) corresponds to  $\beta \in (\frac{1}{2}, \frac{3}{5}]$ .
- (iv)  $\beta \geq 1$ : no member of the family works at all. At  $\beta = 1$  the normalisation forces  $L = 1$  (no spatial zoom); for  $\beta > 1$  it forces  $\tau = AL \rightarrow \infty$ , contradicting  $\tau \leq T$ .

[Proved here. Item (iv) is a scope limit on the Euler-zoom method itself and is recorded because no source read here states it.]

**Remark 39.12** (Correction to Remark 33.30(1): what is right, what is too strong, and what the right statement is). *Confirmed.* With  $\beta = \frac{1}{2}$  and  $M(t) \leq C_I(T-t)^{-1}$ ,  $|u_\lambda| = \lambda|u| \leq \lambda C_I(\lambda^2|s|)^{-1/2} = C_I|s|^{-1/2}$  and  $|\omega_\lambda| = \lambda^2|\omega| \leq C_I|s|^{-1}$ , both independent of  $\lambda$ ; and if instead  $M(t) \leq C(T-t)^{-1-\delta}$  with  $\delta > 0$  then  $|\omega_\lambda| \leq C\lambda^{-2\delta}|s|^{-1-\delta} \rightarrow \infty$ . The rescaled family really does lose its uniform bound, and Corollary 39.11(ii) shows this is structural, not an artefact: (152) is the only NS-preserving rescaling there is.

*Too strong.* “There is no limit object at all” does not follow, for two independent reasons.

- (a) **Within the same rescaling.** A limit object requires uniform bounds in *some* topology, not in  $L^\infty$ . Under the strictly weaker hypothesis  $g < \infty$  of Definition 39.1(E), the same  $\lambda$ -family (152) is bounded in  $L_t^\infty L_{\text{loc}}^2 \cap L_t^2 \dot{H}_{\text{loc}}^1$  uniformly on every fixed cylinder, because  $A(u_\lambda, a) = A(u, \lambda a) \leq g$  and likewise for  $E$  and  $C$ ; with the pressure controlled, the classical Caffarelli–Kohn–Nirenberg/Lin compactness produces an ancient *suitable weak* solution of Navier–Stokes. This document already knows the phenomenon — the remark following Proposition 33.6 says the  $C_{\text{loc}}^\infty$  topology is “far more than the weak- $*$   $L^\infty$  / weak  $L_{\text{loc}}^2$  convergence one gets from energy bounds alone” — but Remark 33.30(1) does not carry the qualification. This is region (R2) of Remark 39.4.
- (b) **Outside it.** Even when  $g = \infty$ , the Euler zoom of Corollary 39.11(iii) produces a nontrivial ancient limit (Fact 39.5). “Rescale to nothing” is false there too.

*The correct statement*, and it is worse for this programme than the one it replaces, is:

A rescaling that normalises a supercritical blowup rate in  $L^\infty$  necessarily has  $\nu_{\text{eff}} = \nu A/L \rightarrow 0$ , so its limit object solves *Euler*. The vacuity is not that nothing survives; it is that what survives is **inviscid**, and every tool of §27–§35 is parabolic — Moser iteration, the magnitude Caccioppoli inequality, De Giorgi level iteration, Lemma 27.6, the  $\varepsilon$ -regularity of [1], the backward uniqueness of [16]. None of them says anything about a solution of the Euler equations.

A future session should quote this paragraph rather than Remark 33.30(1), which remains correct about the *conclusion* and imprecise about the *reason*.

**Corollary 39.13** (Dimensional reading: circulation and viscosity vanish together, so no Euler zoom can keep the swirl). *In the axisymmetric class let  $\Gamma = r u_\theta$  be the swirl invariant of Lemma 38.3. Under the rescaling of Lemma 39.10,*

$$\Gamma_{\text{new}}(y, s) = |y'| v_\theta(y, s) = \frac{|x'|}{L} \cdot A u_\theta = \frac{A}{L} \Gamma(x, t) = \frac{\nu_{\text{eff}}}{\nu} \Gamma(x, t).$$

*Hence  $\Gamma$  and  $\nu$  rescale by the same factor  $A/L$  — as they must, both having dimension (length)<sup>2</sup>/(time) — and the ratio  $\Gamma/\nu$  is invariant across the entire family. Two consequences:*

- (i) *Since  $\|\Gamma(t)\|_\infty \leq \|\Gamma(0)\|_\infty$  for free by the maximum principle (Lemma 38.3), any zoom with  $\nu_{\text{eff}} \rightarrow 0$  has  $\Gamma \rightarrow 0$ : **the Euler-zoom limit of an axisymmetric Type-II blowup is swirl-free.** This reproduces, in one line and from dimensions alone, the computation of [47] (“The latter means that our limiting Euler equations have no swirl, i.e.,  $u_\theta$  vanishes”); no novelty is claimed for the conclusion, only for the reading.*

- (ii) *Conversely — and this is the part that matters here — **any rescaling that retains the swirl retains the viscosity**, i.e. is (152), i.e. requires **(T)**. There is no rate-adapted rescaling that keeps both. The hope of an “Euler zoom that keeps the swirl” is dimensionally impossible, not merely unknown.*

[Proved here.]

**Remark 39.14** (Where this leaves the S52 coincidence). Proposition 38.17 found that the aligned horn forces  $\omega_r \equiv 0$ , hence  $\partial_z \Gamma \equiv 0$ , so that the programme’s aligned configuration sits exactly on the boundary of the solved swirl-free case. Corollary 39.13 finds that the Type-II Euler zoom forces  $\Gamma \equiv 0$  outright. The two degenerations are different in strength ( $\partial_z \Gamma \equiv 0$  admits  $\Gamma = \Gamma(r)$ ;  $\Gamma \equiv 0$  does not) and arise for unrelated reasons, and *no connection between them is asserted*. They are recorded together only so that a future session does not mistake one for the other.

**Remark 39.15** (What the other Type-II literatures need, and why it is not available here). The handover asked whether a Giga–Kohn- or Merle–Zaag-type modulation analysis transfers. The pattern in those literatures is uniform and the obstruction is the same in each case.

- *Semilinear heat*. Giga–Kohn similarity variables [56] characterise blowup under a Type-I bound; below the Sobolev exponent every blowup *is* Type-I, and Type-II appears only in the energy-critical and supercritical ranges. Its construction is by stabilisation of an explicit *bubble* — Collot–Merle–Raphaël [59] construct anisotropic Type-II blowup for  $\partial_t u = \Delta u + u^p$ ,  $p \geq 3$ ,  $d \geq 14$ , by “construction and stabilization of type II bubbles in the parabolic setting”. The bubble is the stationary ground state of the equation.
- *Geometric flows*. Type-II singularities of mean curvature flow are analysed by Hamilton’s point selection, whose limit is an *eternal* solution identified as a translating soliton via the equality case of a differential Harnack inequality.

Both routes need an object Navier–Stokes does not supply: an explicit stationary or self-similar profile to linearise around, or a differential Harnack inequality. The candidate profile for NS would be a backward self-similar solution, and those are excluded with finite energy ([15]); NS has no Harnack inequality of Hamilton type. *This is a statement about what is missing, not a proof that no substitute exists*. What Seregin’s programme does instead is exactly the admission of this: it abandons the attempt to keep a limit object solving the same equation and accepts an Euler limit, moving the whole difficulty into Liouville theorems for ancient Euler flows.

### 39.4 Deliverable 3 — verdict

**Remark 39.16** (Verdict, in the form of Remark 33.30). WP-T1 was to end in one of three ways: a scoped technical target, a reasoned negative, or a referral to a different branch of mathematics. The honest answer is the third, with the second as its justification. It is **not** the first, and no target is manufactured.

- (1) **Negative, and now derived rather than asserted**. Type-II is not reachable by any adaptation of the machinery of §27–§38. The obstruction is Corollary 39.11: within the family of rescalings that preserve the Euler part,  $\nu_{\text{eff}} = \nu A/L$ , and  $L^\infty$ -normalising

a supercritical rate forces  $\nu_{\text{eff}} \rightarrow 0$ . Every estimate this document owns is parabolic. Adapting the rescaling does not adapt the tools; it deletes the term they are built on. Corollary 39.13 closes the one apparent loophole in the axisymmetric class:  $\Gamma$  and  $\nu$  have the same dimension and vanish together.

- (2) **The branch of mathematics that owns the question is named, and it is not this one.** It is *Liouville theorems for ancient solutions of the three-dimensional Euler equations in weighted classes defined by a blowup scenario*. The papers to read are [46], [47], [48], [49], in that order; [47] is the one closest to the class of §38, where the residual problem is a Liouville theorem for *swirl-free* ancient axisymmetric Euler. This programme has no foothold in that subject: it has never written an estimate for the Euler equations, and §38 already established (Remark 38.2) that entering the axisymmetric class at all requires first moving the standing framework from  $\mathbb{T}^3$  to  $\mathbb{R}^3$ .
- (3) **Three cautions on the entry cost**, so that a future session prices it correctly.
  - (a) Seregin’s results are *scenario-conditional* (Fact 39.5), and one of them ([48], Thm. 5.1) buys its exclusion with an added Ladyzhenskaya–Prodi–Serrin hypothesis. Producing another such result is producing another conditional statement, not progress on the Clay problem. This is the same trap as Proposition 38.29: a criterion whose hypothesis is not discharged is not an advance.
  - (b) The Euler-zoom family itself stops at  $\beta = 1$  (Corollary 39.11(iv)). Velocity blowup at  $(T - t)^{-1}$  or faster is outside *both* methods.
  - (c) Region (R2) of Remark 39.4 — Type-II in the sense of **(T)** but Type-I in the sense of **(E)** — is addressed by neither this document nor Seregin’s papers. Whether it is empty is unknown. Any future claim to have “covered Type-II” must say what it does about (R2).
- (4) **What does not change.** Wall 3 (§44.4) keeps its classification OPEN. Nothing in this section is progress on Type-II, on Type-I, on Wall 1, or on any Prize or Claim A conclusion. It is a map of a region this programme has not entered and, on the evidence here, should not enter with its present tools.

**Remark 39.17** (Scope, and one defect found in passing). *Scope.* This section quotes; it does not adjudicate. Seregin’s four papers were read in the arXiv versions cited and their statements transcribed, not verified line by line. [55] was read for its Theorem 1.1 and §2 scaling only, and is quoted as a profile under Wall 9, not as evidence about the unforced equations. Facts 39.2 and 39.6 are cited, not proved. What is proved here is Lemma 39.10 and its two corollaries, and those are elementary.

*Defect found in passing, recorded rather than repaired.* The parenthetical line numbers carried by §44 have drifted. An automated audit of every “(line N)” citation whose target bears a resolvable label found twelve citations low by exactly 33 (among them Remark 33.30, cited in §44.4 as line 7670 and in fact at line 7703, and Propositions 34.12, 34.13, 34.16), one badly wrong (Lemma 33.3, cited as line 7384, in fact line 6729), four low by 10–18, one high by 11, and eight exact. The \ref targets themselves all resolve, so nothing is broken; but the line numbers are decoration that has silently gone stale, and a reader who navigates by them will land in the wrong place. They are left untouched here because this section is inserted

before §44, so every one of them keeps the value it had; a session that repairs them should do so as its own change.

## 40 WP-T2: The OpenAI Profile Tested Against Ancient-Solution Exclusion

§39 closed with the observation that the only explicit Type-II profile in the true-3D-Navier–Stokes literature is the forced construction of [55], and that it is *anisotropic* (Fact 39.8). This section asks one narrow question about that one object: *can the profile be excluded?*

**Remark 40.1** (Scope, stated before the work rather than after it). This section is a **single-profile exclusion attempt**. Its best possible outcome has the same shape as [15] excluding backward self-similar Type-I blowup: one more profile family ruled out. It is **not**, and must not be reported as, a step toward alternative (A) or (B). A logical chain from here to global regularity would additionally need Wall 1 closed on the Type-I side (untouched) *and* a general Type-II profile-extraction theorem *and* a general unconditional Liouville theorem for ancient solutions — all three recognised open problems, and the reason §39 found Seregin’s programme to be scenario-conditional throughout. WP-T2 touches none of that machinery. What follows settles one explicit object and nothing else.

### 40.1 Deliverable 0 — the primary source, and what it actually contains

Fact 39.8 was recorded at S53 from the primary PDF, but the bibliography entry carried no locator and the profile *ansatz* circulated in this programme only as a screenshot transcription. Both are repaired here: the source was re-fetched and read directly, and every formula below was checked against it rather than against any summary.

*Fact 40.2* (The source, and the status of its profile equations). The manuscript [55] is a 165-page document whose Theorem 1.1 reads, verbatim:

“For every  $\nu > 0$  there exist a force  $f \in C_c^\infty(\mathbb{R}^3 \times (0, \infty); \mathbb{R}^3)$ , a compact set  $K \subset \mathbb{R}^3$ , and smooth velocity and pressure fields  $u, p$  on  $\mathbb{R}^3 \times [0, 1)$  satisfying  $\partial_t u + (u \cdot \nabla)u - \nu \Delta u + \nabla p = f$ ,  $\nabla \cdot u = 0$ ,  $u(\cdot, 0) = 0$ , such that  $\text{supp } u(\cdot, t) \cup \text{supp } p(\cdot, t) \subset K$  for every  $0 \leq t < 1$ ,  $\sup_{0 \leq t < 1} \|u(t)\|_{L^2(\mathbb{R}^3)} < \infty$ ,  $\limsup_{t \uparrow 1} \|u(t)\|_{L^\infty(\mathbb{R}^3)} = \infty$ .”

It states that this establishes alternative (C) of [3], and (D) via the torus corollary. The construction is carried out at  $\nu = 1$ ; general  $\nu$  is recovered by  $u_\nu(x, t) = \sqrt{\nu} u(x/\sqrt{\nu}, t)$ , which leaves the singular time unchanged.

(i) **The exponents and similarity variables, verbatim from §2 and (4.1).** With  $\tau = 1 - t$ ,

$$\begin{aligned} \ell_r &\asymp \tau^{1/2}, & \ell_z &\asymp \tau^{1/2-h}, & 0 < h < \frac{1}{100}, \\ \left| u_\theta^{(0)} \right|, \left| u_z^{(0)} \right| &\asymp \tau^{-1/2-h}, & \left| u_r^{(0)} \right| &= O(\tau^{-1/2}), \end{aligned}$$

and the similarity variables are defined by

$$A = \frac{1}{2} + h, \quad D = \frac{1}{2} - h, \quad \tau = q(1 - \eta^2), \quad z = q^D \eta, \quad X = \frac{r^2}{2q}, \quad q > 0, \quad -1 < \eta < 1. \quad (197)$$

Note  $A + D = 1$ ; the source uses this identity explicitly.

**(ii) The leading field, verbatim from (4.3).**

$$u_\theta^{(0)} = q^{-A}E, \quad u_z^{(0)} = q^{-A}U, \quad r u_r^{(0)} = V_0, \quad p^{(0)} = q^{-2A}\Pi, \quad E = C^{-1}\sqrt{2X}\phi, \quad (198)$$

with  $E, U, V_0, \Pi$  functions of  $(X, \eta)$  on the closed profile rectangle  $[0, X_c] \times [-1, 1]$ , and smoothness at the axis imposed on  $\phi$  (equivalently on  $F = \phi/C$ , with  $V_0 = Xv_0$ ).

**(iii) The governing profile equations are stated explicitly.** This is the question Deliverable 0 was posed to settle, and the answer is the favourable one. Incompressibility and the leading radial balance give (4.7),

$$V_0 = \frac{X}{L}(2\eta U - 2D\eta \mathcal{A}_X(U) - d \partial_\eta \mathcal{A}_X(U)), \quad \Pi_X = \frac{E^2}{2X}, \quad (199)$$

with  $d = 1 - \eta^2$ ,  $L = 1 - 2h\eta^2$  and  $\mathcal{A}_X$  the radial average  $\mathcal{A}_X(f) = X^{-1} \int_0^X f(x, \eta) dx$ ; and the vanishing of the leading tangential residual stress is, by the source's Proposition 4.2, the pair (4.13)

$$-2L \frac{X\phi_{XX} + 2\phi_X}{\phi} = S_q, \quad -2L(XU_{XX} + U_X) = S_n, \quad (200)$$

where  $S_q, S_n$  are the explicit first-order sources of (4.9). So Deliverable 2 is a test of an *explicit* system, not a reconstruction from asymptotics.

**(iv) The one structural fact that governs everything below**, stated by the source in its own dimensional analysis (§2.1) and again in its notation guide: *radial viscosity is retained in the leading balance and axial viscosity is not*. The source computes

$$\frac{|u_r|}{\ell_r} = O(\tau^{-1}), \quad \frac{|u_z|}{\ell_z} \asymp \tau^{-1}, \quad \frac{\nu}{\ell_r^2} \asymp \tau^{-1}, \quad \frac{\nu/\ell_z^2}{\nu/\ell_r^2} = \frac{\ell_r^2}{\ell_z^2} \asymp \tau^{2h} \longrightarrow 0,$$

and says of the radial Reynolds number  $\text{Re}_r = |u_r| \ell_r / \nu = O(1)$  that “viscosity continues to compete with radial inflow”. Correspondingly (4.12) defines the leading tangential residual by *adding back*  $\partial_z^2 u_j^{(0)}$ , and the notation guide describes the resulting stress as “retaining radial viscosity and omitting axial viscosity”. *[Fetched and read this session; every displayed formula checked against the source.]*

**Remark 40.3** (What the screenshot transcription got right, and what it omitted). The transcription circulating in this programme before this session —  $u^{(0)} = V_0/r e_r + q^{-A}E e_\theta + q^{-A}U e_z$ ,  $p^{(0)} = q^{-2A}\Pi$ , with (197) — is **correct in every symbol**, and is confirmed by (198). Nothing built on it needs revision. What it omitted is Fact 40.2(iii) and (iv): that the profile equations are explicit, and that the leading balance is *viscous in the radial variable*. Item (iv) is what makes the rest of this section come out the way it does, and it was not on record here.

## 40.2 Deliverable 1 — the anisotropic rescaling, derived by direct substitution

Lemma 39.10 has two parameters, one length and one amplitude. The profile of Fact 40.2 has two lengths with different exponents and two amplitudes with different exponents, so that lemma cannot be applied to it, and its conclusion must not be assumed to transfer. The four-parameter generalisation is derived here from the cylindrical equations directly.

**Lemma 40.4** (The four-parameter anisotropic rescaling: *two effective viscosities*). *Let  $(u, p)$  solve the axisymmetric Navier–Stokes equations ( $\partial_\theta \equiv 0$ ) in cylindrical coordinates. For positive constants  $A_r, A_\theta, A_z, L_r, L_z, T_s$  set*

$$r = L_r \rho, \quad z = L_z \zeta, \quad t = T + T_s s, \quad v_r = A_r u_r, \quad v_\theta = A_\theta u_\theta, \quad v_z = A_z u_z, \quad \tilde{q} = A_p p.$$

*Then continuity becomes*

$$\frac{1}{\rho} \partial_\rho(\rho v_r) + \frac{A_r L_r}{A_z L_z} \partial_\zeta v_z = 0,$$

*so the limit is incompressible iff*

$$\boxed{A_r L_r = A_z L_z =: \Lambda} \quad (201)$$

*Imposing (201) and normalising the advection by  $T_s = \Lambda$ , and writing  $\Delta_\rho^{(m)} := \partial_\rho^2 + \rho^{-1} \partial_\rho - m \rho^{-2}$ , the three momentum equations become*

$$\partial_s v_\theta + v_r \partial_\rho v_\theta + v_z \partial_\zeta v_\theta + \frac{v_r v_\theta}{\rho} = \nu_r \Delta_\rho^{(1)} v_\theta + \nu_z \partial_\zeta^2 v_\theta, \quad (202)$$

$$\partial_s v_z + v_r \partial_\rho v_z + v_z \partial_\zeta v_z + \frac{A_z^2}{A_p} \partial_\zeta \tilde{q} = \nu_r \Delta_\rho^{(0)} v_z + \nu_z \partial_\zeta^2 v_z, \quad (203)$$

$$\partial_s v_r + v_r \partial_\rho v_r + v_z \partial_\zeta v_r + \frac{A_r^2}{A_p} \partial_\rho \tilde{q} - \frac{A_r^2}{A_\theta^2} \frac{v_\theta^2}{\rho} = \nu_r \Delta_\rho^{(1)} v_r + \nu_z \partial_\zeta^2 v_r, \quad (204)$$

*with **two** effective viscosities*

$$\boxed{\nu_r = \nu \frac{A_r}{L_r}, \quad \nu_z = \nu \frac{A_z}{L_z}} \quad (205)$$

[Proved here, by direct substitution into the cylindrical equations; independently re-checked by computer algebra.]

*Proof.*  $\partial_t = T_s^{-1} \partial_s$ ,  $\partial_r = L_r^{-1} \partial_\rho$ ,  $\partial_z = L_z^{-1} \partial_\zeta$ . Substituting  $u_r = v_r/A_r$ ,  $u_\theta = v_\theta/A_\theta$ ,  $u_z = v_z/A_z$ ,  $p = \tilde{q}/A_p$  into  $\frac{1}{r} \partial_r(r u_r) + \partial_z u_z = 0$  and multiplying by  $A_r L_r$  gives the displayed continuity equation. For the azimuthal equation, multiply

$$\partial_t u_\theta + u_r \partial_r u_\theta + u_z \partial_z u_\theta + \frac{u_r u_\theta}{r} = \nu \left( \partial_r^2 u_\theta + \frac{\partial_r u_\theta}{r} - \frac{u_\theta}{r^2} + \partial_z^2 u_\theta \right)$$

by  $A_\theta T_s$ . The time term becomes  $\partial_s v_\theta$ ; the two advection terms and the curvature term acquire  $T_s/(A_r L_r)$  and  $T_s/(A_z L_z)$ , both equal to 1 by (201) and  $T_s = \Lambda$ ; the radial viscous group acquires  $\nu T_s/L_r^2 = \nu \Lambda/L_r^2 = \nu A_r/L_r$  and the axial one  $\nu T_s/L_z^2 = \nu A_z L_z/L_z^2 = \nu A_z/L_z$ . This is (202) with (205). The axial equation is identical after multiplication by  $A_z T_s$ , the pressure term acquiring  $A_z T_s/(A_p L_z) = A_z^2/A_p$ . The radial equation, multiplied by  $A_r T_s$ , acquires  $A_r T_s/(A_p L_r) = A_r^2/A_p$  on the pressure gradient and  $A_r T_s/(A_\theta^2 L_r) = A_r^2/A_\theta^2$  on the centrifugal term.  $\square$

**Remark 40.5** (Consistency with Lemma 39.10). Setting  $A_r = A_\theta = A_z =: A$  and  $L_r = L_z =: L$  makes (201) vacuous, gives  $T_s = \Lambda = AL$  — which is exactly the normalisation  $\tau = AL$  of Lemma 39.10 — and collapses (205) to the single  $\nu_{\text{eff}} = \nu A/L$ . With  $A_p = A^2$  the pressure and centrifugal coefficients in (204) are both 1, matching  $\tilde{p} = (A\tau/L)p = A^2 p$  there. Lemma 39.10 was independently re-derived this session and is correct; Lemma 40.4 contains it.



Now specialise to the profile. This is the computation Deliverable 1 asked for, and the point is that it does *not* reproduce Corollary 39.11(iii).

**Proposition 40.6** (The profile-adapted limit exists, is nontrivial, and is *not* Euler). *Take the exponents of Fact 40.2 and index the family by  $\tau \downarrow 0$ :*

$$L_r = \tau^{1/2}, \quad L_z = \tau^D, \quad A_r = \tau^{1/2}, \quad A_\theta = A_z = \tau^A, \quad A_p = \tau^{2A}.$$

Then:

(i) **Incompressibility is automatic, and it is the source's own identity.**  $A_r L_r = \tau$  and  $A_z L_z = \tau^{A+D} = \tau$ , so (201) holds exactly — because  $A + D = 1$ . The profile's amplitude anisotropy is precisely conjugate to its length anisotropy; this is not a coincidence but incompressibility, and the source uses  $A + D = 1$  to derive the first identity of (199). Hence  $T_s = \Lambda = \tau$ : the natural time unit is the remaining time itself.

(ii) **Radial viscosity is preserved exactly; axial viscosity is lost.** By (205),

$$\nu_r = \nu \frac{\tau^{1/2}}{\tau^{1/2}} = \nu \quad (\text{independent of } \tau), \quad \nu_z = \nu \frac{\tau^A}{\tau^D} = \nu \tau^{2h} \longrightarrow 0.$$

This reproduces, from the rescaling alone, the source's own dimensional statement quoted in Fact 40.2(iv).

(iii) **The radial momentum equation degenerates to a constraint.** The coefficients in (204) are  $A_r^2/A_p = A_\theta^2/A_\theta^2 = \tau^{1-2A} = \tau^{-2h} \rightarrow \infty$ , while  $A_z^2/A_p = 1$  in (203). Multiplying (204) by  $\tau^{2h}$  and letting  $\tau \downarrow 0$  kills every inertial and viscous term and leaves the cyclostrophic balance

$$\partial_\rho \tilde{q} = \frac{v_\theta^2}{\rho}. \quad (206)$$

(iv) **The limiting system is therefore** (206) **together with**

$$\begin{aligned} \partial_s v_\theta + v_r \partial_\rho v_\theta + v_z \partial_\zeta v_\theta + \frac{v_r v_\theta}{\rho} &= \nu \left( \partial_\rho^2 + \frac{\partial_\rho}{\rho} - \frac{1}{\rho^2} \right) v_\theta, \\ \partial_s v_z + v_r \partial_\rho v_z + v_z \partial_\zeta v_z + \partial_\zeta \tilde{q} &= \nu \left( \partial_\rho^2 + \frac{\partial_\rho}{\rho} \right) v_z, \quad \frac{1}{\rho} \partial_\rho (\rho v_r) + \partial_\zeta v_z = 0. \end{aligned}$$

It is closed:  $v_r$  is recovered from continuity,  $\tilde{q}$  from (206) up to a free function  $\tilde{q}_0(\zeta, s)$ , and  $v_\theta, v_z$  evolve. It is nontrivial: the rescaled leading field converges to the profile itself, since  $A_\theta q^{-A} \rightarrow |s|^{-A}$ ,  $X \rightarrow \rho^2/(2|s|)$  and  $\eta \rightarrow \zeta |s|^{-D}$ , giving the explicit self-similar ancient solution  $v_\theta = |s|^{-A} E$ ,  $v_z = |s|^{-A} U$ .

(v) **Consequently the limit is not a solution of the Euler equations**, and Corollary 39.11(iii) does not transfer: it is derived for one length and one amplitude, and the ratio it drives to zero,  $\nu_{\text{eff}} = \nu A/L$ , splits here into two ratios of which only one degenerates.

[Proved here. Items (i)–(iii) are exact; (iv)'s convergence statement is the formal limit of the leading field and is asserted at that level, not as a compactness theorem.]

**Remark 40.7** (What the limiting system is, and why “well-posed” is not claimed for it). The system of Proposition 40.6(iv) has the exact structural signature of a **boundary-layer (Prandtl-type) reduction**: the momentum equation in the thin direction has degenerated into an algebraic constraint fixing the pressure up to a free outer function; the velocity component in that direction is recovered from continuity rather than from its own evolution equation; diffusion survives *only* in the thin direction; and the surviving equations are advected in a direction in which they have no diffusion at all. Here the thin direction is the radial one ( $\ell_r/\ell_z \asymp \tau^h \rightarrow 0$ ), and the free outer function is visible in the source itself as the axis pressure datum  $\Pi(0, \eta)$  of its (4.31), which enters  $\Pi = \Pi(0, \eta) + C_p$  exactly as an outer pressure gradient enters Prandtl’s system.

Two cautions, both material.

- (a) **No source read here writes this system.** The steady, inviscid “quasi-cylindrical” approximation for slender vortices is classical but is an *Euler* reduction and is not this. The closest verified statement is that the inner viscous core of a slender vortex is treated “by making scaling assumptions of the usual boundary-layer type” (Hall 1961; Stewartson–Hall 1963), as reported in [58]. The identification above is therefore *structural*, made from the derived equations, and is offered as such.
- (b) **Deliverable 1 asked whether the limit is well-posed, and the honest answer is that it is well-defined but not known to be well-posed — with specific reason for doubt.** Prandtl-type systems are linearly ill-posed in Sobolev spaces: Gérard-Varet–Dormy [57] prove exactly this for the Prandtl equation, by constructing unstable quasimodes at high tangential frequency around shear flows with a non-degenerate critical point, and remark that “the strong instability is due to viscosity”. Their result is not transported to the present system here, and no claim is made that it applies; it is recorded because the structural features their proof exploits — viscosity in the thin variable, none in the advected variable — are precisely the features Proposition 40.6 produces.

**Corollary 40.8** (If one insists on an Euler zoom, anisotropy forces a choice, and both branches fail). *The Liouville theorems of [46, 47, 48, 49] apply to ancient solutions of Euler, hence require the isotropic zoom of Corollary 39.11(iii), with a single length  $L$  and single amplitude  $A$ . Since  $\sup |u| \asymp \tau^{-A}$  with  $A = \frac{1}{2} + h \in (\frac{1}{2}, 1)$ , that corollary does formally apply and gives  $A = \tau^A$ ,  $\nu_{\text{eff}} = \nu A/L \rightarrow 0$ . But one length cannot match two, and the two available choices fail in opposite ways:*

- (i)  $L \asymp \ell_r = \tau^{1/2}$ : then  $\nu_{\text{eff}} = \nu \tau^h \rightarrow 0$ , and the core has rescaled radial extent  $O(1)$  but rescaled axial extent  $\ell_z/L = \tau^{-h} \rightarrow \infty$ , so the limit is  $\zeta$ -independent. Moreover  $A|u_r| \asymp \tau^A \tau^{-1/2} = \tau^h \rightarrow 0$ : the radial component is annihilated. Finally  $T_s = AL = \tau^{1+h}$ , so  $T_s/\tau = \tau^h \rightarrow 0$  and the limit is steady. The limit object is thus a steady,  $\zeta$ -independent columnar flow  $v = v_\theta(\rho)e_\theta + v_z(\rho)e_z$  — which solves Euler exactly, for any profiles, with pressure given by (206). It is nontrivial, but it is  $\zeta$ -independent on all of  $\mathbb{R}$  and hence of infinite energy, and it lies in no weighted class in which any of the four cited Liouville theorems operates.
- (ii)  $L \asymp \ell_z = \tau^D$ : then the rescaled radial extent is  $\ell_r/L = \tau^h \rightarrow 0$ , the core collapses onto the axis, and the limit is trivial. Nontriviality — the hypothesis every Liouville argument

contradicts — fails outright.

So the Euler-zoom route to excluding this profile is unavailable in both branches, and for reasons that are consequences of the anisotropy alone. [Proved here, at the level of scaling; (i) and (ii) are exponent computations, not compactness theorems.]

### 40.3 Deliverable 2 — the profile *is* excluded, and the Liouville machinery is not what excludes it

Deliverable 2 was posed as: test the extracted ancient solution against the hypotheses of the four Seregin papers. Proposition 40.6 and Corollary 40.8 say that test cannot be run — the profile-adapted limit is not an Euler solution, and the Euler-adapted limit is either trivial or of infinite energy. The correct answer to Deliverable 2 was nevertheless available the whole time, from a lemma this document already had, and it does not need an ancient solution at all.

**Proposition 40.9** (The profile has unbounded swirl, and is therefore impossible for unforced axisymmetric Navier–Stokes). *Let  $\Gamma = ru_\theta$  be the swirl of Lemma 38.3. For the leading profile (198),*

$$\Gamma^{(0)} = r u_\theta^{(0)} = \sqrt{2qX} \cdot q^{-A} \cdot C^{-1} \sqrt{2X} \phi = \frac{2X\phi}{C} q^{\frac{1}{2}-A} = q^{-h} H(X, \eta), \quad (207)$$

where  $H := \sqrt{2X} E = \frac{2X\phi}{C},$

*which is the source’s own identity (“the angular equation is simplest in angular momentum  $ru_\theta^{(0)} = q^{-h}H$ ”, in the proof of its Proposition 4.2). Since  $q \asymp \tau$  on the core ( $|\eta| \leq \eta_0 < 1$ ) and  $H \asymp 1$  at the core edge  $X \asymp X_b,$*

$$\sup_x \left| \Gamma^{(0)}(x, t) \right| \gtrsim \tau^{-h} \longrightarrow \infty \quad (\tau \downarrow 0),$$

*and the same rate follows independently from the source’s exterior swirl  $K = r^{-1-2h} H_{\text{ext}}(\tau/r^2),$  for which*

$$rK = r^{-2h} H_{\text{ext}}(\tau/r^2) \asymp \tau^{-h} \quad \text{at } r \asymp \tau^{1/2}.$$

*Consequently: no smooth axisymmetric solution of the unforced Navier–Stokes equations with  $\|\Gamma(\cdot, 0)\|_{L^\infty} < \infty$  can have this profile as its blowup asymptotics, because Lemma 38.3 gives  $\|\Gamma(\cdot, t)\|_{L^\infty} \leq \|\Gamma(\cdot, 0)\|_{L^\infty}$  throughout the smooth interval, and  $\tau^{-h} \rightarrow \infty$  for every  $h > 0$ . (Schwartz data, as in [3], have  $\|ru_\theta(\cdot, 0)\|_{L^\infty} \leq \| |x| u_0 \|_{L^\infty} < \infty$ .) [Proved here. (207) is re-derived from (198) and agrees with the source’s stated identity.]*

**Corollary 40.10** (The obstruction in geometric form: a Type-I core radius caps the swirl at Type-I). *For unforced axisymmetric flow the maximum principle gives the pointwise bound  $|u_\theta| \leq \Gamma_0/r$  with  $\Gamma_0 = \|\Gamma(\cdot, 0)\|_{L^\infty}$  — i.e. the swirl component automatically satisfies the spatial bound Definition 39.1(S). Hence on a core of radius  $\ell_r,$*

$$|u_\theta| \lesssim \frac{\Gamma_0}{\ell_r}.$$

*A core with the Type-I radial geometry  $\ell_r \asymp \tau^{1/2}$  therefore has  $|u_\theta| \lesssim \tau^{-1/2}$ : **Type-I swirl**. Type-II swirl  $|u_\theta| \asymp \tau^{-1/2-h}$  in the unforced axisymmetric class requires the strictly thinner*

core  $\ell_r \lesssim \tau^{1/2+h}$ . The OpenAI profile has  $\ell_r \asymp \tau^{1/2}$  together with  $|u_\theta| \asymp \tau^{-1/2-h}$ , and so violates the cap by exactly the factor  $\tau^{-h}$  that makes it *Type-II*. [Proved here.]

**Remark 40.11** (Exactly what is and is not excluded). Proposition 40.9 is outcome (i) of the Deliverable 2 trichotomy — an exclusion by a named theorem, with its hypotheses checked — but its reach is small and must be stated at its true size.

- (a) **It is confined to the axisymmetric class**, because Lemma 38.3 is. It says nothing about a non-axisymmetric flow with the same radial and axial scales.
- (b) **It is confined to the unforced equations**. That is the point of the exercise, but it also means it is *not* a statement about [55]: their flow is both forced *and* non-axisymmetric (the oscillatory pulses carry nonzero azimuthal harmonics), so Lemma 38.3 does not apply to it, and no contradiction with their Theorem 1.1 is asserted or implied. Wall 9 of §44 governs, as before.
- (c) **It excludes one profile family, not Type-II**. Any axisymmetric core with  $\ell_r \lesssim \tau^{1/2+h}$  escapes Corollary 40.10 entirely. Region (R2) of Remark 39.4 is untouched, Wall 1 is untouched, and Wall 3 keeps its classification.
- (d) **It is elementary**. The swirl maximum principle is classical and already in this document; the content here is the *check*, and the check is one line once (207) is in hand. No credit is claimed for machinery.

**Remark 40.12** (Two corrections and one clarification to §39). All of S53’s displayed derivations were re-checked from scratch this session;  $\nu_{\text{eff}} = \nu A/L$ , the four-way trichotomy including the  $\beta \geq 1$  scope limit, and the transcribed profile data are all **correct**. Three refinements are nevertheless needed.

- (1) **Corollary 39.13(i) has a hypothesis that this profile violates, and the profile shows the hypothesis is not decorative**. That corollary concludes “the Euler-zoom limit of an axisymmetric Type-II blowup is swirl-free” from  $\Gamma_{\text{new}} = (\nu_{\text{eff}}/\nu)\Gamma$  together with  $\|\Gamma(t)\|_\infty \leq \|\Gamma(0)\|_\infty$ . The second ingredient is Lemma 38.3, which requires the *unforced* equations. For the profile of Fact 40.2,  $\Gamma \asymp \tau^{-h} \rightarrow \infty$  by (207), and  $\Gamma_{\text{new}} = \tau^h \cdot \tau^{-h} = O(1) \neq 0$ : the Euler-zoom limit of *this* object retains its swirl. The corollary is correct as written; but the phrase “any zoom with  $\nu_{\text{eff}} \rightarrow 0$  has  $\Gamma \rightarrow 0$ ” must always be read with its unforced hypothesis attached, and this profile is a concrete witness that it cannot be dropped.
- (2) **Corollary 39.11(iii) does not apply to anisotropic profiles**. It is derived for a single length and a single amplitude. Proposition 40.6 shows that for two of each the conclusion changes qualitatively:  $\nu_r$  survives undiminished while  $\nu_z \rightarrow 0$ , and the limit is not Euler. Nothing in §39 claimed otherwise, but the trichotomy is stated without an anisotropy caveat and could be read as more general than it is.
- (3) **Clarification: the anisotropy is the obstruction to *rescaling*, not to *exclusion***. Fact 39.8 identifies  $\ell_r/\ell_z \rightarrow 0$  as “the sharper point” and it is right about that — Corollary 40.8 is exactly the quantitative form of it. But the anisotropy is not what kills the profile. Corollary 40.10 is a single-scale statement: a profile with  $\ell_r = \ell_z = \tau^{1/2}$  and  $|u_\theta| \asymp \tau^{-1/2-h}$  would be excluded by the same argument. The two length scales matter for Deliverable 1 and are irrelevant to Deliverable 2.

## 40.4 Verdict

**Remark 40.13** (WP-T2 verdict, in the narrow terms the question deserves). Five statements, in the order the deliverables were posed.

- (1) **Deliverable 0: the source exists, was read, and states explicit profile equations.** They are the pair (200), closed by the identities (199) and by the sources of the source's own (4.9). The screenshot transcription this programme had been carrying is correct in every symbol. The one thing it omitted — that the leading balance retains radial viscosity — is what determines Deliverable 1.
- (2) **Deliverable 1: the anisotropic rescaling produces a well-defined, nontrivial, ancient limit, and the limit is not Euler.** Lemma 40.4 replaces the single  $\nu_{\text{eff}}$  by the pair  $(\nu A_r/L_r, \nu A_z/L_z)$ , subject to the incompressibility constraint  $A_r L_r = A_z L_z$  — which the profile satisfies exactly, because  $A + D = 1$ . For the profile's exponents  $\nu_r = \nu$  identically and  $\nu_z = \nu \tau^{2h} \rightarrow 0$ , and the radial momentum equation degenerates to cyclostrophic balance. The limit is a Prandtl-type anisotropic system, not known to be well-posed, and with Gérard-Varet–Dormy giving specific reason to doubt that it is.
- (3) **Deliverable 2: the profile is excluded — outcome (i) — but not by anything in the Liouville literature.** The ancient-Euler route is unavailable in both of its branches (Corollary 40.8). The exclusion is Proposition 40.9: the profile carries  $\Gamma = ru_\theta \asymp \tau^{-h} \rightarrow \infty$ , which the swirl maximum principle forbids for unforced axisymmetric flow. Equivalently (Corollary 40.10): a core of Type-I radius cannot carry Type-II swirl.
- (4) **What this is worth.** One profile family, in the axisymmetric class, for the unforced equations. It is **not** progress toward (A) or (B); it is **not** a statement about [55], whose flow is forced and non-axisymmetric; it is **not** a constraint on Type-II blowup in general, since Corollary 40.10 is escaped by any core with  $\ell_r \lesssim \tau^{1/2+h}$ . Wall 1, Wall 3 and region (R2) are exactly where §39 left them.
- (5) **The transferable lesson, and the only thing here a future session should reuse.** The Liouville programme was the wrong tool for this question and two independent computations say so: the correct rescaling does not produce an Euler solution, and the wrong one produces either nothing or an infinite-energy shear flow. What *did* settle it was a conservation law the axisymmetric class supplies for free. Before importing profile-extraction machinery against a candidate profile, check the candidate against Corollary 40.10 first: it is one line, and for this candidate it was decisive.

## 41 WP-T3: A Quantitative Floor on the Forcing in Wall 9 (Session S55)

Wall 9 (§44.10) records, as a qualitative finding, that the forcing in [55] is *load-bearing and not decorative*. This section turns that into a number. The whole of it is **documentation-sharpening**: it makes Wall 9 more precise and more citable, and it does nothing else. It is *not* a step toward Fefferman's alternatives (A) or (B), *not* a challenge to (C) or (D), and *not* a statement about whether Theorem 1.1 of [55] is correct. Wall 9's classification — PROVED

IMPOSSIBLE as an import route — is unchanged, and Wall 1, Wall 3 and the S54 exclusion (Proposition 40.9) are untouched. The scope is restated at its true size in Remark 41.11, which should be read with any quotation of this section.

The lever that makes the sharpening tractable is structural and is proved first: in the axisymmetric class the swirl equation carries no pressure, and a forcing term enters it *only* through its azimuthal component.

#### 41.1 Deliverable 1 — the forced maximum principle, as a standalone lemma

**Lemma 41.1** (The forced swirl equation: only  $f_\theta$  enters, and it enters as  $rf_\theta$ ). *Let  $(u, p)$  be axisymmetric ( $\partial_\theta \equiv 0$ ) and smooth on an open set of  $\mathbb{R}^3 \times I$ , with  $\operatorname{div} u = 0$ , and define the momentum residual*

$$f := \partial_t u + (u \cdot \nabla)u - \nu \Delta u + \nabla p.$$

*Then  $f$  is axisymmetric, and  $\Gamma = ru_\theta$  satisfies*

$$\partial_t \Gamma + u_r \partial_r \Gamma + u_z \partial_z \Gamma = \nu \left( \Delta - \frac{2}{r} \partial_r \right) \Gamma + r f_\theta, \quad f_\theta := f \cdot e_\theta. \quad (208)$$

*Neither  $f_r$  nor  $f_z$  appears, and no pressure term appears.* [Proved here, from the momentum equations, not taken from the WP-T3 handover.]

*Proof.* Axisymmetry of  $f$  is immediate: every operator on the right of its definition preserves  $\partial_\theta \equiv 0$ . For the equation, take the  $e_\theta$  component of the definition of  $f$ . Three ingredients, each of which is where a forced version could go wrong and each of which is checked:

- (a) **Pressure.**  $(\nabla p)_\theta = r^{-1} \partial_\theta p = 0$ , because  $p$  is axisymmetric. This is the one clean lever: the swirl equation is pressure-free *whether or not* the flow is forced.
- (b) **Advection.**  $((u \cdot \nabla)u)_\theta = u_r \partial_r u_\theta + \frac{u_\theta}{r} \partial_\theta u_\theta + u_z \partial_z u_\theta + \frac{u_r u_\theta}{r}$ , and the  $\partial_\theta$  term vanishes.
- (c) **Vector Laplacian.**  $(\Delta u)_\theta = \Delta u_\theta - \frac{u_\theta}{r^2} + \frac{2}{r^2} \partial_\theta u_r = \Delta u_\theta - \frac{u_\theta}{r^2}$ , again by axisymmetry.

Hence

$$\partial_t u_\theta + u_r \partial_r u_\theta + u_z \partial_z u_\theta + \frac{u_r u_\theta}{r} = \nu \left( \Delta u_\theta - \frac{u_\theta}{r^2} \right) + f_\theta,$$

which is the equation in the proof of Lemma 38.3 with  $+f_\theta$  appended. Multiplying by  $r$  and using the two identities established there — that  $r$  times the left side is  $\partial_t \Gamma + u_r \partial_r \Gamma + u_z \partial_z \Gamma$  (the  $r^{-2}$  terms cancelling exactly), and that  $r(\Delta u_\theta - u_\theta/r^2) = (\Delta - \frac{2}{r} \partial_r) \Gamma$  — gives (208). The added term is  $r \cdot f_\theta$  and nothing else.  $\square$

The maximum principle now has to be run at a moving point, and the two places where such an argument goes wrong — whether the maximum is attained and whether  $t \mapsto \max \Gamma$  is differentiable — are handled explicitly rather than assumed away.

**Lemma 41.2** (Forced maximum principle at the running maximiser). *Let  $\nu > 0$  and let  $(u, p)$  be as in Lemma 41.1 on  $\mathbb{R}^3 \times [t_0, t_1]$ , with residual  $f$ . Assume:*

- (H1) (regularity)  $u$  is  $C^1$  in  $t$  and  $C^2$  in  $x$  and  $p$  is  $C^1$  in  $x$  on  $\mathbb{R}^3 \times [t_0, t_1]$ ; consequently  $\Gamma$ ,  $\partial_t \Gamma$ ,  $\nabla \Gamma$ ,  $D^2 \Gamma$  and  $f_\theta$  are continuous there.

(H2) (attainment, with uniform compactness) For each compact  $J \subset [t_0, t_1]$  there is a compact  $K_J \subset \mathbb{R}^3$  with  $\sup_{x \notin K_J} \Gamma(x, t) < \sup_x \Gamma(x, t)$  for all  $t \in J$ , and  $\sup_x \Gamma(x, t) < \infty$ . (This holds in particular if  $\Gamma(\cdot, t)$  is compactly supported in a fixed compact set, or if  $\Gamma(x, t) \rightarrow 0$  as  $|x| \rightarrow \infty$  uniformly for  $t \in J$ , together with  $\sup_x \Gamma > 0$ .)

(H3) (nondegeneracy)  $M(t) := \max_x \Gamma(x, t) > 0$  for  $t$  in the interval considered.

Write  $\Sigma(t) := \{x : \Gamma(x, t) = M(t)\}$ , which by (H1)–(H2) is nonempty and compact. Then:

(i) **Every maximiser is off the axis.**  $\Gamma$  vanishes on  $\{r = 0\}$  (because  $u_\theta$  is bounded there), so by (H3) every  $x_m \in \Sigma(t)$  has  $r_m > 0$ . In particular the singular coefficient  $-2\nu/r$  of (208) is never evaluated at  $r = 0$ , and no exhaustion  $\{r \geq \varepsilon\}$  of the kind used in Lemma 38.3 is needed.

(ii) **Pointwise differential inequality.** For every  $t$  and every  $x_m \in \Sigma(t)$ ,

$$\partial_t \Gamma(x_m, t) \leq (rf_\theta)(x_m, t). \quad (209)$$

(iii)  **$M$  is locally Lipschitz, with one-sided derivatives everywhere.** On each compact  $J \subset [t_0, t_1]$ ,  $M$  is Lipschitz, and for every  $t$  the one-sided derivatives exist and are given by

$$\frac{d^+ M}{dt}(t) = \max_{x \in \Sigma(t)} \partial_t \Gamma(x, t), \quad \frac{d^- M}{dt}(t) = \min_{x \in \Sigma(t)} \partial_t \Gamma(x, t). \quad (210)$$

Hence  $M$  is differentiable at  $t$  precisely when these agree, and (by Rademacher) at almost every  $t$ .

(iv) **The forced maximum principle.** Combining (ii) and (iii),

$$\frac{d^+ M}{dt}(t) \leq \max_{x \in \Sigma(t)} (rf_\theta)(x, t) \quad \text{for every } t, \quad (211)$$

and, with no differentiability needed anywhere, the integrated form

$$M(t) - M(s) \leq \int_s^t \max_{x \in \Sigma(\sigma)} (rf_\theta)(x, \sigma) d\sigma, \quad t_0 \leq s \leq t < t_1. \quad (212)$$

When  $f \equiv 0$  this returns  $M(t) \leq M(s)$ , i.e. the maximum principle of Lemma 38.3: (211) is its exact forced generalisation.

(v) **The form for  $\|\Gamma\|_{L^\infty}$ , with its sign factor.** Let  $M_\Gamma(t) := \|\Gamma(\cdot, t)\|_{L^\infty} > 0$  and let  $\Sigma_\pm(t)$  be the sets where  $\Gamma = \pm M_\Gamma(t)$ . Then

$$\frac{d^+ M_\Gamma}{dt}(t) \leq \max_{x \in \Sigma_+(t) \cup \Sigma_-(t)} \operatorname{sgn}(\Gamma(x, t)) (rf_\theta)(x, t) \leq \max_{x \in \Sigma_+(t) \cup \Sigma_-(t)} |rf_\theta|(x, t). \quad (213)$$

The sign factor is not removable: see Remark 41.4.

[Proved here.]

*Proof.* (i) Smoothness of  $u$  across the axis forces  $u_\theta = O(r)$ , so  $\Gamma = ru_\theta = O(r^2)$  and in particular  $\Gamma \equiv 0$  on  $\{r = 0\}$ . By (H3),  $M(t) > 0$ , so no point of  $\Sigma(t)$  lies on the axis. Nonemptiness and compactness of  $\Sigma(t)$  follow from (H1)–(H2).

(ii) Fix  $t$  and  $x_m \in \Sigma(t)$ , with cylindrical coordinates  $(r_m, z_m)$ ,  $r_m > 0$  by (i). Because  $\Gamma(\cdot, t)$  is axisymmetric, it is a function of  $(r, z)$  on the open half-plane  $\{r > 0\}$ , and  $(r_m, z_m)$  is an *interior* global maximum of it there. Therefore

$$\partial_r \Gamma(x_m, t) = \partial_z \Gamma(x_m, t) = 0, \quad \partial_r^2 \Gamma(x_m, t) \leq 0, \quad \partial_z^2 \Gamma(x_m, t) \leq 0,$$

the last two being the diagonal entries of a negative semidefinite Hessian. Two consequences at  $x_m$ :

- the drift term vanishes,  $u_r \partial_r \Gamma + u_z \partial_z \Gamma = 0$ , whatever  $u_r, u_z$  are — so *no* bound on the drift is needed, which is why the lemma requires nothing of  $u_r, u_z$  beyond smoothness;
- the diffusion term is nonpositive,

$$\nu \left( \Delta - \frac{2}{r} \partial_r \right) \Gamma = \nu \left( \partial_r^2 \Gamma - \frac{1}{r_m} \partial_r \Gamma + \partial_z^2 \Gamma \right) = \nu (\partial_r^2 \Gamma + \partial_z^2 \Gamma) \leq 0,$$

the singular first-order coefficient being annihilated by  $\partial_r \Gamma(x_m, t) = 0$ . **This is the entire content of why the unforced maximum principle holds**, and it is unaffected by adding a forcing term.

Evaluating (208) at  $x_m$  gives  $\partial_t \Gamma(x_m, t) = 0 + (\leq 0) + (rf_\theta)(x_m, t)$ , which is (209).

(iii) This is the Danskin/envelope argument, written out because it is the step the statement of the deliverable was posed to make explicit. Fix a compact  $J = [a, b] \subset [t_0, t_1]$  and the compact  $K_J$  of (H2); enlarge  $K_J$  to a compact  $K$  with  $K_J \subset \text{int } K$ . By (H1),  $\Gamma$  and  $\partial_t \Gamma$  are bounded on  $K \times J$ , say  $|\partial_t \Gamma| \leq \Lambda$ . Since for  $t \in J$  the supremum of  $\Gamma(\cdot, t)$  is attained in  $K_J$ , for  $s, t \in J$  we have  $M(t) - M(s) \leq \max_{x \in K_J} (\Gamma(x, t) - \Gamma(x, s)) \leq \Lambda |t - s|$  and symmetrically, so  $M$  is Lipschitz on  $J$  with constant  $\Lambda$ ; in particular  $M$  is continuous and, by Rademacher, differentiable a.e.

For the right derivative, fix  $t \in J$ ,  $t < b$ .

*Lower bound.* For any  $x_m \in \Sigma(t)$  and  $s > 0$  small,  $M(t + s) \geq \Gamma(x_m, t + s) = \Gamma(x_m, t) + s \partial_t \Gamma(x_m, t) + o(s) = M(t) + s \partial_t \Gamma(x_m, t) + o(s)$ . Dividing by  $s$  and letting  $s \downarrow 0$ ,  $\liminf_{s \downarrow 0} \frac{M(t+s) - M(t)}{s} \geq \partial_t \Gamma(x_m, t)$ , and taking the maximum over the compact set  $\Sigma(t)$  gives  $\liminf \geq \max_{\Sigma(t)} \partial_t \Gamma$ .

*Upper bound.* Choose  $s_k \downarrow 0$  realising the lim sup, and  $x_k \in \Sigma(t + s_k)$ . By (H2) all  $x_k$  lie in the compact  $K_J$ , so along a subsequence  $x_k \rightarrow \bar{x} \in K_J$ . Joint continuity of  $\Gamma$  and continuity of  $M$  give  $\Gamma(\bar{x}, t) = \lim \Gamma(x_k, t + s_k) = \lim M(t + s_k) = M(t)$ , so  $\bar{x} \in \Sigma(t)$ . Then

$$M(t + s_k) - M(t) \leq \Gamma(x_k, t + s_k) - \Gamma(x_k, t) = s_k \partial_t \Gamma(x_k, t + \vartheta_k s_k)$$

for some  $\vartheta_k \in (0, 1)$ , by the mean value theorem in  $t$ . Dividing by  $s_k$  and using uniform continuity of  $\partial_t \Gamma$  on  $K \times J$ , the right side tends to  $\partial_t \Gamma(\bar{x}, t)$ . Hence  $\limsup \leq \partial_t \Gamma(\bar{x}, t) \leq \max_{\Sigma(t)} \partial_t \Gamma$ .

The two bounds coincide, so the right derivative exists and equals  $\max_{\Sigma(t)} \partial_t \Gamma$ . The left derivative is identical with the inequalities reversed, yielding the minimum.



(iv) (211) is (iii) combined with (ii) applied at each  $x \in \Sigma(t)$ . For (212):  $M$  is Lipschitz on  $[s, t]$ , hence absolutely continuous, hence  $M(t) - M(s) = \int_s^t M'(\sigma) d\sigma$  over the full-measure set where  $M'$  exists; at each such  $\sigma$ ,  $M'(\sigma) = \frac{d^+ M}{dt}(\sigma)$  is bounded by the integrand of (212). The integrand is measurable, being the maximum of a continuous function over an upper semicontinuous compact-valued multifunction, and is bounded on compact subintervals; the inequality follows. *This form assumes no differentiability of  $M$  at any point*, which is the point of recording it.

(v) Where  $\Gamma(x, t) = +M_\Gamma(t)$ ,  $x$  is a maximiser of  $\Gamma$  and (ii) applies verbatim. Where  $\Gamma(x, t) = -M_\Gamma(t)$ ,  $x$  is a global *minimiser* of  $\Gamma$ ; there  $\nabla\Gamma = 0$  and the Hessian is positive semidefinite, so the diffusion term is  $\geq 0$  and (208) gives  $\partial_t \Gamma(x, t) \geq (rf_\theta)(x, t)$ , i.e.  $\partial_t(-\Gamma)(x, t) \leq -(rf_\theta)(x, t)$ . Applying (iii) to  $M_\Gamma = \max_x \max\{\Gamma(x, \cdot), -\Gamma(x, \cdot)\}$  — a maximum over the disjoint union of two copies of the same compact set, to which the same Danskin argument applies unchanged — yields (213).  $\square$

**Remark 41.3** (What is pointwise and what is uniform — read this before citing the floor). Lemma 41.2 is a statement **at a single moving point**  $x_m(t)$ , a set of measure zero in space-time. Three consequences, two negative and one positive, are worth separating.

- (a) **No positive-measure statement follows.** Nothing here shows that  $|f_\theta|$  is large on *any* set of positive measure. A forcing consisting of a single tall spike, tracking the maximiser and supported on a set of measure  $\varepsilon$ , satisfies the conclusion.
- (b) **No  $L^q$  bound follows for  $q < \infty$ .** The same spike has  $\|f\|_{L^q} = O(\varepsilon^{1/q}) \rightarrow 0$ . Any statement about integrated forcing needs a separate argument.
- (c) **The  $L^\infty$  statement does follow, and is the weaker one.**  $\|rf_\theta\|_{L^\infty} \geq (rf_\theta)(x_m(t), t)$  trivially, so (211) implies a floor on the sup-norm. The direction matters: the pointwise statement is the strong one, and the sup-norm statement is what one gets by throwing information away. The *uniform* statement one might want — “ $|f_\theta|$  is bounded below throughout the core” — is a genuinely different and stronger claim and is **not proved here**.

Everything in §41.2 and §41.3 is therefore stated pointwise at  $x_m(t)$ , and should be quoted that way.

**Remark 41.4** (A correction to the WP-T3 handover’s statement of the target). The handover posed Deliverable 1 as: with  $M_\Gamma(t) := \|\Gamma(\cdot, t)\|_{L^\infty}$ , prove  $\frac{d}{dt}M_\Gamma \leq (rf_\theta)(x_m(t), t)$ . **As written for the absolute-value norm this is false.** If the norm is attained at a point where  $\Gamma < 0$ , that point is a global *minimum* of  $\Gamma$ , the diffusion term there is  $\geq 0$  rather than  $\leq 0$ , and the correct inequality is  $\frac{d}{dt}M_\Gamma \leq -(rf_\theta)(x_m, t)$  — the opposite sign. The statement is correct as posed only for the *signed* maximum  $M(t) = \max_x \Gamma$ , which is Lemma 41.2(ii)/(iv); the norm version needs the sign factor of (213). Nothing downstream changes, because §41.2 works with a signed maximum and reports a floor on  $|rf_\theta|$ , but the lemma had to be stated correctly to be usable, and the erroneous form should not be propagated.

## 41.2 Deliverable 2 — the floor for the OpenAI leading field

**Proposition 41.5** (Explicit power-law floor on the azimuthal residual). *Let  $(u^{(0)}, p^{(0)})$  be the leading axisymmetric field (198) of Fact 40.2, with the similarity variables (197) and  $\nu = 1$*

(the source's own normalisation), and let  $f^{(0)} := \partial_t u^{(0)} + (u^{(0)} \cdot \nabla) u^{(0)} - \Delta u^{(0)} + \nabla p^{(0)}$  be its momentum residual — which is exactly what [55] calls the force, by its (3.1). Then, at the running maximiser  $x_m(t)$  of  $\Gamma^{(0)}$ ,

$$\left| (r f_\theta^{(0)})(x_m(t), t) \right| \geq h c_* \tau^{-(1+h)} = h c_* \tau^{-(A+\frac{1}{2})}, \quad (214)$$

$$\left| f_\theta^{(0)}(x_m(t), t) \right| \geq \frac{h c_*}{c_r} \tau^{-(\frac{3}{2}+h)} = \frac{h c_*}{c_r} \tau^{-(A+1)}, \quad (215)$$

with  $c_*, c_r > 0$  the fixed profile constants of the proof, and  $r(x_m(t)) = c_r \tau^{1/2}$ . [Proved here.]

**Proof. Step 1: the swirl of the leading field, re-derived.** From (198),  $r = \sqrt{2qX}$  and  $E = C^{-1} \sqrt{2X} \phi$ ,

$$\Gamma^{(0)} = r u_\theta^{(0)} = \sqrt{2qX} \cdot q^{-A} \cdot C^{-1} \sqrt{2X} \phi = \frac{2X\phi}{C} q^{\frac{1}{2}-A} = q^{-h} H(X, \eta), \quad H = \frac{2X\phi}{C},$$

using  $\frac{1}{2} - A = -h$ . This is (207), re-derived here independently of S54, and it is also the source's own identity (“the angular equation is simplest in angular momentum  $r u_\theta^{(0)} = q^{-h} H$ ”).

**Step 2: the exact time law of the maximum.** By (197),  $\tau = q(1-\eta^2)$ , so  $q = \tau/(1-\eta^2)$  and

$$\Gamma^{(0)} = \tau^{-h} (1-\eta^2)^h H(X, \eta).$$

At fixed  $t$  the pair  $(X, \eta)$  ranges over the whole profile domain as  $(r, z)$  ranges over the half-plane:  $\eta$  and  $q$  are fixed by  $(z, \tau)$  and then  $X = r^2/(2q)$  sweeps  $(0, \infty)$  with  $r$ . Hence

$$M(t) := \sup_x \Gamma^{(0)}(x, t) = c_* \tau^{-h}, \quad c_* := \sup_{(X, \eta)} (1-\eta^2)^h H(X, \eta), \quad (216)$$

and  $c_*$  is a *constant*, independent of  $t$ : the leading field is exactly self-similar, so the maximum is an exact power, not merely  $\asymp \tau^{-h}$ . This is what makes the floor exact rather than heuristic.

**Step 3: the supremum is attained, off the axis and at finite radius.**  $H(0, \eta) = 0$  since  $H = 2X\phi/C$ ;  $H$  is bounded and  $H \rightarrow 0$  as  $X \rightarrow \infty$ , since the exterior power of the source is  $H_{\text{pow}} = \sqrt{2X} c_\infty X^{-A} = \sqrt{2} c_\infty X^{-h}$ ; and the factor  $(1-\eta^2)^h$  vanishes at  $\eta = \pm 1$  where  $H$  stays bounded (smoothness on the closed rectangle). So  $(1-\eta^2)^h H$  attains its supremum at some  $(X_*, \eta_*)$  with  $0 < X_* < \infty$  and  $|\eta_*| < 1$ , and  $c_* \in (0, \infty)$ . Hypotheses (H1)–(H3) of Lemma 41.2 hold for  $\Gamma^{(0)}$  on any  $[t_0, 1)$ .

**Step 4: differentiate.**  $\tau = 1 - t$ , so  $d\tau/dt = -1$  and, from (216),  $M$  is smooth in  $t$  with

$$\frac{dM}{dt}(t) = c_* (-h) \tau^{-h-1} \cdot (-1) = h c_* \tau^{-(1+h)} > 0. \quad (217)$$

(Here the differentiability subtlety of Lemma 41.2(iii) does not bite —  $M$  happens to be smooth — but the lemma was stated so that it would not need to.)

**Step 5: the radius of the maximiser is an output, not an input.** The maximiser sits at  $(X_*, \eta_*)$ , hence at  $q_* = \tau/(1-\eta_*^2)$  and

$$r(x_m(t)) = \sqrt{2q_* X_*} = \underbrace{\left( \frac{2X_*}{1-\eta_*^2} \right)^{1/2}}_{=: c_r > 0} \tau^{1/2}. \quad (218)$$

So  $\ell_r \asymp \tau^{1/2}$  is not assumed here: the maximiser tracks the core radius as a consequence of Steps 1–3, and  $c_r$  is explicit.

**Step 6: apply the lemma.** By (211) and (217),

$$h c_* \tau^{-(1+h)} = \frac{dM}{dt}(t) \leq \max_{x \in \Sigma(t)} (r f_\theta^{(0)})(x, t),$$

so at some (hence, by (218), at a radius  $c_r \tau^{1/2}$ ) maximiser  $x_m(t)$  we get (214). Dividing by  $r(x_m(t)) = c_r \tau^{1/2}$  gives (215). Finally  $A = \frac{1}{2} + h$  gives  $A + \frac{1}{2} = 1 + h$  and  $A + 1 = \frac{3}{2} + h$ , which is the second form of each display.  $\square$

**Remark 41.6** (Calibration: the floor sits exactly at the size of every other term). The exponent  $\tau^{-(A+1)}$  of (215) is not an arbitrary power. Every individual term of the leading field’s own  $\theta$ -momentum equation has the same size on the core, using  $|u_\theta^{(0)}| \asymp \tau^{-A}$ ,  $|u_r^{(0)}| = O(\tau^{-1/2})$ ,  $\ell_r \asymp \tau^{1/2}$ ,  $\ell_z \asymp \tau^D$  from Fact 40.2:

$$\begin{aligned} \left| \partial_t u_\theta^{(0)} \right| &\asymp \frac{\tau^{-A}}{\tau} = \tau^{-(A+1)}, & \left| u_r^{(0)} \partial_r u_\theta^{(0)} \right| &\asymp \frac{\tau^{-1/2} \tau^{-A}}{\tau^{1/2}} = \tau^{-(A+1)}, \\ \left| \partial_r^2 u_\theta^{(0)} \right| &\asymp \frac{\tau^{-A}}{\tau} = \tau^{-(A+1)}, \end{aligned}$$

whereas the *axial* viscous term is strictly smaller,  $\left| \partial_z^2 u_\theta^{(0)} \right| \asymp \tau^{-A} \tau^{-2D} = \tau^{-(A+1)+2h}$ , by a factor  $\tau^{2h}$ . So the floor says: **the azimuthal residual of the axisymmetric leading field is of full leading order in its own momentum equation.** It is not a perturbation, it is not lower order, and it cannot be made so by any choice of the free profile data — the swirl maximum principle fixes it. That is the quantitative content of Wall 9(b)’s “load-bearing, not decorative”. As a by-product the same computation reproduces, from the floor’s side, the source’s own structural statement (Fact 40.2(iv)) that radial viscosity is retained in the leading balance and axial viscosity is not.

**Corollary 41.7** (The swirl maximum is forced into the active annulus). *Keep the hypotheses of Proposition 41.5, and let  $R_\theta^{(0)} := R(u^{(0)}, p^{(0)}) \cdot e_\theta + \partial_z^2 u_\theta^{(0)}$  be the source’s leading tangential residual (its (4.12)). Then*

$$r R_\theta^{(0)} = \left( \partial_t + u_r^{(0)} \partial_r + u_z^{(0)} \partial_z \right) \Gamma^{(0)} - \left( \partial_r^2 - \frac{1}{r} \partial_r \right) \Gamma^{(0)}, \quad (219)$$

and consequently: **if  $R_\theta^{(0)}$  vanishes on an open set  $\mathcal{U}$ , then  $x_m(t) \notin \mathcal{U}$ .** Since [55] makes  $R_\theta^{(0)}$  minus a cylindrical divergence of the stress  $T = q^{-A-1/2} T_0$  (its Proposition 4.2) and makes  $T_0$  vanish for  $X \leq X_a$  and for  $X \geq X_b$  (its Theorem 4.6(ii)), it follows that

$$X_* \in [X_a, X_b]:$$

the swirl maximum of the leading profile must lie in the closed active annulus where the pulses act. This is a consequence of the swirl maximum principle, not a design choice. [Proved here; the consistency of  $X_* \in [X_a, X_b]$  with the source’s profile is discussed, and left open, in Remark 41.10(d).]

*Proof.* For (219): at  $\nu = 1$ , Lemma 41.1 gives  $D_t \Gamma^{(0)} = (\partial_r^2 - \frac{1}{r} \partial_r + \partial_z^2) \Gamma^{(0)} + r f_\theta^{(0)}$  where  $D_t$  is the material derivative along  $(u_r^{(0)}, u_z^{(0)})$ , while  $r R_\theta^{(0)} = r f_\theta^{(0)} + r \partial_z^2 u_\theta^{(0)} = r f_\theta^{(0)} + \partial_z^2 \Gamma^{(0)}$  (as  $r$  does not depend on  $z$ ). Adding gives (219).

Now suppose  $x_m(t) \in \mathcal{U}$  with  $R_\theta^{(0)} \equiv 0$  on  $\mathcal{U}$ . By (219),  $D_t \Gamma^{(0)} = (\partial_r^2 - \frac{1}{r} \partial_r) \Gamma^{(0)}$  there — *radial* diffusion only. At  $x_m(t)$  the gradient vanishes (Lemma 41.2(ii)), so the drift contributes nothing and the first-order radial term is annihilated, leaving

$$\frac{dM}{dt}(t) = \partial_t \Gamma^{(0)}(x_m, t) = \partial_r^2 \Gamma^{(0)}(x_m, t) \leq 0,$$

which contradicts  $dM/dt = h c_* \tau^{-(1+h)} > 0$  from (217).  $\square$

**Remark 41.8** (An independent check of (208) against the source’s own (4.14)). The forced swirl equation can be tested directly against a formula [55] states for a different reason, and it passes. On  $X \leq X_a$  the profiles satisfy the source’s (4.13), i.e.  $-2L(X\phi_{XX} + 2\phi_X)/\phi = S_q$ , and its (4.14) reads

$$\begin{aligned} (\partial_t + u_r^{(0)} \partial_r + u_z^{(0)} \partial_z)(q^{-h} H) &= -\frac{q^{-h-1}}{L} H S_q \stackrel{(4.13)}{=} 2q^{-h-1} \frac{H}{\phi} (X\phi_{XX} + 2\phi_X) \\ &= \frac{4X}{C} q^{-h-1} (X\phi_{XX} + 2\phi_X), \end{aligned}$$

using  $H = 2X\phi/C$ . Independently, (219) with  $R_\theta^{(0)} = 0$  (which holds there,  $T_0$  being zero for  $X \leq X_a$ ) says the same material derivative equals  $(\partial_r^2 - \frac{1}{r} \partial_r) \Gamma^{(0)}$ . Computing that from  $\Gamma^{(0)} = q^{-h} H(X)$  with  $X = r^2/(2q)$  and  $\partial_r = (r/q) \partial_X$  at fixed  $(z, t)$ :

$$\left(\partial_r^2 - \frac{1}{r} \partial_r\right) \Gamma^{(0)} = q^{-h} \frac{2X}{q} H_{XX} = 2X q^{-h-1} \cdot \frac{2}{C} (2\phi_X + X\phi_{XX}) = \frac{4X}{C} q^{-h-1} (X\phi_{XX} + 2\phi_X).$$

**The two agree identically.** This simultaneously checks the source’s (4.14), this document’s (208), and the identity (219) on which Corollary 41.7 rests.

### 41.3 Deliverable 3 — the floor against what the source states

*Fact 41.9* (What [55] states about the size of its forcing). Re-fetched and read this session from the locator carried by the bibliography entry [55]. Four groups of statements are relevant, and they are quoted rather than paraphrased.

- (i) **The force is defined as the residual.** §2: “For any incompressible flow  $u$  and pressure  $p$ , we can always define the external force  $f$  to be the residual in (1.1).” At  $\nu = 1$  its (3.1) is  $R(u, p) = \partial_t u + (u \cdot \nabla)u - \Delta u + \nabla p$ . This is Wall 9(b)’s point, in the source’s own words.
- (ii) **Qualitatively, the background’s residual diverges.** §2: “For the background alone, the momentum residual becomes unbounded as  $t \uparrow 1$ , so it cannot supply the smooth force required by the theorem.” And §2.2: “... needed to sustain the background becomes unbounded as  $t \uparrow 1$ , so it cannot serve as the smooth external force required by the construction.” *No rate is attached in either sentence* — this is the qualitative statement the present section quantifies.

- (iii) **Quantitatively, the rate is stated — in §4, as a stress exponent.** After its (4.11): “In the notation of Section 3, the physical stress representation to be proved is  $T = q^{-A-1/2}T_0$ .” Its Proposition 4.2: “The residuals  $R_\theta^{(0)}, R_z^{(0)}$  in (4.12) are minus the cylindrical divergences  $\partial_r + 2/r$  and  $\partial_r + 1/r$  of  $q^{-A-1/2}T_0$ .” Its Theorem 4.6(iv) makes  $T_0$  nondegenerate on the annulus, “ $|T_0| \geq c\zeta$ ,  $|\partial^\alpha T_0| \leq C_\alpha \zeta \delta^{-m_\alpha}$ ”, with  $\zeta > 0$  exactly on  $X_a < X < X_b$ . In the proof of its Proposition 4.2 the azimuthal case is written out: “The angular equation is simplest in angular momentum  $ru_\theta^{(0)} = q^{-h}H$ ”, with the material derivative  $-q^{-h-1}HS_q/L$ , and “The axial material derivative plus the pressure gradient is similarly  $-q^{-A-1}S_n/L$ .” Its Proposition 5.5 carries the same exponents to the full axisymmetric background: its (5.41) is  $R(u_B, p_B) = -(\partial_r + 2/r)T_{\text{phys},\theta}e_\theta - (\partial_r + 1/r)T_{\text{phys},z}e_z + E_B$  with  $|\partial^\alpha E_B| \leq C_{\alpha,M}q^M$  for every  $M$ , and its (5.43) fixes  $\hat{T} = q^{A+1/2}T_{\text{phys}}$  with  $|D^I(\hat{T} - T_0)| \leq C_I q^{2h} \zeta \delta^{-N_I}$ .
- (iv) **The final force, by contrast, is flat.** Its Theorem 3.1(iii): “ $|\partial_x^\alpha \partial_t^b R(u, p)| \leq C_{\alpha,b,N,X_1} q^N$ ” for every  $N$  on  $0 \leq X \leq X_1$ , and “For  $X \geq X_{\text{ext}}$  the residual is identically zero.” This is a statement about the *full, non-axisymmetric* field, to which Lemma 41.2 does not apply.

[Primary source fetched and read this session; every quotation above is verbatim from it.]

**Remark 41.10** (The comparison, reported exactly as it comes out). (a) **Match the objects first.** The floor of Proposition 41.5 applies to any axisymmetric divergence-free field whose swirl maximum grows like  $\tau^{-h}$ . Two objects in [55] qualify: the leading field  $u^{(0)}$ , and the corrected axisymmetric background  $u_B$  of its Proposition 5.5 — the latter because its (5.42) gives  $q^A u_{\theta,B} = E_0 + O(q^{2h})$ , so that  $\Gamma_B = q^{-h}H(1 + O(q^{2h}))$  and  $M_B(t) = c_* \tau^{-h}(1 + o(1))$ . Neither of the two is the object of Theorem 1.1, which is non-axisymmetric; see item (e) below.

- (b) **Converting the source’s number into the floor’s variables.** By Fact 41.9(iii),  $f_{\theta,B} = -(\partial_r + 2/r)T_{\text{phys},\theta} + (E_B)_\theta$  with  $T_{\text{phys},\theta} = q^{-A-1/2}(T_{0,\theta} + O(q^{2h}))$  and  $E_B$  flat. Hence  $r f_{\theta,B} = -r^{-1} \partial_r(r^2 T_{\text{phys},\theta}) + \text{flat}$ ; and since  $q$  depends on  $(z, t)$  only,  $\partial_r = (r/q) \partial_X$  at fixed  $(z, t)$  and  $r^2 = 2qX$ , so

$$r f_{\theta,B} = -2 q^{-A-1/2} \partial_X (X T_{0,\theta}) + O(q^{-A-1/2+2h}) + \text{flat} \asymp q^{-A-1/2} = q^{-(1+h)} \asymp \tau^{-(1+h)}$$

on the active annulus, where  $q \asymp \tau$  and  $|T_0| \geq c\zeta > 0$ . Dividing by  $r \asymp \tau^{1/2}$ ,  $|f_{\theta,B}| \asymp \tau^{-(A+1)} = \tau^{-(3/2+h)}$ .

- (c) **Verdict: the source meets the floor, and the match is exponent-exact with no slack.** The floor demanded  $|rf_\theta| \gtrsim \tau^{-(A+1/2)}$  and  $|f_\theta| \gtrsim \tau^{-(A+1)}$  ((214), (215)); the source states  $q^{-A-1/2}$  for the stress and  $q^{-A-1}$  for the axial material derivative, i.e. the *same two exponents*. This is a real consistency check, and it is not circular: the floor comes from the swirl maximum principle and the single exponent  $\Gamma^{(0)} \asymp \tau^{-h}$ , while the source’s exponent comes from radially integrating its residual. Two different routes, one answer. Wall 9’s “load-bearing, not decorative” can now be quoted with a number:  $\tau^{-(A+1)}$  **pointwise at the swirl maximum, i.e. full leading order in the  $\theta$ -momentum equation** (Remark 41.6).

- (d) **One derived requirement the source does not state, recorded as a fact to check and *not* as a discrepancy.** Corollary 41.7 forces  $X_* \in [X_a, X_b]$ . Partial independent evidence from the source, in both directions:  $H(0, \eta) = 0$  and  $H \sim \sqrt{2}c_\infty X^{-h}$  at large  $X$ , so  $X_*$  is interior; but its (4.30) gives  $E = c_{\text{patch}}(1 + \eta^2)^{-1} X^{-1/2-\lambda}$  with  $\lambda > 0$  (its Theorem 4.6) on the reserved intervals  $I_{\text{pos}}, I_{\text{mean}} \subset (X_a, X_v)$ , so  $H = \sqrt{2X}E \propto X^{-\lambda}$  is already *decreasing* there, placing  $X_* \leq \inf I_{\text{pos}}$ . The two together pin  $X_*$  to the narrow window  $[X_a, \inf I_{\text{pos}}]$  near the inner edge of the annulus. **Nothing read this session confirms or contradicts  $X_* \geq X_a$ ;** settling it needs the inner-profile construction of the source’s Appendix B, which was not re-derived here. It is recorded as a requirement the leading profile must satisfy for its own stated swirl growth to be possible — checkable in one line by anyone holding the profile — and explicitly not as a defect: the construction’s whole design puts the momentum transport in that annulus, so the requirement is the expected one.
- (e) **No tension with Theorem 1.1, and none is implied.** The final force of [55] is flat (Fact 41.9(iv)), and the floor says nothing against that, for the reason Wall 9 already records: the full field is *not* axisymmetric — the oscillatory pulses “have zero angular average” as cylindrical velocity components but nonzero momentum fluxes — so Lemma 41.2 does not apply to it at all. What the floor quantifies is the size of the momentum source the *axisymmetric* background needs; the construction supplies it internally, through the pulses’ averaged quadratic products, rather than through  $f$ . Read the other way round, the floor says that supply cannot be small: **any mechanism that carries the axisymmetric background’s swirl from  $O(1)$  to  $\tau^{-h}$  must inject azimuthal momentum at full leading order**, which is precisely why breaking axisymmetry is not an optional convenience of the construction.
- (f) **Constants are not compared.** The floor’s constant is  $hc_*$ , and  $h < 1/100$ , so the floor is weak at the level of constants; the source states no numerical value for  $T_0$  at any particular  $X$ . **The comparison in (c) is exponent-level only**, and should be quoted that way.

#### 41.4 Verdict, at its true size

**Remark 41.11** (Scope — read with any quotation of §41). (1) **This is documentation-sharpening.** It makes Wall 9 more precise and more citable. That is all it is for.

- (2) **It is not a step toward (A) or (B).** Nothing here bears on global regularity for the unforced equations. It proves a lower bound on a forcing, which is the opposite direction of travel.
- (3) **It is not a challenge to (C) or (D).** It asserts no error in [55] and finds none; where a derived requirement is not settled by the text read, it is flagged as a fact to check (Remark 41.10(d)), not as a refutation. That construction is not independently re-verifiable from outside in one session, and this section does not attempt it.
- (4) **Wall 9’s classification is unchanged** (§44.10): PROVED IMPOSSIBLE as an import route. Wall 1, Wall 3, Region (R2) and Proposition 40.9 are untouched.

- (5) **The class is axisymmetric and the statement is pointwise.** Lemma 41.2 is confined to the axisymmetric class, and what it proves holds at one moving point (Remark 41.3). No uniform, integrated or non-axisymmetric version is proved.
- (6) **What is genuinely new here is small and worth naming exactly:** the forced generalisation (211) of Lemma 38.3 (elementary, but it did not exist in this document and the handover's form of it was wrong, Remark 41.4); the exponent (215); and Corollary 41.7, which is the only statement here that says something the source does not already say about itself.

## 42 WP-A: An External Log-Weighted BMO Hypothesis on the Vorticity Direction, Priced Against $\sigma^*$ -Decay (Session S57)

### 42.1 Scope, and what is and is not imported

Two preprints of Z. Grujić [60, 61] propose a conditional route to global regularity that operates on *this document's own direction equation* (142), at *this document's own critical scale*  $r^* = M^{-1/2}$ , and attacks the factor  $\mathcal{L}$  at exactly the point Wall 1(d) (§44.2) declares permitted: the gradient of  $u$  is never recovered by an  $L^\infty$  Calderón–Zygmund bound. The hypothesis carrying the route is that the vorticity direction lies in a logarithmically weighted space of bounded mean oscillation. This section prices that hypothesis against the no-concentration family this document already owns —  $\sigma^*$ -decay (Definition 26.3, Corollary 31.5), **(AX)** (Proposition 38.12) and saturation (§37) — under the S50 standard: *a new target must be proved cheaper, not asserted cheaper*.

**Remark 42.1** (Standing: what is a claim and what is a proof). Both sources are **unrefereed single-author preprints** ([60]: v1 9 Jul 2026, v3 9 Sep 2026; [61]: v1 4 Sep 2026, v2 9 Sep 2026; no journal reference on either listing). **Nothing in this document depends on either of them being correct.** They are quoted as claims in Fact 42.2 and nowhere used as hypotheses. Everything labelled *[Proved]* below — Lemma 42.4, Lemma 42.6, Propositions 42.8, 42.9, 42.10 and Theorem 42.11 — is proved here from Definition 26.3 and Poincaré's inequality, and stands whether or not [60, 61] do. Remark 42.15 records a verification of one imported technique and is the only place where a source proof is audited.

### 42.2 The sources, and what each load-bearing step actually is

*Fact 42.2* (The two preprints, classified step by step). Read in full from the primary arXiv sources at S57. Write  $\xi = \omega/|\omega|$  for the direction (this document's  $\hat{e}$ ),  $\phi(r) = 1/|\log r|$ , and

$$\|f\|_{\text{bmo}_\phi} = \|f\|_{L^\infty} + \sup_{x \in \mathbb{R}^3, 0 < r < 1/2} \frac{1}{\phi(r)} \frac{1}{|B_r(x)|} \int_{B_r(x)} |f - c_{B_r(x)}| dy. \quad (220)$$

- (a) **Assumed** ([60, Def. 2.1], [61, Def. 3.1]): a *critical point singularity* —  $\omega(x, t) = \Phi(x, t)|x|^{-2}$  with  $\Phi$  bounded above (and, in [61], below by  $c_0 > 0$ ),  $|\nabla \Phi| \lesssim |x|^{-1}$ ,  $\Phi$  scale-invariant or log-periodic at the core, with the stated consequence that the super-level set  $A_\lambda(t) = \{\omega > \lambda\}$  is *contained* in a ball  $B_R$ ,  $R \leq C\lambda^{-1/2}$ . [61] adds the critical direction bound  $|\nabla \xi| \leq C_1/\max(|x|, \ell(t))$  and an inner scale  $\ell(t) \rightarrow 0$ .

- (b) **Proved**, and it is Constantin–Fefferman’s cancellation [13] in commutator dress: with  $S = \mathcal{T}(\omega\xi)$  a matrix of Calderón–Zygmund operators, the unidirectional part  $\xi(x) \cdot \mathcal{T}(\omega\xi(x))(x) \xi(x)$  vanishes identically, so  $\alpha = \xi \cdot S\xi = \xi \cdot [\mathcal{T}, \xi](\omega) \xi$ .
- (c) **Proved from (a) and the bmo hypothesis** ([60, Thm. 4.1]): if  $\omega \in L_t^\infty L^{3/2, \infty}$  under (a) and  $\xi \in L_t^\infty \text{bmo}_\phi$ , then  $\|\alpha(\cdot, t)\|_{L^{3/2, \infty}(B_R)} \leq C_0/|\log R|$  for all small  $R$ , uniformly in  $t$ . Verified line by line at S57; see Remark 42.15.
- (d) **Proved from (c)** ([60, §5–§6]): a Lorentz-interpolated truncated-enchrophy argument upgrades the distribution function to  $|\{\omega > \lambda\}| \leq C_\omega \lambda^{-3/2} (\log \lambda)^{-3/2}$  and, by O’Neil’s convolution lemma,  $|\{|u| > \lambda\}| \leq C_u \lambda^{-3} (\log \lambda)^{-3}$ .
- (e) **Proved from (d)** ([60, Thm. 7.4], the headline claim): the improved velocity distribution forces the 1D sparseness scale  $r_s \sim \|u\|_\infty^{-1} (\log \|u\|_\infty)^{-1}$  below the analyticity radius  $\rho_s \sim \nu \|u\|_\infty^{-1}$ ; the harmonic-measure maximum principle then contradicts the escape-time property, so  $T^*$  is not singular.
- (f) **Assumed, four times over** ([61, Thm. 5.7, Cor. 5.8, Def. 7.7, Thm. 7.9]): the companion does *not* propagate the hypothesis. It proves a conditional *transfer* theorem — a logarithmic modulus in time of the direction at the inner scale becomes a logarithmic modulus in space — under four hypotheses, none of which is established for Navier–Stokes solutions: an inflow condition (Hk) that is a smallness condition on the inward radial velocity relative to  $\nu$  ( $\sup_A |x|(u_r)_- + 2\nu C_\Phi \leq (3-k)\nu$ ); a tangential strain three logarithms below critical,  $|P_{\xi^\perp} S\xi| \leq \Lambda |x|^{-2} |\log |x||^{-3}$ ; a cap condition on the boundary data; and a core-history modulus  $\omega_{core}(\sigma; t) \leq \Lambda_0/|\log \sigma|$ .
- (g) **Proved, and negative** ([61, Prop. 7.10, Rem. 7.11, Prop. 7.12, Rem. 7.13]): the companion itself exhibits two natural critical cores on which its own hypotheses fail. For a scale-invariant  $e_*$ -axisymmetric tube core  $\omega = \Phi(\vartheta)|x|^{-2}e_*$  the tangential strain is exactly critical,  $|F_{tan}| = \Phi_0|x|^{-2}|\sin 2\vartheta|$  in the spherical case, so the strain hypothesis of (f) fails at order one and “the direction winds logarithmically as  $\ell(t) \rightarrow 0$ ”. For a near-Beltrami core the twist is critical and  $\|\xi\|_{\text{bmo}_\phi} \gtrsim |\log r|$ , so the hypothesis of (e) fails outright.

**Remark 42.3** (Two defects in the sources found at S57, recorded as found). Neither is used below; both are recorded so that no later session repeats the reading.

- (i) **The two papers do not meet.** [60, Thm. 7.4] and its restatement [61, Thm. 7.1] both require  $\xi \in L_t^\infty \text{bmo}_{1/|\log r|}(\mathbb{R}^3)$  — *globally in space*, the supremum in (220) being over all  $x \in \mathbb{R}^3$ . What [61, Cor. 5.8, Thm. 7.9] delivers is  $\|\xi(\cdot, t)\|_{\text{bmo}_{1/|\log r|}(B_{\rho_1})} \leq C$ , a statement on one small ball. The gap is bridgeable — the proof of [60, Thm. 4.1] uses the hypothesis only on balls contained in  $B_{2R^{1/2}}$  — but the local form of Theorem 7.4 that Theorem 7.9 invokes is stated nowhere.
- (ii) **The global form is close to vacuous.** By Proposition 42.9 below,  $\xi$  fails (220) at every scale near any transversal zero of  $\omega$ . A hypothesis on all of  $\mathbb{R}^3$  therefore also asserts that  $\omega$  has no such zero anywhere, which is a statement about the flow far from the singularity and has nothing to do with the mechanism.



### 42.3 Exact scale identification

**Lemma 42.4** (The source's critical regime is this document's critical scale, and its direction bound is this document's borderline). *Let  $\omega$  satisfy Fact 42.2(a) with inner scale  $\ell(t)$  and  $c_0 \leq \Phi \leq C_0$ , and let  $M(t) = \|\omega(\cdot, t)\|_{L^\infty}$ ,  $r^* = M^{-1/2}$  as in Definition 26.3.*

- (i)  $c_0 \ell^{-2} \leq M \leq C_0 \ell^{-2}$ , hence  $C_0^{-1/2} \ell \leq r^* \leq c_0^{-1/2} \ell$ : the source's inner scale and this document's  $r^*$  agree up to fixed constants, and the criticality  $M \cdot (r^*)^2 = 1$  of Remark 26.7 is the concentration rate  $\omega \sim |x|^{-2}$ .
- (ii) The critical direction bound  $|\nabla \xi| \leq C_1 / \max(|x|, \ell)$  of [61, Def. 3.1(6)] implies  $\sigma^* \leq \frac{4\pi}{3} C_1^2 c_0^{-3/2} C_0$ , and no better: it is the assertion that  $\sigma^*$  is bounded, which is the borderline of Remark 26.7 and the Regime-B stalemate of Proposition 31.4. It carries no logarithm, and the source says so.

[Proved.]

*Proof.* (i) On the annulus  $\ell < |x| < R_0$  one has  $c_0 |x|^{-2} \leq \omega \leq C_0 |x|^{-2}$ , and inside  $|x| \leq \ell$  the source's own bound is  $\omega \leq C_0 \ell^{-2}$ ; the supremum is therefore attained at  $|x| \asymp \ell$  and is comparable to  $\ell^{-2}$ . Take  $-1/2$  powers. (ii) On  $B_{r^*}(x^*)$  we have  $|x| \lesssim r^* \asymp \ell$ , so  $|\nabla \xi| \leq C_1 / \ell \leq C_1 C_0^{1/2} / r^* \cdot (C_0^{1/2} c_0^{-1/2})$ ; hence  $\int_{B_{r^*}} |\nabla \hat{e}|^2 \mathbf{1} \leq C_1^2 c_0^{-1} C_0 (r^*)^{-2} \cdot \frac{4\pi}{3} (r^*)^3$  and  $\sigma^* = (r^*)^{-1} \int \leq \frac{4\pi}{3} C_1^2 c_0^{-1} C_0$ , absorbing the comparability constant. The calibration of Remark 26.7 ( $|\nabla \hat{e}| \sim 1/r^*$  on  $B_{r^*}$  gives  $\sigma^* \sim 1$ ) shows the bound is attained, so nothing sharper follows.  $\square$

### 42.4 The Poincaré translation: what is true

Definition 26.3 measures  $|\nabla \hat{e}|^2$  in  $L^2$  on one ball, at one scale, *restricted to*  $\{|\omega| \geq M/4\}$ ; (220) measures  $|\hat{e} - \hat{e}_B|$  in  $L^1$  on every ball, at every scale, with no restriction. The following definition isolates the object that a Poincaré inequality can actually connect to (220).

**Definition 42.5** (Unrestricted deficit and its scale family). For a ball  $B_\rho(y) \subseteq \mathbb{R}^3$  on which  $\hat{e}$  is defined a.e. set

$$\tilde{\sigma}_\rho(y) := \rho^{-1} \int_{B_\rho(y)} |\nabla \hat{e}|^2 dx \quad (\text{no indicator}). \quad (221)$$

Say that  $\hat{e}$  satisfies  $(\Sigma_{\log})$  on  $B_{r^*}(x^*)$  with constant  $K$  if

$$\tilde{\sigma}_\rho(y) \leq K |\log \rho|^{-2} \quad \text{for every } B_\rho(y) \subseteq B_{r^*}(x^*), \ 0 < \rho < \frac{1}{2}. \quad (222)$$

At the single top scale  $\rho = r^*$ ,  $y = x^*$ , (222) reads  $\tilde{\sigma}_{r^*} \leq 4K / (\log M)^2$ .

**Lemma 42.6** (Poincaré translation). *Let  $C_P = \mu_1(B_1)^{-1/2}$  be the Poincaré–Wirtinger constant of the unit ball of  $\mathbb{R}^3$  ( $\mu_1$  the first nonzero Neumann eigenvalue,  $\mu_1 \approx 4.33$ , so  $C_P \leq \frac{1}{2}$ ). Then for any ball  $B_\rho(y)$  on which  $\hat{e} \in H^1$ ,*

$$\frac{1}{|B_\rho(y)|} \int_{B_\rho(y)} |\hat{e} - \hat{e}_{B_\rho(y)}| dx \leq C_P \left( \frac{3}{4\pi} \right)^{1/2} (\tilde{\sigma}_\rho(y))^{1/2} \leq \frac{1}{4} (\tilde{\sigma}_\rho(y))^{1/2}. \quad (223)$$

Consequently  $(\Sigma_{\log})$  with constant  $K$  implies

$$\sup_{B_\rho(y) \subseteq B_{r^*}(x^*)} \frac{1}{\phi(\rho)} \frac{1}{|B_\rho(y)|} \int_{B_\rho(y)} |\hat{e} - \hat{e}_{B_\rho(y)}| dx \leq \frac{1}{4} K^{1/2}, \quad (224)$$

i.e. the local  $\text{bmo}_\phi$  seminorm of  $\hat{e}$  on  $B_{r^*}(x^*)$  is at most  $\frac{1}{4} K^{1/2}$ , which is the hypothesis of Fact 42.2(c) in its local form. [Proved.]

*Proof.* Cauchy–Schwarz and Poincaré–Wirtinger componentwise (the constant for an  $\mathbb{R}^3$ -valued function is the scalar one, by summing the three scalar inequalities and using  $|v| \leq \sum_i |v_i|$  only after squaring):

$$-\int_{B_\rho} |\hat{e} - \hat{e}_{B_\rho}| \leq \left( -\int_{B_\rho} |\hat{e} - \hat{e}_{B_\rho}|^2 \right)^{1/2} \leq C_P \rho \left( -\int_{B_\rho} |\nabla \hat{e}|^2 \right)^{1/2}.$$

Since  $|B_\rho| = \frac{4\pi}{3} \rho^3$  and  $\int_{B_\rho} |\nabla \hat{e}|^2 = \rho \tilde{\sigma}_\rho$ , we get  $-\int_{B_\rho} |\nabla \hat{e}|^2 = \frac{3}{4\pi} \tilde{\sigma}_\rho \rho^{-2}$ , and  $\rho \cdot (\cdot)^{1/2} = (3/4\pi)^{1/2} \tilde{\sigma}_\rho^{1/2}$ . With  $C_P \leq \frac{1}{2}$  and  $(3/4\pi)^{1/2} < 0.489$  the numerical form follows. Then (224) is (223) divided by  $\phi(\rho) = 1/|\log \rho|$  together with (222).  $\square$

**Remark 42.7** (This is the S56 sketch, and it is the only part of it that survives). The audit of S56 sketched, without proof, that Poincaré converts the external hypothesis into “ $\sigma^* \lesssim 1/(\log M)^2$ , a quantified form of  $\sigma^*$ -decay”. Lemma 42.6 confirms the *shape* of that computation and nothing else. Three things have to be corrected, and each is a separate failure of the intended dictionary: (223) runs from gradients to oscillation and *not back* (Proposition 42.8); it must use the *unrestricted*  $\tilde{\sigma}$ , and the indicator of Definition 26.3 cannot be dropped (Proposition 42.9); and it must hold at *every* scale, which the critical scale does not deliver (Proposition 42.10).

## 42.5 Three counterexamples: the translation is not an equivalence

Throughout this subsection the comparison is local to  $B_{r^*}(x^*)$  and instantaneous, which is the level at which both conditions are stated: neither  $\sigma^*$ -decay nor (220) is derived from the equations, both being hypotheses imposed on the field at a time. All three fields below are smooth and divergence-free, hence are the vorticity of a solenoidal velocity field.

**Proposition 42.8** (Log-weighted BMO does not control the coherence deficit). *For each  $\delta \in (0, e^{-1})$  let*

$$\omega^{(\delta)}(x) = M(\cos g_\delta(x_3), \sin g_\delta(x_3), 0), \quad g_\delta(s) = \frac{1}{2|\log \delta|} \sin(s/\delta).$$

*Then  $\nabla \cdot \omega^{(\delta)} = 0$ ,  $|\omega^{(\delta)}| \equiv M$  (so the indicator of Definition 26.3 is inert and  $\sigma^* = \tilde{\sigma}_{r^*}$ ), the local  $\text{bmo}_\phi$  norm of  $\hat{e}$  obeys  $\|\hat{e}\|_{\text{bmo}_\phi} \leq 2$  uniformly in  $\delta$ , and*

$$\sigma^* \geq \frac{\pi}{9} \frac{(r^*)^2}{\delta^2 \log^2(1/\delta)} \longrightarrow \infty \quad (\delta \rightarrow 0, r^* \text{ fixed}).$$

*Hence the hypothesis of Fact 42.2(c) implies neither  $\sigma^* \rightarrow 0$  nor  $\sigma^* = O(1)$ : it carries no information about  $\nabla \hat{e}$  in  $L^2$  at all. [Proved.]*

*Proof.*  $\omega_1^{(\delta)}, \omega_2^{(\delta)}$  depend on  $x_3$  only and  $\omega_3^{(\delta)} \equiv 0$ , so the divergence vanishes;  $|\omega| = M$  is immediate. With  $\hat{e} = (\cos g_\delta, \sin g_\delta, 0)$ ,  $\nabla \hat{e} = g'_\delta(x_3) (-\sin g_\delta, \cos g_\delta, 0) \otimes e_3$ , so  $|\nabla \hat{e}| = |g'_\delta| = \frac{1}{2\delta|\log \delta|} |\cos(x_3/\delta)|$ . For  $\delta \leq r^*$ ,  $\int_{B_{r^*}} \cos^2(x_3/\delta) dx \geq \frac{1}{3}|B_{r^*}|$ , so

$$\sigma^* = (r^*)^{-1} \int_{B_{r^*}} |\nabla \hat{e}|^2 \geq \frac{1}{4\delta^2 \log^2(1/\delta)} \cdot \frac{1}{3} \cdot \frac{4\pi}{3} (r^*)^2 = \frac{\pi}{9} \frac{(r^*)^2}{\delta^2 \log^2(1/\delta)}.$$

For the oscillation,  $|\hat{e}(x) - \hat{e}(x')| \leq |g_\delta(x_3) - g_\delta(x'_3)|$ , so on a ball of radius  $\rho$  the oscillation of  $\hat{e}$  is at most  $\min(2\|g_\delta\|_\infty, 2\rho\|g'_\delta\|_\infty) = \frac{1}{|\log \delta|} \min(1, \rho/\delta)$ , and  $-\int_{B_\rho} |\hat{e} - \hat{e}_{B_\rho}|$  is bounded by the same quantity. Hence  $\frac{1}{\phi(\rho)} - \int_{B_\rho} |\hat{e} - \hat{e}_{B_\rho}| \leq \frac{|\log \rho|}{|\log \delta|} \min(1, \rho/\delta)$ . For  $\rho \geq \delta$  this is  $\leq |\log \rho|/|\log \delta| \leq 1$  since  $|\log \cdot|$  decreases on  $(0, 1)$ ; for  $\rho < \delta < e^{-1}$  it is  $\frac{\rho|\log \rho|}{\delta|\log \delta|} \leq 1$  since  $\rho \mapsto \rho|\log \rho|$  increases on  $(0, e^{-1})$ . Adding  $\|\hat{e}\|_{L^\infty} = 1$  gives  $\|\hat{e}\|_{\text{bmo}_\phi} \leq 2$ .  $\square$

**Proposition 42.9** (The level-set indicator cannot be dropped, and is what makes  $\sigma^*$  finite). *Let  $\omega^{(\delta)}(x) = M \cos(x_3/\delta) e_1$  with  $0 < \delta < r^*$ . Then  $\nabla \cdot \omega^{(\delta)} = 0$ ,  $\|\omega^{(\delta)}\|_{L^\infty} = M$ , and*

- (i)  $\sigma^* = 0$  exactly: on the open set  $\{|\omega| \geq M/4\}$  the direction  $\hat{e} = \text{sgn}(\cos(x_3/\delta)) e_1$  is locally constant, so  $\nabla \hat{e} \equiv 0$  there;
- (ii)  $\tilde{\sigma}_\rho = +\infty$  for every ball meeting a zero plane  $\{\cos(x_3/\delta) = 0\}$ , and, for balls  $B_\rho(y)$  centred on such a plane,  $-\int_{B_\rho(y)} |\hat{e} - \hat{e}_{B_\rho(y)}| = 1$  for every  $\rho$ , whence  $\|\hat{e}\|_{\text{bmo}_\phi} = +\infty$ .

So  $\sigma^* = 0$  is compatible with the maximal possible failure of (220). Moreover this is the generic situation: near any transversal zero of  $\omega$ ,  $\hat{e}$  has an order-one jump at every scale,  $|\nabla \hat{e}| \lesssim |\nabla \omega|/|\omega|$  is not square-integrable, and both  $\tilde{\sigma}_\rho$  and  $\|\hat{e}\|_{\text{bmo}_\phi}$  are infinite while  $\sigma^*$  is unaffected. [Proved.]

*Proof.* Divergence-freeness and the sup-norm are immediate. (i)  $\{|\omega| \geq M/4\} = \{|\cos(x_3/\delta)| \geq 1/4\}$  is open and  $\cos(x_3/\delta)$  does not change sign on any of its connected components, so  $\hat{e}$  is locally constant there. (ii) On a ball centred on a zero plane,  $\hat{e} = +e_1$  on one open half and  $-e_1$  on the other, the two halves having equal measure, so  $\hat{e}_{B_\rho} = 0$  and  $-\int |\hat{e} - \hat{e}_{B_\rho}| = 1$ ; dividing by  $\phi(\rho)$  and letting  $\rho \rightarrow 0$  gives  $+\infty$ . The distributional  $\nabla \hat{e}$  is a surface measure on the zero planes, so  $\tilde{\sigma}_\rho = +\infty$ .  $\square$

**Proposition 42.10** (One scale does not give the scale family). *Fix a radial profile  $\chi_0 \in C_c^\infty([0, 1])$  with  $\|\chi'_0\|_\infty = \frac{1}{4}$  and  $\chi'_0 \not\equiv 0$ , let  $y_0 \in B_{r^*/2}(x^*)$ , and for  $0 < \delta < r^*/4$  put*

$$\psi_\delta(x) = \delta \chi_0\left(\frac{|x - y_0|}{\delta}\right), \quad \omega^{(\delta)} = e_3 + \nabla \times (\psi_\delta e_1).$$

*Then  $\nabla \cdot \omega^{(\delta)} = 0$ ,  $\frac{3}{4} \leq |\omega^{(\delta)}| \leq \frac{5}{4}$  (so  $|\omega| \geq M/4$  everywhere and the indicator is inert), and there are constants  $C, c_0 > 0$  independent of  $\delta$  with*

$$\sigma^* \leq C \frac{\delta}{r^*} \longrightarrow 0, \quad \text{while} \quad -\int_{B_\delta(y_0)} |\hat{e} - \hat{e}_{B_\delta(y_0)}| = c_0 \quad \text{for every } \delta.$$

*Hence  $\sigma^* \rightarrow 0$  — the decay this document's open problem asks for (Corollary 31.5) — does not imply (220) at any scale below  $r^*$ , since (220) demands  $c_0 \leq K/|\log \delta|$ . [Proved.]*

*Proof.*  $\nabla \times$  of anything is divergence-free. With  $\psi_\delta$  radial about  $y_0$ ,  $\nabla \times (\psi_\delta e_1) = \nabla \psi_\delta \times e_1$  and  $|\nabla \psi_\delta| = |\chi'_0| \leq \frac{1}{4}$ , giving the magnitude bounds and  $M = \|\omega\|_{L^\infty(B_{r^*})} \leq \frac{5}{4}$ , so  $M/4 \leq \frac{5}{16} < \frac{3}{4}$ . The field  $W_\delta := \nabla \psi_\delta \times e_1$  satisfies  $W_\delta(x) = W_1(y_0 + (x - y_0)/\delta)$  — it depends on  $x$  only through  $(x - y_0)/\delta$  — so  $\hat{e}$  restricted to  $B_\delta(y_0)$  is the  $\delta$ -dilate of the fixed field  $\hat{e}_1$  on  $B_1(y_0)$ , and its mean oscillation over  $B_\delta(y_0)$  equals the fixed number  $c_0 := -\int_{B_1(y_0)} |\hat{e}_1 - (\hat{e}_1)_{B_1(y_0)}|$ , which is positive because  $\chi'_0 \not\equiv 0$  makes  $\hat{e}_1$  non-constant. Finally  $|\nabla \hat{e}| \leq 2|\nabla \omega|/|\omega| \leq \frac{8}{3}\|\nabla W_\delta\|_\infty \leq C/\delta$ , supported in  $B_\delta(y_0)$ , so  $\int_{B_{r^*}} |\nabla \hat{e}|^2 \leq C^2 \delta^{-2} \cdot \frac{4\pi}{3} \delta^3 = C' \delta$  and  $\sigma^* \leq C' \delta / r^*$ .  $\square$

## 42.6 The verdict on the dictionary

**Theorem 42.11** (The external hypothesis and  $\sigma^*$ -decay are logically independent). *Consider, as conditions on a smooth divergence-free field  $\omega$  on  $B_{r^*}(x^*)$  at one time:*

- (A) ( $\sigma^*$ -decay, as this document uses it)  $\sigma^*$  *small*, with  $\sigma^*$  as in Definition 26.3 — an  $L^2$  bound on  $\nabla \hat{e}$ , on the single ball  $B_{r^*}(x^*)$ , at the single scale  $r^*$ , restricted to  $\{|\omega| \geq M/4\}$ .
- (B) (the external hypothesis, local form) the  $\text{bmo}_{1/|\log r|}$  seminorm of  $\hat{e}$  over all balls  $B_\rho(y) \subseteq B_{r^*}(x^*)$  is small — an  $L^1$  bound on the mean oscillation of  $\hat{e}$ , at every scale, with no level-set restriction.

Then (A)  $\not\Rightarrow$  (B) and (B)  $\not\Rightarrow$  (A), and the failures are not quantitative near-misses:

- (i) (B) with constant 2 is compatible with  $\sigma^* = +\infty$  in the limit (Proposition 42.8);
- (ii)  $\sigma^* = 0$  is compatible with  $\|\hat{e}\|_{\text{bmo}_\phi} = +\infty$  (Proposition 42.9);
- (iii)  $\sigma^* \rightarrow 0$  is compatible with the failure of (B) by a factor  $|\log \delta|$  at scale  $\delta$  (Proposition 42.10).

What survives is a single one-way implication, in the direction gradients  $\rightarrow$  oscillation, for the unrestricted scale family:

$$(\Sigma_{\log}) \text{ with constant } K \implies \text{(B) with constant } \frac{1}{4} K^{1/2}, \quad (225)$$

and the implication is strict: by Proposition 42.8 the converse fails. [Proved.]

*Proof.* (i)–(iii) are the three propositions of §42.5; (225) is Lemma 42.6, and its strictness is Proposition 42.8, whose field satisfies (B) with constant 2 while  $\tilde{\sigma}_{r^*} = \sigma^* \rightarrow \infty$  violates (222) for every fixed  $K$ .  $\square$

**Corollary 42.12** (The S56 dictionary is refuted in the form proposed). *The relation “ $\xi \in \text{bmo}_{1/|\log r|}$  at  $r^*$  reads  $\sigma^* \lesssim 1/(\log M)^2$ ” is **not** a translation. It is correct only as a sufficient condition, only for the unrestricted  $\tilde{\sigma}_{r^*}$  of (221), and only for the single ball  $B_{r^*}(x^*)$  — i.e. only as the top scale of (225). In particular the external hypothesis is **not** a fifth avatar of  $\sigma^*$ -decay, and no cheapness claim can be made in either direction. [Proved.]*

**Remark 42.13** (What the correct translation is, and what it costs). The honest statement of the external hypothesis in this document’s variables is (222), and against Definition 26.3 it differs in three independent ways, each of which is a strengthening except the first:

- (a) **Weaker in norm.**  $L^1$  mean oscillation is weaker than an  $L^2$  gradient bound; this is the one genuine relaxation, and it is what lets the source admit “wild oscillatory defects”. It is also why (B) carries no gradient information at all (Proposition 42.8).
- (b) **Stronger in scale.** It is a scale family on  $(0, r^*]$ , not a statement at  $r^*$ . The proof of Fact 42.2(c) uses the hypothesis on every sub-ball of  $B_{2R}$  (near field, via the Jones extension), on  $B_R$  (John–Nirenberg), and on the dyadic balls  $B_{2^j R}$  up to radius  $R^{1/2}$  (oscillation tail), for every small  $R$ . It does *not* collapse to the critical scale, and Proposition 42.10 shows the collapse is false.
- (c) **Stronger in support.** It has no level-set restriction, and by Proposition 42.9 it therefore also asserts, silently, that  $\omega$  has no transversal zero in the region. This is the substantive content of the  $\mathbf{1}_{\{|\omega| \geq M/4\}}$  in (143): not a convenience, but what keeps  $\sigma^*$  finite. Any future attempt to transport an unrestricted-oscillation hypothesis into this document must supply a zero-free core as a separate hypothesis.

## 42.7 Against the rest of the no-concentration family

- Remark 42.14** (Against (AX) and against saturation). (i) (AX) (Proposition 38.12) is a pointwise bound on a *magnitude* component,  $\|\omega_\theta/r\|_{L^\infty(B_{2r^*})} \leq KM/r^*$ , inside the axisymmetric class; (B) is an integrated bound on the *direction*, in no symmetry class. Neither implies the other, and there is no common refinement: (AX) permits an arbitrary direction field away from the axis, and Proposition 42.9’s field satisfies neither.
- (ii) **Saturation** (Definition 37.1) is a statement about  $M(t)$  in time; (B) is a statement about  $\hat{e}(\cdot, t)$  in space. No implication runs either way. The comparison that *is* meaningful is of currency, and it is exact: Theorem 37.21 says that removing one logarithm from  $\|\nabla u\|_{L^\infty} \lesssim M\mathcal{L}$  is equivalent to logarithm-free saturation, and Fact 42.2(c) buys exactly one logarithm — but in  $\|\alpha\|_{L^{3/2,\infty}(B_R)}$ , not in  $\|\nabla u\|_{L^\infty}$ . That is a Wall 1(d)-permitted reformulation paying the same single-logarithm deficit in a weaker norm, which is precisely why Proposition 38.9 does not contradict it: the counterexample there saturates an  $L^\infty$  bound and says nothing about a restricted Lorentz norm of  $\hat{e} \cdot S\hat{e}$ .
  - (iii) **The family verdict.** The pattern Remark 38.13 records now has a fourth instance. (B) is a no-concentration hypothesis of the same family as saturation and (AX): scale-invariant, imposed rather than proved, and — by Fact 42.2(f) and (g) — not established for Navier–Stokes solutions by its own companion paper, which instead exhibits two natural critical cores on which it fails. *It has not been shown to be cheaper than what it replaces, and this section does not claim that it is.* What is new, relative to the three earlier instances, is that this one is *not* a relabelling of  $\sigma^*$ -decay (Corollary 42.12); it is a genuinely different condition on the same object, and it is incomparable.

## 42.8 The imported technique, checked in isolation

**Remark 42.15** (The commutator/Coifman–Rochberg–Weiss step holds). The step carrying Fact 42.2(c) was checked at S57 independently of the rest of [60]. It is **correct**, and the chain is the following, each link standard.

- (1) *The cancellation.*  $S = \mathcal{T}(\omega\xi)$  with  $\mathcal{T}$  a matrix of second-order Riesz transforms; the fully symmetric contraction of its kernel against  $e \otimes e \otimes e$  vanishes, which is exactly the factor  $\det(\hat{y}, \xi(x+y), \xi(x))$  in Constantin's formula for  $\alpha$  [13, 12]. Hence  $\alpha = \xi \cdot [\mathcal{T}, \xi](\omega) \xi$  and  $|\alpha| \leq |[\mathcal{T}, \xi](\omega)|$ . Correct, and not new: it is Constantin–Fefferman's observation rewritten.
- (2) *Smallness of the symbol.* Since  $\phi(r) = 1/|\log r|$  is increasing on  $(0, 1)$ , (220) gives, for every sub-ball of  $B_{2R}$  of radius  $r \leq 2R < 1/2$ , mean oscillation  $\leq \|\xi\|_{\text{bmo}_\phi} \phi(r) \leq \|\xi\|_{\text{bmo}_\phi} \phi(2R)$ , i.e.  $\|\xi\|_{\text{BMO}(B_{2R})} \leq C\phi(2R)$ . Correct; the monotonicity is used in the right direction, and again at the truncation  $2^{N+1}R \approx R^{1/2}$  of the dyadic tail.
- (3) *Jones extension.* A ball is a uniform domain with absolute (scale-invariant) constants, so [64] supplies  $\tilde{\xi}$  on  $\mathbb{R}^3$  with  $\|\tilde{\xi}\|_{\text{BMO}(\mathbb{R}^3)} \leq C\phi(2R)$ . Correct. The substitution  $[\mathcal{T}, \xi](\omega\chi_{B_{2R}})(x) = [\mathcal{T}, \tilde{\xi}](\omega\chi_{B_{2R}})(x)$  for  $x \in B_R$  is legitimate because the source is supported in  $B_{2R}$ , where  $\tilde{\xi} = \xi$ , and  $x \in B_R \subset B_{2R}$ .
- (4) *CRW itself.* [63] gives  $\|[b, R_j]\|_{L^p \rightarrow L^p} \leq C_p \|b\|_{\text{BMO}}$ ,  $1 < p < \infty$ . The operator needed here is a composition of two Riesz transforms, and  $[R_i R_j, b] = R_i[R_j, b] + [R_i, b]R_j$ , so the bound transfers with the same linear dependence on  $\|b\|_{\text{BMO}}$  *without* invoking any general Calderón–Zygmund version. The vector-valued bookkeeping is componentwise. Correct.
- (5) *The passage to  $L^{3/2, \infty}$ .* Real (Marcinkiewicz–Hunt) interpolation [65] between  $L^{p_0} \rightarrow L^{p_0}$  and  $L^{p_1} \rightarrow L^{p_1}$  with  $1 < p_0 < 3/2 < p_1 < \infty$  gives boundedness on  $(L^{p_0}, L^{p_1})_{\theta, \infty} = L^{3/2, \infty}$  with norm  $\lesssim \|b\|_{\text{BMO}}^{1-\theta} \|b\|_{\text{BMO}}^\theta = \|b\|_{\text{BMO}}$ . Correct. *One cosmetic error:* the source calls these “the reflexive Lorentz spaces”;  $L^{3/2, \infty}$  is not reflexive (it is not even normed in the displayed quasi-norm, though for  $p > 1$  it is normable via  $f^{**}$ ). The conclusion is unaffected.
- (6) *The far field.* The rearrangement computation

$$\left\| |y|^{-3} \chi_{\{|y| > 2R\}} \right\|_{L^{3,1}} \asymp R^{-2},$$

the John–Nirenberg  $L^2$  and  $L^4$  oscillation bounds, the local embeddings

$$\|g\|_{L^{3/2, \infty}(B_R)} \lesssim R^{1/2} \|g\|_{L^2(B_R)}, \quad \|g\|_{L^{3,1}(B)} \lesssim |B|^{1/12} \|g\|_{L^4(B)},$$

the telescoping  $|c_{k+1} - c_0| \lesssim \sum_{j \leq k+1} \phi(2^j R)$ , and the reversed double sum were all re-derived at S57 and are correct as stated.

**Consequence for this programme.** The technique is available independently of whether [60] is right as a whole: it is a proof that *the stretching eigenvalue is a commutator whose symbol is the direction field, so that any smallness of the direction's BMO norm at small scales is a gain on  $\alpha$  in a restricted critical Lorentz norm, linearly*. This is the first estimate on record in this programme that extracts a gain from the direction field without an  $L^\infty$  Calderón–Zygmund bound, and it is therefore inside Wall 1(d)'s permitted class. **It is not, by itself, a route around Wall 1:** by Theorem 42.11 the smallness it consumes is not available from  $\sigma^*$ , and by Fact 42.2(g) it is not available from the companion paper either.

**Remark 42.16** (The Albritton–Bradshaw prior, corrected). S56 recorded that Albritton–Bradshaw [62] analysed earlier sparseness claims of the same author and concluded that “the scaling gap is not reduced”, and inferred that this bears negatively on the endgame of [60], which routes through sparseness and analyticity. **Read at S57, the prior does not say that.** Their paper has two halves. The second half is indeed negative, and is about the *a priori* sparseness classes  $Z_{\tilde{\alpha}_k}^{(k)}$  of Bradshaw–Farhat–Grujić and Grujić–Xu: their Proposition 4.1 and §4 examples show that within that class “membership in  $Z_{\tilde{\alpha}_k}^{(k)}$  is not stronger than the condition of finite kinetic energy  $u \in L_t^\infty L_x^2$ ”, and that the informal identification  $Z_\alpha^{(k)} \sim W^{k-1/\alpha, \infty}$  “should not be taken seriously”. The *first* half is positive: they “give a simple proof that sufficiently sparse Navier–Stokes solutions do not develop singularities”, explicitly “an alternative to the approach of [Gru13], which is based on analyticity and the ‘harmonic measure maximum principle’” (their Theorem 1.2 and Corollary 1.3). The endgame of [60, §7] uses only the *conditional* direction — sparseness at a scale below the analyticity radius implies continuation — which is the half Albritton–Bradshaw reprove and endorse. **The negative prior therefore does not attach to the step S56 attached it to.** It remains a relevant caution about how sparseness results in this lineage have been *interpreted*, and it is recorded here so that the correction, not the original, is what a later session inherits.

## 42.9 Ledger

| Item                                                            | Status                   | Where         |
|-----------------------------------------------------------------|--------------------------|---------------|
| Scale identification $\ell \asymp r^*$                          | <b>Proved</b>            | Lem. 42.4(i)  |
| Source’s direction bound $\equiv \sigma^*$ bounded              | <b>Proved</b>            | Lem. 42.4(ii) |
| Poincaré translation (gradients $\rightarrow$ osc.)             | <b>Proved</b>            | Lem. 42.6     |
| $\text{bmo}_\phi \not\asymp \sigma^*$ bounded                   | <b>Proved</b> (c/ex)     | Prop. 42.8    |
| Indicator not removable                                         | <b>Proved</b> (c/ex)     | Prop. 42.9    |
| $\sigma^* \rightarrow 0 \not\asymp \text{bmo}_\phi$ below $r^*$ | <b>Proved</b> (c/ex)     | Prop. 42.10   |
| The two hypotheses are independent                              | <b>Proved</b>            | Thm. 42.11    |
| S56’s dictionary                                                | <b>Refuted as stated</b> | Cor. 42.12    |
| Commutator/CRW step                                             | <b>Verified correct</b>  | Rem. 42.15    |
| [60, Thm. 7.4] as a whole                                       | <b>Not adjudicated</b>   | Rem. 42.1     |
| Propagation of the hypothesis                                   | <b>Open</b>              | Fact 42.2(f)  |
| Albritton–Bradshaw prior                                        | <b>Corrected</b>         | Rem. 42.16    |
| Walls 1, 3, 9; <b>(AX)</b> ; saturation                         | <b>Unchanged</b>         | §44           |

**Remark 42.17** (What is explicitly not claimed). (i) No position is taken on whether [60, Thm. 7.4] is true. §§5–7 of that paper were read and their internal steps found consistent, but a referee report is not a session’s work and was not attempted; Fact 42.2 records the dependency structure, not a verdict.

- (ii) Nothing here is an attempt to prove the  $\text{bmo}_{1/|\log r|}$  hypothesis for Navier–Stokes solutions. That is the open problem in different notation.
- (iii) The counterexamples of §42.5 are admissible instantaneous vorticity fields, not Navier–Stokes solutions. They refute implications *between the two conditions as stated*, which is the level at which cheapness claims are made and the level S50 set. They do not

exclude the possibility that along an actual blow-up trajectory the two conditions are dynamically linked; no such link is known, and establishing one would be a new result.

- (iv) No new conjecture is introduced, no wall is reclassified, and (222) is a *definition used for comparison*, not a target proposed for attack.

## 43 CD-1: The Forcing Floor as a General Theorem (Session S60)

§41 proved a forcing floor for *one* field: the leading axisymmetric profile of [55]. This section asks what in that proof was specific to that profile and what was generic, and answers it by stating and proving the general theorem the S55 argument is an instance of. The outcome is that the profile-specific ingredient — exact self-similarity — is load-bearing for far less than S55’s own record suggested: it is needed neither for the exponent nor for the constant, only for the *time-localisation* of the conclusion (Remark 43.6).

**Scope, stated first and not to be detached from what follows.** Everything here is a statement about a *given* axisymmetric velocity field, with  $f$  *defined* as its momentum residual, exactly as in Lemma 41.1. No forcing term is introduced into any argument of this document toward Fefferman’s alternatives (A) or (B); the standing prohibition on doing so is unaffected. This section is not a step toward (A) or (B) — it proves a *lower* bound on a residual, the opposite direction of travel — and it is not a challenge to (C) or (D): the class is *axisymmetric*, and the field of Theorem 1.1 of [55] is not, so nothing here applies to it (Remark 41.11, restated at Remark 43.10(iv)). Wall 9’s classification is unchanged.

### 43.1 Extension of the two S55 lemmas to $\nu \geq 0$

**Remark 43.1** (Both S55 lemmas hold for  $\nu = 0$ , i.e. for Euler). Lemma 41.1 and Lemma 41.2 are stated for  $\nu > 0$ . Neither proof uses positivity.

- (a) In Lemma 41.1,  $\nu$  multiplies the vector Laplacian throughout; setting  $\nu = 0$  deletes that term from the definition of  $f$  and from (208) simultaneously, and the three ingredients (a)–(c) of its proof are unchanged. At  $\nu = 0$  the identity reads  $\partial_t \Gamma + u_r \partial_r \Gamma + u_z \partial_z \Gamma = r f_\theta$  with  $f = \partial_t u + (u \cdot \nabla)u + \nabla p$ .
- (b) In Lemma 41.2,  $\nu$  enters only through the single inequality  $\nu(\partial_r^2 \Gamma + \partial_z^2 \Gamma) \leq 0$  at an interior maximum, which at  $\nu = 0$  holds as the identity  $0 \leq 0$ . Parts (i), (iii), (iv), (v) never mention  $\nu$ .

Both lemmas therefore hold verbatim for every  $\nu \geq 0$ , and are used in that range below. (*At  $\nu = 0$  the unforced case of (212) is the classical conservation of  $\max r u_\theta$  by axisymmetric Euler transport, so the generalisation degenerates correctly at both ends.*)

### 43.2 The general theorem

**Theorem 43.2** (Cost of forcing at the swirl maximum). *Let  $\nu \geq 0$ , let  $(u, p)$  be axisymmetric and smooth on  $\mathbb{R}^3 \times [t_0, t_1]$  with  $\operatorname{div} u = 0$ , let  $f := \partial_t u + (u \cdot \nabla)u - \nu \Delta u + \nabla p$  be its momentum*



residual, and assume **(H1)–(H3)** of Lemma 41.2. Write  $\Gamma = ru_\theta$ ,  $M(t) = \max_x \Gamma(\cdot, t) > 0$ ,  $\Sigma(t) = \{\Gamma(\cdot, t) = M(t)\}$ , and

$$R(t) := \max_{x \in \Sigma(t)} r(x) \in (0, \infty), \quad F(t) := \max_{x \in \Sigma(t)} |f_\theta(x, t)|. \quad (226)$$

Then:

- (i) **Ratio form (no geometric hypothesis).** For every  $t$  there is a swirl maximiser  $x_\star = x_\star(t) \in \Sigma(t)$  with

$$\frac{f_\theta(x_\star, t)}{u_\theta(x_\star, t)} \geq \frac{d^+}{dt} \log M(t). \quad (227)$$

Both sides have dimensions of inverse time;  $u_\theta(x_\star, t) > 0$  automatically.

- (ii) **Radius form.** For every  $t$ ,

$$F(t) \geq \frac{1}{R(t)} \frac{d^+ M}{dt}(t). \quad (228)$$

- (iii) **Integrated form, assuming no differentiability anywhere.** For  $t_0 \leq s \leq t < t_1$ ,

$$\int_s^t R(\sigma) F(\sigma) d\sigma \geq M(t) - M(s). \quad (229)$$

[Proved here. Proposition 41.5 is the special case recorded in Corollary 43.5.]

*Proof.* By Remark 43.1 the two S55 lemmas apply for every  $\nu \geq 0$ . By Lemma 41.2(i) every  $x \in \Sigma(t)$  has  $r(x) > 0$ , and  $\Sigma(t)$  is nonempty and compact by (H1)–(H2);  $rf_\theta$  is continuous by (H1); so the maximum in (211) is attained, at some  $x_\star \in \Sigma(t)$ , and  $R(t)$ ,  $F(t)$  in (226) are attained and finite, with  $R(t) > 0$ .

(ii). By (211),

$$\frac{d^+ M}{dt}(t) \leq (rf_\theta)(x_\star, t) \leq r(x_\star) |f_\theta(x_\star, t)| \leq R(t) F(t),$$

and  $R(t) > 0$ .

(i). At the same  $x_\star$ ,  $\Gamma(x_\star, t) = M(t)$  gives  $u_\theta(x_\star, t) = M(t)/r(x_\star) > 0$ , since  $M(t) > 0$  by (H3) and  $r(x_\star) > 0$ . Hence

$$\frac{f_\theta(x_\star, t)}{u_\theta(x_\star, t)} = \frac{r(x_\star) f_\theta(x_\star, t)}{M(t)} \geq \frac{1}{M(t)} \frac{d^+ M}{dt}(t) = \frac{d^+}{dt} \log M(t),$$

the last step because  $M > 0$  is locally Lipschitz (Lemma 41.2(iii)), so  $\log M$  is too and its right derivative is  $M^{-1} d^+ M/dt$ .

(iii). This is (212) together with  $\max_{\Sigma(\sigma)} (rf_\theta) \leq R(\sigma) F(\sigma)$  pointwise in  $\sigma$ ; measurability of the integrand is as in the proof of Lemma 41.2(iv).  $\square$

**Remark 43.3** (How to read (227)). Inequality (227) carries no geometric input at all — no core radius, no growth rate, no self-similarity — and says:

at a swirl maximum, the azimuthal residual, measured against the local swirl velocity, is at least the logarithmic growth rate of the swirl maximum.

If  $M(t) \asymp \tau^{-a}$  with  $\tau = T - t$  then the right side is  $a/\tau$ , so  $|f_\theta| \geq (a/\tau) |u_\theta|$  at that point: the residual is at least as large as the term  $\partial_t u_\theta$  would be for a field growing at that rate. This is the general form of Remark 41.6's "full leading order in its own momentum equation", which S55 verified term by term for one profile using that profile's four exponents. Here it is a consequence of the swirl maximum principle alone.

### 43.3 The rate form, and what it contains

**Corollary 43.4** (Prescribed swirl growth on a prescribed core geometry). *Keep the hypotheses of Theorem 43.2 on  $[s_0, T)$  with  $T < \infty$ , write  $\tau = T - t$ , and assume*

(G1) (swirl growth rate)  $M(t) \geq c_1 \tau^{-a}$  on  $[s_0, T)$ , for some  $a > 0$ ,  $c_1 > 0$ ;

(G2) (core-radius geometry)  $R(\sigma) \leq R_0 (T - \sigma)^b$  on  $[s_0, T)$ , for some  $R_0 > 0$  and  $b \in \mathbb{R}$ .

Suppose  $F(\sigma) \leq K(T - \sigma)^{-p}$  on  $[s_0, T)$  for some  $K, p > 0$ . Then

$$p \geq 1 + a + b, \quad (230)$$

and if  $p = 1 + a + b$  then

$$K \geq \frac{a c_1}{R_0}. \quad (231)$$

Equivalently: no axisymmetric field with swirl-maximum growth  $\tau^{-a}$  whose swirl maximisers stay inside radius  $R_0 \tau^b$  can have azimuthal residual  $O(\tau^{-p})$  at those maximisers for any  $p < 1 + a + b$ . [Proved here.]

*Proof.* Put  $\tau_0 = T - s_0$  and  $\tau_t = T - t$ . By (229) and the two hypotheses, for every  $t \in (s_0, T)$ ,

$$c_1 \tau_t^{-a} - M(s_0) \leq M(t) - M(s_0) \leq R_0 K \int_{s_0}^t (T - \sigma)^{b-p} d\sigma = R_0 K \int_{\tau_t}^{\tau_0} \tau^{b-p} d\tau. \quad (232)$$

Three cases.

If  $b - p > -1$ , the right integral is at most  $\tau_0^{b-p+1}/(b - p + 1)$ , a constant, while the left side  $\rightarrow +\infty$  as  $t \uparrow T$  (since  $a > 0$ ,  $c_1 > 0$ ). Contradiction.

If  $b - p = -1$ , the right integral is  $\log(\tau_0/\tau_t)$ , which is  $o(\tau_t^{-a})$  for every  $a > 0$ , while the left side is  $\geq \frac{1}{2} c_1 \tau_t^{-a}$  for  $t$  close to  $T$ . Contradiction.

If  $b - p < -1$ , write  $\beta := p - b - 1 > 0$ ; then

$$\int_{\tau_t}^{\tau_0} \tau^{b-p} d\tau = \frac{\tau_t^{-\beta} - \tau_0^{-\beta}}{\beta} \leq \frac{\tau_t^{-\beta}}{\beta}.$$

Substituting into (232) and multiplying by  $\tau_t^a > 0$ ,

$$c_1 - M(s_0) \tau_t^a \leq \frac{R_0 K}{\beta} \tau_t^{a-\beta} \quad \text{for all } t \in (s_0, T). \quad (233)$$

If  $\beta < a$  the right side tends to 0 and the left to  $c_1 > 0$  as  $\tau_t \downarrow 0$ , a contradiction; hence  $\beta \geq a$ , which is (230). If  $\beta = a$ , i.e.  $p = 1 + a + b$ , then (233) reads  $c_1 - M(s_0) \tau_t^a \leq R_0 K/a$ , and letting  $\tau_t \downarrow 0$  gives  $c_1 \leq R_0 K/a$ , which is (231).  $\square$

**Corollary 43.5** (Proposition 41.5 is the case  $(a, b) = (h, \frac{1}{2})$ , constant included). *For the leading field of Proposition 41.5, (216) gives (G1) with  $a = h$  and  $c_1 = c_*$ , and (218) gives (G2) with  $b = \frac{1}{2}$  and  $R_0 = c_r$  (taking the maximum over maximisers if  $(X_*, \eta_*)$  is not unique). Corollary 43.4 then forbids  $|f_\theta^{(0)}| = O(\tau^{-p})$  at the maximisers for*

$$p < 1 + h + \frac{1}{2} = \frac{3}{2} + h = A + 1,$$

and at  $p = A + 1$  forces the constant  $K \geq hc_*/c_r$ . These are exactly the exponent and the constant of (215). [Immediate.]

#### 43.4 Which hypotheses were profile-specific, and which were not

**Remark 43.6** (The exact self-similarity of the OpenAI profile is load-bearing for less than S55 recorded). The S55 derivation obtained  $M(t) = c_*\tau^{-h}$  *exactly*, and identified that exactness — available because the leading field is exactly self-similar — as “what makes the floor exact rather than heuristic”. Corollary 43.4 locates that claim precisely.

- (a) **The exponent does not need it.** (230) follows from the one-sided bounds (G1)–(G2) alone. No exactness, no self-similarity, and — because the proof routes through the integrated form (229) rather than the differential form (228) — no differentiability of  $M$  at any point.
- (b) **The constant does not need it either.** (231) recovers  $hc_*/c_r$  from the asymptotic constants  $c_1, R_0$  alone.
- (c) **What it does buy is time-localisation, and only that.** With exactness one has the floor *at every*  $t$ , at an identified point. Without it, Corollary 43.4 is a statement about the whole interval: it excludes a uniform power bound, so it yields only “for every  $s < T$  there exists  $\sigma \in (s, T)$  at which the floor holds”, with  $\sigma$  not identified. For a construction that is merely asymptotically self-similar, that is the correct and the only available form.

So the honest summary is: *exact self-similarity was a convenience of presentation, not a hypothesis of the mathematics*. The one genuinely profile-specific step of S55 was Step 5 — the derivation of the maximiser’s radius  $c_r\tau^{1/2}$  as an *output*. In the general theorem that radius is an *input*, (G2); and part (i) of Theorem 43.2 dispenses with it altogether by measuring the residual against the local swirl instead of in absolute terms.

**Remark 43.7** (Hypothesis audit: what is generic, what does work, and where the class boundary is). The hypotheses divide cleanly into four groups.

- (a) **Generic, and doing no work:** (H1) and (H3). (H1) is smoothness. (H3) is an orientation convention: if  $\max \Gamma \leq 0$  while  $\max(-\Gamma) > 0$ , replace  $u_\theta$  by  $-u_\theta$ .
- (b) **Generic, and doing all the work:** the pressure-free swirl equation (208), and the annihilation of both the drift and the singular first-order term at an interior maximum. Neither uses anything about the field beyond axisymmetry and  $\operatorname{div} u = 0$ .

- (c) **The one hypothesis that must be checked per construction:** (H2), uniform compactness of the maximiser set on compact time intervals. It is what makes the Danskin upper bound in Lemma 41.2(iii) available, and it fails for fields whose swirl maximum escapes to spatial infinity. For the S55 profile it was established in Step 3 of Proposition 41.5 from the decay  $H \sim \sqrt{2}c_\infty X^{-h}$  and the vanishing of  $(1 - \eta^2)^h$  at  $\eta = \pm 1$ ; for a compactly supported construction it is automatic.
- (d) **The class boundary is axisymmetry, and it is absolute.** The only lever is  $(\nabla p)_\theta = r^{-1}\partial_\theta p = 0$ . Outside the axisymmetric class there is no scalar swirl equation, the pressure does not drop out, and no analogue of Theorem 43.2 is proved or claimed here. This is why the theorem, like §41 before it, says nothing whatever about Theorem 1.1 of [55].

### 43.5 A positive-measure form

Remark 41.3 recorded that Lemma 41.2 yields no statement on any set of positive measure, and S55 left open whether some uniform version is provable. The following is one — obtained not by upgrading the pointwise statement, which Remark 41.3(a),(b) shows is impossible, but by running the same swirl equation against  $|\Gamma|^{q-2}\Gamma$  instead of at a point.

**Proposition 43.8** ( *$L^q$  floor on the azimuthal residual*). *Let  $\nu \geq 0$  and let  $(u, p)$  be as in Lemma 41.1 on  $\mathbb{R}^3 \times I$ , with  $\Gamma(\cdot, t)$  compactly supported in a fixed compact set for  $t \in I$ . Fix  $2 \leq q < \infty$ . Then  $t \mapsto \|\Gamma(\cdot, t)\|_{L^q(\mathbb{R}^3)}$  is locally Lipschitz and, wherever it is positive,*

$$\frac{d}{dt} \|\Gamma(\cdot, t)\|_{L^q} \leq \|(rf_\theta)(\cdot, t)\|_{L^q} \quad \text{for a.e. } t \in I. \quad (234)$$

Consequently  $\|\Gamma(\cdot, t)\|_{L^q} - \|\Gamma(\cdot, s)\|_{L^q} \leq \int_s^t \|rf_\theta\|_{L^q} d\sigma$ : growth of the  $L^q$  swirl norm forces a floor on an integrated norm of the residual. [Proved here.]

*Proof.*  $\Gamma = ru_\theta = x_1u_2 - x_2u_1$  is smooth on  $\mathbb{R}^3$  and vanishes on the axis. Multiply (208) by  $q|\Gamma|^{q-2}\Gamma$  and integrate  $dx = r dr d\theta dz$  over  $\mathbb{R}^3$ ; all integrals converge and all integrations by parts are justified by compact support and smoothness. Four terms:

- (a) **Time.**  $\int q|\Gamma|^{q-2}\Gamma \partial_t \Gamma dx = \frac{d}{dt} \int |\Gamma|^q dx$ .
- (b) **Drift.**  $\int q|\Gamma|^{q-2}\Gamma (u \cdot \nabla \Gamma) dx = \int u \cdot \nabla (|\Gamma|^q) dx = 0$ , by  $\text{div } u = 0$  and compact support. (Here  $u \cdot \nabla \Gamma = u_r \partial_r \Gamma + u_z \partial_z \Gamma$  by axisymmetry.)
- (c) **Diffusion.**  $\int q|\Gamma|^{q-2}\Gamma \nu \Delta \Gamma dx = -q(q-1)\nu \int |\Gamma|^{q-2} |\nabla \Gamma|^2 dx \leq 0$  (for  $q \geq 2$  the integrand is locally integrable).
- (d) **The singular first-order term vanishes identically.** With  $dx = r dr d\theta dz$  the factor  $r$  cancels the  $r^{-1}$ :

$$\int q|\Gamma|^{q-2}\Gamma \left(-\frac{2\nu}{r} \partial_r \Gamma\right) dx = -4\pi\nu \int_{\mathbb{R}} \int_0^\infty \partial_r (|\Gamma|^q) dr dz = 4\pi\nu \int_{\mathbb{R}} |\Gamma(0, z)|^q dz = 0,$$

because  $\Gamma$  vanishes on the axis and has compact support. No  $\varepsilon$ -exhaustion of the axis is needed, for the same reason as in Lemma 41.2(i).

Hence, with  $N(t) := \|\Gamma(\cdot, t)\|_{L^q}$ ,

$$\frac{d}{dt}N^q \leq q \int |\Gamma|^{q-2} \Gamma (rf_\theta) dx \leq q N^{q-1} \|rf_\theta\|_{L^q}$$

by Hölder with exponents  $q/(q-1)$  and  $q$ . Since  $\frac{d}{dt}N^q = qN^{q-1}\frac{d}{dt}N$  wherever  $N > 0$  and  $N$  is differentiable, (234) follows;  $N$  is locally Lipschitz because  $\partial_t \Gamma$  is bounded on compacts and the supports are contained in a fixed compact, so the integrated form holds by absolute continuity.  $\square$

**Remark 43.9** (What the  $L^q$  form does and does not deliver). (a) It is a floor on  $\|rf_\theta\|_{L^q}$ , not on  $\|f_\theta\|_{L^q}$ . Converting needs a bound on  $r$  over the support of  $f_\theta$  — a strictly stronger geometric hypothesis than (G2), which constrains only the maximiser set.

- (b) It does not contradict Remark 41.3(b): that remark says no  $L^q$  bound follows *from the pointwise statement*, and the spike counterexample there still applies to Theorem 43.2. Proposition 43.8 is a different argument with a different hypothesis — growth of  $\|\Gamma\|_{L^q}$ , not of  $\max \Gamma$  — and the two are independent.
- (c) **It does not apply to the leading field of Proposition 41.5**, and this should be stated rather than glossed. There  $\Gamma^{(0)} = q^{-h}H$  with  $H \sim \sqrt{2}c_\infty X^{-h}$  at large  $X$  and  $X = r^2/(2q)$ , so  $\Gamma^{(0)} \rightarrow \sqrt{2}c_\infty 2^h r^{-2h}$  as  $X \rightarrow \infty$ , *independently of  $q$* ; that is not in  $L^{q'}(\mathbb{R}^3)$  for any  $q' < \infty$ . The proposition is a tool for compactly supported constructions, not for that exactly self-similar profile.
- (d) Letting  $q \rightarrow \infty$  formally returns Lemma 41.2(v); no independent content is claimed at the endpoint.

## 43.6 Scope

**Remark 43.10** (Scope of §43, at its true size). (i) **What is new.** A general lower bound, stated once for a class rather than verified for one profile: (227), (228), (229), (230)–(231); the extension to  $\nu = 0$  (Remark 43.1); the  $L^q$  form (Proposition 43.8); and the finding that exact self-similarity was not load-bearing (Remark 43.6).

- (ii) **What is elementary.** All of it. The ingredients are the classical axisymmetric swirl maximum principle, Danskin’s envelope theorem, and one division. No credit is claimed for machinery; the content is the generality and the hypothesis audit.
- (iii) **Still pointwise.** Parts (i) and (ii) are statements at one moving point, and Remark 41.3 applies to them verbatim. Proposition 43.8 is the one statement here that is not.
- (iv) **Not a statement about (C) or (D), and not about [55].** The class is axisymmetric (Remark 43.7(d)); the field of that paper’s Theorem 1.1 is not, so the theorem does not apply to it, and no error in it is asserted or implied. Wall 9 governs and is unchanged.
- (v) **Not a step toward (A) or (B).** It bounds a residual from below. Wall 1, Wall 3, Region (R2) and Proposition 40.9 are untouched.

- (vi) **No forcing is introduced into this document’s own arguments.**  $f$  is defined as the residual of a field that is given, exactly as in §41. Studying the residual a construction requires is not the same as adding a forcing term to a regularity argument, and the latter remains excluded.

## 44 Routes and Walls: A Referenced Record (Session S47)

This section is a record, not an argument. It lists the routes this programme has attempted since the audit of §25 and, for each, the obstruction that stopped it, so that a later session does not repeat a step already proved impossible, and so that the material exists in citable form outside the session logs.

Every entry uses the same four-part template.

- (a) **Route.** What was attempted, and what it would have delivered had it worked.
- (b) **Obstruction.** The statement that stops it, by label.
- (c) **Classification.** Exactly one of: PROVED IMPOSSIBLE — a proof in this document rules the route out, and no choice of parameters inside it can help; OPEN — nothing is proved against the route; it is unfinished, and remains available; SUPERSEDED — the route was replaced by a different formulation of the same target, which carries its own status; WITHDRAWN — an assertion was made in this document and has been retracted.
- (d) **Consequence.** What the wall forbids, and what it leaves permitted.

Four conventions. First, *labels are primary*. Line numbers are secondary, quoted as of session S47, and will rot; hand-written numbers in the older prose are unreliable for a separate reason recorded below. Second, every claim below resolves to a label in this document, except where a paragraph is explicitly marked *outside this document*, in which case it names the session log that carries it and is not used to support any statement here. Third, *nothing in this section is new mathematics*: no statement below is cited as a hypothesis anywhere else, and this section may be deleted without altering any proof.

Fourth, a caution the first successful compilation of this document (S47) exposed. Parts of §20–§22 carry *hand-written* numbers in their prose and subsection headings — “Lemma 20.1”, “Proposition 20.3”, “Theorem 20.4” — which are off by one from the numbers L<sup>A</sup>T<sub>E</sub>X assigns: those three statements live in the section L<sup>A</sup>T<sub>E</sub>X numbers 21, and Theorem 21.4 prints as Theorem 21.4, not 20.4. The title of §25 and the defect headings inside it use the hand-written scheme. This record uses \ref throughout, so its numbers are the printed ones; when they disagree with a quoted phrase from the older prose, the \ref is right. The drift is pre-existing, is not repaired here, and is noted so that a reader is not sent to the wrong section.

### 44.1 Wall 0 — The S31 audit: the $H^1$ Prize claim of §20–§22

- (a) **Route.** A vorticity scale-recursion (Proposition 21.3) iterated down scales to give the uniform enstrophy cascade bound  $Z(r, z_0) \leq D r^{2/3}$  for all  $u_0 \in H^1(\mathbb{T}^3)$  (Theorem 21.4, line 4407), from which Corollary 21.5 (line 4451) and Corollary 21.6 (line 4467, “Prize for  $u_0 \in H^1$ ”) were drawn.

- (b) **Obstruction.** §25 (line 5218) records *eight* independent defects, not one:
- D1. the induction of Theorem 21.4 requires  $D$  small and its base case requires  $D$  large; the stated choice satisfies neither, so the recursion collapses to the CKN small-energy regime;
  - D2. an invalid Hölder triple ( $\frac{1}{3} + \frac{1}{2} + \frac{1}{2} > 1$ ) in Lemma 20.2 Step 3, which destroys the claimed  $5/3$  superlinearity;
  - D3. a reversed Jensen inequality in Lemma 20.2 Step 5 — the true relation runs the other way, and no  $L_t^\infty$  information is available for Leray–Hopf solutions to repair it;
  - D4. regularity circularity (a local *enstrophy* inequality is assumed where only a local *energy* inequality holds) together with global/local norm mixing;
  - D5. Theorem 22.6 is conditional on  $\int_0^T M dt < \infty$ , which is the Beale–Kato–Majda criterion, so it assumes an equivalent of its conclusion;
  - D6. the two-regime dichotomy needs  $\nu > C_1 C/c$ , a condition on the given physical viscosity that NS scaling cannot arrange;
  - D7. the Regime II conclusion is false:  $\dot{M} = CM^{5/4}$  blows up in finite time;
  - D8. unproved  $|\nabla^2 \omega|^2$ -coercivity in Lemma 22.4, whose stretching estimate also uses  $\|\nabla u\|_{L^\infty} \leq CM$  — Calderón–Zygmund operators are unbounded on  $L^\infty$  and the correct bound carries a logarithm.
- (c) **Classification.** WITHDRAWN. Theorem 21.4, Corollary 21.5 and Corollary 21.6 are withdrawn. Global regularity for general  $u_0 \in H^1(\mathbb{T}^3)$  is **open** in this document.
- (d) **Consequence.** What survives is listed in “What survives the audit” (§25): the Route C bootstrap (Theorem 3.3), Claim A for shear (§11) and near-shear (§12) data,  $\lambda_{\max} = 0$  for shear flows, the orthogonal split (Lemma 22.1) and the exact scale invariance of  $\sigma = A_{\text{loc}}/M^{3/2}$  — the audit’s own list, which names  $\sigma$  and not the later localised  $\sigma^*$  of Proposition 26.4. §20 remains usable only as a *conditional small-data* statement, and only after D2–D4 are repaired and the superlinearity exponent re-derived. Defect D8 is the first appearance of Wall 1 below; it was already flagged in the audit and is the reason the logarithm is not a later regression.

#### 44.2 Wall 1 — The Calderón–Zygmund logarithm

- (a) **Route.** Every enstrophy-budget estimate of §27–§35 controls the vortex stretching by a sup-norm of  $\nabla u$ , recovered from  $\omega$  by Biot–Savart. Closing any of them requires that sup-norm to cost  $O(M)$ .
- (b) **Obstruction.** It costs  $M\mathcal{L}$ . The factor  $\mathcal{L} = \log(e + \|u\|_{H^3}/M)$  enters at Lemma 27.6 (line 5701, definition on line 5712) through  $\|\nabla u\|_{L^\infty} \lesssim M\mathcal{L}$ , which is the  $L^\infty$  endpoint failure of the Biot–Savart operator — a Calderón–Zygmund operator, unbounded on  $L^\infty$ . This is the correction Defect D8 of §25 had already flagged as unavoidable, so the factor is not a later regression but the audit’s finding carried forward explicitly. Every budget downstream inherits it, and it appears at three distinct closure attempts:

- (i) **Crude  $K$ -absorption.** Proposition 28.4 (line 5903): the coupling term re-enters at top order multiplied by  $\mathcal{L}$  — “the crude route misses closing the loop by exactly one logarithm”.
  - (ii) **The annulus core.** Remark 34.20 (line 8395): absorption requires  $C\mathcal{L} \leq \delta\nu$ , which fails for fixed  $\nu$  as  $\mathcal{L} \rightarrow \infty$ ; the shortfall is “exactly the same single logarithm”.
  - (iii) **The corrected band route.** Proposition 35.15 (line 9209): both Conjecture 28.5 and the  $Y$ -half of Conjecture 34.22 close if and only if the correlation constant *decays* like  $C_1/\mathcal{L}$ ; boundedness is not enough, because the hypothesis is used precisely to cancel the  $\mathcal{L}$  carried by the budget (159).
  - (iv) **Saturation of the Type-I bound.** *Added at S50.* Theorem 37.21: the statement  $M(t) \geq c(T-t)^{-1}$  near a putative Type-I first singularity is *equivalent*, with reciprocal constants, to  $\|\nabla u\|_{L^\infty} \leq KM$  — the bound item (a) above requires. It is a fourth face of this wall, not a route around it.
- (c) **Classification.** OPEN, and structural. No proof in this document says the logarithm cannot be removed; what the items above establish is that it is not a bookkeeping artefact of any one estimate, since it is paid once, at the point where a sup-norm of a Calderón–Zygmund image is taken, and then inherited. *Amended at S50:* item (iv) widens the scope of the entry without changing its classification. Theorem 37.21 is an equivalence, not an impossibility proof: it says the saturation route and the Calderón–Zygmund route have the same content, and says nothing about whether either can be closed. A proof that Wall 1 *cannot* be removed would be a new result and is not in this document.
- (d) **Consequence.** *Forbidden:* presenting any of (i)–(iv) as a separate obstacle to be attacked separately; they are one obstacle seen four times. *Forbidden (added at S50):* presenting logarithm-free saturation of the Type-I vorticity bound as a cheaper, sharper or better-defined target than this wall, as Remark 36.21 proposed; by Theorem 37.21 it is the same target, and by Corollary 37.22 the logarithm-carrying form that *is* provable (Theorem 37.19) returns only the estimate the document already had. *Forbidden:* closing any of them with a *boundedness* hypothesis on a correlation constant — by Proposition 35.15 the required hypothesis is decay. *Permitted:* any reformulation in which  $\nabla u$  is *not* recovered by an  $L^\infty$  Calderón–Zygmund bound. Two such are on record: the profile reformulation of §33, whose status is Wall 2 — WITHDRAWN at S48, the reformulation having been shown in §36 never to incur the logarithm at all; and a symmetry-reduced setting in which  $\nabla u$  is recovered by lower-dimensional elliptic problems, which is untested here and must be settled by a written estimate before anything is built on it.

#### 44.3 Wall 2 — The profile route was thought to relocate Wall 1 into hypothesis (R); it does not

*Revised at S48. This entry was written at S47 and classified OPEN; §36 settled (R) against the literature and the entry is now WITHDRAWN. The S47 text is retained in (a)–(b) because the record is the point of this section; (c) and (d) are rewritten.*

- (a) **Route.** Type-I blowup-profile compactness (§33): pass to a rescaled limit  $U$  and work on the profile, where Proposition 33.8(d) (line 6941) states that “on the profile the



logarithmic factor  $\mathcal{L}$  may be replaced by an absolute constant”. If that were unconditional, Wall 1 would be removed for every estimate transferred to  $U$ .

- (b) **Obstruction, as recorded at S47.** It was held not to be unconditional. Proposition 33.8(d) was labelled *Proved-modulo*-(R), and its proof states the mechanism plainly: “the CZ step is replaced by the interior estimate (R), which is logarithm-free”. Hypothesis (R) (line 6785) *is* an assumed log-free gradient bound — scale-uniform interior regularity  $\|\nabla^k u_\lambda\|_{L^\infty} \leq C_k(c_I)(1+R)S_1^{-(k+1)/2}$  — carried in the S39 ledger as **Conditional** (Serrin-type). On that reading the barrier was relocated, not removed.
- (c) **Classification.** WITHDRAWN. The assertion in (b) that the barrier is “relocated, not removed” is retracted. Theorem 36.16 proves (R), and Corollary 36.17 proves Proposition 33.8(d) unconditionally. The reason the S47 reading was wrong is recorded in Remark 36.2: the hypothesis actually available on the rescaled family is an  $L_t^\infty L_x^\infty$  *velocity* bound (line 5565 assumes it outright, alongside the vorticity bound), and the pair  $s' = s = \infty$  satisfies  $2/s' + 3/s = 0 < 1$  — the *strict*, subcritical range of Serrin’s criterion [7, 24], where boundedness suffices and no smallness is required. The S47 entry looked for (R) on the  $\varepsilon$ -regularity shelf ([1, 19, 18]), where the smallness-versus-boundedness objection is real; the statement is not on that shelf. Its quantitative, scale-invariant form comes from the  $L^\infty$ -based local theory of [17, §4] and [22, Thm. 2.4], neither of which uses a local energy, a pressure norm, or a smallness hypothesis. Remark 36.18 corrects the mechanism stated inside Proposition 33.8(d): the logarithm is not *removed* on the profile, it is never *incurred*, because the profile route does not pass through the Calderón–Zygmund operator.
- (d) **Consequence.** *Permitted, and now correct:* describing the profile route as logarithm-free without carrying (R). *Forbidden:* reading this withdrawal as removing Wall 1. Wall 1 concerns  $u$  on  $\mathbb{T}^3$  measured against  $M(t)$ , and stands; the closest the present result comes to it is Observation 36.19, which is recorded there as an open item and used nowhere. *Forbidden:* reading it as progress on the Prize. Wall 3 is unaffected: the ceiling of §33 is still “no Type-I singularity”, and (N), (N’), (A-up) and Conjecture 33.25 are all exactly where S47 left them. *The one live successor:* the saturation proviso inside Proposition 33.8(d), which Proposition 33.8(e) uses and which (R) never covered — residue (1) of §36.5. *Also forbidden, permanently:* attempting to weaken the standing hypotheses of §27 to a vorticity-only Type-I bound. Proposition 36.12 shows the velocity rate cannot be recovered from  $\|\omega\|_{L^\infty}$  and the energy: the interpolation exponent is forced to 3/5, not 1/2.

#### 44.4 Wall 3 — The ceiling of the S32–S45 line is below the Prize

- (a) **Route.** The alignment pincer of §26–§33, closed at both horns.
- (b) **Obstruction.** Remark 33.30 (line 7670) states the ceiling: if (A-up) and Conjecture 33.25 were proved and (R), (N), (N’) discharged, the conclusion would be *no Type-I singularity* — “strictly less than the Clay problem”. The reason is not a gap but the construction: every estimate in §27–§33 uses the Type-I rates, and the profile extraction uses them *essentially* — without  $M(t) \lesssim (T-t)^{-1}$  the rescaled family has

no uniform bound and there is no limit object. “A Type-II singularity would rescale to nothing, and the entire reframe would be vacuous against it.” The same remark adds two further reservations: **(N)** restricts the conclusion to non-degenerate points, and Conjecture 33.25 must beat the constant 1 exactly, which is the marginal-target pattern the S31 audit identified as where errors hide.

- (c) **Classification.** OPEN, and outside the present methods. Type-II exclusion is carried in the S39 ledger as “Open (outside all present methods)”.
- (d) **Consequence.** *Forbidden:* reading any part of §27–§35 as progress on Type-II; the document says so in terms. *Forbidden:* reporting a closed pincer as the Prize. *Permitted:* closing the pincer as a Type-I exclusion result, provided it is stated as such — and, in a symmetry class where Type-I is already excluded in the literature, only if the prior exclusion is cited rather than re-derived.

#### 44.5 Wall 4 — Level iteration for the annulus residual

- (a) **Route.** Remark 30.6 proposed absorbing the annulus residual by a nested family of level cutoffs  $\chi_k$  with Young parameters  $\delta_k = \delta_0 8^{-k}$ , on the “geometric beats geometric” principle.
- (b) **Obstruction.** Three propositions in §34, each independent of the others:
  - (i) Proposition 34.12 (line 8156):  $\Lambda_k \geq 2^8 4^k \geq 2^8 > 1$  for *every* sequence  $(\delta_k)$  — the recursion never contracts. The proposed  $\delta_k = \delta_0 8^{-k}$  makes it strictly worse ( $32^k$ ); Young’s inequality has no purchase on the  $2^8 4^k$  term because that term does not involve  $\delta_k$ ; and the summability  $\sum_k \delta_k \leq \frac{1}{2}$  that absorption independently requires forces the two geometric factors onto the same side, where they reinforce.
  - (ii) Proposition 34.13 (line 8191): the unrolled inequality is implied by  $Y_0 \leq Y_0$  and carries no information for every  $N$ ; letting  $N \rightarrow \infty$  does not help.
  - (iii) Proposition 34.16 (line 8270): by a co-area identity the residual produced by *any* admissible cutoff is a level density of the measure  $\mu_F$ , bounded above only by  $(\theta_+ - \theta_-)^{-1} \leq 16$  times a total mass and bounded below not at all. No cleverer choice of levels exists to be found.
- (c) **Classification.** PROVED IMPOSSIBLE for the route, and SUPERSEDED as a target: Conjecture 34.22 (line 8435) is stated as “the replacement for Remark 30.6” and is itself OPEN.
- (d) **Consequence.** *Forbidden:* any further attempt to close the annulus residual by choosing levels, cutoffs or Young parameters. Remark 34.21 states what would actually have to be proved, and stresses that the absorbed term must carry *the same* cutoff  $\eta$  as the left-hand side — “this, not the value of any constant, is what defeats the nested scheme” — and that by Lemma 34.2 one admissible level pair suffices, so the difficulty is analytic and not combinatorial. This wall is what reclassified the annulus residual  $\mathcal{A}_{\text{ann}}$  — H1’s last gap — from *open-mechanical* to *open*, at the close of §34 and in the S44 ledger of §35.

## 44.6 Wall 5 — $c_K$ and band absorption are not the same hypothesis

- (a) **Route.** Remarks 34.24 and 34.23 described Conjecture 28.5 (the  $c_K$  decorrelation hypothesis, line 5917) and Conjecture 34.22 (line 8435) as “of the same analytic type”, with the hope that proving one would deliver the other, or that the apparently absolute constants of the band statement made it the weaker target.
- (b) **Obstruction.** §35 settles both hopes negatively.
  - (i) *Disjoint supports.* Remark 35.4 (line 8738):  $E = Q(r^*) \cap \{m \geq M/4\}$  and  $\text{Band} \subseteq \{m \leq M/4\}$  meet only in a null level set; neither hypothesis constrains any integral supported in the other’s region.
  - (ii) *Counterexamples both ways.* Proposition 35.8 (line 8846): a family with both band left-hand sides vanishing identically — so Conjecture 34.22 holds with  $C = 0$  — while  $c_K \geq |E|/(8\ell^5) \rightarrow \infty$ . Proposition 35.10 (line 8947): a family with  $c_K = 0$ , the strongest possible form of Conjecture 28.5, for which the constant required in (168) diverges.
  - (iii) *The strength claim was inverted.* Remark 35.3 (line 8708) records that the first draft of §35 stated Lemma 35.2 with the reciprocal volume factor. Three conclusions drawn from it — that  $\Xi$ -boundedness is weaker than  $c_K$ -boundedness, that Conjecture 28.5 could be weakened to an  $O(1)$  statement without loss, and that the measured  $c_K \approx 1$  was direct evidence for  $\Xi = O(1)$  — are **withdrawn**. Corrected,  $\Xi[\mathbf{1}_E] = c_K |Q(r^*)|/|E| \geq c_K$ : the volume-normalised statistic is *stronger*, not weaker.
- (c) **Classification.** PROVED IMPOSSIBLE for the implication, in a precisely bounded sense, and WITHDRAWN for the strength claim. The bound on the sense matters and is stated in the document, not added here: by Remark 35.7 (line 8822) an admissible test configuration (Definition 35.6) need not come from a solution of Navier–Stokes, so Propositions 35.8 and 35.10 do *not* establish that the two conjectures are independent as statements about Navier–Stokes solutions — “that question is not decidable by any argument available here”. What is proved is that no implication can be derived from the a priori budgets and pointwise structure (A1)–(A3) in terms of which both are phrased.
- (d) **Consequence.** *Forbidden:* treating a proof of either conjecture as progress on the other; *forbidden:* the reading of “same analytic type” as “one covers the other”. *Permitted:* Conjecture 35.20 (line 9412), the single hypothesis (LC) that covers both  $Y$ -halves (Proposition 35.21). Note its exact position, which is easy to misstate: (LC) is *verbatim* the  $c_K$  hypothesis — same normalisation, same  $1/\mathcal{L}$  decay — asserted on the *larger* region  $\{m \geq M/16\}$ ; by Remark 35.22 it implies neither Conjecture 28.5 nor conversely, and is “a genuinely new hypothesis of an old form”. It is weaker than the alternative common hypothesis (JLE-1, Conjecture 35.23) in the regime that matters, not weaker than Conjecture 28.5. Neither the angular half (169) (Remark 35.26) nor the constant condition  $CC_1 \leq \delta\nu$  (Remark 35.16) is covered by any of them.

## 44.7 Wall 6 — The $c_K$ numerics cannot discriminate

- (a) **Route.** Measure  $c_K$  in direct numerical simulation and use the result as evidence for or against Conjecture 28.5. Four sessions did so (S34, S35, S42–43, S45).
- (b) **Obstruction.** The measurement is uninformative for a logical reason, independent of what any run reports.
  - (i) Conjecture 28.5 and Conjecture 35.20 are conditional statements: each asserts a bound *near a putative Type-I singularity*, in the limit  $\mathcal{L} \rightarrow \infty$ . A solver that remains globally regular is never in that regime, so neither the conjecture nor its negation predicts anything about the numbers such a solver produces. Both predict the same thing — nothing — and a measurement that both sides of a disjunction predict cannot separate them.
  - (ii) Proposition 32.1 (line 6514) removes the one reading under which a regular run could still have served as a proxy. It proves that at a genuine blowup time  $\limsup_{t \rightarrow T^-} \Omega(t) = \infty$ ; hence bounded-ensrophy escape (Regime A) is impossible, and  $\sigma^*$ -decay is, given H1, *necessary* rather than one route among several. A resolved simulation is a bounded-ensrophy object throughout. So the regime the conjectures speak about is not merely one the runs did not visit; it is not of the type a resolved run represents, and the decay is not something such a run could exhibit.
  - (iii) Independently of (i)–(ii), the in-document assessment Remark 35.19 (line 9363) states that the  $O(1)$  measurements of S34–S35 are “consistent with, but do not establish” either conjecture, that a measured  $c_K \approx 1$  does *not* support  $\Xi = O(1)$  after the correction of Remark 35.3, and that the discrepancy is worst exactly in the thin, filamentary configurations expected near a singularity.
- (c) **Classification.** PROVED IMPOSSIBLE as a test of the conjectures. The ground is (i)–(ii): an observation about the *form* of the statements — each is conditional on a regime the solver cannot enter, and Proposition 32.1 is what ties that regime to genuine blowup. No new mathematics is required for it, and it does not depend on resolution, viscosity or initial condition. The diagnostic suite itself is OPEN and valid for other purposes.
- (d) **Consequence.** *Forbidden:* measuring  $c_K$ ,  $c_K^{\text{band}}$  or the band ratios again *as evidence* for or against Conjecture 28.5, Conjecture 34.22 or Conjecture 35.20; they may be computed as context. *Permitted:* measurements whose answer differs between competing mechanisms *in a regular flow* — questions about a mechanism, not proxies for blowup. Also permitted, and still outstanding, are the three quantities Remark 35.19(iii) names as not reported in the form needed: the volume fractions  $|E|/|Q(r^*)|$  and  $|\text{Band}|/|Q(r^*)|$  at the working scale, the band correlation constant  $c_K^{\text{band}}$ , and the  $\mathcal{L}$ -dependence of both correlation constants.

*Outside this document.* Session S45 measured all three (SESSION-LOG/2026-09-10-S45-scale-diagnostics-jet-sweep.md; EXP-L4-SCALE-\*, EXP-L4-JET\*) and found the  $\mathcal{L}$ -slope of  $\log c_K$  to be  $-0.196$  ( $r^2 = 0.05$ ) on one pool and  $+0.088$  ( $r^2 = 0.02$ ) across all twelve runs, against the  $-1$  a clean  $C/\mathcal{L}$  law would require, over  $\mathcal{L} \in [1.35, 4.68]$ . That is a negative signal for Proposition 35.15’s hypothesis and nothing more: by (i) above it

does not refute the conjecture, and it is recorded here only because it is the measurement Remark 35.19(iii) asked for. No statement in this document rests on it.

#### 44.8 Wall 7 — The S35 second good term does not exist

- (a) **Route.** Proposition 29.2 (line 6026, *provisional* as stated) recorded the identity (149) with *two* good terms on the left: the  $K$ -density  $\nu \iint |\nabla m|^2 w \eta^2$  and, additionally,  $\nu \iint m^2 w^2 \eta^2$  — the critical quadratic with the favourable weight, presented as reinforcing Proposition 29.1.
- (b) **Obstruction.** Lemma 30.1 (line 6166): the  $+\nu w \hat{e}$  term of (142) contributes exactly  $+\nu w^2$  to  $\frac{1}{2} D_t w$ , which cancels the sink term  $-\nu \iint m^2 w^2 \eta^2$  *identically*. The second left-hand term is removed; (149) must be read without it.
- (c) **Classification.** WITHDRAWN, and repaired. The  $K$ -control conclusion stands with amended constants; the critical quadratic is handled solely by Proposition 29.1, which suffices, so no downstream conclusion depended on the cancelled term.
- (d) **Consequence.** *Forbidden:* citing (149) in its original two-term form, or invoking the  $w^2$  term as an independent source of smallness. This is the smallest of the walls and is recorded because it is one of three instances — with Wall 5(iii) and the corrigendum of §28 — of the same failure mode: a favourable statement written down before the computation producing it was closed. Note also Remark 29.4, which states what would follow if couplings C1–C2 verified in full: Conjecture 28.5 would become *unnecessary*. That conditional remains conditional.

#### 44.9 Wall 8 — The localized Constantin–Fefferman machinery does not transfer to profiles

- (a) **Route.** Carry the H2 / Regime-B analysis of §31, whose engine is Proposition 31.2 (line 6381), across the rescaling limit onto the Type-I profile  $U$ , so that the S39 reframe inherits it.
- (b) **Obstruction.** Remark 33.9(i) (line 7031): the far-field step of Proposition 31.2 bounds  $\int_{|y| \geq \rho'} |\omega(x^* - y)| |y|^{-3} dy$  by  $C(\rho')^{-3/2} \Omega^{1/2}$  with  $\Omega = \|\omega\|_{L^2}^2$ , and for  $U \in \mathcal{A}(c_I)$  on  $\mathbb{R}^3$  there is no reason for  $\|\omega_U(\cdot, s)\|_{L^2(\mathbb{R}^3)}$  to be finite; the naive substitute  $|\omega_U| \leq M_U$  makes the integral logarithmically divergent at large  $|y|$ . The document states the consequence in terms: “the entire H2/Regime-B analysis of §31 is unavailable on the profile as written”.
- (c) **Classification.** PROVED IMPOSSIBLE as written — the obstruction is a divergence, not a missing estimate. Whether some localized substitute exists on  $\mathbb{R}^3$  is OPEN and untried.
- (d) **Consequence.** *Forbidden:* “Any future session that uses Proposition 31.2 on a profile is in error” (Remark 33.29(4)). The companion prohibition of Remark 33.9(ii) is equally binding and is easier to violate by accident: the Bridge Lemma (Conjecture 26.8) may *not* be pulled back, because Lemma 33.10 gives semicontinuity in the wrong direction and  $\sigma^*(U) = 0$  does not give  $\sigma^*(u_{\lambda_j}) \rightarrow 0$ . Any argument that routes a profile conclusion

back through H1 on the approximating sequence is invalid; the S39 dichotomy is built to obtain its contradiction entirely on  $U$  for exactly this reason.

#### 44.10 Wall 9 — Forced blowup constructions cannot be de-forced

*This entry is recorded outside the mathematics of this document.* It carries no label here because no part of this document uses it; it is included because it bounds what future sessions may import. The source is `SESSION-LOG/2026-09-09-S40-openai-comparison-aniso-collapse.md`.

- (a) **Route.** Import the mechanism of a recently published finite-time blowup construction for the *forced* Navier–Stokes equations — an axisymmetric anisotropic self-similar collapsing vortex core, with  $\ell_r \asymp \tau^{1/2}$ ,  $\ell_z \asymp \tau^{1/2-h}$  and  $\text{Re}_\theta \rightarrow \infty$  while  $\text{Re}_r = O(1)$  — by deleting the forcing term, and read the result against the Prize equations.
- (b) **Obstruction.** The forcing is load-bearing at three separate points of that construction, not decorative: an exponentially small external force seeds each oscillatory pulse; what remains of the momentum residual after pulse cancellation *becomes* part of  $f$ ; and the compact-support cutoff introduces a further force term. The forcing is *defined* as the Navier–Stokes residual of the constructed field. Setting  $f \equiv 0$  is therefore not a simplification of the construction but a different question, which the source does not claim to answer. In Fefferman’s formulation [3] the result establishes alternatives (C) and (D), not (A)/(B).
- (c) **Classification.** PROVED IMPOSSIBLE as an import route — there is nothing to de-force, since the residual is the forcing by construction. The underlying blowup question for the unforced equations is of course OPEN; that is the Prize.
- (d) **Consequence.** *Forbidden:* forcing terms, modified equations and “weak internal forces” as mechanisms anywhere in this programme; *forbidden:* reading a forced blowup result as evidence about the unforced equations in either direction. *Permitted:* the *geometry* of such constructions as a source of adversarial initial data for the unforced solver, which is how `layer3/aniso_collapse_IC.py` was built (fed to an unmodified  $f \equiv 0$  solver). The one substantive observation, recorded as an observation and not as evidence: sustaining that anisotropic collapse against dissipation required an externally injected, precisely tuned force.
- (e) **Quantitative addendum (S55).** Item (b)’s “load-bearing, not decorative” now carries a number, proved in §41: for the *axisymmetric* leading field and background of that construction, the swirl maximum principle alone forces the azimuthal momentum residual to satisfy  $|rf_\theta| \gtrsim \tau^{-(A+1/2)}$ , hence  $|f_\theta| \gtrsim \tau^{-(A+1)} = \tau^{-(3/2+h)}$ , pointwise at the running swirl maximum (Proposition 41.5) — full leading order in its own momentum equation. The source’s stated stress scaling  $T = q^{-A-1/2}T_0$  meets this floor with the same exponents (Remark 41.10). The addendum sharpens the wording of this entry and changes neither its classification nor its consequences, and it asserts nothing about the correctness of that construction’s Theorem 1.1.

#### 44.11 Status table

| #     | Route                               | Classification                                                                                                   | Primary reference      |
|-------|-------------------------------------|------------------------------------------------------------------------------------------------------------------|------------------------|
| 0     | §20–§22 $H^1$ Prize claim           | <b>Withdrawn</b> (8 grounds)                                                                                     | §25                    |
| 1     | Sup-norm of a CZ image; crude $K$   | <b>Open</b> (structural)                                                                                         | Prop. 28.4             |
| 1'    | same log, annulus core              | <b>Open</b>                                                                                                      | Rem. 34.20             |
| 1''   | same log, corrected band            | <b>Open</b> ; needs <i>decay</i>                                                                                 | Prop. 35.15            |
| 2     | Profile route as log-free           | <b>Withdrawn</b> at S48; (R) proved                                                                              | Thm. 36.16, Cor. 36.17 |
| 3     | Pincer closed $\Rightarrow$ Prize   | <b>Open</b> ; ceiling is Type-I only                                                                             | Rem. 33.30             |
| 3'    | rate-adapted rescaling as an escape | <b>Proved impossible</b> for these tools: $\nu_{\text{eff}} = \nu A/L \rightarrow 0$ , so the limit solves Euler | Cor. 39.11             |
| 3''   | an Euler zoom that keeps the swirl  | <b>Proved impossible</b> ( $\Gamma$ and $\nu$ share a dimension)                                                 | Cor. 39.13             |
| 3'''  | Type-II itself                      | <b>Open</b> ; owned by Liouville theory for ancient Euler                                                        | Rem. 39.16             |
| 4     | Nested level iteration              | <b>Proved impossible</b>                                                                                         | Prop. 34.12            |
| 4'    | unrolled form                       | <b>Proved impossible</b> (vacuous)                                                                               | Prop. 34.13            |
| 4''   | any level selection                 | <b>Proved impossible</b> (co-area)                                                                               | Prop. 34.16            |
| 4'''  | replacement target                  | <b>Superseded; Open</b>                                                                                          | Conj. 34.22            |
| 5     | $c_K \Rightarrow$ band absorption   | <b>Proved impossible</b> over (A1)–(A3)                                                                          | Prop. 35.10            |
| 5'    | band absorption $\Rightarrow c_K$   | <b>Proved impossible</b> over (A1)–(A3)                                                                          | Prop. 35.8             |
| 5''   | scope of 5, 5'                      | Test configurations, <i>not</i> NS solutions                                                                     | Rem. 35.7              |
| 5'''  | band is the weaker target           | <b>Withdrawn</b> (lemma inverted)                                                                                | Rem. 35.3              |
| 5'''' | common hypothesis (LC)              | <b>Open</b> ; covers both $Y$ -halves                                                                            | Conj. 35.20            |
| 6     | $c_K$ numerics as evidence          | <b>Proved impossible</b> (logical)                                                                               | Prop. 32.1             |
| 6'    | in-document assessment              | Consistent with, does not establish                                                                              | Rem. 35.19             |
| 7     | S35 second good term                | <b>Withdrawn</b> (exact cancellation)                                                                            | Lem. 30.1              |
| 8     | Localized CF on profiles            | <b>Proved impossible</b> ( $\Omega = \infty$ )                                                                   | Rem. 33.9(i)           |
| 8'    | Pull-back of H1 to sequence         | <b>Forbidden</b> (no reverse inequality)                                                                         | Rem. 33.9(ii)          |
| 9     | De-forcing a forced blowup          | <b>Proved impossible</b> as an import                                                                            | S40 log (outside doc)  |
| 9'    | how much forcing, quantitatively    | <b>Floor proved</b> : $ f_\theta  \gtrsim \tau^{-(A+1)}$ ; source's stated exponent agrees                       | Prop. 41.5             |
| 10    | Axisymmetric-with-swirl pivot (S51) | Wall 1 <b>survives and is sharp</b> in this class                                                                | Prop. 38.9             |
| 10'   | its Type-I branch                   | <b>Closed by others</b> (Seregin–Šverák 2009)                                                                    | Fact 38.24             |
| 10''  | its Type-II branch                  | <b>Open</b> ; outside all present methods                                                                        | Rem. 38.32             |

**Remark 44.1** (Two corrections to the S46 handover, recorded here). The execution plan `HANDOVER-S46-OPUS.md` that commissioned this section states two things that the document does not support, and the document governs.

- (i) It refers to “the S31 audit (all seven grounds)”. §25 lists **eight** defects, D1–D8. D8

(unproved coercivity in Lemma 22.4, together with the  $L^\infty$  Calderón–Zygmund bound) is the one most easily dropped from a summary and is also the first appearance of Wall 1, so the omission is not neutral. Wall 0 above lists all eight.

- (ii) It describes Conjecture 35.20 as “a common *weaker* target”. By its own statement (LC) is the  $c_K$  hypothesis asserted on the *larger* region  $\{m \geq M/16\}$ , with the same normalisation and the same  $1/\mathcal{L}$  decay; by Remark 35.22 it neither implies nor is implied by Conjecture 28.5. It is weaker than the *other* common hypothesis (JLE-1, Conjecture 35.23) in the regime that matters — which is what §35.6 says — not weaker than Conjecture 28.5.

A third statement of the handover is correct but is routinely rounded off in summary: the S44 non-implications are proved over the a priori structure (A1)–(A3) of Definition 35.6, *not* for Navier–Stokes solutions (Remark 35.7). “Proved independent” without that qualifier overstates them.

**Remark 44.2** (What this section does not do). It does not assess whether any wall can be broken, and it contains no encouragement to try. Walls 1, 2, 3 and 5''' are open questions with no proof against them; Walls 4, 6, 8 and 9 carry proofs, and the proofs are in the propositions cited, not here. Nothing in this section bears on H2, on  $\sigma^*$ -decay, on hypothesis **(R)**, on gap **(A-up)**, on the withdrawn material of §20–§22, or on any Prize or Claim A conclusion, all of which retain the status they had before it was written.

## References

- [1] L. Caffarelli, R. Kohn, L. Nirenberg, Partial regularity of suitable weak solutions of the Navier–Stokes equations, *Comm. Pure Appl. Math.*, 35(6):771–831, 1982.
- [2] Ph. Bénéilan, H. Brezis, M. G. Crandall, A semilinear equation in  $L^1(\mathbb{R}^N)$ , *Ann. Scuola Norm. Sup. Pisa*, 2(4):523–555, 1975.
- [3] C. L. Fefferman, *Existence and Smoothness of the Navier–Stokes Equation*, Clay Mathematics Institute, 2000.
- [4] M. Giaquinta, E. Giusti, On the regularity of the minima of variational integrals, *Acta Math.*, 148:31–46, 1982.
- [5] F. Lin, A new proof of the Caffarelli–Kohn–Nirenberg theorem, *Comm. Pure Appl. Math.*, 51(3):241–257, 1998.
- [6] G. Prodi, Un teorema di unicità per le equazioni di Navier–Stokes, *Ann. Mat. Pura Appl.*, 48:173–182, 1959.
- [7] J. Serrin, On the interior regularity of weak solutions of the Navier–Stokes equations, *Arch. Rat. Mech. Anal.*, 9:187–195, 1962.
- [8] E. M. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton Univ. Press, 1970.



- [9] E. W. Stredulinsky, Higher integrability from reverse Hölder inequalities, *Indiana Univ. Math. J.*, 29:407–413, 1980.
- [10] N. V. Krylov, *Lectures on Elliptic and Parabolic Equations in Hölder Spaces*, Graduate Studies in Mathematics, Amer. Math. Soc., 1996.
- [11] Ph. Bénilan, H. Brézis, M. G. Crandall, A semilinear equation in  $L^1(\mathbb{R}^N)$ , *Ann. Sc. Norm. Sup. Pisa*, 2(4):523–555, 1975.
- [12] P. Constantin, C. Fefferman, Direction of vorticity and the problem of global regularity for the Navier–Stokes equations, *Indiana Univ. Math. J.*, 42(3):775–789, 1993.
- [13] P. Constantin, Geometric Statistics in Turbulence, *SIAM Review*, 36(1):73–98, 1994.
- [14] H. Kozono, Y. Taniuchi, Bilinear estimates in BMO and the Navier–Stokes equations, *Math. Z.*, 235(1):173–194, 2000.
- [15] J. Nečas, M. Růžička, V. Šverák, On Leray’s self-similar solutions of the Navier-Stokes equations, *Acta Math.*, 176:283–294, 1996.
- [16] L. Escauriaza, G. Seregin, V. Šverák,  $L_{3,\infty}$ -solutions of Navier-Stokes equations and backward uniqueness, *Uspekhi Mat. Nauk*, 58:3–44, 2003.
- [17] G. Koch, N. Nadirashvili, G. Seregin, V. Šverák, Liouville theorems for the Navier–Stokes equations and applications, *Acta Math.*, 203(1):83–105, 2009.
- [18] G. Seregin, A certain necessary condition of potential blow up for Navier–Stokes equations, *Comm. Math. Phys.*, 312(3):833–845, 2012.
- [19] O. A. Ladyzhenskaya, G. A. Seregin, On partial regularity of suitable weak solutions to the three-dimensional Navier–Stokes equations, *J. Math. Fluid Mech.*, 1(4):356–387, 1999.
- [20] M. Struwe, On partial regularity results for the Navier–Stokes equations, *Comm. Pure Appl. Math.*, 41(4):437–458, 1988.
- [21] S. Takahashi, On interior regularity criteria for weak solutions of the Navier–Stokes equations, *Manuscripta Math.*, 69(3):237–254, 1990.
- [22] G. Seregin, V. Šverák, On type I singularities of the local axi-symmetric solutions of the Navier–Stokes equations, *Comm. Partial Differential Equations*, 34(2):171–201, 2009. (arXiv:0804.1803; Definition 2.3, Theorem 2.4, Theorem 2.8, Remark 2.9 are the statements used in §36.)
- [23] D. Albritton, T. Barker, On local type I singularities of the Navier–Stokes equations and Liouville theorems, *J. Math. Fluid Mech.*, 21(3):Art. 43, 2019. (arXiv:1811.00502.)
- [24] R. M. Chen, G. P. Galdi, B. Poggi, A. Schikorra, On Serrin interior regularity criterion for Navier–Stokes equations, arXiv:2606.24733, 2026. (Quoted in §36 for the exact statement of Serrin’s criterion and its attribution.)

- [25] H. Kwon, The role of the pressure in the regularity theory for the Navier–Stokes equations, arXiv:2104.03160, 2021.
- [26] E. Giusti, *Direct Methods in the Calculus of Variations*, World Scientific, 2003. (Iteration Lemma 6.1; hole-filling.)
- [27] O. A. Ladyzhenskaya, N. N. Ural'tseva, *Linear and Quasilinear Elliptic Equations*, Academic Press, 1968. (Chapter II: De Giorgi level truncation.)
- [28] J. T. Beale, T. Kato, A. Majda, Remarks on the breakdown of smooth solutions for the 3-D Euler equations, *Comm. Math. Phys.*, 94(1):61–66, 1984. (The Theorem on p. 62 and inequality (15) on p. 64 are the statements quoted in §37.3. Inequality (15) is the source of the factor  $\mathcal{L}$  of Lemma 27.6, which had carried no citation before S50. The paper states, p. 62: “Similar statements hold for the Navier–Stokes equations; the arguments below apply equally well in this case.”)
- [29] J. Leray, Sur le mouvement d’un liquide visqueux emplissant l’espace, *Acta Math.*, 63(1):193–248, 1934. (Origin of the  $L^p$  lower blow-up rates, stated there without proof; see [32, §1].)
- [30] Y. Giga, Solutions for semilinear parabolic equations in  $L^p$  and regularity of weak solutions of the Navier–Stokes system, *J. Differential Equations*, 62(2):186–212, 1986. (Proof, by semigroup theory, of the rates Leray stated.)
- [31] J. C. Robinson, W. Sadowski, R. P. Silva, Lower bounds on blow up solutions of the three-dimensional Navier–Stokes equations in homogeneous Sobolev spaces, *J. Math. Phys.*, 53(11):115618, 15 pp., 2012.
- [32] J. C. Cortissoz, J. A. Montero, Lower bounds for possible singular solutions for the Navier–Stokes and Euler equations revisited, *J. Math. Fluid Mech.*, 2016; arXiv:1503.03063v2. (Theorem 2.1 is quoted verbatim in Fact 37.9; §1 is quoted for the classical Leray–Giga rates.)
- [33] B. Ingimarnson, I. Kukavica, Lower bounds on the blowup rate of vorticity in the Euler equations, arXiv:2603.17431v2, 2026. (Theorems 2.1 and 2.5 and the proof of Theorem 2.1 are quoted in Fact 37.12. This is the state of the art for a lower bound on  $\|\omega\|_{L^\infty}$ , and it is a lim sup carrying a logarithm.)
- [34] A. Cheskidov, M. Dai, S. Palasek, Instantaneous Type I blow-up and non-uniqueness of smooth solutions of the Navier–Stokes equations, arXiv:2511.09556v2, 2026. (§2.2 is quoted in Remark 37.2 for the definition of Type I as saturation of the velocity lower bound; Theorem 1.1, Properties 2 and 4, in Fact 37.13.)
- [35] O. A. Ladyzhenskaya, Unique global solvability of the three-dimensional Cauchy problem for the Navier–Stokes equations in the presence of axial symmetry, *Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)*, 7:155–177, 1968 (Russian).
- [36] M. R. Ukhovskii, V. I. Yudovich, Axially symmetric flows of ideal and viscous fluids filling the whole space, *J. Appl. Math. Mech.*, 32:52–61, 1968.

- [37] C.-C. Chen, R. M. Strain, T.-P. Tsai, H.-T. Yau, Lower bound on the blow-up rate of the axisymmetric Navier–Stokes equations, *Int. Math. Res. Not.*, 2008, Art. ID rnn016, 31 pp. (Theorem 1.1 is quoted verbatim in Fact 38.22. Part II: *Comm. Partial Differential Equations*, 34(1–3):203–232, 2009.)
- [38] H. Chen, D. Fang, T. Zhang, Regularity of 3D axisymmetric Navier–Stokes equations, *Discrete Contin. Dyn. Syst.*, 37(4):1923–1939, 2017. (Source of the scalar  $J = \omega_r/r$  and of the criterion  $|u_\theta| \leq Cr^{-1+\varepsilon}$  quoted in Fact 38.28(a).)
- [39] Z. Lei, Q. S. Zhang, Criticality of the axially symmetric Navier–Stokes equations, *Pacific J. Math.*, 289(1):169–187, 2017; arXiv:1505.02628. (The Form Boundedness Condition, the main theorem and the  $|\ln r|^{-2}$  corollary are quoted in Fact 38.28(b).)
- [40] D. Wei, Regularity criterion to the axially symmetric Navier–Stokes equations, *J. Math. Anal. Appl.*, 435(1):402–413, 2016; arXiv:1508.03318. (The  $|\ln r|^{-3/2}$  improvement, quoted in Fact 38.28(c). This is the state of the art for a swirl-only criterion.)
- [41] Q. S. Zhang, A review of results on axially symmetric Navier–Stokes equations, with addendum by X. Pan and Q. S. Zhang, arXiv:2101.04905, 2021. (§2.2 is quoted in Fact 38.27 for the  $(J, \Omega, \omega_z)$  system and the criticality observation.)
- [42] J. Neustupa, M. Pokorný, An interior regularity criterion for an axially symmetric suitable weak solution to the Navier–Stokes equations, *J. Math. Fluid Mech.*, 2(4):381–399, 2000.
- [43] D. Chae, J. Lee, On the regularity of the axisymmetric solutions of the Navier–Stokes equations, *Math. Z.*, 239(4):645–671, 2002.
- [44] I. Kukavica, M. Ziane, One component regularity for the Navier–Stokes equations, *Nonlinearity*, 19(2):453–469, 2006.
- [45] C. Cao, E. S. Titi, Global regularity criterion for the 3D Navier–Stokes equations involving one entry of the velocity gradient tensor, *Arch. Ration. Mech. Anal.*, 202(3):919–932, 2011.
- [46] G. Seregin, Remarks on Type II blowups of solutions to the Navier–Stokes equations, *Commun. Pure Appl. Anal.*, 23(10):1389–1406, 2024; arXiv:2304.04045v1, 8 April 2023.
- [47] G. Seregin, A note on potential Type II blowups of axisymmetric solutions to the Navier–Stokes equations, arXiv:2402.13229v3, 8 August 2026.
- [48] G. Seregin, A note on certain scenarios of Type II blowups of suitable weak solutions to the Navier–Stokes equations, arXiv:2507.08733, submitted 11 July 2025, revised 3 January 2026.
- [49] G. Seregin, On potential Type II blowups for the Navier–Stokes equations, arXiv:2606.29468v1, 28 June 2026.
- [50] G. Seregin, W. Zajączkowski, A sufficient condition of local regularity for the Navier–Stokes equations, *Zap. Nauchn. Sem. POMI*, 336:46–54, 2006.

- [51] G. Seregin, V. Šverák, Regularity criteria for Navier–Stokes solutions, in Y. Giga, A. Novotný (eds.), *Handbook of Mathematical Analysis in Mechanics of Viscous Fluids*, Springer, Cham, pp. 829–867.
- [52] T. Tao, Quantitative bounds for critically bounded solutions to the Navier–Stokes equations, arXiv:1908.04958v2; in *Nine Mathematical Challenges* (A. Kechris et al., eds.), Proc. Sympos. Pure Math.
- [53] T. Barker, C. Prange, From concentration to quantitative regularity: a short survey of recent developments for the Navier–Stokes equations, *Vietnam J. Math.*, 2023; arXiv:2211.16215.
- [54] T. Y. Hou, Potentially singular behavior of the 3D Navier–Stokes equations, *Found. Comput. Math.*, 2022; arXiv:2107.06509v2 (“The potentially singular behavior of the 3D Navier–Stokes equations”), 26 May 2022.
- [55] OpenAI, Finite time blowup for Navier–Stokes, manuscript, 2026; <https://cdn.openai.com/pdf/32d9f210-8b73-45e0-91bc-82a30aef8a9a/navier-stokes.pdf>. [*S53 read Theorem 1.1 and §2. Re-fetched and read further at WP-T2 (§40): §3.1, §4.1–4.2 including the similarity variables (4.1), the leading ansatz (4.3), the profile identities (4.7) and the profile equations (4.13). A **forced** construction:  $f \in C_c^\infty$ , and the full flow is not axisymmetric — the oscillatory pulses carry nonzero azimuthal harmonics. Quoted throughout as a profile, under Wall 9 of §44, not as evidence about the unforced equations. Re-fetched and read again at WP-T3 (§41) for the quantitative forcing statements: §2 and §2.2 (the force is the residual; the background’s residual diverges), (3.1), Theorem 3.1(iii) and its (3.4)–(3.5), (4.11)–(4.14), Proposition 4.2, Theorem 4.6(ii)/(iv)/(vi) and its (4.27), (4.30), Proposition 5.5 with its (5.41)–(5.44), and Lemma 4.9 with its (4.32); all quotations in Fact 41.9 are verbatim.*]
- [56] Y. Giga, R. V. Kohn, Characterizing blowup using similarity variables, *Indiana Univ. Math. J.*, 36(1):1–40, 1987.
- [57] D. Gérard-Varet, E. Dormy, On the ill-posedness of the Prandtl equation, *J. Amer. Math. Soc.*, 23:591–609, 2010; arXiv:0904.0434. [*Abstract fetched and quoted at §40: “We prove here that it is linearly ill-posed in Sobolev type spaces ... the strong instability is due to viscosity”. Cited only for the structural caution of Remark 40.7; **not** applied to the system derived there.*]
- [58] E. W. Mayer, Long’s vortex revisited, arXiv:1303.1212, 5 March 2013. [*Read at §40 for one statement only: that the inner viscous core of a slender vortex is treated by Hall (1961) and Stewartson–Hall (1963) “by making scaling assumptions of the usual boundary-layer type”.*]
- [59] C. Collot, F. Merle, P. Raphaël, On strongly anisotropic type II blow up, arXiv:1709.04941, 14 September 2017.
- [60] Z. Grujić, Logarithmic depletion of vortex stretching and singularity evasion in the 3D Navier–Stokes equations, arXiv:2607.08866; v1 9 July 2026, v3 9 September 2026. [*Unrefereed preprint, no journal reference. Read in full from the primary source at*

*S57 (§42): Definition 2.1, the  $\text{bmo}_\phi$  definition (2), Remarks 2.2–2.3, the commutator identity (7) of §3, Theorem 4.1 and its complete proof, §5.1–5.3 including (23)–(40), §6 including (41)–(49), and §7 including Definition 7.1, Theorem 7.3, Theorem 7.4 and its proof. Quoted as a claim only; **no statement in this document depends on it** (Remark 42.1). The commutator/Coifman–Rochberg–Weiss step is independently verified in Remark 42.15.]*

- [61] Z. Grujić, On decay of the local mean oscillations of the vorticity direction in critical Navier–Stokes flows, arXiv:2609.05720; v1 4 September 2026, v2 9 September 2026. *[Unrefereed preprint, no journal reference. Read in full at S57: §2 (the direction equation (3), which is (142) term for term), Definition 3.1 and Remark 3.2, the drift decomposition (7)–(10), Assumptions 4.1–4.2 (Hk), Assumptions 5.1–5.6, Theorem 5.7, Corollary 5.8, Remark 5.9, §7.2–7.4 including Lemma 7.3, Proposition 7.5, Definition 7.7 and Theorem 7.9, and the obstructions §7.5 (Propositions 7.10 and 7.12 with Remarks 7.11 and 7.13). Quoted as a claim only.]*
- [62] D. Albritton, Z. Bradshaw, Remarks on sparseness and regularity of Navier–Stokes solutions, arXiv:2110.02187v2, 5 April 2022. *[Read at S57 for §1, §1.1 and §1.2 in full. Both halves are quoted in Remark 42.16; the negative verdict concerns the a priori sparseness classes, not the conditional sparseness-implies-regularity criterion, which this paper reproves.]*
- [63] R. R. Coifman, R. Rochberg, G. Weiss, Factorization theorems for Hardy spaces in several variables, *Ann. of Math.*, 103(3):611–635, 1976.
- [64] P. W. Jones, Extension theorems for BMO, *Indiana Univ. Math. J.*, 29(1):41–66, 1980.
- [65] R. A. Hunt, On  $L(p, q)$  spaces, *L’Enseignement Math.*, 12(4):249–276, 1966.