Indiana University Health

Work/School Release Form

Employee/Student Information

Name: Reid, Peyton
Address: 2513 W WELLINGTON DR
MUNCIE, IN 473041270
Sex: Male
Date of Birth: 02/06/2015
Phone: (765) 729-0063

Release Details

This notice verifies that the above identified individual was seen in this facility.

Date seen: 02/08/25 22:54:1710:54 pm EST

He/she may return to work/school after this date: 2/10/25

They will return to work with the following restrictions:

__ None
__ No heavy lifting over __ lbs
__ No prolonged standing greater than __ minutes
__ Desk work only
Other:____

Restrictions apply through this date:___

After the date above, they should be able to participate full in all duties and activities.

Provider Name: Carl Pafford, MD Date: 02/08/25 22:54:17 10:54 pm EST

Signature Line
Electronically Signed on 02/08/25 22:54 EST

Pafford, Carl M, MD

Indiana University Health

Reid, Peyton

DOB: 02/06/2015
MRN: 74813166
Visit Date: 02/08/2025

Discharge Instructions

We would like to thank you for allowing us to assist you with your healthcare needs. The following includes patient education materials and information regarding your injury/illness.

Your ED Provider

Time Seen:
Pafford, Carl M, MD / 02/08/2025 21:08

Tests Performed

Radiology
CT Head W/O IV Contrast 02/08/2025 21:59 EST
XR Shoulder 2 View Min Left 02/08/2025 21:49 EST

Diagnosis from Today's Visit

Influenza A
Vasovagal syncope
Scalp hematoma
Contusion of left shoulder
Laceration of left eyebrow

What to Do Next

You were treated today on an emergency basis; it may be wise to contact your primary care provider to notify them of your visit today. You may have been referred to your regular doctor or a specialist, please follow up as instructed. If your condition worsens or you can't get in to see the doctor, contact the Emergency Department.

You Need to Schedule the Following Appointments

Follow Up with Westergren, Eric S, DO, FAM
When: Within 5 to 7 days

Where: IU Health Ball Memorial Physicians - Yorktown 1420 S Pilgrim Blvd
Yorktown, IN 47396-
(765) 759-4068

Scheduled Follow-Up Appointments

Appointment Type	When	With	Where	Contact Information	Status
BMH OT Continuing Treatment	02/13/2025 03:10 PM EST	Mccaslin, Mary A, OT	RehabP CB Muncie 200 N Tillotson Ave Ste A Muncie, IN 47304-3988	(765) 254-9717	Confirmed

Appointment Type	When	With	Where	Contact Information	Status
BMH OT Continuing Treatment	02/20/2025 03:10 PM EST	Mccaslin, Mary A, OT	RehabP CB Muncie 200 N Tillotson Ave Ste A Muncie, IN 47304-3988	(765) 254-9717	Confirmed
BMH OT Continuing Treatment	02/27/2025 03:10 PM EST	Mccaslin, Mary A, OT	RehabP CB Muncie 200 N Tillotson Ave Ste A Muncie, IN 47304-3988	(765) 254-9717	Confirmed
BMH OT Continuing Treatment	03/06/2025 03:10 PM EST	Mccaslin, Mary A, OT	RehabP CB Muncie 200 N Tillotson Ave Ste A Muncie, IN 47304-3988	(765) 254-9717	Confirmed
BMH OT Continuing Treatment	03/13/2025 03:10 PM EDT	Mccaslin, Mary A, OT	RehabP CB Muncie 200 N Tillotson Ave Ste A Muncie, IN 47304-3988	(765) 254-9717	Confirmed
BMH OT Continuing Treatment	03/20/2025 03:10 PM EDT	Mccaslin, Mary A, OT	RehabP CB Muncie 200 N Tillotson Ave Ste A Muncie, IN 47304-3988	(765) 254-9717	Confirmed
BMH OT Continuing Treatment	03/27/2025 03:10 PM EDT	Mccaslin, Mary A, OT	RehabP CB Muncie 200 N Tillotson Ave Ste A Muncie, IN 47304-3988	(765) 254-9717	Confirmed
BMH OT Continuing Treatment	04/03/2025 03:10 PM EDT	Mccaslin, Mary A, OT	RehabP CB Muncie 200 N Tillotson Ave Ste A Muncie, IN 47304-3988	(765) 254-9717	Confirmed
BMH OT Continuing Treatment	04/10/2025 03:10 PM EDT	Mccaslin, Mary A, OT	RehabP CB Muncie 200 N Tillotson Ave Ste A Muncie, IN 47304-3988	(765) 254-9717	Confirmed
BMH OT Continuing Treatment	04/17/2025 03:10 PM EDT	Mccaslin, Mary A, OT	RehabP CB Muncie 200 N Tillotson Ave Ste A Muncie, IN 47304-3988	(765) 254-9717	Confirmed
BMH OT Continuing Treatment	04/24/2025 03:10 PM EDT	Mccaslin, Mary A, OT	RehabP CB Muncie 200 N Tillotson Ave Ste A Muncie, IN 47304-3988	(765) 254-9717	Confirmed
BMH OT Continuing Treatment	05/01/2025 03:10 PM EDT	Mccaslin, Mary A, OT	RehabP CB Muncie 200 N Tillotson Ave Ste A Muncie, IN 47304-3988	(765) 254-9717	Confirmed
BMH OT Continuing Treatment	05/08/2025 03:10 PM EDT	Mccaslin, Mary A, OT	RehabP CB Muncie 200 N Tillotson Ave	(765) 254-9717	Confirmed

Appointment Type	When	With	Where	Contact Information	Status
			Ste A Muncie, IN 47304-3988		
BMH OT Continuing Treatment	05/15/2025 03:10 PM EDT	Mccaslin, Mary A, OT	RehabP CB Muncie 200 N Tillotson Ave Ste A Muncie, IN 47304-3988	(765) 254-9717	Confirmed

Education Materials

Hematoma

A hematoma is a collection of blood trapped outside of a blood vessel. It's what we think of as a bruise or a contusion. It's often seen under the skin as a black and blue spot on your arm or leg, or a bump on your head after an injury. It can be almost anywhere on or in your body. It can also occur in an internal organ. This can be more serious.

A hematoma is caused by an injury with damage to small blood vessels. This causes blood to leak into the tissues. Blood forms a pocket under the skin that swells and looks like a purplish patch. Hematomas sometimes form under the skin from bleeding during childbirth and can be particularly serious. Another serious form of hematoma forms after a fall on the head, called a subdural hematoma.

Gradually the blood in the hematoma is absorbed back into the body. The swelling and pain of the hematoma will go away. This takes from 1 to 4 weeks, depending on the size of the hematoma. The skin over the hematoma may turn bluish then brown and yellow as the blood is dissolved and absorbed. Usually, this only takes a couple of weeks but can last months.

Home care

- Limit motion of the joints near the hematoma. If the hematoma is large and painful, avoid sports and other vigorous physical activity until the swelling and pain goes away.
- Apply an ice pack over the injured area for 20 minutes every 1 to 2 hours the first day. Continue with ice packs 3 to 4 times a day for the next 2 days. To make an ice pack, put ice cubes in a plastic bag that seals at the top. Wrap the bag in a thin towel or cloth. Don't put ice or an ice pack directly on the skin. Continue the use of ice packs to ease pain and swelling as needed.
- Take acetaminophen for pain relief, unless you were given a different pain medicine to use. Talk with your provider before using this medicine if you:
    - Have chronic liver or kidney disease
    - Have had a stomach ulcer or digestive tract bleeding
    - Are taking blood-thinner medicines.

Follow-up care

Follow up with your healthcare provider, or as advised. If X-rays or a CT scan were done, you'll be told if there is a change in the reading, especially if it affects treatment.

When to get medical advice

Call your healthcare provider right away if any of the following occur:

- Redness around the hematoma
- Increase in pain or warmth in the hematoma
- Increase in size of the hematoma
- Fever of 100.4°F (38°C) or higher, or as directed by your provider
- If the hematoma is on the arm or leg, watch for:
    - More swelling or pain in the extremity
    - Numbness or tingling or blue color of the hand or foot

© 2000-2024 The StayWell Company, LLC. All rights reserved. This information is not intended as a substitute for professional medical care. Always follow your healthcare professional's instructions.

Upper Extremity Bruise

You have a bruise (contusion) on your arm, wrist, hand, or fingers. Symptoms include pain, swelling, and skin discoloration. No bones are broken. This injury may take from a few days to a few weeks to heal. During that time, the bruise may change from a red color to purple- blue, to green-yellow, to yellow-brown.

Home care

- Unless another medicine was prescribed, you can take acetaminophen, ibuprofen, or naproxen to control pain. Talk with your healthcare provider before using these medicines if you have chronic liver or kidney disease or ever had a stomach ulcer or digestive bleeding.
- Raise the injured area to reduce pain and swelling. As much as possible, sit or lie down with the injured area raised about the level of your heart. This is especially important during the first 48 hours.
- Ice the injured area to help reduce pain and swelling. To make an ice pack, put ice cubes in a plastic bag that seals at the top. Wrap the bag in a thin towel. Don't put an ice pack directly on the skin. Apply to the bruised area for 20 minutes every 1 to 2 hours the first day. Continue this 3 to 4 times a day until the pain and swelling goes away.
- If a sling was provided, you may remove it to shower or bathe. To prevent joint stiffness, don't wear it for more than 1 week. Also, make sure your hand is slightly above the level of the elbow in the sling to prevent your hand from becoming swollen and discolored by gravity.

Follow-up care

Follow up with your healthcare provider, or as advised. Call if you're not improving in the next 1 to 2 weeks.

When to get medical advice

Call your healthcare provider right away if any of these occur:


- Increased pain or swelling
- Hand or fingers become cold, blue, numb, or tingly
- Signs of infection: warmth, drainage, or increased redness or pain around the injury
- Inability to move the injured body part, the hand, or individual fingers.
- Frequent bruising for unknown reasons

© 2000-2024 The StayWell Company, LLC. All rights reserved. This information is not intended as a substitute for professional medical care. Always follow your healthcare professional's instructions.

You can wash and pat dry your facial laceration. Do not rub it or put any lotions over it. Keep your face out of the sun to decrease the pigment uptake. Watch for signs of infection such as fever, redness, pus leaking from the wound

LACERATION, Face (Dermabond)

A laceration is a cut through the skin. A laceration on your face has been closed with Dermabond skin adhesive, a type of skin glue.



HOME CARE

Medications: Acetaminophen (Tylenol) or ibuprofen (Motrin, Advil) may be taken for pain, unless another pain medicine was prescribed. NOTE: If you have chronic liver or kidney disease or ever had a stomach ulcer or GI bleeding, talk with your doctor before using these medications.

General Care:

- Keep the wound clean and dry. You may shower or bathe as usual, but do not use soaps, lotions, or ointments on the wound area. Do not scrub the wound. After bathing, pat the wound dry with a soft towel.
- Do not scratch, rub, or pick at the film. Do not place tape directly over the film.
- Do not apply liquids (such as peroxide), ointments, or creams to the wound while the film is in place.
- Most facial skin wounds heal without problems. However, an infection sometimes occurs despite proper treatment. Therefore, watch for the signs of infection listed below.

FOLLOW UP as directed by the doctor or our staff. The Dermabond film will fall off naturally in 5 to 10 days.

GET PROMPT MEDICAL ATTENTION if any of the following occur:

- Signs of infection:
  Fever over 100.0°F (37.8°C)
  Increasing pain in the wound
  Increasing redness or swelling
  Pus coming from the wound
- Wound bleeds more than a small amount or bleeding doesn't stop

Wound edges come apart

© 2000-2011 Krames StayWell, 780 Township Line Road, Yardley, PA 19067. All rights reserved. This information is not intended as a substitute for professional medical care. Always follow your healthcare professional's instructions.

Your Medications

	What	How Much	When	Instructions	Next Dose
Unchanged	cetirizine (ZyrTEC Children's Allergy 1 mg/ mL oral syrup)	2.5 Milliliter By mouth	Once a day		

Allergies

NKA

Additional Information

General Information: We are happy to serve you. The examination and treatment you have received have been on an emergency basis only. Call the Emergency Department for any questions or concerns (listed on front page). An emergency is an illness or injury so serious that you need to get care right away to keep from getting worse. If you do not have an emergency, you should go to your doctor's office. For help finding a doctor, visit www.iuhealth.org and search "Find a Doctor". We are very concerned about your general health and safety. Because of this, when driving or riding in a vehicle, we encourage you to always wear your seat belt, and if you have an infant or child, always use the appropriate infant/child car seat. You were treated today on an emergency basis; it may be wise to contact your primary care provider to notify them of your visit today. You may have been referred to your regular doctor or a specialist, please follow up as instructed.

Medication Instructions: If you need a prescription refill, you must see your doctor. Take all medications only as directed. Do not take a medication you are allergic to. If you have been prescribed new medications check with your pharmacist to make sure there are no problems with the medications you are taking.
To help ensure your safety, keep a copy of this discharge instruction to share with your health care provider at your next visit. It contains a copy of your medication list, diagnoses made in our emergency department and information about treatment you received here.

Referral Instructions: You were treated today on an emergency basis; it may be wise to contact you primary care provider to notify them of your visit today. You may have been referred to your regular doctor or a specialist, please follow up as instructed. If your condition worsens or you can't get in to see the doctor, contact the Emergency Department.

Insurance Information: When you sign in, it is your responsibility to give correct and complete billing information. This is important in case there is a need to contact you with test results. Federal law requires us to provide you a medical screening exam and stabilizing treatment for any emergent medical condition found regardless of your insurance or ability to pay. Your insurance company may require pre-authorization or notification of your visit, both of which are your responsibility. We will make every effort to properly bill your insurance company on your behalf. If you need financial assistance, please call the number listed on front page.

Diagnostic Studies: If you had a diagnostic study performed, we will notify you of any lab, EKG or X-ray findings that would require a change in treatment. A specialist reviews most diagnostic studies after your visit. Some test results are not available for 24-72 hours.

If you have been given medications or a prescription in the Emergency Department: Many medications can make you feel sleepy, impair your ability to drive, or effect your ability to make good decisions. Some of these medicines can even make you stop breathing when you go to sleep, particularly if several different medicines are combined. For this reason, we ask that you arrange for someone else to give you a ride home, do not operate machinery or make important decisions, and do not take any sleeping pills, pain pills, or drink alcohol for at least 4-6 hours from the time you leave the emergency department. It is also important to know that the use of narcotic medications can be habit forming.

Crisis Hotlines:
USA National Suicide Hotline (Toll free and 24/7): Dial 988

Antibiotics Are Not Always the Answer: Antibiotics only treat bacterial infections; they are not needed for viral illnesses. When an antibiotic is not prescribed, ask your healthcare professional for tips on how to relieve symptoms and feel better. For more information on when antibiotics may be helpful, visit www.cdc.gov/getsmart.

Quit Smoking for Health: Smoking has been proven to be hazardous to your health. If you would like to quit smoking, call 1-800-QUIT-NOW (1-800-784-8669) for free counseling and two weeks of free Nicotine Replacement Therapy. Helpful information can also be found online at www.iuhealth.org or the Quitline can also provide information about the closest smoking cessation program to you.

My IU Health: My IU Health allows convenient, electronic access to manage your health. Sign up for a secure account at www.myiuhealth.org.