



# Timing Noise Analysis of NANOGrav pulsars

Delphine Perrodin

Franklin & Marshall College  
Osservatorio Astronomico di Cagliari

IPTA meeting @ Kiama  
June 26, 2012






LEGGENDO  
METROPOLITANO

IL TEMPO È PRESENTE

GIUGNO 2012  
QUARTIERE CASTELLO  
leggendometropolitano.it



This slide contained a picture of Delphine's Drivers License and Italian Identity Card. For the sake of privacy they have been removed. Cheers! Imagine a funny drivers license picture.





Perrodin, Jenet, Lommen, Finn, Demorest,  
Ferdman, Gonzalez, Nice, Ransom, Stairs (2012)

# Outline



Perrodin, Jenet, Lommen, Finn, Demorest,  
Ferdman, Gonzalez, Nice, Ransom, Stairs (2012)

# Outline

- Autocorrelation of pulsar timing residuals



Perrodin, Jenet, Lommen, Finn, Demorest,  
Ferdman, Gonzalez, Nice, Ransom, Stairs (2012)

# Outline

- Autocorrelation of pulsar timing residuals
- “nearest-neighbor” autocorrelation



Perrodin, Jenet, Lommen, Finn, Demorest,  
Ferdman, Gonzalez, Nice, Ransom, Stairs (2012)

# Outline

- Autocorrelation of pulsar timing residuals
- “nearest-neighbor” autocorrelation
- delta-function statistic





Perrodin, Jenet, Lommen, Finn, Demorest,  
Ferdman, Gonzalez, Nice, Ransom, Stairs (2012)

# Outline

- Autocorrelation of pulsar timing residuals
- “nearest-neighbor” autocorrelation
- delta-function statistic
- exponential fit



Perrodin, Jenet, Lommen, Finn, Demorest,  
Ferdman, Gonzalez, Nice, Ransom, Stairs (2012)

# Outline

- Autocorrelation of pulsar timing residuals
- “nearest-neighbor” autocorrelation
- delta-function statistic
- exponential fit
- Red noise in NANOGrav pulsars




Perrodin, Jenet, Lommen, Finn, Demorest,  
Ferdman, Gonzalez, Nice, Ransom, Stairs (2012)

# Outline

- Autocorrelation of pulsar timing residuals
- “nearest-neighbor” autocorrelation
- delta-function statistic
- exponential fit
- Red noise in NANOGrav pulsars
- Observing strategies

# NANOGrav residuals


## NANOGrav 5-year timing results overview:



(plot: D. Nice)

Demorest et al. (2012)

# NANOGrav residuals



Demorest et al. (2012)





# Characterizing (red) timing noise

# Characterizing (red) timing noise

- Useful for GW detection pipelines; need to separate out noise from GW signature at low frequencies

# Characterizing (red) timing noise

- Useful for GW detection pipelines; need to separate out noise from GW signature at low frequencies
- Data non-uniformly sampled. Easier to study noise in time-domain.

# Characterizing (red) timing noise

- Useful for GW detection pipelines; need to separate out noise from GW signature at low frequencies
- Data non-uniformly sampled. Easier to study noise in time-domain.
- We calculated the autocorrelation function of timing residuals:

# Characterizing (red) timing noise

- Useful for GW detection pipelines; need to separate out noise from GW signature at low frequencies
- Data non-uniformly sampled. Easier to study noise in time-domain.
- We calculated the autocorrelation function of timing residuals:

$$\hat{R}(k) = \frac{1}{(n-k)\sigma^2} \sum_{t=1}^{n-k} (X_t - \mu)(X_{t+k} - \mu)$$





# Method 1:

# Method 1:

- Based on TEMPO<sub>2</sub>'s “CheckWhite” plugin

# Method 1:

- Based on TEMPO2's "CheckWhite" plugin
- Examines correlations of adjacent observations

# Method 1:

- Based on TEMPO2's "CheckWhite" plugin
- Examines correlations of adjacent observations
- Good for first examination (but does not find scale)

# Method 1:

- Based on TEMPO2's "CheckWhite" plugin
- Examines correlations of adjacent observations
- Good for first examination (but does not find scale)

Did not find any non-white  
correlation with this method.





## Method 2:

## Method 2:

- Autocorrelation function (as a function of time lag) of “white” data should be delta-function (zero-lag correlation corresponds to amplitude of white noise).

## Method 2:

- Autocorrelation function (as a function of time lag) of “white” data should be delta-function (zero-lag correlation corresponds to amplitude of white noise).
- Can test the “whiteness” of timing residuals by measuring deviations from a delta-function.

## Method 2:

- Autocorrelation function (as a function of time lag) of “white” data should be delta-function (zero-lag correlation corresponds to amplitude of white noise).
- Can test the “whiteness” of timing residuals by measuring deviations from a delta-function.

$$\chi^2 = \frac{1}{N-1} \sum_{i=1}^N \frac{Y_i^2}{\sigma_i^2}$$

## Method 2:

- Autocorrelation function (as a function of time lag) of “white” data should be delta-function (zero-lag correlation corresponds to amplitude of white noise).
- Can test the “whiteness” of timing residuals by measuring deviations from a delta-function.

$$\chi^2 = \frac{1}{N-1} \sum_{i=1}^N \frac{Y_i^2}{\sigma_i^2}$$

Results? High  $\chi^2$  in J1643-1224 (112), J1910+1256 (382) and J1640+2224 (654).





## Method 3:

## Method 3:

- Based on TEMPO2's "SpectralModel" plugin method for weak red noise.

## Method 3:

- Based on TEMPO2's "SpectralModel" plugin method for weak red noise.
- Model for timing noise in frequency domain:

## Method 3:

- Based on TEMPO2's "SpectralModel" plugin method for weak red noise.
- Model for timing noise in frequency domain:

$$P(f) = \frac{A}{\left(1 + \left(\frac{f}{f_c}\right)^2\right)^{\alpha/2}} + B$$

## Method 3:

- Based on TEMPO2's "SpectralModel" plugin method for weak red noise.
- Model for timing noise in frequency domain:

$$P(f) = \frac{A}{\left(1 + \left(\frac{f}{f_c}\right)^2\right)^{\alpha/2}} + B$$

Fourier transform of autocorrelation function (alpha=2):

## Method 3:

- Based on TEMPO2's "SpectralModel" plugin method for weak red noise.
- Model for timing noise in frequency domain:

$$P(f) = \frac{A}{\left(1 + \left(\frac{f}{f_c}\right)^2\right)^{\alpha/2}} + B$$

Fourier transform of autocorrelation function (alpha=2):

$$C(d) = Ce^{-d/\tau} + \sqrt{2\pi}B\delta(d)$$





## Method 3:

## Method 3:

Results? Significant effect seen in  
J1910+1256 , J2145-0750 and J1643-1224.

# Method 3:

Results? Significant effect seen in J1910+1256 , J2145-0750 and J1643-1224.

Pulsar	Freq(MHz)	$C(10^{-13}s^2)$
J1910+1256	1400	$9.9 \pm 1.8$
J2145-0750	1400	$9.8 \pm 1.3$
J1643-1224	1400	$11 \pm 7$

- Compare amplitude of non-white noise with amplitude of white noise
- Variance of white noise expected to decrease as  $1/N$

$$\sigma_w^2 = \frac{\sigma_0^2}{N}$$

- Improve rms until level of white noise down to level of red noise:

$$\sigma_w^2 = \frac{\sigma_0^2}{N} = \sigma_r^2$$

- Define:


$$\text{Time Factor} = N = \frac{\sigma_0^2}{\sigma_r^2}$$

Can increase number of observations up to N times current number and still expect rms to go down as  $1/\sqrt{N}$


J1910+1256 and J1643-1224 exhibit the lowest factors:


Pulsar	Freq(MHz)	Obs	$\chi^2$	$C(10^{-13}s^2)$	Factor
J1910+1256	1400	AO	382	$9.9 \pm 1.8$	7
J1643-1224	1400	GBT	112	$11 \pm 7$	18
J1640+2224	1400	AO	654	–	40
J2145-0750	1400	GBT	37	$9.8 \pm 1.3$	66

Even then, can increase the cadence or integration time  
6-fold and still not reach the level of red noise ->  
incentive to increase observation time in order to  
achieve lower rms











# Validity of methods?

# Validity of methods?

- “Nearest-neighbor” method not sufficient, does not probe time scales

# Validity of methods?

- “Nearest-neighbor” method not sufficient, does not probe time scales
- Delta-function statistic useful for identifying deviations from white noise

# Validity of methods?

- “Nearest-neighbor” method not sufficient, does not probe time scales
- Delta-function statistic useful for identifying deviations from white noise
- Exponential fit not adequate for weak red noise in nanograv pulsars. Shape of correlation function changed by parameter fitting (Levin & van Haasteren; Ellis et al.)



# Validity of methods?

- “Nearest-neighbor” method not sufficient, does not probe time scales
- Delta-function statistic useful for identifying deviations from white noise
- Exponential fit not adequate for weak red noise in nanograv pulsars. Shape of correlation function changed by parameter fitting (Levin & van Haasteren; Ellis et al.)
- Try fitting more general functions?

# Validity of methods?

- “Nearest-neighbor” method not sufficient, does not probe time scales
- Delta-function statistic useful for identifying deviations from white noise
- Exponential fit not adequate for weak red noise in nanograv pulsars. Shape of correlation function changed by parameter fitting (Levin & van Haasteren; Ellis et al.)
- Try fitting more general functions?
- Bayesian methods are being explored (van Haasteren, Ellis)



# Red noise in NANOGrav pulsars?

# Red noise in NANOGrav pulsars?

- Evidence for weak red noise in J1910+1256 and J1643-1256 (borderline case: J1640+2224)

# Red noise in NANOGrav pulsars?

- Evidence for weak red noise in J1910+1256 and J1643-1256 (borderline case: J1640+2224)
- However: J1910+1256 was only observed at one frequency and DM variations were not included -> not necessarily a “bad” pulsar

# Red noise in NANOGrav pulsars?

- Evidence for weak red noise in J1910+1256 and J1643-1256 (borderline case: J1640+2224)
- However: J1910+1256 was only observed at one frequency and DM variations were not included -> not necessarily a “bad” pulsar
- J1640+2224 has orbital period close to  $\frac{1}{2}$  year, can lead to poor orbital sampling





# Conclusions

# Conclusions

- NANOGrav pulsars are white noise-dominant

# Conclusions

- NANOGrav pulsars are white noise-dominant
- Can freely increase the cadence or integration time of our observations in order to achieve lower rms and better sensitivity to GW

# Conclusions

- NANOGrav pulsars are white noise-dominant
- Can freely increase the cadence or integration time of our observations in order to achieve lower rms and better sensitivity to GW
- DM variation method successful at removing red noise

# Conclusions

- NANOGrav pulsars are white noise-dominant
- Can freely increase the cadence or integration time of our observations in order to achieve lower rms and better sensitivity to GW
- DM variation method successful at removing red noise
- At IPTA level: should test DM variation algorithms on same data sets