

Stochastic background searches using the optimal statistic

Xavi Siemens

Sydney Chamberlin, Jolien Creighton, Paul Demorest,
Justin Ellis, Larry Price, Joe Romano


Anholm et al. 2009, van Haasteren et al. 2009, Demorest et al 2012

The pipeline




Justin
Ellis

Look at SNR
and decide
whether we
have made a
detection



The pipeline




Look at SNR
and decide
whether we
have made a
detection

Find $\Omega_{95\%}, A_{95\%}$: Values
of injected BG for which
95% of the time $\hat{\Omega} > \hat{\Omega}^*$



Justin
Ellis



Sydney
Chamberlin

Optimal statistic

Start from likelihood (standard Gaussian multivariate)

$$p(r|\Omega) = \frac{1}{\sqrt{\det(2\pi\Sigma_r)}} \exp\left(-\frac{1}{2}r^T\Sigma_r^{-1}r\right)$$

Residuals $\mathbf{r} = \begin{bmatrix} \mathbf{r}_1 \\ \mathbf{r}_2 \\ \vdots \\ \mathbf{r}_l \end{bmatrix}$

Covariance matrix for residuals $\Sigma_r = \langle \mathbf{r}\mathbf{r}^T \rangle = \begin{bmatrix} \mathbf{P}_1 & \Omega\mathbf{S}_{12} & \cdots & \Omega\mathbf{S}_{1l} \\ \Omega\mathbf{S}_{21} & \mathbf{P}_2 & \cdots & \Omega\mathbf{S}_{2l} \\ \vdots & \vdots & \ddots & \vdots \\ \Omega\mathbf{S}_{l1} & \Omega\mathbf{S}_{l2} & \cdots & \mathbf{P}_l \end{bmatrix}$

$$\Sigma_r = \begin{bmatrix} \mathbf{P}_1 & \Omega\mathbf{S}_{12} & \cdots & \Omega\mathbf{S}_{1l} \\ \Omega\mathbf{S}_{21} & \mathbf{P}_2 & \cdots & \Omega\mathbf{S}_{2l} \\ \vdots & \vdots & \ddots & \vdots \\ \Omega\mathbf{S}_{l1} & \Omega\mathbf{S}_{l2} & \cdots & \mathbf{P}_l \end{bmatrix} \quad \textbf{covariance matrix}$$

Auto-correlation matrices (diagonal elements of covariance matrix)

$$P_I = \langle r_I r_I^T \rangle_{ij} = R_I \left[\int_{-\infty}^{\infty} df e^{2\pi i f (t_i - t_j)} P_r(f) + \sigma_I^2 \delta_{ij} \right] R_I^T$$

Red noise power spectrum $P_r(f) = A f^{-\gamma}$

Red and white noise parameters modeled and measured from individual residual pulsar data (see poster by Justin Ellis)

Timing model $R_I = I - A_I (A_I^T C_I^{-1} A_I)^{-1} A_I^T C_I^{-1}$

$$\Sigma_r = \begin{bmatrix} \mathbf{P}_1 & \Omega \mathbf{S}_{12} & \cdots & \Omega \mathbf{S}_{1l} \\ \Omega \mathbf{S}_{21} & \mathbf{P}_2 & \cdots & \Omega \mathbf{S}_{2l} \\ \vdots & \vdots & \ddots & \vdots \\ \Omega \mathbf{S}_{l1} & \Omega \mathbf{S}_{l2} & \cdots & \mathbf{P}_l \end{bmatrix} \quad \text{covariance matrix}$$

Cross-correlation (off-diagonal elements of covariance matrix)

$$\Omega_\beta S_{IJ} = \langle r_I r_J^T \rangle_{ij} = R_I \left[\frac{\chi_{IJ}}{2} \int_{-\infty}^{\infty} df e^{2\pi i f (t_i - t_j)} P_g(f) \right] R_J^T$$

Hellings-Downs curve χ_{IJ}

GW power spectrum $P_g(f) = \frac{H_0^2}{16\pi^4} \Omega_\beta f^{\beta-5} = \frac{A_g^2}{24\pi^2} \left(\frac{f}{f_{1\text{yr}}} \right)^{2\alpha} f^{-3}$

Timing model $R_I = I - A_I (A_I^T C_I^{-1} A_I)^{-1} A_I^T C_I^{-1}$

Optimal statistic is obtained by maximizing the likelihood over GW amplitude (but fixed spectrum)

Pair-wise optimal statistic

$$Q_{IJ} = N_{IJ} r_I^T P_I^{-1} S_{IJ} P_J^{-1} r_J$$

Normalization

$$N_{IJ} = \left(\text{Tr} \left[P_I^{-1} S_{IJ} P_J^{-1} S_{IJ}^T \right] \right)^{-1}$$

Pairwise standard deviation

$$\sigma_{IJ} \approx \sqrt{N_{IJ}}$$

Normalization is chosen so that

$$\langle Q_{IJ} \rangle = \Omega_\beta$$

**PTA optimal
statistic**

$$Q_{\text{opt}} = \frac{\sum_{IJ} Q_{IJ} \sigma_{IJ}^{-2}}{\sum_{IJ} \sigma_{IJ}^{-2}}$$

**Pairwise standard
deviation**


$$\sigma_{IJ} \approx \sqrt{N_{IJ}}$$

Normalization is chosen so that

$$\langle Q_{\text{opt}} \rangle = \Omega_{\beta}$$


$$P_I = \langle r_I r_I^T \rangle_{ij} = R_I \left[\int_{-\infty}^{\infty} df e^{2\pi i f (t_i - t_j)} P_r(f) + \sigma_I^2 \delta_{ij} \right] R_I^T$$

Autocorrelation matrix for single pulsar with quadratic subtraction




$$P_I^{-1}$$

SVD inverse of Autocorrelation matrix for single pulsar with quadratic subtraction




$$\Omega_{\beta} S_{IJ} = \langle r_I r_J^T \rangle_{ij} = R_I \left[\frac{\chi_{IJ}}{2} \int_{-\infty}^{\infty} df e^{2\pi i f (t_i - t_j)} P_g(f) \right] R_J^T$$

S_{IJ} cross-correlation matrix



$$Q_{IJ} = N_{IJ} r_I^T P_I^{-1} S_{IJ} P_J^{-1} r_J$$

Optimal filter: $N_{IJ} P_I^{-1} S_{IJ} P_J^{-1}$



Open Mock Data Challenge

Dataset 1: 36 pulsars, 100ns white noise, SMBBH spectrum, $A = 5 \times 10^{-14}$

Dataset 2: 36 pulsars, different white noises for pulsars (given), SMBBH spectrum, $A = 5 \times 10^{-14}$

Dataset 3: 36 pulsars, different white noises for pulsars (given), SMBBH spectrum, $A = 10^{-14}$, and additional red noise

Open Mock Data Challenge

Dataset 1: 36 pulsars, 100ns white noise, SMBBH spectrum, $A = 5 \times 10^{-14}$

FOUND: $A = (4.9 \pm 0.19) \times 10^{-14}$

SNR = 13

Dataset 2: 36 pulsars, different white noises for pulsars (given), SMBBH spectrum, $A = 5 \times 10^{-14}$

Dataset 3: 36 pulsars, different white noises for pulsars (given), SMBBH spectrum, $A = 10^{-14}$, and additional red noise

Open Mock Data Challenge

Dataset 1: 36 pulsars, 100ns white noise, SMBBH spectrum, $A = 5 \times 10^{-14}$

$$\text{FOUND: } A = (4.9 \pm 0.19) \times 10^{-14}$$

$$\text{SNR} = 13$$

Dataset 2: 36 pulsars, different white noises for pulsars (given), SMBBH spectrum, $A = 5 \times 10^{-14}$

$$\text{FOUND: } A = (4.7 \pm 0.27) \times 10^{-14}$$

$$\text{SNR} = 8.8$$

Dataset 3: 36 pulsars, different white noises for pulsars (given), SMBBH spectrum, $A = 10^{-14}$, and additional red noise

Open Mock Data Challenge

Dataset 1: 36 pulsars, 100ns white noise, SMBBH spectrum, $A = 5 \times 10^{-14}$

$$\text{FOUND: } A = (4.9 \pm 0.19) \times 10^{-14}$$

$$\text{SNR} = 13$$

Dataset 2: 36 pulsars, different white noises for pulsars (given), SMBBH spectrum, $A = 5 \times 10^{-14}$

$$\text{FOUND: } A = (4.7 \pm 0.27) \times 10^{-14}$$

$$\text{SNR} = 8.8$$


Dataset 3: 36 pulsars, different white noises for pulsars (given), SMBBH spectrum, $A = 10^{-14}$, and additional red noise

$$\text{FOUND: } A = (1.2 \pm 0.07) \times 10^{-14}$$

$$\text{SNR} = 8.7$$


Noise estimation bias

Noise estimation biases SNR




Noise estimation bias

Noise estimation biases amplitude estimation



Noise estimation bias

Two wrongs that make a right



Very preliminary, rough, upper limit estimates*



CAUTION: Based on variance of estimators, not injections

NANOGrav: $A_{95\%} = 6.4 \times 10^{-15}$ compare with 7.2×10^{-15}
Demorest et al. 2012

EPTA (5 pulsar dataset): $A_{95\%} = 6.5 \times 10^{-15}$ compare with 6×10^{-15}
van Haasteren et al. 2011

Agreement between three independent techniques!!

Working on combining both data sets

Working on PPTA data...

*Your mileage may vary

The end

How do we derive the likelihood for the residuals?

- Start from Gaussian likelihood for GWs

$$p(y|\Omega) = \frac{1}{\sqrt{\det(2\pi\Sigma_y)}} \exp\left(-\frac{1}{2}y^T\Sigma_y^{-1}y\right)$$

$$\mathbf{y} = \begin{bmatrix} \mathbf{y}_1 \\ \mathbf{y}_2 \\ \vdots \\ \mathbf{y}_l \end{bmatrix} \quad \Sigma_y = \begin{bmatrix} \mathbf{P}_1 & \Omega\mathbf{S}_{12} & \cdots & \Omega\mathbf{S}_{1l} \\ \Omega\mathbf{S}_{21} & \mathbf{P}_2 & \cdots & \Omega\mathbf{S}_{2l} \\ \vdots & \vdots & \ddots & \vdots \\ \Omega\mathbf{S}_{l1} & \Omega\mathbf{S}_{l2} & \cdots & \mathbf{P}_l \end{bmatrix} \quad \begin{aligned} \Omega S_{IJ} &= \langle y_I y_J^T \rangle \\ P_I &= \langle y_I y_I^T \rangle \end{aligned}$$

$$y_I \rightarrow r_I = R_I y_I \quad R_I = I - A_I (A_I^T C_I^{-1} A_I)^{-1} A_I^T C_I^{-1}$$

$$p(y|\Omega) \rightarrow p(r|\Omega) = \frac{1}{\sqrt{\det(2\pi\Sigma_r)}} \exp\left(-\frac{1}{2}r^T\Sigma_r^{-1}r\right)$$

$$\Sigma_r = R\Sigma_y R^T$$