

Chapter 5

Later

Theorem 5.0.1. *The following are equivalent:*

1. *The Intermediate Value Theorem holds for X : for any continuous map $f : X \rightarrow \mathbb{R}$, and any c between $f(x_1)$ and $f(x_2)$ for some $x_1, x_2 \in X$, there exists some $x \in X$ such that $f(x) = c$.*
2. *There is no continuous surjection from X onto a discrete two-point space $\{0, 1\}$.*
3. *X cannot be partitioned into two nonempty disjoint open sets.*
4. *X cannot be partitioned into two nonempty disjoint closed sets.*
5. *Every proper subset of X that is both open and closed is either \emptyset or X itself.*

Remark If X is path-connected, then it is connected. The converse is not necessarily true.

Example 5.0.1. *Let $X = \{(x, y) | y = \sin(x)\} \cup \{(0, y) | y \in [-1, 1]\} \subset \mathbb{R}^2$ with the Euclidean metric. Then X is connected but not path-connected.*

Proof

First we prove that X is path-connected. Let $p_1, p_2 \in X$ such that $P_1 = (0, 0)$ and $P_2 = (1, \sin(1))$. Let $\gamma : [0, 1] \rightarrow X$ with $\gamma(t) = (x(t), y(t))$ such that $\gamma(0) = P_1$ and $\gamma(1) = P_2$. Since γ is continuous, both $x(t)$ and $y(t)$ are continuous.

Let $U = \{t | x(t) = 0\} \subseteq [0, 1]$. Since U is a preimage of a closed set under a continuous map, U is closed. Thus, $t_0 = \sup U$ exists. Note that $t_0 < 1$ since $x(1) = 1$.

Let $\Phi(t) = \frac{1}{x(t)}$ which is well-defined on for $t > t_0$. Notice that $\Phi(t) \rightarrow +\infty$ as $t \rightarrow t_0^+$. Then, for $t > t_0$, $y(t) = \sin(\Phi(t))$. Take the interval $(t_0, t_0 + \epsilon)$, then $y(t)$ oscillates between -1 and 1 infinitely many times. So $y(t)$ is not continuous at t_0 . This is a contradiction. Thus, no such path γ exists and X is not path-connected.

Lemma 5.0.1. *The closure of a connected set is connected.*

Proof Let A be a connected set and \bar{A} be its closure. Suppose $\bar{A} = U_1 \cup U_2$ where U_1 and U_2 are nonempty disjoint open sets in \bar{A} . Then $A = (A \cap U_1) \cup (A \cap U_2)$ is non-trivial: let $U_1 \cap A = \emptyset$, then $U_1 \subseteq \bar{A} \setminus A$ which is impossible.

Second, we prove that X is connected. Let $X = Y \cup \{(0, y) | y \in [-1, 1]\}$. We know that $\bar{Y} = X$ and Y is path-connected (thus connected). Then using the property that the closure of a connected set is connected, we have X is connected.

Remark Also, we can prove it using 2 in the ??.

5.0.1 Connected Components

For any $x, y \in X$, we say that $x \sim y$ if $\exists \gamma : [0, 1] \rightarrow X$ continuous such that $\gamma(0) = x$ and $\gamma(1) = y$. It is easy to check that \sim is an equivalence relation. We call the equivalence classes the connected components of X .

Exercise Define connected components for the notion of connected (not just path-connected) spaces.

Observation The number of connected components is preserved under homeomorphisms.

Example 5.0.2. $(0, 1) \cup (2, 3)$ is not homeomorphic to $(0, 1)$ since the former has 2 connected components while the latter has 1 connected component.

Example 5.0.3. The interval $[0, 1]$ is not homeomorphic to $[0, 1] \times [0, 1]$. Removing a point (not in the boundary) from $[0, 1]$ results in a disconnected space, while removing a point from $[0, 1] \times [0, 1]$ still results in a connected space. Thus, they are not homeomorphic.

Example 5.0.4. $(0, 1) \approx [0, 1] \approx [0, 1)$. Removing a boundary point from $[0, 1]$ still results in a connected space, while removing a boundary point from $(0, 1)$ results in a disconnected space. Thus, they are not homeomorphic.

Example 5.0.5. We can split the alphabet(capital letter) into nine homeomorphism classes:

- A
- B
- $C, I, J, L, M, N, S, U, V, W, Z$
- D, O
- E, F, G, T, Y
- H
- K, X, Z
- P
- Q, R

Example 5.0.6 (First homology group).

$$2026 \rightsquigarrow 2$$

$$2025 \rightsquigarrow 1$$

$$1949 \rightsquigarrow 2$$

$$1982 \rightsquigarrow 3$$

$$1988 \rightsquigarrow 5$$

Property 5.0.1. $U \subseteq \mathbb{R}^2$ open in one of the metrics defined by the norms listed below iff it is open in the others:

- Euclidean norm: $\|(x, y)\|_2 = \sqrt{x^2 + y^2}$
- Supremum norm: $\|(x, y)\|_\infty = \max\{|x|, |y|\}$
- Diamond norm: $\|(x, y)\|_{1/2} = \sqrt{|x|} + \sqrt{|y|}$

In other words, these metrics define the same topology. So they have the same open sets.

Definition 5.0.1. Two norms on V $\|\cdot\|_a$ and $\|\cdot\|_b$ are called equivalent if $\exists c_1, c_2$ such that:

$$c_1\|v\|_a \leq \|v\|_b \leq c_2\|v\|_a \quad (5.1)$$

Exercise

1. This is an equivalent relation.
 2. Equivalent norms define the same topology.
 3. $\|\cdot\|_1$, $\|\cdot\|_2$, and $\|\cdot\|_\infty$ on \mathbb{R}^n are equivalent.
- * All norms on \mathbb{R}^n are equivalent.