

Chapter 19

Lecture 19: Countability. Sorgenfrey line and Sorgenfrey plane. Different notions of separability.

19.1 Countability

Definition 19.1.1. X is first countable if for every $x \in X$, there exists a countable basis at x . That is, there exists $\mathcal{B} = \{B_n : n \in \mathbb{N}\}$ a collection of open sets containing x such that for any open set U containing x , there exists $B_n \in \mathcal{B}$ such that $B_n \subseteq U$.

Definition 19.1.2. X is second countable if there exists a countable basis $\mathcal{B} = \{B_n : n \in \mathbb{N}\}$ for the topology of X . That is, for every x and every open set U containing x , there exists $B_n \in \mathcal{B}$ such that $x \in B_n \subseteq U$.

Exercise $\mathbb{R}^n$ with standard topology, $B_n = \{U_\epsilon(x) | x \in \mathbb{Q}^n, \epsilon \in \mathbb{Q}_{>0}\}$. B_n is a countable basis. So $\mathbb{R}^n$ is second countable. Show the details.

Exercise Show that if X_n 's are first(second) countable, then $\prod X_n$ with product topology is first(second) countable.

Theorem 19.1.1. Let X be second countable. Then

1. Every open cover of X has a countable subcovering (X is Lindelöf space).
2. There is a countable subset $A \subseteq X$ such that $\bar{A} = X$ ($A \subseteq X$ is called dense and X is called separable).

Proof to 1

Let $\mathcal{A}$ be an open cover of X . Let $\mathcal{B}$ be a countable basis for X . For each positive integer n for which it is possible, choose an element $A_n \in \mathcal{A}$ such that $B_n \subseteq A_n$. The collection $\mathcal{A}'$ of all such possible A_n is countable since it is indexed with a subset J of the positive integers.

Furthermore, $\mathcal{A}'$ covers X . To see this, let $x \in X$. Since $\mathcal{A}$ is a covering of X , there exists $A \in \mathcal{A}$ such that $x \in A$. Since $\mathcal{B}$ is a basis for the topology of X , there exists $B_m \in \mathcal{B}$ such that $x \in B_m \subseteq A$. By construction of $\mathcal{A}'$, there exists $A_m \in \mathcal{A}'$ such that $B_m \subseteq A_m$. Thus, $x \in A_m$. So $\mathcal{A}'$ is a countable subcovering of $\mathcal{A}$.

Proof to 2

From each non-empty basis element B_n , choose a point $x_n \in B_n$. Let D be the set of all such points x_n . Then D is countable.

Given any point $x \in X$, every basis element containing x contains a point of D , so x is in the closure of D . Thus, $\bar{D} = X$. ■

Property 19.1.1. A second countable space is:

1. First countable.
2. Separable.
3. Lindelöf.

Example 19.1.1 (non-example). $\mathbb{R}_l$ with lower limit topology (Sorgenfrey line) is first countable, Lindelöf, separable, but not second countable.

First-countable: for any $x \in \mathbb{R}_l$, $\{[x, x + \frac{1}{n}) : n \in \mathbb{N}\}$ is a countable local basis at x .

Assume the contrary that $\mathbb{R}_l$ is second countable. Let $\mathcal{B} = \{B_n : n \in \mathbb{N}\}$ be a countable basis for $\mathbb{R}_l$. For each $x \in \mathbb{R}$, there exists $B_{n_x} \in \mathcal{B}$ such that $x \in B_{n_x} \subseteq [x, x+1)$. So $\min B_{n_x} = x$. This implies that the map $f : \mathbb{R} \rightarrow \mathcal{B}, f(x) = B_{n_x}$ is injective. However, $\mathbb{R}$ is uncountable while $\mathcal{B}$ is countable, which is a contradiction. So $\mathbb{R}_l$ is not second countable.

Separable: $\mathbb{Q} \subseteq \mathbb{R}_l$ is countable and dense in $\mathbb{R}_l$.

Lindelöf: See the below property.

Property 19.1.2 (Lindelöf property of $\mathbb{R}_l$). Any open cover of $\mathbb{R}_l$ has a countable subcover.

Proof

Let $\mathcal{A}$ be a covering of $\mathbb{R}_l$ by $[a_\alpha, b_\alpha)$'s. We need to show that there exists a countable subcovering of $\mathcal{A}$ covering $\mathbb{R}_l$.

Let $C = \bigcup (a_\alpha, b_\alpha)$ such that $\mathbb{R} \setminus C$ is not empty. Let $x \in \mathbb{R} \setminus C$. Then $x = a_\beta$ for some β such that $x \notin (a_\beta, b_\beta)$. Take $q_x \in (a_\beta, b_\beta) \cap \mathbb{Q}$ thus $x < q_x$.

Take $x, y \in \mathbb{R}$ with $x < y$. Then $q_x < q_y$ or we make a contradiction(why?). The map $x \mapsto q_x$ is injective from $\mathbb{R} \setminus C$ to $\mathbb{Q}$. So $\mathbb{R} \setminus C$ is countable.

Now we show that some countable subcollection of $\mathcal{A}$ covers $\mathbb{R}$. To begin, choose for each element $\mathbb{R} \setminus C$ an interval $[a_\alpha, b_\alpha)$ of $\mathcal{A}$ that contains it. Then we have chosen a countable subcollection $\mathcal{A}'$ of $\mathcal{A}$ covering $\mathbb{R} \setminus C$.

Now take the set C and topologize it as a subspace of $\mathbb{R}$ (C is open in $\mathbb{R}$, topologizing means taking the intersection of C with open sets in $\mathbb{R}$ as the open sets of C). Then C is second countable(since $\mathbb{R}$ is second countable). Now C is covered by the open intervals (a_α, b_α) which are open in $\mathbb{R}$ and hence open in C . So there exists a countable subcovering of C by (a_α, b_α) 's. Suppose this subcollection is indexed by $\alpha = \alpha_1, \alpha_2, \dots$. Then the collection

$$\mathcal{A}'' = \{[a_\alpha, b_\alpha) : \alpha = \alpha_1, \alpha_2, \dots\} \quad (19.1)$$

is a countable subcollection of $\mathcal{A}$ that covers C .

Hence the countable collection $\mathcal{A}' \cup \mathcal{A}''$ is a countable subcollection of $\mathcal{A}$ that covers $\mathbb{R}_l$. ■

Property 19.1.3. $\mathbb{R}_l \times \mathbb{R}_l$ is not Lindelöf. The space $\mathbb{R}_l \times \mathbb{R}_l$ with the product topology is called the Sorgenfrey plane. (The proof can be shown by a picture not drawn here.)

Proof The space $\mathbb{R}_l^2$ has as basis all sets of the form $[a, b) \times [c, d)$ where $a < b$ and $c < d$. To show that $\mathbb{R}_l^2$ is not Lindelöf, we consider the subspace

$$L = \{(x, -x) | x \in \mathbb{R}_l\} \subseteq \mathbb{R}_l^2. \quad (19.2)$$

It is easy to check that L is closed in $\mathbb{R}_l^2$. Let's cover $\mathbb{R}_l^2$ by the open set $\mathbb{R}_l^2 \setminus L$ and by all basis elements of the form

$$[a, b) \times [-a, d) \quad (19.3)$$

Each of these open sets intersects L in at most one point. Since L is uncountable, no countable subcollection covers $\mathbb{R}_l^2$. Thus, $\mathbb{R}_l^2$ is not Lindelöf. ■

Example 19.1.2. Subspace of Lindelöf space is not necessarily Lindelöf. The ordered square I_0^2 is compact thus is Lindelöf. But the subspace $A = I_0 \times (0, 1)$ is not Lindelöf. For A is the union of disjoint sets $U_x = \{x\} \times (0, 1)$, each of which is open in A . This collection of sets $\{U_x | x \in I_0\}$ is uncountable, and no proper subcollection covers A .

19.2 Separation Axioms

Topological space X can satisfy the following separability axioms:

1. T_1 (Frechet): Given two distinct points $x, y \in X$, there exists an open set U such that $x \in U$ and $y \notin U$. (Equivalently, all singleton sets are closed. Easy exercise)
2. T_2 (Hausdorff): Given two distinct points $x, y \in X$, there exist open sets U, V such that $x \in U, y \in V$ and $U \cap V = \emptyset$.
3. T_3 (Regular): X is T_1 and given a closed set $F \subseteq X$ and a point $x \notin F$, there exist open sets U, V such that $x \in U, F \subseteq V$ and $U \cap V = \emptyset$.
4. T_4 (Normal): X is T_1 and given two disjoint closed sets $F_1, F_2 \subseteq X$, there exist open sets U, V such that $F_1 \subseteq U, F_2 \subseteq V$ and $U \cap V = \emptyset$.

Lemma 19.2.1. *Let X be a T_1 topological space. Then*

1. *X is regular if and only if for every point $x \in X$ and every neighborhood U of x , there exists a neighborhood V of x such that $\bar{V} \subseteq U$.*
2. *X is normal if and only if for every closed set $F \subseteq X$ and every neighborhood U of F , there exists a neighborhood V of F such that $\bar{V} \subseteq U$.*

Proof

$\Rightarrow$ Let X be regular.

If $U = X$, there is nothing to prove.

Suppose $U \neq X$. Then $F = X \setminus U$ is closed and $x \notin F$. By regularity of X , there exist open sets V, W such that $x \in V, F \subseteq W$ and $V \cap W = \emptyset$. Now we need to show $\bar{V} \subseteq U$.

If $y \in F$, then $y \in W$. Since $V \cap W = \emptyset$, $y \notin \bar{V}$ because we find a neighborhood W of y such that $W \cap V = \emptyset$. Thus, $\bar{V} \subseteq X \setminus F = U$.

$\Leftarrow$ Suppose that the point x and the closed set B not containing X are given. Let $U = X \setminus B$. By hypothesis, there is a neighborhood V of x such that $\bar{V} \subseteq U$. The open sets V and $X \setminus \bar{V}$ then separate x and B .

The proof of 2. is similar. ■

Theorem 19.2.1. 1. *X is Hausdorff, then $Y \subseteq X$ with subspace topology is Hausdorff.*

2. *X_α is Hausdorff, then $\prod X_\alpha$ with product topology is Hausdorff.*

3. *X is regular, then $Y \subseteq X$ with subspace topology is regular.*

4. *X_α is regular, then $\prod X_\alpha$ with product topology is regular.*

Proof

(a) is obvious.

(b) Let $x = (x_\alpha), y = (y_\alpha) \in \prod X_\alpha$ with $x \neq y$. Then there exists β such that $x_\beta \neq y_\beta$. Since X_β is Hausdorff, there exist open sets $U_\beta, V_\beta \subseteq X_\beta$ such that $x_\beta \in U_\beta, y_\beta \in V_\beta$ and $U_\beta \cap V_\beta = \emptyset$. Then $\pi_\beta^{-1}(U_\beta), \pi_\beta^{-1}(V_\beta)$ are open in $\prod X_\alpha$, $x \in \pi_\beta^{-1}(U_\beta), y \in \pi_\beta^{-1}(V_\beta)$ and $\pi_\beta^{-1}(U_\beta) \cap \pi_\beta^{-1}(V_\beta) = \emptyset$. So we are done.

(c) (It is easy to verify the subspace of T_1 space is T_1 .) Let $Y \subseteq X$ with subspace topology. Let $y \in Y$ and B be a closed set in Y such that $y \notin B$. Then there exists a closed set F in X such that $B = F \cap Y$. Since $y \notin B, y \notin F$. Since X is regular, there exist open sets $U, V \subseteq X$ such that $y \in U, F \subseteq V$ and $U \cap V = \emptyset$. Then $U \cap Y, V \cap Y$ are open in $Y, y \in U \cap Y, B \subseteq V \cap Y$ and $(U \cap Y) \cap (V \cap Y) = \emptyset$. So we are done.

(d) Let $\{X_\alpha\}$ be a family of regular spaces. Let $X = \prod X_\alpha$. X_α is regular implies that X_α is Hausdorff. By (b), X is Hausdorff thus is T_1 .

Let $x = (x_\alpha) \in X$ and let U be a neighborhood of x in X . Choose a basis element $\prod U_\alpha \subseteq U$ containing x . By the previous lemma, we can choose, for each α , a neighborhood V_α of x_α in X_α such that $\bar{V}_\alpha \subseteq U_\alpha$. If it happens that $U_\alpha = X_\alpha$ for some α , we choose $V_\alpha = X_\alpha$. Then $V = \prod V_\alpha$ is a neighborhood of x in X . Recall from lecture 11, that $\bar{V} = \prod \bar{V}_\alpha$. It follows at once that $\bar{V} \subseteq \prod U_\alpha \subseteq U$. By the previous lemma, X is regular. ■

Remark X is normal does not imply that $Y \subseteq X$ with subspace topology is normal. X_1, X_2 are normal does not imply that $X_1 \times X_2$ with product topology is normal.

Exercise Show that $\mathbb{R}_l$ is normal.

Solution It is immediate that one-point sets are closed in $\mathbb{R}_l$, since the topology of $\mathbb{R}_l$ is finer than the standard topology on $\mathbb{R}$. Let A and B are disjoint closed sets in $\mathbb{R}_l$. For each $a \in A$, since B is closed and $a \notin B$, there exists a basis element $[a, x_a)$ such that $[a, x_a) \cap B = \emptyset$. For each $b \in B$, since A is closed and $b \notin A$, there exists a basis element $[b, y_b)$ such that $[b, y_b) \cap A = \emptyset$. Then the open sets

$$U = \bigcup_{a \in A} [a, x_a), \quad V = \bigcup_{b \in B} [b, y_b) \quad (19.4)$$

are disjoint open sets containing A and B respectively. So we are done. ■

Fact (Without Proof, Efforts are required to show this) $\mathbb{R}_l^2$ is not normal. But it is regular (as a product of regular spaces).

Exercise $\mathbb{R}_K$ is a topological space with basis given by all open intervals (a, b) and all sets of the form $(a, b) \setminus K$ where $K = \{1/n \mid n \in \mathbb{N}\}$. $\mathbb{R}_K$ is Hausdorff. But $\mathbb{R}_K$ is not regular.

The set K is closed in $\mathbb{R}_K$ and it does not contain the point 0. Suppose that there exist disjoint open sets U and V such that $0 \in U$ and $K \subseteq V$. Choose a basis element containing 0 and lying in U . It must be a basis

element of the form $(a, b) \setminus K$, since each basis element of the form (a, b) contains intersects K . Choose $1/n \in K$ such that $1/n < b$. Then there exists a basis element of the form (c, d) such that $1/n \in (c, d) \subseteq V$. Finally, the basis elements $(a, b) \setminus K$ and (c, d) intersect, since $c < 1/n < b$. So U and V are not disjoint. ■