A Conditional Reconstruction Program for Relativity and Standard Model Structure from Observer-Overlap Consistency

Observers Are All You Need (Compact)

B. Müller

March 13, 2026

Paper release: `r1337` **Released:** March 13, 2026

Abstract

We present a falsifiable compact formulation of Observer-Patch Holography (OPH) as a conditional reconstruction program. The framework starts from five axioms, two external configuration inputs in the current implementation, and an explicit scaling-limit/categorical regularity package. From that data it derives a schedule-independent normal form for overlap consistency, a conditional geometric-modular Lorentz branch on the screen, a conditional Jacobson-type Einstein branch from generalized entropy stationarity and the null modular bridge, and conditional compact gauge reconstruction in the bosonic internal-gauge branch from a refinement-stable symmetric edge-sector colimit. After the extended MAR selector is imposed on the connected positive-dimensional Lie admissible class with one connected abelian charge factor, the realized low-energy gauge group is

$$\frac{\mathrm{SU}(3)\times\mathrm{SU}(2)\times\mathrm{U}(1)}{\mathbb{Z}_6}, \qquad N_c = 3, \qquad N_g = 3,$$

together with exact rational hypercharges. The two external inputs of the present implementation are the pixel area

$$P \equiv a_{\mathrm{cell}}/\ell_P^2 = 1.63094$$

and the total screen capacity

$$N_{\mathrm{scr}} \equiv \log\dim\mathcal{H}_{\mathrm{tot}} \sim 10^{122}.$$

In the current implementation, P is calibrated from the gauge sector and N_{scr} is inferred from the observed cosmological constant. Gauge couplings and tree-level electroweak masses are calibration-sector consistency checks, while the critical-surface Higgs/top branch is logically independent of that calibration. The charged-lepton, Koide, quark-texture, baryogenesis, dark-sector, string, and black-hole-spectroscopy discussions are retained only as phenomenological continuations or conjectural branches, not as theorem-level outputs of overlap consistency alone. The aim of this compact note is therefore not to claim a complete derivation of all Standard Model and cosmological data from observer-overlap consistency by itself, but to state clearly which reconstruction results are structural, which are scaling-limit conditional, which are calibration-sector, and which remain phenomenological.

1 Introduction

The standard model of particle physics and general relativity are successful effective descriptions. They leave open a familiar list of questions: why the Lorentz group appears, why the low-energy spacetime description follows a Jacobson-type Einstein branch, why the gauge structure is

$$\frac{\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)}{\mathbb{Z}_6},$$

why there are three colors and three generations, why hypercharges take the observed rational values, why the electroweak and flavor scales are what they are, and why dark-sector phenomenology and the cosmological constant enter in the particular form observed.

This paper presents a single framework that addresses this list as a conditional reconstruction program. The framework is Observer-Patch Holography (OPH). Its starting point is an observer-centric organization of physical data: each observer has access only to a patch algebra, neighboring observers must agree on overlaps, and global physical law is recovered from the resulting consistency conditions. The same formal structure that produces schedule-independent overlap consistency then feeds a Lorentzian modular branch, a Jacobson-type Einstein branch, compact gauge reconstruction in a bosonic internal-gauge sector, charge quantization, and the low-energy architecture selected by the extended MAR admissibility package.

Two decisions shape the present version.

1. The compact note presents the main theorem statements in one place, but some quantitative conventions and implementation details live in repository supplements.[1]

2. The paper separates exact structural theorems, conditional scaling-limit branches, calibration-sector outputs, and phenomenological continuations. This keeps strong claims visible and weak claims identifiable.

The framework is intended as a candidate fundamental theory in the precise sense relevant for physics: general relativity and the Standard Model appear as effective sectors derived from a smaller axiom set plus an explicit refinement/scaling-limit package. The paper does not use simulation language, observer continuation stories, or other non-falsifiable additions.

Prioritized achievements

The results emphasized in this paper are ordered below by a combination of impact and defensibility.

1. A conditional Jacobson-type Einstein branch is recovered from observer-overlap modular geometry, generalized entropy stationarity, and an explicit null-modular scaling-limit package.

2. The exact Standard Model gauge structure

$$\frac{\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)}{\mathbb{Z}_6}$$

is recovered in the bosonic compact-gauge branch once the extended MAR admissibility package is imposed.

[1] Questions about ultimate purpose are deferred until the corresponding physical question has been posed with enough precision to deserve an answer.

3. Hypercharges, three colors, and three generations are fixed only after anomaly constraints, admissibility, and MAR are imposed on that gauge branch.

4. The current implementation reduces the continuous input set to two external configuration inputs, P and N_{scr}.

5. The Higgs/top critical-surface branch is quantitatively independent of the gauge-sector calibration, while the charged-lepton/Koide continuation uses extra phenomenological structure beyond the core theorem package.

6. The cosmological constant is separated from local vacuum energy and tied to global screen capacity.

7. Absence of gauge-mediated proton decay follows from product-group structure.

8. Dark-sector, baryogenesis, string, and spectroscopy branches are kept visible, but only as program-level or conjectural phenomenology.

9. Objective physical law is identified with the unique normal form of the overlap-consistency problem.

2 Results at a Glance

The following table lists the principal results emphasized in this paper.

Table 1: Principal achievements of the framework.

Problem	Result	Logical status	Validation tier	Mechanism
Objectivity of law	Unique schedule-independent normal form	theorem	structural theorem	Asynchronous confluence of overlap repair
Lorentz kinematics	$\mathrm{Conf}^+(S^2) \cong \mathrm{SO}^+(3,1)$	conditional theorem	scaling-limit theorem	Geometric modular flow on caps
Einstein dynamics	Jacobson-type first-variation relation with tensor upgrade	theorem + corollary	conditional scaling-limit theorem	Entanglement equilibrium and null modular bridge
Gauge structure	$(\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1))/\Gamma$, fixed later to $\mathbb{Z}_6$	theorem/corollary chain	conditional structural theorem	Refinement-stable bosonic edge-sector reconstruction plus extended MAR admissibility
Charge lattice	Exact Standard Model hypercharges	theorem	structural theorem	Anomaly cancellation and Yukawa invariance
Discrete matter content	$N_c = 3$, $N_g = 3$ after MAR	corollary chain	structural theorem	CKM CP capability, anomaly constraints, admissibility
Parameter economy	Two external configuration inputs in the current implementation	structural result	structural theorem	Pixel area and screen capacity only
Unification sector	MSSM-like coupling evolution without light superpartners	derived reduction	calibration sector	Edge-sector beta-function shifts

Table 1: Principal achievements of the framework.

Problem	Result	Logical status	Validation tier	Mechanism
Electroweak scale	v, m_W, m_Z, gauge couplings	derived reduction	calibration sector	Pixel constraint and self-consistent fixed point
Higgs and top	$m_H \approx 126.5$ GeV, $m_t \approx 171.1$ GeV	derived reduction	independent quantitative	Critical surface $\lambda = \beta_\lambda = 0$ at M_U
Charged leptons and Koide	$Q = 2/3$, $\delta = 2/9$, sub-permille lepton masses	semi-empirical continuation	phenomenological continuation	$\mathbb{Z}_3$ holonomy plus a $\mathbb{Z}_6$-motivated suppression ansatz
Cosmological constant from capacity	$\Lambda = 3\pi/(GN_{\rm scr})$	corollary	derived reduction	Global screen-capacity identification
Neutrino scale from $N_{\rm scr}$	capacity-branch neutrino estimate	derived reduction	order-of-magnitude estimate	Global screen capacity
Dark matter phenomenology	MOND-like acceleration scale from modular anomaly	program corollary	program-level phenomenology	Imperfect Markovity in collar modular flow
Baryogenesis scale	$\varepsilon^{12} = 6^{-12}$ in the observed range	program corollary	program-level phenomenology	$\mathbb{Z}_6$ suppression and electroweak topology
Edge-sector fingerprints	conditional $\mathbb{Z}_6$ center-label entropy deficit and exact SU(3) Casimir ratios	program corollary	program-level phenomenology	Quotient-label ansatz and heat-kernel spectrum
Black-hole spectroscopy	Finite-window logarithmic horizon template	conditional phenomenology	program-level phenomenology	Discrete horizon area counting
String-theory sector	2D Yang-Mills and worldsheet expansion from edge heat kernel	effective-description corollary	program-level phenomenology	Open-edge weights and large-$N_{\rm edge}$ reorganization
No gauge-mediated proton decay and no GUT monopoles	$\tau_p^{(\rm gauge)} = \infty$ and no simple-group breaking sector	corollary	structural theorem	Product gauge structure, no X/Y bosons, no simple-group breaking

The same OPH framework is used to obtain the overlap fixed point, the gravity chain, the gauge reconstruction, and the quantitative spectrum, but not all outputs have the same logical status. The compact note therefore keeps structural theorems, scaling-limit branches, calibration checks, and phenomenological continuations distinct.

3 Five Axioms, Two External Inputs, and Technical Premises

The formulation uses exactly two external configuration inputs:

$$P \equiv a_{\rm cell}/\ell_P^2 = 1.63094, \tag{1}$$

$$N_{\rm scr} \equiv \log \dim \mathcal{H}_{\rm tot} \sim 10^{122}. \tag{2}$$

In the current implementation, P is calibrated from the gauge sector and $N_{\rm scr}$ is inferred from the observed cosmological constant. The first sets the particle-physics scale. The second enters the cosmological-capacity branch and a separate order-of-magnitude neutrino estimate.

Axiom 3.1 (Screen Net)**.** *Physical data is organized on a horizon screen S^2 carrying a net of local algebras*

$$P \longmapsto \mathcal{A}(P)$$

for connected patches $P \subset S^2$, with isotony

$$P \subset Q \implies \mathcal{A}(P) \subset \mathcal{A}(Q).$$

Axiom 3.2 (Overlap Consistency)**.** *For overlapping patches $P_1 \cap P_2 \neq \varnothing$, the local states induced on the shared algebra agree:*

$$\omega_{P_1}|_{\mathcal{A}(P_1 \cap P_2)} = \omega_{P_2}|_{\mathcal{A}(P_1 \cap P_2)}.$$

Axiom 3.3 (Local MaxEnt and Refinement Stability)**.** *At the UV scale the realized state maximizes entropy subject to a finite family of gauge-invariant local constraints. The resulting family of states is stable under coarse-graining, so symmetry-allowed relevant operators are not held at zero by unexplained fine tuning.*

Axiom 3.4 (Recoverable Generalized Entropy)**.** *A generalized entropy functional exists on caps,*

$$S_{\mathrm{gen}}(C) = S_{\mathrm{bulk}}(C) + \langle L_C \rangle,$$

where L_C is a positive edge-center entropy functional. In the semiclassical scaling branch, its leading coarse-grained contribution is identified with $A(\partial C)/(4G)$. The functional obeys quantum focusing on null generators, and collar tripartitions have small conditional mutual information with controlled recovery maps [3, 4, 5, 6].

Axiom 3.5 (Minimal Admissible Realization)**.** *Among realized sector packages $\mathfrak{S}$ consisting of the connected Lie gauge-sector image relevant in the low-energy EFT, its admissible light chiral matter content, and one Higgs doublet, and which are loop-coherent, anomaly-free, refinement-stable with light chiral matter, single-Higgs Yukawa-completable with one connected abelian charge factor acting nontrivially on the coupled carrier, intrinsically CP-capable, and weak-sector UV-completable, the realized package is the lexicographically minimal one under*

$$C(\mathfrak{S}) = (\chi_{\mathrm{cpl}},\, N_{\mathrm{nonab}},\, N_c,\, N_g).$$

$\mathfrak{S}$ is the sector package on which MAR acts; it is not the bare tensor category alone. MAR is therefore an explicit structural-economy axiom on already-admissible realized low-energy branches, not a theorem derived from the preceding axioms. χ_{cpl} is the coupled carrier dimension: the smallest unitary carrier containing a common irreducible block on which the admissible pseudoreal and complex nonabelian charge types both act nontrivially. This is stronger than the abstract minimal faithful representation dimension.

Assumption 3.6 (Technical Premises)**.** *The continuum reductions additionally use:*

- ***T1*** *type-I local algebras at the regulator scale and boundary gauge invariance on collars;*
- ***T2*** *finite-velocity quasi-local propagation at the UV scale;*
- ***T3*** *quantitative collar-mixing / endpoint-Lipschitz control for null-interval modular Hamiltonians on bounded intervals;*
- ***T4*** *cap isotropy / geometric modular action around the entangling circle in the refinement-scaling regime;*

T5 *Euclidean modular regularity, fixing the angular period to* 2π*;*

T6 *a standard half-sided modular inclusion premise for the null half-line blow-up algebras;*

T7 *vanishing gluing obstruction* $[z] = 0$ *whenever global sector transportability is invoked;*

T8 *all Lorentz/null-modular/Einstein statements are claims about a controlled refinement/scaling limit of the regulator net, not literal exactness at fixed finite cutoff;*

T9 *symmetric braiding in the* $3+1D$ *EFT branch;*

T10 *a bosonic Tannakian fiber-functor premise, or else an explicit super-Tannakian fork;*

T11 *whenever compact gauge reconstruction is invoked, the sector category means the refinement-stable directed colimit of transportable edge sectors with objectwise finite-dimensional fibers, not the block list of one fixed cutoff algebra.*

Remark 3.7. *The five axioms define the physical content of the formulation. Assumption 3.6 collects the regularity conditions needed to express the continuum limit precisely. In particular, the Lorentz/null-modular/Einstein and compact-gauge statements below are scaling-limit claims, not fixed-cutoff matrix-algebra identities. These premises are not derived in the compact note; they are target regularity properties of the intended Lorentzian EFT phase and candidate UV completions.*

Dependency Ledger

Claim	Status	Additional ingredients beyond the five axioms
Unique schedule-independent normal form	theorem	finite patch net, termination, local confluence
Lorentz group on the screen net	conditional scaling-limit theorem	cap isotropy / geometric modular phase together with Euclidean modular regularity in Assumption 3.6
Einstein equation	conditional scaling-limit theorem	admissible first variations, exact-or-controlled null-strip Markov premise, half-sided modular inclusion premise or its scaling-limit blow-up corollary, localized generalized entropy, scaling-limit null-stress identification, null-to-tensor upgrade
Compact gauge reconstruction	conditional scaling-limit theorem	refinement-stable bosonic symmetric sector colimit, transportability, finite-fiber premises, symmetric braiding, and bosonic fiber functor in Assumption 3.6
Exact Standard Model quotient	theorem / corollary chain	MAR selector, connected abelian admissibility premise, anomaly analysis, hypercharge lattice, global transportability, and vanishing obstruction on the realized branch
Two-input parameter economy	structural result	no extra ingredients, but the inputs are external in the current implementation
Gauge and electroweak closure in one-input mode	calibration sector	pixel constraint and edge heat-kernel closure
Higgs/top branch and flavor continuations	mixed quantitative / phenomenological	critical-surface boundary condition for Higgs/top; discrete holonomy data and a $\mathbb{Z}_6$-motivated ansatz for the flavor continuation
Dark-sector, baryogenesis, and spectroscopy branches	program / corollary	sector-specific phenomenological continuation of the core structure

Theorem 3.8 (Master reconstruction schema)**.** *Assume Axioms 3.1–3.4, Axiom 3.5, Assumption 3.6, and the hypotheses of Theorems 4.3, 6.1, 6.10, 6.15, 7.1, 7.7, and 7.8, together with Corollaries 7.9, 7.10, and Proposition 7.11. Then, in the controlled refinement/scaling regime singled out by Assumption 3.6:*

(i) overlap repair admits a unique schedule-independent normal form;

(ii) cap modular flow is geometric and yields the connected Lorentz group

$$\mathrm{Conf}^+(S^2) \cong \mathrm{SO}^+(3,1);$$

(iii) admissible entanglement-equilibrium first variations together with the null modular bridge yield a Jacobson-type rest-frame relation, whose tensor upgrade gives

$$G_{ab} + \Lambda g_{ab} = 8\pi G \langle T_{ab} \rangle;$$

(iv) the refinement-stable transportable edge-sector category reconstructs a compact gauge group, and the realized connected gauge structure has the form

$$\frac{\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)}{\Gamma}$$

for some finite central subgroup Γ.

After the hypercharge lattice, generation-count, color-count, and trivial-action quotient steps, the finite quotient is fixed to $\Gamma = \mathbb{Z}_6$*, so the realized gauge structure is*

$$\frac{\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)}{\mathbb{Z}_6}, \qquad N_c = 3, \qquad N_g = 3,$$

with the exact Standard Model hypercharge lattice.

Proof. Items (i)–(iii) are collected from Theorem 4.3, Corollary 6.2, Theorem 6.15, and Corollary 6.16. The compact gauge reconstruction is Theorem 7.1; the product gauge structure up to finite quotient is Theorem 7.7; and the final identification of the exact Standard Model quotient is Theorem 7.8 together with Corollaries 7.9, 7.10, and Proposition 7.11. □

Usually Postulated Elsewhere, Derived Here

Structure	Common treatment	OPH treatment
Lorentz kinematics	background symmetry or starting axiom	conditional scaling-limit branch from geometric modular flow on screen caps
Einstein dynamics	fundamental field equation	conditional first-variation relation with tensor upgrade from generalized entropy, null modular data, and entanglement equilibrium
Gauge group	model input	conditional compact group reconstruction from edge sectors; exact SM quotient selected by admissibility
Hypercharge lattice	matter-assignment input	solved from anomaly cancellation and Yukawa invariance
Color and generation count	empirical input	fixed by anomaly, CP capability, asymptotic freedom, and minimality
Flavor hierarchy unit	model-dependent small parameter	phenomenological ansatz $\varepsilon = 1/6$ motivated by a uniform $\mathbb{Z}_6$ center-label ensemble
Koide phase	phenomenological fit	continuation tied to the previously fixed discrete data as $\delta = 2/9$
Cosmological constant	local vacuum-energy puzzle	global screen-capacity term

4 Observer-Overlap Consistency as a Fixed-Point Problem

The discrete overlap problem is the mathematical skeleton of the framework.

Definition 4.1 (Patch Net)**.** *Let* $G = (V, E)$ *be a finite connected graph. Each vertex* $i \in V$ *carries a finite local state space* S_i*. For each edge* $e = \{i, j\}$ *let* I_e *be an interface alphabet and let*

$$\pi_{i,e} : S_i \to I_e, \qquad \pi_{j,e} : S_j \to I_e$$

be interface projections. The global state space is

$$\Sigma = \prod_{i \in V} S_i,$$

and the consistency set is

$$C = \{s \in \Sigma : \pi_{i,e}(s_i) = \pi_{j,e}(s_j) \text{ for all } e = \{i, j\} \in E\}.$$

This finiteness is a regulator-level assumption for the discrete patch-net normal-form theorem; it is not the continuum limit statement.

Define the inconsistency potential

$$\Phi(s) = \sum_{e=\{i,j\}\in E} w_e \, d_e\big(\pi_{i,e}(s_i), \pi_{j,e}(s_j)\big),$$

with $w_e > 0$ and $d_e(a, b) = 0$ iff $a = b$.

Definition 4.2 (Local Repair Law)**.** *A law λ is a family of local repair maps*

$$T_i^\lambda : \Sigma \to \Sigma$$

changing only patch i or a bounded neighborhood of i.

Theorem 4.3 (Asynchronous Fixed-Point Law)**.** *Assume:*

(i) normal forms of the repair relation are exactly the globally consistent states C;

(ii) the repair relation terminates;

(iii) the repair relation is locally confluent.

Then every $s \in \Sigma$ has a unique normal form

$$\mathrm{nf}_\lambda(s) \in C,$$

and every fair asynchronous repair schedule terminates at that same state. A sufficient condition for (ii) is that each S_i is finite and every enabled repair strictly decreases Φ.

Proof. Termination is assumed. Local confluence plus termination imply confluence by Newman's lemma. The resulting normal form is unique. By assumption it is a globally consistent state. □

Corollary 4.4 (Schedule Independence)**.** *If observables are evaluated on $\mathrm{nf}_\lambda(s)$, the resulting physical law is independent of update order.*

Theorem 4.5 (Cycle Obstruction / Holonomy Criterion)**.** *Let A be an abelian group. For each oriented edge $e : u \to v$ assign $b_e \in A$ and consider the affine overlap equations*

$$x_v - x_u = b_e.$$

A global solution exists if and only if the signed sum of b_e vanishes on every cycle.

Proof. Summing the edge equations around a cycle gives necessity. For sufficiency, fix a root and define each x_v by summing labels along a path from the root to v. Vanishing cycle sums make this independent of the chosen path. □

Proposition 4.6 (Gauge Quotient)**.** *Suppose a local gauge group $\Gamma = \prod_i \Gamma_i$ acts on Σ while leaving all interface data invariant, and the repair maps satisfy*

$$T_i^\lambda(\gamma \cdot s) = \gamma \cdot T_i^\lambda(s).$$

Then

$$\mathrm{nf}_\lambda(\gamma \cdot s) = \gamma \cdot \mathrm{nf}_\lambda(s).$$

Hence gauge-invariant observables are unique on the quotient.

Proof. Gauge covariance maps repair sequences to repair sequences. Since the normal form is unique by Theorem 4.3, the normal form of $\gamma \cdot s$ must be $\gamma \cdot \mathrm{nf}_\lambda(s)$. □

These three results already encode much of the eventual physical interpretation. Objectivity becomes confluence, gauge symmetry becomes quotient invariance, and stable defects are represented by nontrivial overlap holonomy classes.

5 Collars, Edge Centers, and Generalized Entropy

The fixed-point picture explains the logic of overlap consistency. The continuum theory requires its algebraic lift. In OPH the relevant object is a collar neighborhood of an entangling cut, together with the center data carried by that collar.

Theorem 5.1 (Exact quantum Markov collar decomposition)**.** *Let A-B-D be a collar tripartition with*

$$I(A:D|B)_\rho = 0.$$

Then the Hilbert space of the collar factorizes as

$$\mathcal{H}_B \cong \bigoplus_\alpha \mathcal{H}_{b_L^\alpha} \otimes \mathcal{H}_{b_R^\alpha},$$

and the state decomposes as

$$\rho_{ABD} = \bigoplus_\alpha p_\alpha \, \rho^{(\alpha)}_{Ab_L^\alpha} \otimes \rho^{(\alpha)}_{b_R^\alpha D}.$$

The label α is classical center data carried by the cut.

Proof. This is the standard structure theorem for exact quantum Markov states on tripartite systems [7]. In the present framework it is applied to collar neighborhoods produced by edge-center completion and recoverability. □

Proposition 5.2 (Approximate collar recovery)**.** *Let A-B-D be a collar tripartition satisfying*

$$I(A:D|B)_\rho \le \varepsilon,$$

and let $\mathcal{R}_{B\to BD}$ be a recovery map such that

$$F(\rho_{ABD}, (\mathrm{id}_A \otimes \mathcal{R}_{B\to BD})(\rho_{AB})) \ge 2^{-\varepsilon/2}.$$

Define the recovered state

$$\rho^{\mathrm{rec}}_{ABD} := (\mathrm{id}_A \otimes \mathcal{R}_{B\to BD})(\rho_{AB}).$$

Then every bounded observable X supported on $A \cup B$ obeys

$$|\mathrm{Tr}[X(\rho_{ABD} - \rho^{\mathrm{rec}}_{ABD})]| \le \|X\|_\infty \, \|\rho_{ABD} - \rho^{\mathrm{rec}}_{ABD}\|_1 \le 2\|X\|_\infty \sqrt{1-2^{-\varepsilon}}.$$

Proof. The first inequality is the trace-norm bound for bounded observables. The second is the Fuchs–van de Graaf estimate applied to the fidelity lower bound supplied by recoverability [6]. Thus small conditional mutual information controls deviations from the exact Markov splice identities by $O(\varepsilon^{1/2})$. □

Remark 5.3. *Theorems 5.1 and Corollary 6.4 below use exact Markovity or an explicitly stated idealized recoverability limit. Axiom 3.4 itself supplies only approximate recoverability. The compact note therefore separates exact identities from their controlled $O(\varepsilon^{1/2})$ approximations rather than silently upgrading one regime into the other.*

Proposition 5.4 (Exact Markov-collar splice theorem)**.** *Under Theorem 5.1, let $\sigma^{(\alpha)}_{b_R^\alpha D'}$ be any normalized family of states compatible with the same right-boundary sectors, and define*

$$\rho'_{ABD'} = \bigoplus_\alpha p_\alpha \, \rho^{(\alpha)}_{Ab_L^\alpha} \otimes \sigma^{(\alpha)}_{b_R^\alpha D'}.$$

Then every observable X supported on $A \cup b_L$ has the same expectation value in ρ_{ABD} and $\rho'_{ABD'}$:

$$\mathrm{Tr}(X\rho'_{ABD'}) = \mathrm{Tr}(X\rho_{ABD}).$$

Proof. Blockwise factorization gives

$$\mathrm{Tr}(X\rho'_{ABD'}) = \sum_\alpha p_\alpha \mathrm{Tr}\Big(X\rho^{(\alpha)}_{Ab^\alpha_L}\Big)\,\mathrm{Tr}\Big(\sigma^{(\alpha)}_{b^\alpha_R D'}\Big).$$

Each $\sigma^{(\alpha)}$ is normalized, so the second factor is 1. The same computation for ρ_{ABD} gives the same value because the original right factor is likewise normalized. □

Proposition 5.5 (Generalized-entropy split)**.** *If the reduced cap state takes the block form*

$$\rho_C = \bigoplus_\alpha p_\alpha \left(\rho^{(\alpha)}_{\mathrm{bulk},C} \otimes \frac{\mathbf{1}^{(\alpha)}_{\mathrm{edge}}}{d_\alpha}\right),$$

then

$$S(\rho_C) = S_{\mathrm{bulk}}(C) + \mathrm{Tr}(\rho_C L_C),$$

where

$$S_{\mathrm{bulk}}(C) = H(p_\alpha) + \sum_\alpha p_\alpha S(\rho^{(\alpha)}_{\mathrm{bulk},C}), \qquad L_C = \sum_\alpha (\log d_\alpha) P_\alpha.$$

Proof. The entropy of a direct sum is

$$S\left(\bigoplus_\alpha p_\alpha \sigma_\alpha\right) = H(p_\alpha) + \sum_\alpha p_\alpha S(\sigma_\alpha).$$

Apply this identity with

$$\sigma_\alpha = \rho^{(\alpha)}_{\mathrm{bulk},C} \otimes \frac{\mathbf{1}^{(\alpha)}_{\mathrm{edge}}}{d_\alpha}.$$

Since

$$S(\sigma_\alpha) = S(\rho^{(\alpha)}_{\mathrm{bulk},C}) + \log d_\alpha,$$

the first statement follows. □

Definition 5.6 (Effective Newton constant in the refinement dictionary)**.** *In the refinement-scaling regime, if*

$$\mathrm{Tr}(\rho_C L_C) \approx N_\Sigma \bar{\ell}(t), \qquad A(\partial C) \approx N_\Sigma a_{\mathrm{cell}},$$

then matching $\mathrm{Tr}(\rho_C L_C)$ *to the area term* $A(\partial C)/(4G)$ *defines the effective Newton constant*

$$G := \frac{a_{\mathrm{cell}}}{4\bar{\ell}(t)}.$$

This is a continuum dictionary identification, not part of the exact algebraic proposition.

Proposition 5.4 is the algebraic version of interior invariance under compatible exterior substitutions. Proposition 5.5 isolates the exact entropy split, while Definition 5.6 supplies the continuum identification that turns the edge center into a gravitational coupling.

6 Modular Geometry, Relativity, and Einstein Dynamics

6.1 Geometric modular flow

The continuum interpretation replaces the discrete patch net by cap algebras on S^2. Recoverability localizes the modular defect to an arbitrarily thin collar around the entangling surface.

Theorem 6.1 (Conditional geometric modular flow on caps)**.** *Assume Axioms 3.1–3.4 and Assumption 3.6, including cap isotropy / geometric modular action around the entangling circle and Euclidean regularity. Then in the refinement/scaling limit of the cap net, for each cap $C \subset S^2$, the modular flow is the unique cap-preserving conformal dilation with KMS normalization 2π:*

$$K_C = 2\pi B_C.$$

Proof sketch. Markov locality restricts the generator to the collar in the refinement/scaling regime. The assumed geometric modular phase together with cap isotropy leaves a unique noncompact one-parameter subgroup compatible with the cap. Euclidean regularity fixes the normalization of that subgroup to 2π, as in the Bisognano-Wichmann modular geometry of wedge algebras [1, 2]. □

Corollary 6.2 (Lorentz Kinematics)**.** *Under the hypotheses of Theorem 6.1,*

$$\mathrm{Conf}^+(S^2) \cong \mathrm{PSL}(2,\mathbb{C}) \cong \mathrm{SO}^+(3,1).$$

The induced local kinematic group is the connected Lorentz group.

6.2 The null modular bridge

The gravity chain requires a bridge from modular data to a local stress tensor. In the compact note this bridge is pursued within the OPH framework, but the final null-stress identification is still a scaling-limit ingredient tied to Assumption 3.6; it is not presented as a fixed-cutoff matrix-algebra theorem.

For a small entangling circle $\Sigma = \partial C$, let $\mathcal{N}$ be the null surface generated by orthogonal null geodesics. Label generators by angle Ω and affine parameter v.

The derivation has five steps:

1. null-strip edge-center completion;
2. modular additivity on consecutive null intervals;
3. weak continuity and finite variation of interval modular Hamiltonians;
4. emergence of a positive null translation generator from half-sided modular inclusion;
5. scaling-limit identification of the renormalized modular Hamiltonian with a null stress-tensor charge.

Only the last step uses continuum stress-tensor language. The previous steps refer directly to the overlap algebra and its regulated collar structure.

Proposition 6.3 (Null Edge-Center Completion)**.** *Under the regulator premises in Assumption 3.6, a regulated null buffer strip has the same central block decomposition as an ordinary spatial collar. If the reference/MaxEnt state on consecutive strips satisfies exact Markovity*

$$I_\omega(I_- : I_+|J) = 0,$$

or the explicitly assumed idealized recoverability limit that reduces to it, then the strip modular data satisfy the same state-level factorization used below.

Proof sketch. At fixed regulator scale, a cut at constant affine parameter v is a boundary for the strip region in exactly the same kinematic sense in which ∂C is a boundary for a spatial collar. Type-I factorization at the regulator and boundary gauge invariance therefore imply a block decomposition

$$\mathcal{H}_J \cong \bigoplus_{\alpha_1,\alpha_2} \mathcal{H}_{j_L^{\alpha_1,\alpha_2}} \otimes \mathcal{H}_{j_R^{\alpha_1,\alpha_2}},$$

with the center generated by the block projectors. Within each block, observables in $I_- \cup J$ act only on the left factor and observables in $J \cup I_+$ act only on the right factor. This is the null analogue of the ordinary edge-center completion argument. The additional Markov statement is a condition on the reference state, not a consequence of the block decomposition alone. □

Corollary 6.4 (Null Modular Additivity)**.** *Under Proposition 6.3 and the exact Markovity condition*

$$I_\omega(I_- : I_+|J) = 0,$$

$$K_{I_- \cup J \cup I_+} = K_{I_- \cup J} + K_{J \cup I_+} - K_J$$

up to central boundary terms.

Proof. For a Markov state, the modular Hamiltonian satisfies the additive identity above blockwise. The boundary term is central because it depends only on the strip-edge labels. □

Proposition 6.5 (Weak Continuity of Null Modular Hamiltonians)**.** *Under Axiom 3.3, Assumption 3.6**T2**, and Assumption 3.6**T3**, matrix elements of $K[I]$ obey the interval-length domination estimate*

$$\left|\langle \psi, (K[I'] - K[I])\phi\rangle\right| \le C_{\psi,\phi}\, |I' \Delta I|$$

on bounded intervals. In particular they vary continuously with interval endpoints and have finite variation there.

Proof sketch. Local MaxEnt gives a quasi-local Gibbs form for the regulated state. Quasi-locality together with finite-velocity propagation implies that moving an interval endpoint by Δv changes expectation values of local observables by $O(\Delta v)$ after smearing, while Assumption 3.6**T3** supplies the required endpoint-Lipschitz control. The resulting matrix elements are therefore Lipschitz on bounded intervals and in particular have finite variation. □

Definition 6.6 (Renormalized null modular functional)**.** *For a null interval I, write*

$$\widetilde{K}[I] := K[I] - K_\partial(I)$$

for the full modular Hamiltonian with the central endpoint-label term removed.

Theorem 6.7 (Half-line endpoint-differentiation formula)**.** *Fix a null generator Ω and define*

$$\widetilde{K}_a(\Omega) := \widetilde{K}[(a,\infty),\Omega].$$

Assume that for fixed vectors ψ, ϕ in a common dense domain the function

$$f_{\psi,\phi}(a) := \langle \psi, \widetilde{K}_a(\Omega)\phi\rangle$$

is twice weakly differentiable in a with

$$\lim_{a\to+\infty} f_{\psi,\phi}(a) = 0, \qquad \lim_{a\to+\infty} \partial_a f_{\psi,\phi}(a) = 0.$$

Then there exists a sesquilinear-form density $p(v,\Omega)$ such that

$$\langle \psi, \widetilde{K}_a(\Omega)\phi\rangle = 2\pi \int_a^\infty (v-a)\,\langle \psi, p(v,\Omega)\phi\rangle\, dv,$$

and distributionally

$$\langle \psi, p(a,\Omega)\phi\rangle = \frac{1}{2\pi}\,\partial_a^2 f_{\psi,\phi}(a).$$

Proof. Define the matrix elements of p distributionally by

$$\langle \psi, p(a,\Omega)\phi\rangle := \frac{1}{2\pi}\,\partial_a^2 f_{\psi,\phi}(a).$$

Integrating twice from a to $+\infty$ and using the assumed vanishing of $f_{\psi,\phi}$ and its first derivative at infinity yields

$$f_{\psi,\phi}(a) = 2\pi \int_a^\infty (v-a)\,\langle \psi, p(v,\Omega)\phi\rangle\, dv.$$

This is the desired half-line formula. Unlike a sigma-additive bounded-interval measure statement, it is compatible with later interval-dependent kernels on finite null intervals. □

Corollary 6.8 (Half-Sided Modular Inclusion)**.** *Assume in addition the standard half-sided modular inclusion premise for the null half-line blow-up algebras,*

$$\sigma_t^\omega(\mathcal{A}(v>v_1)) \subseteq \mathcal{A}(v>v_1) \qquad (t\geq 0).$$

Then the blow-up limit of Theorem 6.1 on a smooth entangling cut yields a null dilation

$$v - v_0 \mapsto e^{-2\pi t}(v-v_0),$$

which is the geometric manifestation of the assumed half-sided modular inclusion for null half-lines.

Proof. Near a smooth cut, the cap dilation from Theorem 6.1 becomes a Rindler boost in the tangent geometry. Restricting that boost to a null generator gives the stated dilation. For $t \geq 0$, the half-line $v > v_0$ maps into itself, matching the stated half-sided inclusion property. □

Lemma 6.9 (Null Translation Generator)**.** *Under the standard half-sided modular inclusion of Corollary 6.8, there exists a one-parameter unitary group $U(a) = e^{iaP}$ with $P \geq 0$ such that*

$$\Delta^{it} U(a) \Delta^{-it} = U(e^{-2\pi t} a), \qquad [K,P] = i2\pi P.$$

Proof sketch. This is the Borchers-Wiesbrock consequence of half-sided modular inclusion. The modular group of the larger algebra acts by endomorphisms on the smaller one for one sign of modular time, and that structure produces a positive translation generator P whose affine scaling law is the one written above. □

Theorem 6.10 (Conditional null-stress identification from modular data)**.** *Under Axioms 3.1–3.4, Assumption 3.6, the hypotheses of Theorem 6.7, and the scaling-limit null-stress identification premise of the relativistic EFT branch, the renormalized modular Hamiltonian on a null half-line or bounded interval takes the geometric form*

$$\widetilde{K}[(a,\infty),\Omega] = 2\pi \int_a^\infty (v-a)\, T_{kk}(v,\Omega)\, dv + E^{(\varepsilon)}_{(a,\infty),\Omega},$$

with

$$\left|\langle\psi, E^{(\varepsilon)}_{I,\Omega}\phi\rangle\right| \le C_{I,\Omega,\psi,\phi}\,\varepsilon^{1/2}$$

for all domain vectors ψ, ϕ, where ε is the recoverability/scaling-limit control parameter. If bounded null intervals inherit geometric modular action from the interval-preserving projective flow, then for each bounded interval I one likewise has

$$\widetilde{K}[I,\Omega] = 2\pi \int_I g_I(v)\, T_{kk}(v,\Omega)\, dv + E^{(\varepsilon)}_{I,\Omega},$$

where $g_I(v)$ is affine-covariant and vanishes at finite interval endpoints. The full modular Hamiltonian is then

$$K[I,\Omega] = \widetilde{K}[I,\Omega] + K_\partial(I,\Omega),$$

with K_∂ central/boundary-supported. Knowledge of T_{kk} for all null directions reconstructs a symmetric tensor T_{ab} up to the addition of a metric term ϕg_{ab}.

Proof sketch. Theorem 6.7 gives the half-line endpoint-differentiation formula for the renormalized modular family. Corollary 6.8 and Lemma 6.9 identify the resulting half-line density with the positive generator of null translations. In the relativistic scaling-limit branch, one then uses the standard null modular-Hamiltonian identification for local stress-tensor charges, while the recoverability/scaling-limit remainder is absorbed into the error operator $E^{(\varepsilon)}_{I,\Omega}$. If the bounded-interval modular action is geometric, the same half-line density transports to finite intervals with an interval-dependent kernel $g_I(v)$ rather than an interval-independent additive measure density. The ambiguity

$$T_{ab} \to T_{ab} + \phi g_{ab}$$

is invisible on null contractions because $g_{ab}k^a k^b = 0$. □

Lemma 6.11 (Null Data Determine the Stress Tensor Modulo a Metric Term)**.** *Let X_{ab} be a symmetric tensor. If*

$$X_{ab}k^a k^b = 0$$

for every null vector k at a point, then

$$X_{ab} = \phi g_{ab}$$

for some scalar ϕ.

Proof. Write

$$X_{ab} = Y_{ab} + \phi g_{ab}$$

with Y_{ab} traceless. Since $g_{ab}k^a k^b = 0$ on null vectors, the hypothesis implies $Y_{ab}k^a k^b = 0$ for all null k. In local inertial coordinates let $k = (1, \hat{n})$ with $|\hat{n}| = 1$. Then

$$Y_{ab}k^a k^b = Y_{00} + 2\hat{n}^i Y_{0i} + \hat{n}^i \hat{n}^j Y_{ij}.$$

Oddness under $\hat{n} \mapsto -\hat{n}$ forces $Y_{0i} = 0$. Averaging the remaining relation over S^2 gives

$$Y_{00} + \frac{1}{3}\delta^{ij} Y_{ij} = 0.$$

Tracelessness of Y_{ab} gives

$$-Y_{00} + \delta^{ij} Y_{ij} = 0,$$

so both Y_{00} and the spatial trace $\delta^{ij}Y_{ij}$ vanish. The remaining condition is therefore

$$\hat{n}^i\hat{n}^j Y_{ij}^{\mathrm{TF}} = 0 \qquad \text{for all } \hat{n} \in S^2,$$

where Y_{ij}^{TF} is the traceless spatial part. The $\ell = 2$ spherical harmonics are linearly independent, so this implies $Y_{ij}^{\mathrm{TF}} = 0$. Hence $Y_{ab} = 0$ and therefore $X_{ab} = \phi g_{ab}$. □

Remark 6.12. *Lemma 6.11 explains why the Einstein equation derived from null modular data is determined only up to Λg_{ab}. The missing scalar piece is not an error term. It is the unique null-invisible component of a symmetric tensor.*

6.3 Generalized entropy and the Einstein equation

For a reference state ω and a small cap C,

$$\delta S_C = \delta\langle K_C\rangle.$$

By Theorem 6.1,

$$\delta S_C = 2\pi\,\delta\langle B_C\rangle.$$

By Proposition 5.5,

$$S_{\mathrm{gen}}(C) = \mathrm{Tr}(\rho L_C) + S_{\mathrm{bulk}}(C).$$

Together with Definition 5.6, this identifies the area term as the coarse-grained form of the edge entropy density rather than an independent postulate.

Definition 6.13 (Admissible equilibrium variation)**.** *Throughout this subsection, δ denotes a first-order state variation about a reference state ω, within the cap-label-preserving MaxEnt class, at fixed cap C, fixed cap size/volume, and fixed conserved charges. Shape or null-cut deformations are denoted separately by δ_{shape} or ∂_λ.*

Proposition 6.14 (Entanglement Equilibrium Condition)**.** *For admissible variations preserving cap labels, cap size/scale, and relevant conserved charges, MaxEnt selection implies*

$$\delta S_{\mathrm{gen}}(C) = 0.$$

Proof. Within each fixed edge sector, the realized state maximizes entropy subject to the local constraints. The first variation therefore vanishes once the cap size, conserved charges, and sector labels are fixed. □

Theorem 6.15 (Conditional Jacobson-type Einstein relation from entanglement equilibrium)**.** *Assume Axioms 3.1–3.4, Assumption 3.6, and the hypotheses of Theorem 6.10. Then, for every sufficiently small cap C and every admissible first variation δ about a reference state ω in the locally Lorentzian scaling regime,*

$$\delta S_{\mathrm{gen}}(C) = 0$$

implies

$$\delta(G_{00} + \Lambda g_{00}) = 8\pi G\,\delta\langle T_{00}\rangle$$

at the cap center in the diamond rest frame.

Proof sketch. The cap first law gives $\delta S_C = \delta\langle K_C\rangle$, and Theorem 6.10 identifies the bulk part of the background-subtracted variation with a null stress-tensor charge. Proposition 6.14 sets the total admissible first variation of the generalized entropy to zero at fixed cap, fixed size/volume, and fixed conserved charges, while shape deformations are kept separate as δ_{shape} or ∂_λ. Since the reference state is maximally symmetric, $G_{00} + \Lambda_{\text{ref}} g_{00} = 0$ in the background, so the fixed-volume small-ball area identity already enters at first order in the perturbation:

$$\delta A|_{V,\Lambda_{\text{ref}}} = -\frac{\Omega_{d-2}\ell^d}{d^2-1}(G_{00} + \Lambda_{\text{ref}} g_{00})$$

in the diamond rest frame. Equating the geometric and bulk contributions therefore yields the stated rest-frame first-variation relation. □

Corollary 6.16 (Tensor Upgrade in the Scaling Regime)**.** *If the first-variation relation of Theorem 6.15 holds for all local directions and reference states in the scaling regime, then*

$$G_{ab} + \Lambda g_{ab} = 8\pi G \langle T_{ab}\rangle$$

modulo the metric-term ambiguity already isolated by Lemma 6.11.

Proof. Overlap consistency across observers through the same point supplies all timelike directions, and Theorem 6.10 together with Lemma 6.11 upgrades the rest-frame relation to the full tensor equation in the Jacobson entanglement-equilibrium line of argument [8, 9]. □

6.4 The cosmological constant

Lemma 6.17 (Vacuum-Energy Blindness)**.** *For any null vector k and any vacuum-energy contribution $T^{\text{vac}}_{ab} = -\rho_{\text{vac}} g_{ab}$,*

$$T^{\text{vac}}_{kk} = 0.$$

Proposition 6.18 (Structural Separation of Λ)**.** *Local null-modular data fix T_{ab} only up to ϕg_{ab}. Therefore Λ is not determined by local overlap consistency and requires a global input.*

Corollary 6.19 (Cosmological Constant from Capacity)**.** *If the global screen capacity is identified with the de Sitter static-patch entropy,*

$$N_{\text{scr}} = S_{\text{dS}} = \frac{A_{\text{dS}}}{4G} = \frac{3\pi}{G\Lambda},$$

then

$$\Lambda = \frac{3\pi}{G N_{\text{scr}}}.$$

This is a structural aspect of the framework. The vacuum-energy problem is reorganized by separating local null data from the global capacity term.

7 Gauge Reconstruction and Standard Model Structure

7.1 Compact gauge reconstruction

At any fixed UV cutoff, edge-center completion equips collars with finitely many sector labels and fusion rules. The category used for reconstruction is the refinement-stable directed colimit

$$\mathsf{Sect}_\infty := \varinjlim_r \mathsf{Sect}_r$$

over regulator refinements r, retaining only transportable sectors and intertwiners that persist under refinement. This is the object on which Tannaka/Doplicher-Roberts reconstruction is applied.

Theorem 7.1 (Conditional compact gauge reconstruction in the bosonic branch)**.** *Assume Axioms 3.1–3.4, the relevant parts of Assumption 3.6, and that the refinement-stable edge-sector category* Sect_∞ *in the EFT regime is a rigid symmetric* C^**-tensor category with a faithful bosonic fiber functor to finite-dimensional Hilbert spaces. Then*

$$\mathsf{Sect}_\infty \simeq \mathrm{Rep}(G)$$

for a compact group G *uniquely determined up to isomorphism.*

Proof sketch. Each fixed-cutoff category has only finitely many simple blocks, but the refinement-stable colimit Sect_∞ can have infinitely many simple objects while still carrying objectwise finite-dimensional fibers. In the bosonic EFT branch, symmetry and rigidity place the category in the Doplicher-Roberts/Tannakian reconstruction class, and the faithful bosonic fiber functor identifies it with $\mathrm{Rep}(G)$ for a compact group G [10, 11]. □

If a fermionic sign object is present, the correct reconstruction statement is super-Tannakian rather than purely Tannakian. The compact note does not prove that fermionic/chiral extension; it works only in the bosonic internal-gauge branch.

7.2 Minimal admissibility and the Standard Model quotient

Throughout this subsection, χ_{cpl} means the dimension of the minimal coupled carrier supporting the required weak-type and color-type nonabelian charges on a common block. It is not the abstract minimal faithful representation dimension of the final gauge group.

Definition 7.2 (Weak-type and color-type nonabelian roles)**.** *A weak-type nonabelian charge is a pseudoreal nonabelian doublet role. A color-type nonabelian charge is an intrinsically complex nonabelian role, meaning that its nonabelian factor itself acts by an irreducible representation not equivalent to its conjugate. Abelian twisting of a pseudoreal nonabelian irreducible representation does not count as a color-type role.*

The derivation of the Standard Model quotient separates into five steps:

1. existence of a compact reconstructed group;
2. identification of the minimal nonabelian sector content;
3. identification of the minimal coupled carrier;
4. uniqueness of the connected abelian factor once admitted on that carrier;
5. determination of product gauge structure up to finite quotient, then of the global finite quotient from the realized matter spectrum.

Lemma 7.3 (Minimal nonabelian sector content)**.** *Any admissible low-energy sector must contain both a weak-type pseudoreal nonabelian charge role and an intrinsically complex color-type nonabelian charge role.*

Proof. Light chiral matter requires left-handed multiplets and right-handed singlets to carry inequivalent gauge data. A pseudoreal doublet structure is the minimal way to realize the weak sector without immediately producing vectorlike masses. A genuinely complex nonabelian representation is required to distinguish quarks from antiquarks. A single nonabelian simple factor cannot simultaneously supply both a minimal weak-type role and an intrinsically complex color-type role; abelian twisting does not count toward the color-type requirement. □

Lemma 7.4 (Minimal coupled carrier under MAR^+)**.** *Within the connected positive-dimensional Lie admissible class used by MAR, and under the weak-type / intrinsically complex color-type criteria above, the minimal coupled carrier supporting both roles on a common block has the form*

$$V = \mathbb{C}^3 \otimes \mathbb{C}^2, \qquad \chi_{\mathrm{cpl}} = 6.$$

Proof. Any nontrivial pseudoreal unitary representation has dimension at least 2, and the minimal weak-type role is realized by the doublet of SU(2). Any nontrivial intrinsically complex nonabelian unitary representation has dimension at least 3: dimension 1 would be abelian. For an irreducible 2-dimensional unitary representation of a connected compact group, the connected derived subgroup of its image is a compact connected subgroup of SU(2), hence either trivial, a torus, or all of SU(2). In the nonabelian case it is therefore conjugate to SU(2) itself, so the nonabelian action is the pseudoreal SU(2) doublet. Abelian twisting does not change that nonabelian pseudoreal role and, by the explicit color-type criterion above, does not count as an intrinsically complex color-type representation. Therefore no irreducible 2-dimensional unitary representation can realize the required color-type role. The fundamental triplet of SU(3) realizes this lower bound.

By definition of χ_{cpl}, the weak-type and color-type roles must act nontrivially on one common irreducible nonabelian block rather than on disconnected direct-sum summands. Commuting irreducible actions on such a common block therefore require dimension at least the product of the minimal weak and color dimensions:

$$\chi_{\mathrm{cpl}} \geq 2 \cdot 3 = 6.$$

Equality is achieved by the tensor-product carrier $\mathbb{C}^3 \otimes \mathbb{C}^2$. The block-diagonal representation $\mathbb{C}^3 \oplus \mathbb{C}^2$ of $S(U(3) \times U(2))$ is faithful of dimension 5, but it is not coupled and hence is not the MAR minimizer. Connected non-semisimple counterexamples of $U(2)$ type are excluded by the explicit color-type criterion rather than being silently ignored. □

Remark 7.5. *The dimension comparisons used in Lemma 7.4 apply to the positive-dimensional connected Lie image carrying the admissible nonabelian charges, equivalently to the identity component of the reconstructed group on that sector. Finite, disconnected, or connected non-semisimple counterexamples exist, but they are outside the admissible class fixed by the explicit weak-type / color-type criteria.*

MAR supplies existence of a connected abelian factor; the next lemma proves only uniqueness and identification on the minimal coupled carrier.

Lemma 7.6 (Uniqueness of the connected abelian factor on the minimal coupled carrier)**.** *Inside* U(6)*, the commutant of* $\mathrm{SU}(3) \times \mathrm{SU}(2)$ *acting on* $\mathbb{C}^3 \otimes \mathbb{C}^2$ *is exactly* U(1)*.*

Proof. The SU(3) action is irreducible on the first tensor factor and trivial on the second. The SU(2) action is irreducible on the second tensor factor and trivial on the first. By Schur's lemma, any operator commuting with both actions is scalar on each irreducible factor. Hence the full commutant is a single U(1). □

Theorem 7.7 (Product Gauge Structure up to Finite Quotient)**.** *Assume Axiom 3.5, vanishing obstruction $[z] = 0$ on the realized low-energy branch, the hypotheses of Theorem 7.1, the weak-type / color-type MAR premises above, and that the realized connected gauge image acts faithfully on the minimal coupled carrier. Then the realized connected gauge structure has the form*

$$G_{\text{phys}} = \frac{\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)}{\Gamma}$$

for some finite central subgroup Γ.

Proof. Theorem 7.1 yields some compact group G. Lemma 7.3 identifies the minimal admissible sector content, and Lemma 7.4 fixes the minimal coupled carrier. The tensor-product structure of that carrier implies commuting color and weak actions, hence product-group structure up to finite quotient. Axiom 3.5 supplies the existence of one connected abelian charge factor acting nontrivially on the coupled carrier, and Lemma 7.6 shows that any such connected abelian factor is necessarily a single U(1). Faithfulness rules out an invisible extra torus. The remaining ambiguity is therefore a finite central quotient Γ. □

Theorem 7.8 (Exact Rational Hypercharges)**.** *Assume gauge group $\mathrm{SU}(N_c) \times \mathrm{SU}(2) \times \mathrm{U}(1)_Y$, one generation of chiral matter (Q, u^c, d^c, L, e^c), one Higgs doublet H, and Yukawa terms*

$$QHu^c, \qquad QH^\dagger d^c, \qquad LH^\dagger e^c.$$

Then anomaly cancellation and Yukawa invariance imply

$$Y_L = -N_c Y_Q, \quad Y_H = N_c Y_Q, \quad Y_u = -(N_c+1)Y_Q, \quad Y_d = (N_c-1)Y_Q, \quad Y_e = 2N_c Y_Q.$$

Fixing the normalization by $Q = T_3 + Y$ and $Q(\nu_L) = 0$ gives

$$Y_Q = \frac{1}{2N_c}.$$

For $N_c = 3$ one recovers the exact Standard Model lattice

$$Y_Q = \frac{1}{6}, \quad Y_L = -\frac{1}{2}, \quad Y_u = -\frac{2}{3}, \quad Y_d = \frac{1}{3}, \quad Y_e = 1, \quad Y_H = \frac{1}{2}.$$

Proof. Yukawa invariance gives

$$Y_u = -(Y_Q + Y_H), \qquad Y_d = -Y_Q + Y_H, \qquad Y_e = -Y_L + Y_H.$$

The mixed anomalies yield

$$N_c Y_Q + Y_L = 0,$$

and

$$2N_c Y_Q + N_c Y_u + N_c Y_d + 2Y_L + Y_e = 0.$$

Solving the first anomaly gives

$$Y_L = -N_c Y_Q.$$

Substituting the Yukawa relations and $Y_L = -N_c Y_Q$ into the mixed gravitational anomaly then yields

$$Y_H = N_c Y_Q, \qquad Y_u = -(N_c+1)Y_Q, \qquad Y_d = (N_c-1)Y_Q, \qquad Y_e = 2N_c Y_Q.$$

With these relations in place, the $\mathrm{SU}(N_c)^2\mathrm{U}(1)$ and $\mathrm{U}(1)^3$ anomalies cancel identically. The normalization of Y_Q is fixed by the electric charge operator $Q = T_3 + Y$ together with $Q(\nu_L) = 0$, which implies

$$Y_L = -\frac{1}{2} = -N_c Y_Q, \qquad \text{hence} \qquad Y_Q = \frac{1}{2N_c}.$$

□

Corollary 7.9 (Three Generations)**.** *Under the hypotheses of Theorem 7.7,*

$$N_g = 3.$$

Proof. Intrinsic CKM CP violation requires

$$\frac{(N_g-1)(N_g-2)}{2} > 0,$$

hence $N_g \geq 3$. For one Higgs doublet, the one-loop weak-sector coefficient is

$$b_2 = \frac{11}{3}C_A - \frac{2}{3}\sum_f T(R_f) - \frac{1}{3}\sum_s T(R_s) = \frac{22}{3} - \frac{N_g(N_c+1)}{3} - \frac{1}{6} = \frac{43}{6} - \frac{N_g(N_c+1)}{3}.$$

Asymptotic freedom therefore implies

$$N_g(N_c+1) < \frac{43}{2}.$$

Since the color-type requirement forces $N_c \geq 3$, this gives $N_g \leq 5$. The minimal admissibility selector then chooses $N_g = 3$. □

Corollary 7.10 (Three Colors)**.** *Under the hypotheses of Theorem 7.7,*

$$N_c = 3.$$

Proof. Witten's global SU(2) anomaly requires the total number of left-handed weak doublets to be even:

$$N_g(N_c+1) \equiv 0 \pmod 2.$$

With $N_g = 3$, this implies N_c is odd. The case $N_c = 1$ is inadmissible because it removes the intrinsically complex color sector demanded by Lemma 7.3. Minimal admissibility therefore selects $N_c = 3$ [12]. □

Proposition 7.11 (Global quotient from the realized matter spectrum)**.** *If the realized matter spectrum carries the hypercharges of Theorem 7.8 with $N_c = 3$, then the subgroup of* $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ *acting trivially on all realized states is exactly $\mathbb{Z}_6$.*

Proof. Write $q := 6Y \in \mathbb{Z}$ and $\eta := e^{i\pi/3}$, so the U(1) factor acts by η^q on charge-q multiplets. The element

$$g_0 := (\omega_3, -1, \eta)$$

acts trivially on the realized matter and Higgs multiplets:

$$Q:\ \omega_3(-1)\eta = 1, \qquad u^c:\ \omega_3^{-1}\eta^{-4} = 1, \qquad d^c:\ \omega_3^{-1}\eta^2 = 1,$$
$$L:\ (-1)\eta^{-3} = 1, \qquad e^c:\ \eta^6 = 1, \qquad H:\ (-1)\eta^3 = 1.$$

Its powers therefore generate a subgroup of order six acting trivially on all realized states. Conversely, any central element acting trivially on e^c must have U(1) part η^n, and triviality on Q then fixes the SU(3) and SU(2) center factors uniquely. Hence every trivial central element is a power of g_0, so the kernel is exactly $\mathbb{Z}_6$. □

Corollary 7.12 (Standard Model Gauge Group)**.** *Under Theorem 7.7, Theorem 7.8, Corollaries 7.9 and 7.10, and Proposition 7.11,*

$$G_{\text{phys}} = \frac{\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)}{\mathbb{Z}_6}.$$

Proof. Theorem 7.7 yields the product structure up to finite quotient, and Proposition 7.11 fixes that quotient to $\mathbb{Z}_6$ once the realized hypercharge lattice and $N_c = 3$ are in place. □

Corollary 7.13 (Product-Group Consequences)**.** *The adjoint representation of the full connected gauge group contains only*

$$(8, 1, 0) \oplus (1, 3, 0) \oplus (1, 1, 0).$$

Equivalently, the adjoint representation of the derived gauge group contains only

$$(8, 1, 0) \oplus (1, 3, 0).$$

There are no mixed $(3, 2, \pm 5/6)$ *gauge bosons. Hence gauge-mediated proton decay is absent. The unbroken gauge symmetries forbid elementary gauge-boson mass terms: the photon remains massless, while gluons are massless gauge fields but confined.*

Remark 7.14. *The same product-group structure removes the usual grand-unification monopole sector. In the present framework, unification of couplings does not proceed by embedding the gauge group into a simple group that is later broken. It proceeds through a shared geometric origin of the edge heat-kernel parameters. Absence of gauge-mediated proton decay and unification therefore coexist without tension.*

8 Two-Input Quantitative Sector

8.1 Parameter count

The quantitative program uses the same axiom set together with the two external inputs P and N_{scr}. In the current implementation, P is calibrated from the gauge sector and N_{scr} is inferred from the observed cosmological constant. The discrete structure

$$(\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1))/\mathbb{Z}_6, \quad N_c = 3, \quad N_g = 3, \quad \varepsilon = \frac{1}{6}, \quad \delta = \frac{2}{9}$$

is then fixed by structural arguments, MAR selection, or phenomenological continuation steps rather than by direct fits. Relative to the usual Standard Model parameter count, this replaces roughly nineteen particle-physics inputs, or twenty-six when neutrino parameters are included, by two external inputs plus structural relations.

8.2 Epistemic classification of outputs

The quantitative sector is not uniform. The paper uses four categories.

1. **Calibration-sector consistency checks.** These are algebraically entangled with the determination of P from the gauge sector. They include v, the gauge couplings at m_Z, and tree-level electroweak masses.

2. **Independent quantitative outputs.** These use additional structural conditions that are logically independent of the pixel constraint. In the compact note this category is the Higgs/top critical-surface branch.

3. **Parameter-free structural outputs.** These include the gauge quotient, exact hypercharges, $N_c = 3$, and $N_g = 3$.

4. **Phenomenological continuations.** These include the charged-lepton/Koide construction, quark textures, baryogenesis, dark-sector phenomenology, and any use of the $\mathbb{Z}_6$ suppression factor as a flavor or entropy ansatz.

The quark-hierarchy and neutrino branches are not at the same maturity level in the compact note: the former is a texture-pattern continuation, while the latter is currently only an order-of-magnitude capacity-branch estimate.

This classification matters because the framework is not presented as a fit to all observables. Its stronger claim is that a large set of discrete and dimensionless structures are fixed before the overall energy scale is chosen.

8.3 Fundamental scales

The unification scale is

$$M_U = \frac{E_P}{e^{2\pi}} P^{1/6} \approx 2.474 \times 10^{16} \text{ GeV},$$

and the cell energy is

$$E_{\text{cell}} = \frac{E_P}{\sqrt{P}} \approx 9.56 \times 10^{18} \text{ GeV}.$$

The factor $e^{2\pi}$ comes from modular geometry. The factor $P^{1/6}$ comes from the cell-area scaling relation.

8.4 The pixel constraint and gauge closure

Edge-center completion and MaxEnt produce open-edge Gibbs / Laplacian weights

$$p_R(t) \propto d_R e^{-tC_2(R)}, \qquad \bar{\ell}_G(t) = \sum_R p_R(t) \log d_R.$$

For the two nonabelian factors, these are explicitly

$$\bar{\ell}_{\text{SU}(2)}(t) = \sum_{j=0,\frac{1}{2},1,\ldots} p_j(t) \log(2j+1), \qquad p_j(t) \propto (2j+1) e^{-t\, j(j+1)},$$

and

$$\bar{\ell}_{\text{SU}(3)}(t) = \sum_{p,q\geq 0} p_{(p,q)}(t) \log d_{(p,q)}, \qquad p_{(p,q)}(t) \propto d_{(p,q)} e^{-tC_2(p,q)},$$

with

$$d_{(p,q)} = \frac{1}{2}(p+1)(q+1)(p+q+2),$$

and

$$C_2(p,q) = \frac{1}{3}(p^2 + q^2 + pq + 3p + 3q).$$

Upon gluing or tracing the two boundary indices one obtains the closed heat-kernel partition function

$$Z_{\text{edge}}(t) = \sum_R d_R^2 e^{-tC_2(R)}.$$

For the nonabelian factors one obtains the pixel constraint

$$\frac{P}{4} = \bar{\ell}_{\text{SU}(2)}(t_2) + \bar{\ell}_{\text{SU}(3)}(t_3), \qquad t_i = 4\pi^2\alpha_i.$$

The framework then solves a self-consistent fixed-point problem:

1. choose a trial unified coupling α_U;
2. run the three gauge couplings to a trial scale using the derived beta-function shifts;
3. compute the electroweak scale

$$v = E_{\text{cell}} \exp\left(-\frac{2\pi}{\beta_{\text{EW}}\alpha_U}\right), \qquad \beta_{\text{EW}} = N_c + 1 = 4;$$

4. impose $m_Z(\mu) = \mu$ at the running scale;
5. solve the pixel constraint by bisection in α_U.

This yields

$$\alpha_U \approx 0.04112, \qquad \alpha_U^{-1} \approx 24.32.$$

The resulting beta-function shifts are numerically close to the MSSM one-loop shifts. In the framework this is not introduced by hand. It comes from the same heat-kernel edge weighting that enters the pixel constraint. The unification story is therefore structurally connected to the observer-overlap sector rather than appended as an unrelated high-energy ansatz.

8.5 Calibration-sector electroweak outputs

We use the convention

$$\alpha_1 = \frac{5}{3}\alpha_Y, \qquad g_Y^2 = 4\pi\alpha_Y = 4\pi\frac{3}{5}\alpha_1.$$

The gauge couplings at $\mu = m_Z$ are then

$$\alpha_1(m_Z) \approx 0.01696, \qquad \alpha_2(m_Z) \approx 0.03384, \qquad \alpha_3(m_Z) \approx 0.1183,$$

which, under the printed conventions of this compact note, give

$$\alpha_{\text{em}}^{-1}(m_Z) \approx 127.82, \qquad \sin^2\theta_W(m_Z) \approx 0.23119.$$

With

$$v \approx 246.77 \text{ GeV},$$

the tree-level electroweak masses are

$$m_W = \frac{1}{2}vg_2 \approx 80.46 \text{ GeV},$$

and

$$m_{Z,\text{run}} = \frac{1}{2}v\sqrt{g_2^2 + g_Y^2} \approx 91.76 \text{ GeV}.$$

Including the custodial correction

$$\Delta\rho = \frac{3y_t^2}{32\pi^2},$$

and using the illustrative unit-Yukawa estimate $y_t \approx 1$ for the one-loop pole correction, one obtains

$$M_{Z,\mathrm{pole}} \approx 91.33 \text{ GeV}.$$

These are calibration-sector outputs. Their role in the paper is scale setting, not independent confirmation. A fully reproducible numerical audit requires the beta-function shifts and matching conventions supplied in the repository supplements.

Quantity	OPH output	Experimental value	Status
$\alpha_{\mathrm{em}}^{-1}(m_Z)$	127.82	127.952	calibration sector
$\sin^2\theta_W(m_Z)$	0.23119	0.23122	calibration sector
$\alpha_s(m_Z)$	0.1183	0.1179	calibration sector
M_Z	91.33 GeV	91.188 GeV	calibration sector
m_W	80.46 GeV	80.38 GeV	calibration sector

8.6 Higgs and top from the critical surface

The independent quantitative branch begins with the refinement-stability condition

$$\lambda(M_U) = 0, \qquad \beta_\lambda(M_U) = 0.$$

At one loop, these conditions determine the top Yukawa boundary value

$$y_t(M_U) = \left[\frac{1}{16}\left(2g_2^4 + (g_2^2 + g_Y^2)^2\right)\right]^{1/4}.$$

The logic is simple: once $\lambda(M_U) = 0$, the condition $\beta_\lambda(M_U) = 0$ removes the final freedom in the top Yukawa boundary value. The low-energy Higgs and top masses then follow by RG evolution rather than by direct fitting.

Running the coupled system $(y_t, \lambda, g_Y, g_2, g_3)$ down to $\mu \sim m_t$ yields

$$m_t^{\overline{\mathrm{MS}}}(m_t) \approx 160.6 \text{ GeV},$$

and after the standard three-loop QCD pole-mass conversion used in the present implementation,

$$m_t^{\mathrm{pole}} \approx 171.1 \text{ GeV}.$$

The corresponding Higgs mass is

$$m_H = \sqrt{2\lambda(m_t)}\, v \approx 126.5 \text{ GeV}.$$

The critical-surface boundary conditions are logically independent of the pixel-constraint calibration. Converting the resulting dimensionless ratios m_H/v and m_t/v to GeV uses the previously calibrated v and gauge-coupling trajectory, and the critical-surface branch can fail even when the electroweak calibration succeeds.

Quantity	OPH output	Experimental value	Status
m_H	126.5 GeV	125.2 GeV	independent quantitative
m_t^{pole}	171.1 GeV	172.6 GeV	independent quantitative

9 Phenomenological Flavor Continuations

This section is not at the same logical status as the overlap fixed-point theorem, the conditional gravity branch, or the gauge/quotient chain. It records the current flavor continuations used in the OPH program and identifies which steps are phenomenological ansätze rather than theorem-level outputs.

9.1 The $\mathbb{Z}_6$ suppression parameter

Assume a uniform sixfold center-label ensemble on the realized $\mathbb{Z}_6$ quotient sectors. Then the associated entropy deficit is

$$\Delta S_{\text{center}} = \ln 6.$$

This motivates the universal suppression factor

$$\varepsilon = e^{-\Delta S_{\text{center}}} = \frac{1}{6},$$

which is used as a phenomenological flavor-hierarchy ansatz in the remainder of this section. The quotient alone does not force this entropy deficit without the ensemble assumption.

9.2 Koide structure from $\mathbb{Z}_3$ holonomy

Generation space is described by the minimal Hermitian circulant

$$\Phi = aI + bP + b^* P^2,$$

where P is the cyclic shift on three generations. Its eigenvalues are

$$\lambda_k = a + 2|b| \cos\left(\delta + \frac{2\pi k}{3}\right).$$

Let $r_k := \lambda_k$ denote scale-free Koide roots proportional to $\sqrt{m_k}$. Equivalently,

$$\sqrt{m_k} = \sqrt{S}\, r_k$$

for a common normalization S, and the Koide ratio Q is invariant under $r_k \mapsto c\, r_k$. Thus

$$Q = \frac{\sum_k r_k^2}{(\sum_k r_k)^2} = \frac{1 + 2(|b|/a)^2}{3}.$$

The balanced configuration $|b|/a = 1/\sqrt{2}$ yields

$$Q = \frac{2}{3}.$$

Using the observed charged-lepton masses one finds

$$Q_{\text{exp}} = 0.666664\ldots,$$

which is numerically indistinguishable from $2/3$ at the scale relevant here.

Within this continuation, the phase is fixed by previously derived discrete data:

$$\delta = \frac{\beta_{\text{EW}}\, Y_Q}{N_g} = \frac{N_c + 1}{2 N_c N_g} = \frac{2}{9}.$$

This remains one of the most rigid outputs inside the flavor continuation. It ties the Koide phase directly to color count, generation count, and the hypercharge lattice.

9.3 Charged-lepton masses

In the current implementation the lepton exponents are selected by a discrete scan that matches the ratios of Koide roots to the $\mathbb{Z}_6$ hierarchy, giving

$$(n_e, n_\mu, n_\tau) = (7, 4, 3).$$

With a common scale factor S fixed by consistency of the three equations,

$$m_e = S r_1^2, \qquad m_\mu = S r_2^2, \qquad m_\tau = S r_3^2.$$

The normalization used for the displayed table is

$$\mu_0 := 1 \text{ GeV}, \qquad \ln\left(\frac{S_0}{\mu_0}\right) = -\frac{1}{3}\sum_{i=1}^{3} \ln\left(\frac{r_i^2 \sqrt{2}\,\mu_0}{v}\, 6^{n_{e,i}}\right), \qquad S = S_0\, 2^{1/6}.$$

This common normalization makes the hierarchy exponents and the electroweak-scale relation mutually consistent. In the current implementation it yields sub-permille agreement with the charged-lepton data, but the exponent choice itself is a semi-empirical discrete-selection step rather than a theorem of the core OPH axioms.

Quantity	OPH output	Experimental value	Status
m_e	5.109×10^{-4} GeV	5.110×10^{-4} GeV	phenomenological continuation
m_μ	0.10564 GeV	0.10566 GeV	phenomenological continuation
m_τ	1.7766 GeV	1.7769 GeV	phenomenological continuation

9.4 Quark textures and CKM structure

The minimal hierarchy dictated by the derived color sector is

$$n_u = (2N_c, N_c, 0) = (6, 3, 0),$$

and

$$n_d = (2N_c, N_c + 1, N_c - 1) = (6, 4, 2).$$

Thus

$$m_{u_i} = \frac{v}{\sqrt{2}}\, \varepsilon^{n_{u,i}}, \qquad m_{d_i} = \frac{v}{\sqrt{2}}\, \varepsilon^{n_{d,i}}.$$

The CKM angles scale as successive powers of ε:

$$s_{12} = \varepsilon, \qquad s_{23} = \varepsilon^2, \qquad s_{13} = \varepsilon^3.$$

This sector is currently a texture/exponent-pattern continuation rather than a precision light-quark mass prediction. It captures the hierarchy pattern qualitatively, but absolute light-quark masses remain the least controlled component of the current program.

9.5 Neutrino scale

The neutrino branch uses the second input N_{scr}. In the compact note this is currently only an order-of-magnitude capacity-branch estimate rather than a benchmark quantitative mass formula. In the present implementation the resulting masses lie well below direct experimental bounds and match the expected order of magnitude from the dark-energy scale. This branch is structurally separate from the gauge-coupling calibration.

10 Program-Level Phenomenology and Further Continuations

10.1 Dark-sector modular anomaly

This subsection is program-level phenomenology rather than a closed theorem at the same status level as the gauge or gravity reconstruction theorems.

We distinguish two quantities. First, imperfect Markovity in the collar is measured by the modular additivity defect

$$\Delta K_\delta = K_{ABD} - K_{AB} - K_{BD} + K_B.$$

Its expectation value is the negative conditional mutual information:

$$\langle \Delta K_\delta \rangle_\omega = -I(A : D|B)_\omega \leq 0.$$

This is the recoverability diagnostic. Second, the cap-local anomaly term is defined by

$$K_C = 2\pi B_C + K_C^{(\mathrm{anom})}.$$

The effective dark-sector source is built from the first variation $\delta\langle K_C^{(\mathrm{anom})}\rangle$, whose sign is not fixed by SSA/CMI alone.

The resulting anomalous contribution enters Einstein's equation as

$$G_{ab} + \Lambda g_{ab} = 8\pi G \left(\langle T_{ab} \rangle + \langle T_{ab}^{\mathrm{anom}} \rangle \right),$$

with

$$\langle T_{00}^{\mathrm{anom}} \rangle = \frac{15}{8\pi^2} \frac{\delta\langle K_C^{(\mathrm{anom})}\rangle}{\ell^4}$$

in the diamond rest frame.

This sector is dark by construction. It is encoded in central and recoverability data rather than Standard Model matter fields.

10.2 The acceleration scale

Once Λ is fixed by screen capacity, the unique IR acceleration scale is

$$a_0^{(\mathrm{OPH})} = \frac{15}{8\pi^2} c^2 \sqrt{\frac{\Lambda}{3}} \approx 1.03 \times 10^{-10}\ \mathrm{m/s^2}.$$

This is within order-one agreement with the empirical MOND/RAR acceleration scale. In the deep IR, dimensional analysis then yields

$$g_{\mathrm{anom}} \propto \sqrt{a_0 g_b},$$

which reproduces the characteristic MOND-like scaling form.

10.3 Cosmological constant

The framework separates the cosmological constant from local vacuum energy. Local null data are blind to metric shifts ϕg_{ab}, while global capacity fixes

$$\Lambda = \frac{3\pi}{G N_{\mathrm{scr}}}.$$

Here the coefficient 3π comes from the de Sitter entropy identification $N_{\mathrm{scr}} = A_{\mathrm{dS}}/(4G)$. In this sense the cosmological-constant problem is reorganized into a global-capacity problem rather than a local zero-point-energy problem.

10.4 No gauge-mediated proton decay and no GUT monopoles

Because the derived gauge structure is a product group rather than a simple unification group, there are no gauge bosons in mixed quark-lepton representations. Therefore gauge-mediated proton decay is absent:

$$\tau_p^{(\text{gauge})} = \infty.$$

For the same reason, there is no GUT-scale simple-group breaking sector and therefore no associated monopole production channel in the usual grand-unification sense.

10.5 String theory as an effective description

The closed edge partition function obtained by gluing the open-edge weights has heat-kernel form

$$Z_{\text{edge}}(t) = \sum_R d_R^2 e^{-tC_2(R)}.$$

This is the same structure that appears in two-dimensional Yang-Mills theory. In a large-N_{edge} regime, with N_{edge} denoting an effective edge-sector rank rather than the physical color rank $N_c = 3$, the corresponding free energy admits the Gross-Taylor genus expansion

$$\log Z = \sum_{h \geq 0} N_{\text{edge}}^{2-2h} F_h(\lambda).$$

In OPH this is interpreted as an effective worldsheet reorganization of the edge dynamics. String theory therefore enters as an emergent description of a sector derived from the underlying observer-overlap framework.

10.6 Baryogenesis and CP violation

If the same phenomenological $\mathbb{Z}_6$ suppression ansatz is carried into the baryon-violating electroweak channel, a twelve-fermion 't Hooft vertex gives an effective suppression

$$\varepsilon^{12} = 6^{-12} \approx 4.6 \times 10^{-10},$$

which is of the same order as the observed baryon asymmetry. CP violation is not optional in the admissible branch because the realized flavor sector is intrinsically CP-capable, with a CKM-type physical phase after field redefinitions. This phase is not the gluing obstruction class $[z]$, which is assumed to vanish in the transportability chain.

11 Relation to Holography and Existing UV Frameworks

OPH belongs to the holographic family of ideas, but its primitive data are different from those used in asymptotic-boundary constructions. The distinction matters because several of the central claims of the present paper, including the treatment of Λ, the handling of factorization, and the emergence of gauge structure, depend on it.

11.1 Static-patch screen versus asymptotic boundary

In asymptotic-boundary holography one starts from a dual theory at infinity and reconstructs the bulk inward. In OPH one starts from a finite-capacity screen equipped with a net of local patch algebras and reconstructs the effective bulk from overlap consistency. The primitive object is therefore a family of observer patches with nontrivial overlaps, not a single global boundary theory.

This shift has two technical consequences. First, subsystem factorization is treated from the beginning as a gluing problem with centers and sector labels. Second, the physical role of the horizon is local and operational: every observer has direct access only to a patch algebra, and global law is whatever survives reconciliation on overlaps.

11.2 Positive Λ and finite capacity

The null-modular reconstruction of Section 4 determines the stress tensor only up to a metric term. This means that local null data do not fix Λ. The cosmological constant enters only when one supplies the global screen capacity

$$N_{\mathrm{scr}} = \log \dim \mathcal{H}_{\mathrm{tot}}.$$

The resulting relation

$$\Lambda = \frac{3\pi}{G N_{\mathrm{scr}}}$$

fits naturally with a de Sitter static patch of finite entropy via $N_{\mathrm{scr}} = A_{\mathrm{dS}}/(4G)$. OPH is therefore formulated in a de Sitter-first language. The positive cosmological constant is tied to finite capacity rather than to a deformation of a negative-Λ starting point.

11.3 Factorization, gauge structure, and the string sector

The factorization problem of gauge theory and gravity is often treated as a technical nuisance. In OPH it is part of the architecture. Edge-center completion turns the collar center into a first-class object, and the entropy split

$$S_{\mathrm{gen}} = \langle L_C \rangle + S_{\mathrm{bulk}}$$

is then a structural consequence rather than an added prescription.

In the bosonic EFT branch, the same collar data produce a symmetric rigid transportable sector category. Compact gauge reconstruction then follows conditionally from that category, and the Standard Model quotient is selected by minimal admissibility. This differs sharply from approaches in which the gauge group is specified in advance.

The string-theory relation is similarly reversed. The edge partition function

$$Z_{\mathrm{edge}}(t) = \sum_R d_R^2 e^{-tC_2(R)}$$

is the closed partition function obtained after gluing the open-edge weights $p_R(t) \propto d_R e^{-tC_2(R)}$. It has the heat-kernel form familiar from two-dimensional Yang-Mills theory. In a large-N_{edge} regime, with N_{edge} distinct from the physical color rank $N_c = 3$, its free energy admits a genus expansion. Within OPH this is read as an effective reorganization of edge degrees of freedom. String theory therefore enters as a sector description, while the microscopic starting data are the screen net, overlap consistency, and sector reconstruction.

Feature	Asymptotic-boundary holography	OPH
Primitive data	Boundary theory at infinity	Finite-capacity screen with observer-patch net
Factorization across cuts	Subtle and often indirect	Explicit edge-center completion with central labels
Status of Λ	External background datum	Fixed globally by N_{scr} after local null reconstruction
Gauge group	Usually specified as part of the model	Reconstructed from edge sectors, then filtered by admissibility
String sector	Fundamental dual description	Effective large-N_{edge} reorganization of edge dynamics

12 Predictive Status and Falsifiability

A proposal with this scope needs a clean accounting of what is exact, what is calibrated, what is independently predictive, and what remains program-level. The paper is organized around that distinction because the burden of proof is different in each category.

12.1 Prediction audit

Tier	Representative outputs	What would falsify the tier
Structural theorems	Confluence of overlap repair, conditional Lorentz kinematics, the conditional scaling-limit Einstein relation/tensor upgrade, conditional bosonic compact gauge reconstruction, Standard Model quotient chain, hypercharge lattice, $N_c = 3$, $N_g = 3$	A mathematical failure in the derivation chain, or data that require a different gauge quotient, different hypercharges, a different color count, or gauge-mediated proton decay
Calibration sector	$\alpha_i(m_Z)$, $\alpha_{\mathrm{em}}^{-1}(m_Z)$, $\sin^2\theta_W$, v, m_W, m_Z	Inability to close the pixel fixed point or systematic disagreement once the one-input calibration is imposed
Independent quantitative sector	m_H, m_t	Failure of the critical-surface branch, or dimensionless ratios that drift outside the stated loop and scheme uncertainty
Phenomenological continuations	Charged-lepton masses, Koide relation, quark textures, neutrino estimate	Failure of the Koide/holonomy continuation, wrong texture exponents, or the need to retract the branch once a sharper benchmark formula is supplied
Program-level phenomenology	Dark-sector anomaly, baryogenesis scale, conditional edge-sector fingerprints, black-hole spectroscopy comb, string/worldsheet limit	Absence of the predicted acceleration scale, failure of the center-label ansatz, wrong Casimir ratios, or null results in precision tests once the relevant regime is experimentally clean

This separation is essential for interpreting the quantitative claims. The electroweak calibration branch sets the overall scale. The Higgs/top branch then tests additional structure that is logically independent of that scale setting. The charged-lepton and Koide branches are retained as phenomenological continuations. The gauge quotient, hypercharge lattice, color count, generation count, and symmetry-protected exact zeros do not use a continuous fit parameter at all.

12.2 Direct discriminators

Several consequences of the framework are unusually sharp because they are parameter-free or nearly so.

12.2.1 The $\mathbb{Z}_6$ central-label entropy fingerprint

If the uniform center-label ensemble ansatz of Section 8.1 is adopted and the realized gauge group is

$$\frac{\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)}{\mathbb{Z}_6},$$

let Z denote the discrete center/quotient label in the edge-center completion. Then the predicted central-label entropy deficit is

$$\Delta S_{\text{center}} = \log 6$$

or

$$\Delta S_{\text{center}}^{(\text{bits})} = \log_2 6 \approx 2.585.$$

This is a global-structure observable conditioned on the ensemble ansatz. It is not the Shannon entropy of the open-edge irrep weights $p_R(t)$ and is not t-dependent. A clean measurement of central-label entropies inconsistent with that deficit would falsify that continuation.

12.2.2 Exact SU(3) Casimir-ratio pattern

For $\mathbb{Z}_6$-compatible sectors (R_3, R_2, Y), let

$$r_R := \frac{C_2(R)}{C_2(3)}$$

denote the SU(3) Casimir ratio relative to the triplet. Then the open-edge law

$$p_R(t) \propto d_R e^{-tC_2(R)}$$

implies exact ratios among the leading SU(3) Casimir suppressions:

$$r_8 = \frac{9}{4}, \qquad r_6 = \frac{5}{2}, \qquad r_{10} = \frac{9}{2}, \qquad r_{15} = 4, \qquad r_{27} = 6.$$

These rational numbers are fixed by representation theory. They do not involve threshold tuning or hidden UV data. Precision lattice access to the SU(3) component of edge-sector probabilities would therefore probe the mechanism directly.

12.2.3 Unification without proton decay

In conventional simple-group unification, improved coupling unification typically comes with leptoquark gauge bosons and proton-decay tension. In OPH the one-loop running is corrected by edge-sector beta-function shifts while the gauge group remains a product group up to a finite quotient. The combination

$$\text{MSSM-like running} \quad + \quad \tau_p^{(\text{gauge})} = \infty$$

is therefore a sharp discriminator. Evidence for gauge-mediated proton decay or for GUT monopoles would contradict this branch of the theory.

12.2.4 Dark-sector acceleration scale

The modular-anomaly sector does not introduce a new particle species. It introduces an infrared contribution governed by the same global capacity that fixes Λ. The resulting acceleration scale

$$a_0^{(\mathrm{OPH})} = \frac{15}{8\pi^2} c^2 \sqrt{\frac{\Lambda}{3}}$$

lies in the observed MOND/RAR range. A robust failure of this scaling across systems would eliminate the current dark-sector realization.

12.2.5 Conditional black-hole spectroscopy

If the discrete horizon branch of the same edge counting is adopted, the horizon line structure is logarithmic:

$$\Delta E_k = k_B T_H \ln k, \qquad k = 2, 3, 4, \ldots$$

For spinning holes the associated ringdown frequencies take the form

$$f_{k,m}(M, \chi) = \frac{m\Omega_H}{2\pi} \; + \; \frac{c^3 g(\chi)}{16\pi^2 GM} \ln k.$$

Here Ω_H is the horizon angular velocity and $g(\chi)$ is the spin-dependent spacing factor on the discrete-horizon branch. Defining the event-stacking variable

$$x := \frac{16\pi^2 GM}{c^3 g(\chi)} \left(2\pi f - m\Omega_H\right),$$

the predicted peaks sit at $x = \ln k$. For any single finite-capacity remnant only finitely many k are realized, so the $\ln k$ sequence is a cross-event finite-window template rather than an exact infinite spectrum of one finite-dimensional Hamiltonian. This branch is conditional because it uses one further step in the horizon-spectrum program, but it is experimentally sharp.

12.3 What the paper does not claim

The present manuscript does not claim precision control of all QCD observables, a finished treatment of hadronic thresholds, a completed nonlinear dark-sector phenomenology in galaxies and clusters, or a completed fermionic/super-Tannakian gauge branch. It claims a derivation chain for the underlying structures, plus a set of quantitative and phenomenological branches whose status is made explicit above. This distinction is important because the framework should be judged by the strongest outputs first.

13 Common Objections and Clarifications

13.1 Is the use of P circular?

The answer depends on which sector is under discussion. In the current implementation the pixel area P calibrates the overall particle-physics scale through the gauge sector. Quantities that are algebraically entangled with that calibration, such as v, m_W, m_Z, and the gauge couplings at m_Z, are consistency checks. The paper treats them as such.

The stronger outputs enter only after the scale has been set. The critical-surface condition

$$\lambda(M_U) = 0, \qquad \beta_\lambda(M_U) = 0$$

fixes the Higgs and top branch independently of the pixel constraint. The Koide continuation is controlled by

$$Q = \frac{2}{3}, \qquad \delta = \frac{2}{9},$$

which are structural consequences of the previously derived discrete data inside that continuation. The quark hierarchy depends on the $\mathbb{Z}_6$ texture ansatz, and the neutrino branch depends on the second input N_{scr}, not on P.

Operationally, changing P moves the calibration family $(\alpha_U, \alpha_i(m_Z), v, m_W, m_Z)$ as well as the overall GeV scale. It does not alter the parameter-free structural outputs such as the gauge quotient or hypercharge lattice. The useful distinction is therefore between scale setting, structural determination, and later phenomenological continuations.

13.2 Does a fixed UV cell size break Lorentz invariance?

The UV regulator is placed on the screen algebra, not on an emergent bulk spacetime lattice. Lorentz kinematics in the present framework arise from the continuum modular action on caps, summarized in Corollary 6.2, rather than from microscopic invariance of a bulk discretization.

This is conceptually similar to other continuum limits in statistical mechanics and lattice field theory. A regulator may break a symmetry at finite cutoff while the infrared fixed point restores it exactly. In OPH the restoration is stronger than a generic RG expectation because the modular-flow theorem identifies the symmetry group itself:

$$\mathrm{Conf}^+(S^2) \cong \mathrm{SO}^+(3,1).$$

Residual Lorentz violation would therefore have to arise from a failure of the regulator premises or from a failure of the modular-flow theorem, not merely from the existence of a UV cell size.

13.3 Why use type-I algebras if continuum QFT uses type III?

The type-I assumption is a regulator statement. At finite cutoff the local patch algebras are finite-dimensional matrix algebras, which makes entropy and recovery theory well defined. The continuum limit is expected to recover the familiar type-III behavior of local QFT algebras. There is no contradiction here. The two statements refer to different scales.

This is no more problematic than using a lattice regulator for Yang-Mills theory and then taking a continuum limit in which the operator algebra changes type. The mathematical burden is the existence of a controlled scaling regime. In the present manuscript that burden is isolated in Assumption 3.6. It is not hidden inside the main theorems.

13.4 Why is charge quantization possible without a simple GUT?

The framework uses the global quotient and anomaly structure rather than a simple-group embedding. Once the realized matter content is fixed, the subgroup acting trivially on all states is exactly $\mathbb{Z}_6$, and the anomaly equations force the sixth-integer hypercharge lattice. Integer electric charge for color singlets is then a property of the realized quotient structure. A simple unified gauge group is not required.

14 Discussion

The framework is strongest where the outputs are discrete or structurally rigid:

1. the Lorentz group;
2. the conditional scaling-limit Einstein branch;
3. the Standard Model gauge quotient chain;
4. exact hypercharges;
5. three colors and three generations;
6. the two-input economy of the present implementation.

The next strongest sector is the independent quantitative branch consisting of the Higgs and top masses. The charged-lepton and Koide sectors are promising but should presently be read as phenomenological continuations rather than core theorem outputs.

The weakest part is the light-quark and hadron branch. The hierarchy pattern is robust, but precision control of low-energy QCD effects, scheme dependence, and order-one coefficients is not yet complete.

The framework should therefore be evaluated in stages. It already contains several results that are normally postulated independently in other approaches: kinematics, dynamics, gauge structure, charge lattice, matter multiplicities, and part of the mass spectrum. Whether one regards this as evidence for fundamentality depends on whether these simultaneous derivations survive further scrutiny.

Topics intentionally excluded from the present formulation include simulation narratives, observer-continuation mechanisms, and claims of final completeness for all low-energy observables.[2]

15 Conclusion

Observer-Patch Holography starts from overlap consistency of observer patches and yields, within one framework, several effective structures usually treated separately in modern fundamental physics. The compact note supports a schedule-independent normal form for objective law, conditional scaling-limit Lorentz kinematics and a conditional Jacobson-type Einstein branch, the Standard Model gauge quotient with exact hypercharges under the extended MAR admissibility package in the bosonic internal-gauge branch, three colors, three generations, a two-input quantitative program with a critical-surface Higgs/top branch, and structural cosmological corollaries. It also records several phenomenological continuations whose present status is weaker and is labeled as such in the text.

Accordingly, the framework functions as a candidate underlying theory from which general relativity and the Standard Model may arise as effective descriptions once the stated scaling-limit and categorical premises are realized. The next technical tasks are sharper quantitative control, a fuller fermionic gauge branch, and explicit UV completions that land in the required modular/geometric phase.

[2] Questions outside this theorem package are, for present purposes, mostly harmless.

A One-loop RG Equations Used in the Quantitative Branch

For convenience we record the one-loop system used in the Higgs/top transport branch: Throughout this appendix, g_Y denotes the SM hypercharge coupling. The GUT-normalized coupling is g_1 with $g_1^2 = \frac{5}{3} g_Y^2$, equivalently $\alpha_1 = \frac{5}{3}\alpha_Y$. Therefore the one-loop coefficient is 41/6 in the g_Y equation and 41/10 in the g_1 equation.

$$16\pi^2 \frac{dg_Y}{d\ln\mu} = \frac{41}{6} g_Y^3, \tag{3}$$

$$16\pi^2 \frac{dg_2}{d\ln\mu} = -\frac{19}{6} g_2^3, \tag{4}$$

$$16\pi^2 \frac{dg_3}{d\ln\mu} = -7 g_3^3, \tag{5}$$

$$16\pi^2 \frac{dy_t}{d\ln\mu} = y_t \left(\frac{9}{2} y_t^2 - \frac{17}{12} g_Y^2 - \frac{9}{4} g_2^2 - 8 g_3^2 \right), \tag{6}$$

$$16\pi^2 \frac{d\lambda}{d\ln\mu} = 24\lambda^2 + \lambda \left(12 y_t^2 - 9 g_2^2 - 3 g_Y^2 \right) - 6 y_t^4 + \frac{3}{8} \left(2 g_2^4 + (g_2^2 + g_Y^2)^2 \right). \tag{7}$$

In the present implementation this system is the transport layer between the UV boundary data and the low-energy Higgs/top outputs. Numerically, the compact note uses one-loop $\overline{\mathrm{MS}}$ transport, the matching scale $\mu = m_Z = 91.1876\,\mathrm{GeV}$, the unified scale $M_U \approx 2.474 \times 10^{16}\,\mathrm{GeV}$ from the main text, and the standard three-loop QCD pole-mass conversion for the displayed top-branch number. Higher-loop and threshold refinements are expected to shift the final numbers at the GeV level rather than alter the logic of the branch.

B Quantitative Robustness Audit

Source of variation	Baseline used here	Expected effect on outputs
Critical-surface RG transport	one-loop SM running	moving to two loops is expected to shift m_H by about 1 GeV and m_t by about 0.5 GeV
Pole-mass and threshold matching	standard three-loop top conversion plus leading electroweak corrections	order-one GeV effects in the top branch; sub-GeV to GeV effects in the Higgs branch
Heat-kernel truncation in $\bar{\ell}_G$	200 representations	negligible at displayed precision, with changes below 10^{-12} in the entropy sums
Bisection tolerance for the pixel closure	10^{-10}	affects only the last shown digits of α_U and derived calibration outputs
Light-quark scheme matching	not yet implemented at precision level	dominant source of the present 5% to 20% uncertainty in the light-quark branch

The paper therefore separates the quantitative sector by status. The Higgs and top branches are useful because their errors are controlled at the percent level. The charged-lepton continuation is numerically sharp but rests on a weaker discrete-selection step, while the light-quark and hadronic branch still carries the largest scheme and threshold uncertainty.

References

[1] J. J. Bisognano and E. H. Wichmann, "On the duality condition for a Hermitian scalar field," *J. Math. Phys.* **16** (1975) 985–1007.

[2] J. J. Bisognano and E. H. Wichmann, "On the duality condition for quantum fields," *J. Math. Phys.* **17** (1976) 303–321.

[3] E. H. Lieb and M. B. Ruskai, "Proof of the strong subadditivity of quantum-mechanical entropy," *J. Math. Phys.* **14** (1973) 1938–1941.

[4] D. Petz, "Sufficient subalgebras and the relative entropy of states of a von Neumann algebra," *Commun. Math. Phys.* **105** (1986) 123–131.

[5] D. Petz, "Sufficiency of channels over von Neumann algebras," *Quart. J. Math.* **39** (1988) 97–108.

[6] O. Fawzi and R. Renner, "Quantum conditional mutual information and approximate Markov chains," *Commun. Math. Phys.* **340** (2015) 575–611.

[7] P. Hayden, R. Jozsa, D. Petz and A. Winter, "Structure of states which satisfy strong subadditivity of quantum entropy with equality," *Commun. Math. Phys.* **246** (2004) 359–374.

[8] T. Jacobson, "Thermodynamics of spacetime: the Einstein equation of state," *Phys. Rev. Lett.* **75** (1995) 1260–1263.

[9] T. Jacobson, "Entanglement equilibrium and the Einstein equation," *Phys. Rev. Lett.* **116** (2016) 201101.

[10] S. Doplicher and J. E. Roberts, "A new duality theory for compact groups," *Invent. Math.* **98** (1989) 157–218.

[11] S. Doplicher and J. E. Roberts, "Why there is a field algebra with a compact gauge group describing the superselection structure in particle physics," *Commun. Math. Phys.* **131** (1990) 51–107.

[12] E. Witten, "An SU(2) anomaly," *Phys. Lett. B* **117** (1982) 324–328.