

A new class of generalized inverses in rings with involution

Liyun Wu*, Jianlong Chen and Huihui Zhu

Southeast University
Southeast University
Hefei University of Technology
E-mail: wlymath@163.com

Let R be a $*$ -ring and let $a, w, v \in R$. This paper contributes to define two new classes of generalized inverses, called the w -core inverse and the dual v -core inverse in R . An element $a \in R$ is w -core invertible if there exists some $x \in R$ such that $awx^2 = x$, $xawa = a$ and $(awx)^* = awx$. Such an x is called a w -core inverse of a , which encompasses several known generalized inverses such as the core inverse, the e -core inverse and the Moore-Penrose inverse. Several characterizations and representations for elements to be w -core invertible are given, based on $\{1, 3\}$ -inverses, inverses along an element, (b, c) -inverses, projections, ideals and units. Also, the connections between the w -core inverse and other generalized inverses are given.

References

- [1] Baksalary, O. M., Trenkler, G. (2010). Core inverse of matrices. *Linear Multilinear Algebra* 58:681–697.
- [2] Mary, X., Patrício, P. (2013). Generalized inverses modulo H in semigroups and rings. *Linear Multilinear Algebra* 61:886–891.
- [3] Penrose, R. (1955). A generalized inverse for matrices. *Proc. Camb. Phil. Soc.* 51:406–413.
- [4] Drazin, M. P. (2012). A class of outer generalized inverses. *Linear Algebra Appl.* 436:1909–1923.