

The n -torsionfreeness of syzygies of the residue field

Yuya Otake

Nagoya University

E-mail: `m21012v@math.nagoya-u.ac.jp`

The notion of n -torsionfree modules was introduced by Auslander and Bridger [1] as a natural generalization of the notion of torsionfree modules over integral domains. The n -torsionfreeness of syzygies plays an important role in Auslander's approximation theory [1, 2]. As a dual theory of approximation theories, the existence of finite projective hulls has been well-studied [2, 4, 5, 6]. We will see that the existence of finite projective hulls is closely related to the $(n + 1)$ -torsionfreeness of n -syzygies. As an application, we consider the n -torsionfreeness of syzygies of the residue field k over a local ring R . Let $t = \text{depth } R$ and $\Omega^t k$ be the t th syzygy of the residue field k . Dey and Takahashi [3] proved that $\Omega^t k$ is $(t + 1)$ -torsionfree, and it is a $(t + 2)$ nd syzygy if and only if the local ring R has type one. Motivated by their results, we study higher torsionfreeness of the syzygy $\Omega^t k$.

References

- [1] M. Auslander; M. Bridger, Stable module theory, Memoirs of the American Mathematical Society 94, American Mathematical Society, Providence, R.I., 1969.
- [2] M. Auslander; R.-O. Buchweitz, The homological theory of maximal Cohen-Macaulay approximations, Colloque en l'honneur de Pierre Samuel (Orsay, 1987), M'ém. Soc. Math. France (N.S.) 38 (1989), 5—37.
- [3] S. Dey; R. Takahashi, On the subcategories of n -torsionfree modules and related modules, Collect. Math. 74 (2023), no. 1, 113—132.
- [4] H.-B. Foxby, Embedding of modules over Gorenstein rings, Proc. Amer. Math. Soc. 36 (1972), 336—340.
- [5] Y. Otake, Ext modules related to syzygies of the residue field, preprint (2023), arXiv:2304.02900.
- [6] R. Takahashi; S. Yassemi; Y. Yoshino, On the existence of embeddings into modules of finite homological dimensions, Proc. Amer. Math. Soc. 138 (2010), no. 7, 2265—2268.