

# The first Hilbert coefficient and the reduction number of stretched ideals

Kazuho Ozeki

Nihon University

E-mail: `ozeki.kazuho@nihon-u.ac.jp`

The homological property of the associated graded ring of an ideal is an important problem in commutative algebra. In this talk we explore the structure of the associated graded ring of stretched  $\mathfrak{m}$ -primary ideals in the case where the reduction number attains almost minimal value in a Cohen-Macaulay local ring  $(A, \mathfrak{m})$ . In the second half of this talk, We also present the structure of stretched  $\mathfrak{m}$ -primary ideals with small first Hilbert coefficient.

## References

- [1] K. Ozeki, The reduction number of stretched ideals, J. Math. Soc. Japan, Vol. 74 (2022), pp.1021-1045.
- [2] K. Ozeki, The first Hilbert coefficient of stretched ideals, Acta Mathematica Vietnamica, Vol. 47 (2022), pp.251-267.
- [3] M. E. Rossi and G. Valla, Stretched  $\mathfrak{m}$ -primary ideals, Beiträge Algebra und Geometrie Contributions to Algebra and Geometry, Vol. 42 (2001), pp.103-122.
- [4] J. D. Sally, Stretched Gorenstein rings, J. Lond. Math. Soc. (2), Vol. 20 (1979), pp.19-26.

---

2020 Mathematics Subject Classification: 13H10, 13D40

Keywords: commutative ring, stretched local ring, stretched ideal, Cohen-Macaulay local ring, associated graded ring, Hilbert function, Hilbert coefficient