Modules of 0-Hecke algebras from posets

Seung-Il Choi*, Young-Hun Kim and Young-Tak Oh

Seoul National University
Seoul National University
Sogang University
E-mail: ignatioschoi@snu.ac.kr

In this talk, our main focus is on exploring a specific class of mathematical objects called right 0-Hecke modules, which arise from partially ordered sets. These modules, also known as poset modules, were introduced by Duchamp, Hivert, and Thibon. Their work involved investigating the intriguing algebraic and combinatorial properties of these modules.

We specifically consider the category of the full subcategory of right 0-Hecke modules, where the objects are direct sums of finitely many isomorphic copies of poset modules. Our first result establishes that the direct sum of the Grothendieck groups of the poset module category is isomorphic to the Hopf algebra of quasisymmetric functions. Furthermore, we demonstrate that every right weak Bruhat interval module can be viewed as a poset module. By introducing a certain functor, we also establish that every left weak Bruhat interval module can be viewed as a poset module.

References

- [1] S.-I. Choi, Y.-H. Kim, S.-Y. Nam, and Y.-T. Oh. The projective cover of tableau-cyclic indecomposable Hn(0)-modules. Transactions of the American Mathematical Society, 375(11):7747–7782, 2022
- [2] W.-S. Jung, Y.-H. Kim, S.-Y. Lee, and Y.-T. Oh. Weak Bruhat interval modules of the 0-Hecke algebra. Math. Z., 301(4):3755–3786, 2022
- [3] G. Duchamp, F. Hivert, and J.-Y. Thibon. Noncommutative symmetric functions. VI. Free quasi-symmetric functions and related algebras. Internat. J. Algebra Comput., 12(5):671–717, 2002
- [4] S.-I. Choi, Y.-H. Kim, and Y.-T. Oh. Poset modules of 0-Hecke algebras and related quasisymmetric power sum expansions, ArXiv (to appear)