

Quantum projective planes finite over their centers and Beilinson algebras

Ayako Itaba

Tokyo University of Science
E-mail: `itaba@rs.tus.ac.jp`

In noncommutative algebraic geometry, a quantum polynomial algebra defined by Artin and Schelter [1] is a basic and important research object, which is a noncommutative analogue of a commutative polynomial algebra. Also, Artin and Schelter [1] gave the classifications of low dimensional quantum polynomial algebras. Moreover, Artin, Tate and Van den Bergh [2] found a nice correspondence between 3-dimensional quantum polynomial algebras and geometric pair (E, σ) , where E is the projective plane or a cubic divisor in the projective plane, and σ is the automorphism of E . So, this result allows us to write a 3-dimensional quantum polynomial algebra A as the form $A = A(E, \sigma)$. For a 3-dimensional quantum polynomial algebra $A = A(E, \sigma)$, Artin, Tate and Van den Bergh [3] showed that A is finite over its center if and only if the order $|\sigma|$ of σ is finite. For a 3-dimensional quantum polynomial algebra $A = A(E, \sigma)$ with the Nakayama automorphism ν of A , the author and Mori [7] proved that the order $|\nu^* \sigma^3|$ of $\nu^* \sigma^3$ is finite if and only if the norm $\|\sigma\|$ of σ introduced by Mori [9] is finite if and only if the noncommutative projective plane in the sense Artin and Zhang [4] is finite over its center. In [6], for a 3-dimensional quantum polynomial algebra A of Type S', the author prove that the following conditions are equivalent; (1) the noncommutative projective plane is finite over its center. (2) The Beilinson algebra of A is 2-representation tame in the sense of Herschend-Iyama-Oppermann [5]. (3) The isomorphism classes of simple 2-regular modules over the Beilinson algebra of A are parametrized by the projective plane. Note that, this result holds for a Type S by Mori [9]. The Beilinson algebra of A is introduced by Minamoto and Mori [8], which is a typical example of 2-representation infinite algebra.

References

- [1] M. Artin and W. Schelter, Graded algebras of global dimension 3, *Adv. Math.* 66 (1987), 171-216.
- [2] M. Artin, J. Tate and M. Van den Bergh, Some algebras associated to automorphisms of elliptic curves, *The Grothendieck Festschrift*, vol. 1, *Progress in Mathematics* vol. 86 (Birkhäuser, Basel, 1990) 33-85.

2020 Mathematics Subject Classification: 16W50, 16S37, 16D90, 16E65, 16G10, 16G60

Keywords: Quantum polynomial algebras, Geometric algebras, Quantum projective planes, Calabi-Yau algebras, Beilinson algebras, Regular modules

- [3] M. Artin, J. Tate and M. Van den Bergh, Module over regular algebras of dimension 3, *Invent. Math.* 106 (1991), no. 2, 335-388.
- [4] M. Artin and J. J. Zhang, Noncommutative projective schemes, *Adv. Math.* 109 (1994), no. 2, 228-287.
- [5] M. Herschend, O. Iyama and S. Oppermann, n-representation infinite algebras, *Adv. Math.* 252 (2014), 292-342.
- [6] A. Itaba, Quantum projective planes and Beilinson algebras of 3-dimensional quantum polynomial algebras for Type S', submitted (2023), arXiv:2304.02242.
- [7] A. Itaba and I. Mori, Quantum projective planes finite over their centers, *Canad. Math. Bull.*, Vol. 66 (2023), Issue 1, 53-67.
- [8] H. Minamoto and I. Mori, The structure of AS-Gorenstein algebras, *Adv. Math.* 226 (2011), no. 5, 4061-4095.
- [9] I. Mori, Regular modules over 2-dimensional quantum Beilinson algebras of Type S, *Math. Z.* 279 (2015), no. 3-4, 1143-1174.