Rings and radicals related to n-primariness

Hongying Chen* and Yang Lee

Pusan National University Yanbian University E-mail: hongyingchen93@gmail.com

An ideal I of a ring R is called n-primary(resp., T-primary) provided that $AB \subseteq I$ for ideals A, B of R implies that (A+I)/I or (B+I)/I is nil of index n(resp., (A+I)/I or (B+I)/I is nil) in R/I, where $n \ge 1$. It is proved that for a proper ideal I of a principal ideal domain R, I is T-primary if and only if I is of the form p^kR for some prime element p and $k \ge 1$ if and only if I is 2-primary, through which we study the structure of matrices over principal ideal domains. We prove that for a T-primary ideal I of a ring R, R/I is prime when the Wedderburn of R/I is zero. In addition we provide a method of constructing strictly descending chain of n-primary radicals from any domain, where the n-primary radical of a ring R means the intersection of all the n-primary ideals of R.

References

- [1] S.A. Amitsur, *Nil radicals; Historical notes and some new results*, In: Colloq. Math. Soc. Janos Bolyai 6: Rings, Modules and Radicals, Keszthely (Hungary) 1971, 47-65
- [2] C. Beddani, W.Messirdi, 2-prime ideals and their applications, J. Algebra Appl. 15 (2016), 1650051, 11 pp.
- [3] G.F. Birkenmeier, H.E. Heatherly, E.K. Lee, *Completely prime ideals and associated radicals*, in: S.K. Jain, S.T. Rizvi (Eds.), Proc. Biennial Ohio State.Denison Conference, 1992, World Scientific, Singapore, 1993, pp. 102-129.
- [4] J.Han, T.K. Kwak, C.I. Lee, Y. Seo, Generalizations of reversible and Armendariz rings, Internat. J. Algebra Comput. 25 (2016), 911-933.
- [5] I.N. Herstein, Topics in Ring Theory, Univ. Chicago Press, Chicago, 1965.
- [6] C.Y. Hong, T.K. Kwak, On minimal strongly prime ideals, Comm. Algebra 28 (2000), 4867-4878.
- [7] S.U. Hwang, Y. C. Jeon and Y. Lee, Structure and topological conditions of NI rings, J. Algebra, **302** (2006),186-199.

²⁰²⁰ Mathematics Subject Classification: 16D25, 16S36, 16U80

Keywords: n-priamry ring, T-primary ring, n-priamry radical, T-primary radical, prime ring, polynomial ring, matrix ring

- [8] S.U. Hwang, N.K. Kim, Y. Lee, On rings whose right annihilators are bounded, Glasgow Math. J. 51 (2009), 539-559.
- [9] Y.C. Jeon, H.K. Kim, Y. Lee, J.S. Yoon, On weak Armendariz rings, Bull. Korean Math. Soc. 46 (2009), 135-146.
- [10] N.K. Kim, Y. Lee, M. Ziembowski, Annihilating properties of ideals generated by coefficients of polynomials and power series, Int. J. Algebra Comput., Vol. 32, No. 02, pp. 237-249 (2022).
- [11] A.A. Klein, Rings of bounded index, Comm. Algebra 12 (1984) 9-21.
- [12] G. Köthe, Die Struktur der Ringe, deren Restklassenring nach dem Radikal vollständig reduzibel ist. Math. Z. 32 (1930), 161-186.
- [13] J. Krempa, Logical connections among some open problems concerning nil rings, Fund. Math. 76 (1972), 121-130.
- [14] T.Y. Lam, A First Course in Noncommutative Rings, Springer-Verlag, New York, 1991.
- [15] G. Marks, On 2-primal Ore extensions, Comm. Algebra 29 (2001), 2113-2123.
- [16] R. Nikandish, M.J. Nikmehr, A. Yassine, More on the 2-primary ideals of commutative rings, Bull. Korean Math. Soc. 57(2020), 117-126.
- [17] L.H. Rowen, Ring Theory, Academic Press, San Diego, 1991.
- [18] A.D. Sands, Radical and Morita contexts, J. Algebra 24 (1973), 335-345.
- [19] G. Shin, Prime ideals and sheaf representation of a pseudo symmetric ring, Trans. Amer. Math. Soc. 184 (1973), 43-60.