

On the set of all generalized Drazin invertible elements in a ring

Fei Peng* and Xiaoxiang Zhang

Southeast University
E-mail: pfmath@163.com

Berkani and Sarihr [Studia Math. (2001) 148: 251-257] showed that the set of all Drazin invertible elements in an algebra over a field is a regularity in the sense of Kordula and Muller [Studia Math. (1996) 119: 109-128]. In this note, the above result is extended to the case of a ring. Counterexamples are provided to show that the set of all generalized Drazin invertible elements in a ring need not be a regularity in general. We determine when the set of all generalized Drazin invertible matrices in the 2×2 full matrix ring over a commutative local ring is a regularity. We also give a sufficient condition for the set of all generalized Drazin invertible elements in a ring to be a regularity.

References

- [1] M. Berkani and M. Sarihr, An Atkinson-type theorem for B-Fredholm operators, Studia Math. 148 (2001) 251-257.
- [2] J. Cui, J.L. Chen, When is a 2×2 matrix ring over a commutative local ring quasipolar? Comm. Algebra 39 (2011) 3212-3221.
- [3] M.P. Drazin, Pseudo-inverses in associative rings and semigroups, Amer. Math. Monthly 65 (1958) 506-514.
- [4] R. Harte, On quasinilpotents in rings, Panamer. Math. J. 1 (1991) 10-16.
- [5] V. Kordula, V. Muller, On the axiomatic theory of spectrum, Studia Math. 119 (1996) 109-128.
- [6] J.J. Koliha, P. Patricio, Elements of rings with equal spectral idempotents, J. Austral. Math. Soc. 72 (2002) 137-152.
- [7] R.A. Lubansky, Koliha-Drazin invertibles form a regularity, Math. Proc. R. Ir. Acad. 107A (2007) 137-41.

2020 Mathematics Subject Classification: 16U90

Keywords: Regularity, Drazin inverse, generalized Drazin inverse, ring