Some Properties of Left r-Clean Bimodules

Dian Ariesta Yuwaningsih*, Indah Emilia Wijayanti and Budi Surodjo

Universitas Gadjah Mada E-mail: dian.ariesta.yuwaningsih@mail.ugm.ac.id

Let R be an associative ring with identity and M an R-bimodule. Ring R is considered clean if each element is the sum of idempotent and unit elements. By generalizing the unit elements to regular elements, the definition of a clean ring has been generalized to an r-clean ring. Previous researchers have developed a clean ring definition and their generalization to the module structure through its endomorphism ring. This research introduces the generalization of r-clean rings, called left r-clean R-bimodules, defined without their endomorphism rings. Furthermore, we present some of their properties and give the sufficient and necessary conditions for an R-bimodule to be r-clean.

References

- [1] Ali, M.M., 2008, Idempotent and Nilpotent Submodules of Multiplication Modules, Communication in Algebra, Vol. 36, No. 1, pp. 4620-4642.
- [2] Ashrafi, N. and Nasibi, E., 2013, On r-Clean Rings, Mathematical Reports, Vol. 15, No. 2, pp. 125-132.
- [3] Ashrafi, N. and Nasibi, E., 2013, Rings in Which Elements are the Sum of An Idempotent and A Regular Element, Bulletin of the Iranian Mathematical Society, Vol. 39, No. 3, pp. 579-588.
- [4] Camillo, V.P., Khurana, D., Lam, T.Y., Nicholson, W.K., and Zhou, Y., 2006, Continuous Modules are Clean, Journal of Algebra, Vol. 304, pp. 94-111.
- [5] Chen, H. and Chen, M., 2003, On Clean Ideals, International Journal of Mathematics and Mathematical Sciences, Vol. 62, pp. 3949-3956.
- [6] Malik, D.S., Mordeson, J.N., and Sen, M.K., 1977, Fundamentals of Abstract Algebra, The McGraw-Hill Companies, Inc., New York.
- [7] Nicholson, W.K., 1977, Lifting Idempotents and Exchange Rings, Transactions of the American Mathematical Society, Vol. 229, pp. 269-278.
- [8] Nicholson, W.K., Vadarajan, K., and Zhou, Y., 2004, Clean Endomorphism Rings, Arch. Math., Vol.83, pp. 340-343.