



Contents lists available at ScienceDirect

Journal of Quantitative Spectroscopy & Radiative Transfer

journal homepage: www.elsevier.com/locate/jqsrt

Note

Broadening of the $J = 3 \leftarrow 2$ spectral line of carbon monoxide by pressure of CO, N₂ and O₂

M.A. Koshelev*, V.N. Markov

Institute for Applied Physics of the Russian Academy of Sciences, 46 Uljanova St., 603950 Nizhnii Novgorod, Russia

ARTICLE INFO

Article history:

Received 23 January 2009

Received in revised form

3 February 2009

Accepted 5 February 2009

PACS:

33.20.Bx

33.70.Jg

Keywords:

Carbon monoxide

Pressure broadening

Line shifting

Line shape studies for the investigation of collisional broadening and shift of spectral lines of molecules are of interest in astrophysics, planetary atmosphere studies, propagation of radiation in the Earth's atmosphere, quantum chemistry, etc. The employment of complementary experimental techniques and methods for such studies is very desirable since it may reveal possible sources of error and can yield very reliable line-shape parameters.

Here we compare results of our investigation on the collisional relaxation for the $J = 3 \leftarrow 2$ rotational line of CO in the ground vibrational state induced by nitrogen and oxygen using a spectrometer with radio acoustic detection of absorption (RAD—spectrometer) [1]. Comparison of our data with the results of Ref. [2], where a video-type sub-mm spectrometer was used for analogous investigations, is presented. The self-broadening study of the line is presented in this note as well.

The RAD spectrometer and method of the line shape investigation were described in detail in Ref. [3]. The length of the absorption cell was about 17 mm and the corresponding optical path-length is about 1.4×10^{-2} (the line intensity was taken from the HITRAN database [4]). All measurements were carried out at 297 ± 1 K. The Voigt profile was used for the line-shape parameter determination from the experimental data. Typical dependences of the collisional line half width at half maximum (HWHM) on pressure of CO, N₂, and O₂ are depicted in Fig. 1. The pressure broadening coefficient was obtained as a slope of the straight line fitted to the corresponding experimental dependence. The results of the measurements of the pressure broadening coefficient of the $J = 3 \leftarrow 2$ spectral line of carbon monoxide by pressure of N₂ and O₂ in comparison with data from Ref. [2] are presented in Table 1. The self-broadening coefficient of the line was measured as 3.135(10) MHz/Torr. Table 1 shows that our results are in good agreement within quoted experimental uncertainties. This agreement obtained by means of different techniques for the line-shape study increases the reliability of

* Corresponding author. Tel: +7 831 4164866; fax: +7 831 4363792.

E-mail address: koma@appl.sci-nnov.ru (M.A. Koshelev).




Fig. 1. Pressure dependence of the $J = 3 \leftarrow 2$ CO line half width for CO (●), N₂ (Δ) and O₂ (■) as buffer gasses. Solid lines are results of linear regression of the experimental data.

Table 1

Pressure broadening coefficient of the $J = 3 \leftarrow 2$ spectral line of carbon monoxide by pressure of N₂ and O₂ in comparison with data from Ref. [2].

Buffer gas	Broadening coefficient MHz/Torr	References
N ₂	2.895(10)	This work
	2.885(6)	Ref. [2]
O ₂	2.570(10)	This work
	2.516(37)	Ref. [2]

Errors in parentheses correspond to one standard uncertainty in unit of the last digit quoted.

the measured data. Our study did not reveal any noticeable shift of the line center within experimental uncertainty ± 5 kHz/Torr. The observed rest frequency of the line center is 345795.987(3) MHz, which is in good agreement with results of the Lamb-dip measurements [5] proving consistency of our data.

The investigations described in this note were supported in parts by Russian Foundation for Basic Research (RFBF) and Deutsch Forschungsgemeinschaft (DFG). The authors express their deep gratitude to these institutions.

References

- [1] Krupnov AF. In: Chantry GW, editor. Modern aspects of microwave spectroscopy. London: Academic Press; 1979. p. 217.
- [2] Priem D, Rohart F, Colmont JM, Włodarczak G, Bouanich JP. Lineshape study of the $J = 3 \leftarrow 2$ rotational transition of CO perturbed by N₂ and O₂. J Mol Struct 2000;517–518:435–54.
- [3] Markov VN, Golubiatnikov GYu, Savin VA, Sergeev DA, Guarnieri A, Maeder H. Line broadening and shifting studies of the $J = 5 \leftarrow 4$ transition of carbon monoxide perturbed by CO, N₂, and O₂. J Mol Spectrosc 2002;212:1–5.
- [4] Rothman LS, Jacquemart D, Barbe A, Benner DC, Birk M, Brown LR, et al. The HITRAN 2004 molecular spectroscopic database. JQSRT 2005;96:139–204.
- [5] Winnewisser G, Belov SP, Klaus Th, Schieder R. Sub-Doppler measurements on the rotational transitions of carbon monoxide. J Molec Spectrosc 1997; 184:468–72.