

LETTER TO THE EDITOR

Microwave Measurements of $J = 2 \leftarrow 1$, $K = 0, 1$ Ammonia Transitions at 1.215 THz

A. F. Krupnov,¹ M. Yu. Tretyakov,¹ M. Bogey, S. Bailleux, A. Walters, B. Delcroix, and S. Civiš²

Laboratoire de Spectroscopie Hertzienne, associe au CNRS, Universite des Sciences et Technologies de Lille, 59655 Villeneuve d'Ascq Cedex, France

Received January 24, 1996




FIG. 1. The submillimeter-wave absorption signal of the $J = 2 \leftarrow 1$, $K = 1$, $a \leftarrow s$ $^{14}\text{NH}_3$ ammonia transition.

The first rotational transition of the ammonia molecule at 572 GHz was previously measured by microwave spectroscopy in (1, 2). In the present letter, measurements of the two second rotational transitions of the ammonia molecule $J = 2 \leftarrow 1$, $K = 0, 1$ at 1214.8 and 1215.2 GHz are reported.

The experimental apparatus was based on the Lille millimeter-wave/submillimeter-wave spectrometer (3, 4), with the addition of a high-frequency Russian-constructed BWO produced by the ISTOK Research and Production Co. This new BWO was phase-locked against an existing Thomson BWO, operating around 405 GHz, using a similar broadband multiplier-mixer to that first used in (5). The Thomson BWO was in turn phase-locked as explained in previous articles (3, 4).

Neither the geometry of the cell nor the InSb bolometer was optimized

¹ Permanent address: Applied Physics Institute of Russian Academy of Sciences, 46 Uljanova Street, 603 600 Nizhnii Novgorod, Russia.

² Permanent address: J. Heyrovský Institute of Physical Chemistry, Academy of Sciences of the Czech Republic, Dolejškova 3, 182 23 Prague 8, Czech Republic.

TABLE 1
Observed and Calculated Frequencies of Ground State $^{14}\text{NH}_3$ Ammonia Transitions

Ammonia Transitions	Obs. Freq. MHz	Calc. Freq. ^a MHz	O - C MHz
$J = 2 \leftarrow 1, K = 0$	1 214 852.874(140)	1 214 852.6	0.3
$J = 2 \leftarrow 1, K = 1$	1 215 245.674(178)	1 215 245.3	0.4

^aRef.6.

for the terahertz frequency region; a 3-m-long cell and nonmagnetically tuned bolometer were used. Nor was any attempt made to reduce power loss by water absorption in the path of the terahertz radiation through the atmosphere. Sample pressures less than 0.5 Pa were used to prevent saturation of the lines.

A typical example of measured spectral lines is presented in Fig. 1.

Observed frequencies in MHz are presented in Table 1 together with previously predicted frequencies calculated from simultaneous analysis of the microwave, submillimeter-wave, far-infrared, and infrared-microwave two-photon transitions between the ground and ν_2 inversion-rotation levels of $^{14}\text{NH}_3$ (6). The calculated frequencies were converted from values quoted in cm^{-1} to seven significant digits and are hence given with eight digits. The observed frequency quoted is the mean of seven independent measurements for each line taken with varying time constants (30–100 msec). No frequency shift correlated to the time constant was observed. The uncertainty quoted for the measured results is the standard deviation of these seven measurements. This standard deviation is comparable to the uncertainty we would predict based on our experiments at lower frequencies. Linewidths observed were close to the Doppler limit at this frequency.

With the addition of these two new transitions measured by microwave techniques the low- J rotational spectrum of ammonia is fixed by three, not one, microwave points for further fits. Although a fit of these microwave transitions alone would not be expected to lead to molecular constants better than those already available, their inclusion in a global fit should give improvements in the quality of future spectral predictions for this molecule. These frequencies may be useful for the calibration of Fourier transform spectrometers in this region.

ACKNOWLEDGMENTS

Studies were funded partly by the French Research Ministry and the CNRS (INSU grant), partly by the Russian Fund for Fundamental Studies

(Grant 94-02-05424-a), and the Ministry of Science and Technical Politics through the State Program on Fundamental Metrology. M.Yu.T. thanks the European Union (PECO Contract CIPDCT 94-0614), A.F.K. thanks the Nord-Pas de Calais Region, and S.C. thanks the CNRS for supporting their stay in Lille.

REFERENCES

1. A. F. Krupnov, V. A. Skvortsov, and L. A. Sinegubko, *Izv. Vyssh. Uchebn. Zaved. Radiofiz.* **11**, 1186–1190 (1968).
2. P. Helminger, F. C. De Lucia, and W. Gordy, *J. Mol. Spectrosc.* **39**, 94–97 (1971).
3. M. Cordonnier, M. Bogey, C. Demyunck, and J.-L. Destombes, *J. Chem. Phys.* **97**, 7984–7989 (1992).
4. S. Bailleux, M. Bogey, C. Demyunck, J.-L. Destombes, and A. Walters, *J. Chem. Phys.* **101**, 2729–2733 (1994).
5. G. Winnewisser, A. F. Krupnov, M. Yu. Tretyakov, M. Liedtke, F. Lewen, A. H. Saleck, R. Schieder, A. P. Shkaev, and S. A. Volokhov, *J. Mol. Spectrosc.* **165**, 294–300 (1994).
6. S. Urban, V. Spirko, D. Papousek, J. Kauppinen, S. P. Belov, L. I. Gershtein, and A. F. Krupnov, *J. Mol. Spectrosc.* **88**, 274–292 (1981).