

Note

Broadening and shifting studies of $J = 5 \leftarrow 4$ carbon monoxide line perturbed by Ne, Ar, and Kr

V.N. Markov^{a,*} and D.A. Sergeev^b

^a Applied Physics Institute, Russian Academy of Science, Uljanova Street 46, Nizhnii Novgorod 603950, Russia

^b Advanced School of General and Applied Physics, NN State University, Uljanova Street 46, Nizhnii Novgorod 603950, Russia

Received 28 April 2003; in revised form 6 October 2003

A growing interest in determination of molecular line parameters—such as pressure broadening, pressure shift, and lineshape makes development of accurate methods of measurements very desirable. Recently, few works were published concerning this problem in the mm and submm region [1–3].

This paper continues our previous lineshift and line broadening studies of carbon monoxide molecule induced by collision with different types of gases [4–7]. Since carbon monoxide (CO) is a simple diatomic molecule, the choice of the rare gases (Ne, Ar, and Kr) was motivated by the development of theories describing collisional line broadening and shifting phenomena, the experimental data are useful for deducing intermolecular potential parameters.

We also desire to compare our data with data [3], where analogous investigation was performed but by using a different method.

All investigations were made at room temperature (295 ± 1) K. A detailed description of the experimental technique and general methods of determination of halfwidth and linecenter frequency is presented [4].

Shortly, for this study the submillimeter RAD spectrometer with backward wave oscillator (BWO) and acoustic cell was used [8]. The BWO frequency was phase stabilized and controlled by a computer via GPIB. The amplitude modulation of submm radiation was used for this investigation. The lineshape experimental data were fitted to the Voigt model function including baseline influence [4].




Fig. 1. Experimental dependence of HWHM of $J = 5 \leftarrow 4$ CO line broadened by Ar and Ne gases.

The example pressure dependence of HWHM of $J = 5 \leftarrow 4$ CO line on the pressure of Ar and Ne gases is depicted in Fig. 1. The dependences of the center frequency of the $J = 5 \leftarrow 4$ CO line on the pressure of rare gases (Ne, Ar, and Kr) are presented in Fig. 2. Comparison of our experimental results with data [3] is presented in Table 1.

As we can see our data have a good coincidence with results in the paper [3], but the accuracy of determination of pressure shift coefficients is much high in our work.

* Corresponding author. Fax: +7-8312-363792.

E-mail address: markov@appl.sci-nnov.ru (V.N. Markov).




Fig. 2. Dependences of center frequency of $J = 5 \leftarrow 4$ CO line on the pressure buffer gases Ne (●), Ar (○), and Kr (■).

Table 1
Pressure broadening coefficient δ and shift coefficient of $J = 5-4$ spectral line CO molecule for Ne, Ar, and Kr

Perturbed gas	γ (MHz/Torr)	δ (kHz/Torr)	Reference
Ne	1.33(2)	23(4)	This work
	1.383(47)	24.8(123)	[3]
Ar	2.04(2)	45(5)	This work
	1.917(31)	54.8(136)	[3]
Kr	2.08(2)	49(5)	This work
	1.935(31)	64.5(165)	[3]

Numbers in parentheses are one standard deviation obtained from least-squares analysis.

Acknowledgments

The investigations described in this paper were supported by the Deutsche Forschungsgemeinschaft (DFG) and Russian Fund of Basics Research (RFBR), Grant 03-02-16125 and the Contract of the Ministry of Industry, Science and Technology of Russian Federation. The authors express their deep gratitude to these institutions. They are also grateful to Professor A.F. Krupnov for his interest in this work and for helpful criticism of various points in an earlier draft.

References

- [1] D. Priem, F. Rohart, J.M. Colmont, G. Woldarczak, J.-P. Bouanich, *J. Mol. Struct.* 517/518 (2000) 435–454.
- [2] C. Pazzarini, L. Dore, G. Cazzoli, *J. Mol. Spectrosc.* 216 (2002) 428–436.
- [3] K.M.T. Yamada, H. Abe, *J. Mol. Spectrosc.* 217 (2003) 87–92.
- [4] H. Mäder, A. Guarnieri, J. Dose, N. Nissen, V.N. Markov, A.M. Shtanyuk, A.F. Andrianov, V.N. Shanin, A.F. Krupnov, *J. Mol. Spectrosc.* 180 (1996) 183–187.
- [5] N. Nissen, J. Dose, A. Guarnieri, H. Mäder, V.N. Markov, G.Yu. Golybyatnikov, I.I. Leonov, V.N. Shanin, A.F. Krupnov, *Z. Naturforsch. A* 54 (1999) 218–224.
- [6] V.N. Markov, H. Mäder, *J. Mol. Spectrosc.* 205 (2001) 350–352.
- [7] V.N. Markov, G.Yu. Golybyatnikov, V.A. Savin, D.A. Sergeev, A. Guarnieri, H. Mäder, *J. Mol. Spectrosc.* 212 (2002) 1–5.
- [8] A.F. Krupnov, in: G.W. Chantry (Ed.), *Modern Aspects of Microwave Spectroscopy*, Academic Press, London, 1979, pp. 217–256.