Journal of Quantitative Spectroscopy & Radiative Transfer 337 (2025) 109379

Contents lists available at ScienceDirect

Journal of Quantitative Spectroscopy and Radiative Transfer

journal homepage: www.elsevier.com/locate/jqsrt

Collisional parameters of the pure rotational R(0) line of CO in N_2: Experiment versus theory

D.S. Makarov *, I.N. Vilkov, T.A. Galanina, E.A. Serov, M.A. Koshelev, M.Yu. Tretyakov

Federal Research Center A.V. Gaponov-Grekhov Institute of Applied Physics of the Russian Academy of Sciences (IAP RAS), 46 Ulyanov Str., 603950 Nizhny Novgorod, Russia

ARTICLE INFO

Keywords:
Microwave spectroscopy
High resolution
Carbon monoxide
Line shape analysis
Collisional parameters
Speed dependent effects

ABSTRACT

Line shape of the first rotational transition of the CO molecule perturbed by N_2 collisions was studied beyond the traditional Voigt profile at pressures from 0.1 up to 1500 Torr and temperatures from 238 up to 353 K using two different spectrometers with complementary abilities. Line intensity, shifting, mixing, broadening, and underlying continuum coefficients were retrieved. The experimental speed dependence (SD) broadening parameters and their temperature dependence were compared to the results of the recent quantum scattering calculations showing fair agreement yet pointing to possible necessity of improvement of the theoretical approach to the CO–N_2 collisional system.

1. Introduction

Carbon monoxide is one of the most frequently studied molecules (near 18 000 search results found by Google Scholar dated by 2024). The necessity and the importance of a deep understanding of its internal dynamics and interaction properties are exhaustively and in detail described in the Introduction section of the recent paper [1], hereinafter referred to as Paper I, which is devoted to the high accuracy *ab initio* study of collisional broadening of the R(0) line of CO molecule in N_2 environment. Earlier experimental studies of the R(0) line by means of a videospectrometer covering wide temperature ranges employ only Lorentz [2] or Voigt [3] line shape model for the analysis of the recorded data. In a joint measurements using Fourier spectrometer and radioacoustic spectrometer [4] Voigt profile was also used. Study performed at room temperature by means of a videospectrometer operating in the modes of radiation power and frequency modulation [5] also considers Galatry and Rautian model profiles together with the Voigt profile, showing the insufficiency of the latter one. In addition, Raman spectra study should be mentioned [6], using the same potential energy surfaces as in Paper I to compare calculated and measured broadening parameters, thus still employing the Voigt model.

The main goal of the present paper is direct experimental verification of the Paper I calculations which were initially planned as part of the joint project by analogy with our previous studies described in [7,8] (the latter is referred to as Paper II). As *ab initio* calculations in Paper I result in a dependence of the collisional cross-section on kinetic energy and corresponding speed-dependence of the broadening coefficient, it is accounted for in the experimental data analysis for the first time.

The paper has the following structure. Section 2 gives a brief overview of the experimental techniques used for the studied line profile recordings under different thermodynamic conditions and approaches used for processing the obtained data. In Section 3 the retrieved values of the line shape parameters are analyzed and compared with the results of *ab initio* calculations. The main findings of the study are summarized in Section 4.

2. Experimental techniques and data processing

2.1. Data acquisition

The profiles of the R(0) line of the most abundant isotopologue $^{12}C^{16}O$ in the ground vibrational state were recorded by means of the radioacoustic detection (RAD) spectrometer [9] and resonator (cavity) spectrometer [10]. The temperature range covered by the RAD spectrometer measurements was from 238 to 353 K, 20 recording batches were obtained at 12 temperature values. The data batches were obtained as follows: the first small amount of CO was taken (140–150 mTorr) and the line profile was recorded in pure CO, then N_2 was added step-by-step, typically by portions of ~300 mTorr up to the pressure near 2 Torr with subsequent line profile recording. The radiation source (BWO, backwards wave oscillator) was phase-locked to the GPS-tracked frequency standard, its power was optimized to reduce the radiation intensity and thus the effect of molecular transition saturation. For each recording, multiple scans (151 frequency points per

* Corresponding author.
E-mail address: dmak@ipfran.ru (D.S. Makarov).

https://doi.org/10.1016/j.jqsrt.2025.109379
Received 20 November 2024; Received in revised form 29 January 2025; Accepted 29 January 2025
Available online 7 February 2025
0022-4073/© 2025 Elsevier Ltd. All rights are reserved, including those for text and data mining, AI training, and similar technologies.



Fig. 1. RAD spectrometer series of step-by-step recordings at 303 K (points, upper panels) and fit residuals (scaled by the factor of 100) for the speed-independent Voigt profile (upper solid traces) and qSDRP for cases of line-by-line and multifit approach (middle and lower solid traces, respectively). Fit quality Q is shown near each residual.

scan, 0.5 s per point) through the frequency range were performed with further averaging of the spectra, typically over 16 scans. The recordings were obtained in the radiation frequency manipulation (FMp) mode with precisely known frequency deviation value and subsequent 1-F synchronous demodulation of the acquired signal (see Paper II for details).

The resonator measurements were carried out at room temperature (297–297.5 K), employing various partial pressures of CO and N_2 with the total pressure up to about 2 atm. The intrinsic part of losses (baseline) was recorded separately with a gas chamber filled with pure argon and subtracted from the data recorded in the CO–N_2 mixture. The calculated contribution of all other (except the R(0)) lines of CO was subtracted from the recordings, including the wings of the subsequent 7 rotational lines as well as the weak lines of less abundant isotopologues located nearby and the higher vibrational state using the approach described in Paper II. The air pressure broadening coefficients were taken from HITRAN [11] and recalculated to N_2 pressure broadening ones based on the values from [12]. A method of the line mixing coefficients calculation for CO–N_2 basing on the values for CO–He [13] is described in detail in Paper II.

2.2. Low pressure data treatment

The recorded spectra were treated using least-square fitting of the model function to the experimental data employing the Levenberg — Marquardt method. The Hartmann — Tran profile (hereinafter referred to as HTP), reduced to the quadratic speed-dependent Rautian profile (qSDRP) by setting the correlation parameter η to zero as in Paper II, was employed as a line shape model [14]. Thus, the parameters of interest were the following: central frequency f_0, speed-averaged collisional broadening Γ_0, quadratic SD broadening parameter Γ_2, and speed-averaged collisional shift of the central frequency Δ_0. The quadratic speed-dependent shift Δ_2 was negligibly small and unobservable with signal-to-noise ratio (SNR) achieved in our experiment. The specified values describe the line profile at definite pressure and temperature. Considering the linear dependence of Γ_0, Γ_2 and Δ_0 on pressure, the corresponding pressure-independent line shape coefficients were denoted as γ_0, γ_2 and δ_0. The speed-independent Voigt profile was also used, which enabled an adequate comparison of the retrieved collisional line broadening coefficient ($\gamma_V = \Gamma_V/P$) and its temperature dependence with the previously reported data.

There are two approaches to the data treatment to obtain pressure-independent coefficients. The first one is "line-by-line", when each recording of the batch is analyzed individually, with the following analysis of the line shape parameters versus expected pressure dependence. The second one is the multispectrum fit approach (or "multifit"), which suggests analysis of the whole recording batch with predefined pressure dependence of the parameters. The results of the model profiles fitting within different approaches are shown in Fig. 1, confirming that using Voigt profile is insufficient at current sensitivity level. The better fit quality in case of line-by-line approach is caused by the influence of extra uncertainties (e.g., pressure and temperature measurement errors) on the fit results in case of the multifit approach.

Besides a proper line shape model, two hardware factors were taken into account: (i) radiation power-versus-frequency dependence was accounted for as a linear with frequency term multiplied to the line profile function, and (ii) the polynomial up to the third-order was added to the model to account for the baseline, i.e. an additional absorption signal caused by the radiation absorbed in the radioacoustic cell elements.

For RAD recordings treatment, the effective (i.e. observed in the experiment) speed-averaged collisional width Γ_0 was considered as

$$\Gamma_0(P_{N_2}) = \sqrt{\left(\Gamma_{self} + \gamma_0^{N_2} \cdot P_{N_2}\right)^2 + \Omega_R^2}, \quad (1)$$

where Γ_{self} is the self-broadening component of the collisional width, $\gamma_0^{N_2}$ is the nitrogen pressure broadening coefficient, P_{N_2} is the partial pressure of nitrogen, and Ω_R is the Rabi frequency, an additional adjusted parameter to describe the molecular transition saturation effect by radiation power (see Paper II for details). Typical dependences of Γ_0, Γ_2 and Δ_0 (with the subtracted self-pressure components derived within current treatment) on P_{N_2} are shown in Fig. 2. The points display the results of the line-by-line treatment, while the lines display pressure dependences for the corresponding temperatures obtained with the multifit treatment. This figure clearly shows the limitations of the line-by-line approach: at higher pressures (and wider frequency scan ranges) the parameters corresponding to the weak effects may noticeably deviate from the expected dependences due to the correlation with the hardware-related parameters.

It was impossible to fit velocity-changing collisions rate parameter ν_{vc} due to its strong correlation with the speed-dependent width parameter Γ_2: at pressures typical for RAD recordings, the manifestations of both effects are similar in line shape in terms of form and magnitude. The Dicke effect is pronounced in the line shape if the Doppler width is greater than the collisional one, which is true for the lowest pressures the measurements were carried out at (below 0.1–0.2 Torr), which makes this effect almost negligible for almost all the recording, contrary to the SD effect which is valuable for all recordings at any pressure. Currently, the most reliable solution seems to use fixed ν_{vc}, starting with the result of some preliminary evaluation (e.g., *ab initio* calculation



Fig. 2. Γ_0, Γ_2 and Δ_0 as a function of N_2 pressure (self-pressure contributions are subtracted) for a number of temperatures, obtained from line-by-line treatment of RAD spectrometer recordings (points) and from multifit (lines) for the same temperatures. Error bars correspond to 1σ fitting uncertainty.

from [7], assuming that collisions with non-polar species Ar and N_2 might be similar) and to vary it slightly looking at the $\Gamma_{0,2}$ vs. P dependence. The smaller the systematic discrepancies of $\Gamma_{0,2}$ from the expected dependence, the closer the ν_{vc} value to the "true" value. Due to the correlation, the varying fixed ν_{vc} value also slightly changes half-width parameters $\Gamma_{0,2}$ and the corresponding broadening coefficients $\gamma_{0,2}$. The derivatives of γ_0 and γ_2 by ν_{vc} in numerical simulation were found as 0.013 and 0.064, respectively, and thus taken into account together with the uncertainty of ν_{vc} to calculate total uncertainty of the studied parameters.

At 303 K, ν_{vc} for CO–N_2 collisions was changed from 0.0 to 0.9 MHz/Torr, which varied γ_0 and γ_2 within about 0.01 and 0.05 MHz/Torr (0.3% and 15%), respectively. In both cases this systematic change exceeded 3σ statistical interval. The most plausible ν_{vc} value at 296 K was estimated as 0.3(1) MHz/Torr, which is quite in line with the values found for the CO–Ar mixture in Paper II and [7] (0.151 and 0.301 MHz/Torr, respectively). In addition, the influence of the temperature dependence of ν_{vc} was estimated using the *ab initio* value of $n_{\nu_{vc}} = 0.9$ (which is in agreement with $n_{\nu_{vc}} = 1$ as supposed in [14]) for CO–Ar collisions from [7] assuming (i) their similarity with CO–N_2 collisions, and (ii) the applicability of the single power law for the temperature dependences:

$$X(T) = X(T_0) \cdot \left(\frac{T_0}{T}\right)^n, \tag{2}$$

where X stands for $\gamma_{0,2}$, δ_0 or ν_{vc}. Note that possible uncertainty in $n_{\nu_{vc}}$ leads to the corresponding error of $\nu_{vc}(T)$ no more than 2%–3% at the edges of the studied temperature range, which is much smaller than 0.1 MHz/Torr (or 33%) uncertainty of the manual adjustment and may be neglected when calculating the total error budget. The same reasoning allows neglecting possible temperature dependence of the correlation between $\nu_{vc}(T)$ and $\gamma_{0,2}(T)$.

Having adjusted ν_{vc} and its temperature dependence, all the low pressure data were processed both within the line-by-line and the multifit approaches. The latter allows not only direct derivation of the pressure-independent coefficients, but also reduces statistical uncertainties and possible correlations between the studied parameters.

2.3. High pressure data treatment

In the recordings taken at a pressure near and above 1 atm, the line mixing effect is highly pronounced. In our case it is accounted for by one additional parameter proportional to pressure in the model function known as the Rozenkranz correction [15]. This correction is also incorporated into the up-to-date version of the HTP used in the HAPI [16] code and HITRAN database [11]. As was shown in Paper II, the manifestation of the speed-dependence of the line mixing effect cannot be observed with the SNR insured by the present-day resonator spectrometer.

In addition, at atmospheric pressures the continuum absorption is noticeable. In the vicinity of the R(0) line, the frequency dependence of the continuum is very close to frequency squared, which is stipulated by the radiation term. The pressure dependence of the continuum within the binary collision approximation (which is safely valid under our experimental conditions) is well known, so the continuum absorption can be modeled as

$$C(f, P_{\mathrm{CO}}, P_{\mathrm{N_2}}) = \left(C_{self} \cdot P_{\mathrm{CO}}^2 + C_{foreign} \cdot P_{\mathrm{CO}} \cdot P_{\mathrm{N_2}}\right) \cdot f^2, \tag{3}$$

where f is the radiation frequency, C_{self} and $C_{foreign}$ are the self- (i.e. provided by CO–CO pairs) and foreign- (CO–N_2 pairs) continua coefficients, P_{CO} is the CO partial pressure. This term is added to the line profile model.

In contrast to RAD data treatment, the spectrometer baseline (cavity losses) is determined experimentally. Thus, no additive terms accounting for the baseline are necessary.

Fig. 3 shows the results of the fitting model profiles with and without speed-dependences of the collisional broadening to the resonator spectrometer recordings. The manifestation of the SD effect is also observed in the fit residuals similar to the RAD spectrometer data. The use of more sophisticated line shape model which takes into account the SD effect demonstrates almost noise-like residuals. Some systematic discrepancies in case of the multifit approach are supposed to have the same origin as for the low-pressure data.

3. Retrieved parameters and their comparison with *ab initio* calculations

Temperature dependences of the broadening coefficients $\gamma_{0,2}$ and shifting coefficient δ_0 are based on the RAD spectrometer data. The multifit approach allowed deriving these coefficients directly, while the line-by-line approach demands analysis of the pressure dependences



Fig. 3. The same as Fig. 1 but for the resonator spectrometer (residuals are scaled by the factor of 100). Note three orders of magnitude difference in the frequency scale.



Fig. 4. Temperature dependence of N_2-pressure shifting coefficient. Error bars are 1σ fit uncertainty. Grey zone is for the 95% confidence interval of the final dependence shown by black solid line.

first (the latter is also used for verification of RAD spectrometer data together with the resonator spectrometer ones). For γ_0 and γ_2, the results of the line-by-line and multifit treatment of the RAD spectrometer data agree within the 3σ interval. We consider the multifit results as more reliable and they will be mostly shown and discussed (if the opposite is not specified).

The difference between the results of the two fitting approaches is demonstrated on an example of the temperature dependence of the pressure shifting coefficient δ_0 in Fig. 4. Absolute shifting coefficient values are small and significantly scattered. In addition, there is no unambiguous physically grounded expression for the temperature dependence of this coefficient (see, e.g., discussion in [17]). In such a situation, the linear dependence assumption seems appropriate. Due to the large uncertainty of the data there is no preference between line-by-line or only multifit results. In this case, parameters of the final linear temperature dependence were derived using both sets of values as independently obtained data. The corresponding expression is

$$\delta_0(T) = -4(5) - 0.044(10) \times (T - 296) \quad \text{kHz/Torr},$$

where T is in Kelvin and the uncertainties are given in the units of the last significant digit.

Thorough verification of the retrieved values of γ_0 and γ_2 can be made using data from the two spectrometers at significantly different (near three orders of magnitude) pressures. Fig. 5 shows all measured



Fig. 5. All retrieved CO–N_2 collisional broadening parameters Γ_0 and Γ_2 recalculated to 296 K. Expected linear pressure dependences (solid and dashed lines in the upper panel) are determined using low-pressure data only and extrapolated to high pressures. Lower panels show corresponding pressure-normalized obs.-minus-calc. values.

Γ_0 and Γ_2 values for the CO–N_2 collisional system recalculated to 296 K using the single power law (2) with temperature-dependent parameters derived in this work from the RAD spectrometer data and self-broadening coefficients from Paper II (numerical values are given in Table 2 of the current study). Plotted linear functions are derived using only RAD data. Despite this, the pressure-normalized observed-minus-calculated values (lower part of the figure) are grouping near the zero level in both pressure ranges demonstrating perfect agreement between the two datasets. This agreement confirms reliability of the results and minimization of possible systematic errors.



Fig. 6. Line mixing parameter (upper panel), frequency-normalized continuum contribution (central panel) and central frequency shift from the unperturbed value 115271.202 MHz (lower panel) normalized to CO partial pressure versus N_2 to CO partial pressures ratio (see Eq. (4) and discussion) at ~297 K. Uncertainty intervals correspond to 1σ.

The line integrated intensity, shifting, mixing and foreign continuum coefficients were retrieved from the high pressure recordings. The following dependences were used for evaluating the line-by-line results:

$$\begin{aligned} \frac{Z}{P_{\mathrm{CO}}} &= z_{self} + z_{foreign}\frac{P_{\mathrm{N_2}}}{P_{\mathrm{CO}}} \\ \frac{C}{P_{\mathrm{CO}}^2 \cdot f^2} &= C_{self} + C_{foreign}\frac{P_{\mathrm{N_2}}}{P_{\mathrm{CO}}} \end{aligned} \tag{4}$$

where Z/z is for either Y/y or Δ_0/δ_0, Z and C are the parameter values obtained from the spectra recordings, while $z_{foreign}$ and $C_{foreign}$ are the required coefficients. The coefficients z_{self} and C_{self} were taken from Paper II and fixed in the multifit procedure.

The results are shown in Fig. 6. The difference between the line-by-line and multifit results is slightly greater than the statistical uncertainty of $y_{foreign}$ and $C_{foreign}$ coefficients obtained from each fitting approach but the results agree within 3σ. Similar to the argon atom, an N_2 molecule has no dipole moment and also has smaller mass. Consequently, it was expected that these parameters have close values for the CO–N_2 and CO–Ar collisional systems. Indeed, the resulting values $y_{foreign} = 7.95(22)$ 1/Torr and $C_{foreign} = 4.80(56) \cdot 10^{-25}$ cm^{-1}/MHz^2 Torr^2 retrieved from the recordings at ~297 K agree within about 10% with the corresponding results from Paper II (which were retrieved at similar temperature). The pressure shifting coefficient $\delta_{foreign} = -4.8(8)$ kHz/Torr also agrees with the RAD spectrometer measurements at the corresponding temperature. The integrated intensity value retrieved from the resonator data is $3.277(9) \cdot 10^{-24}$ cm/molec, which agrees with $3.2814(73) \cdot 10^{-24}$ cm/molec from Paper II and is slightly less than $3.3 \cdot 10^{-24}$ cm/molec given in HITRAN (errors in parentheses correspond to 1σ fit uncertainties).



Fig. 7. Comparison of the Voigt profile collisional broadening parameter obtained in current and in earlier studies.

Table 1
The speed-independent CO–N_2 broadening coefficient γ_V at 296 K and its temperature exponent. Uncertainties of the values retrieved in this study include fit uncertainty combined with errors of pressure and temperature readings like in Paper II.

Source	γ_V, MHz/Torr	n_{γ_V}
This study (RAD)	3.190(7)	0.775(4)
Colmont [3]	3.3(1)	0.86(12)
Connor [2]	3.02(14)	0.74(10)
Nissen FT [4]	3.16(3)	0.74(2)
Nissen RAD [4]	3.22(2)	–
Puzzarini [5]	2.998(98)	–

To compare our results with the ones reported earlier, the collisional broadening coefficient γ_V retrieved from the Voigt profile was considered. Fig. 7 shows that the current result is in a good agreement with the earlier measurements and has a better accuracy. The numerical values of the $\gamma_V(296)$ and its temperature dependence parameter n_{γ_V} for our and previous data can be compared using Table 1.

Details of *ab initio* calculations of the pressure broadening parameters for the R(0) line in the CO–N_2 collisional system are given in Paper I. The speed-dependence of broadening was not evaluated in Paper I because of the absence of related experimental data at the time of publication. The calculation of the γ_0 and γ_2 *ab initio* values and validation of accuracy of these theoretical data is one of the main goals of the current study. To run a "numerical experiment" in a proper way (see fig. 18 in Paper II and the related discussion for more details), $\gamma(v)$ dependence was retrieved using the results of approximating the *ab initio* pressure broadening cross-section $PBXS(E_{kin}, J)$ from Table I and Eqs. (5)–(6) of Paper I. Fig. 8 shows the calculated dependences of γ on the absorbing molecules speed v normalized to their most probable speed $\bar{v} = \sqrt{2k_BT/m}$ (where k_B is the Boltzmann constant and m is the mass of a CO molecule). With these dependences calculated for the number of gas temperatures, full speed-dependent *"ab initio"* profiles were calculated using Eq. (12) from Paper II, simulating the profiles observed in the experiment. These profiles were then analyzed similarly



Fig. 8. *Ab initio* speed dependences of the collisional broadening $\gamma(v/\bar{v})$ of the R(0) line of CO perturbed by N_2 at different temperatures. Dashed line shows Maxwell–Boltzmann distribution.



Fig. 9. Experimental and derived from *ab initio* calculations dependences of the $\gamma_{0,2}$ coefficients of qSDRP versus temperature.

to the experimental data giving theoretical (*"ab initio"*) values of $\gamma_{0,2}$ versus temperature. Fig. 9 shows the comparison of the experimental and theoretical T-dependences of the $\gamma_{0,2}$ parameters.

Measured and calculated values of $\gamma_{0,2}(T)$ are in good general agreement, but from the numerical data given in Table 2 one can find that the difference in $\gamma_0(296)$ values is about 6%, whereas for $\gamma_2(296)$ it is about 25%. The impact of the difference between measured and *ab initio* parameters in the line profile is shown in Fig. 10 together with the noise trace corresponding to the SNR~3000 which is typical for the RAD recordings. The difference between the temperature-dependent exponents is less but still exceeds the joint uncertainty interval. Note that the authors of Paper I "do not consider the pressure shift coefficient" because it is small and "the shape of the line is dominated by the collisional broadening" (see Introduction in Paper I). Thus, a similar comparison of our Δ_0 data with the corresponding *ab initio* values is left for the future studies.



Fig. 10. The impact of difference between measured and *ab initio* parameter values: only Γ_2 parameter and both Γ_0 and Γ_2, calculated for 296 K, 0.1 Torr of CO and 1.0 Torr of N_2. Noise trace corresponds to SNR~3000.

Table 2
Comparison of the measured and *ab initio* values of speed-independent and speed-dependent collisional broadening coefficients (in MHz/Torr) and their temperature-dependent parameters (unitless). Experimental uncertainties include fit uncertainty combined with the errors of pressure and temperature readings, and, for $\gamma_{0,2}$, uncertainties due to correlation with manually adjusted ν_{vc} (see Section 2.2 for details). Uncertainties of *ab initio* values include fit statistical uncertainty and variation due to usage of different interaction potentials from Paper I.

Parameter	Measured	*Ab initio*
$\gamma_V(296)$	3.190(7)	3.395(21)
n_V	0.775(4)	0.726(2)
$\gamma_0(296)$	3.237(7)	3.451(21)
n_0	0.781(4)	0.724(2)
$\gamma_2(296)$	0.296(7)	0.372(2)
n_2	0.971(80)	0.759(3)

The comparison of the currently known *ab initio* and measured line shape parameters [8,18–22] revealed no dependence on the "complexity" of the molecular collisional system. Even the calculations for the simplest systems like hydrogen isotopologue lines perturbed by He demonstrate impressive but still incomplete (with about the same difference as in our case) agreement for different series of transitions. In Paper I, Paper II and some of the above-mentioned studies it is demonstrated that different PES-s produce different values of line shape parameters. Thus, it may be conjectured that the clue to the successful quantum scattering calculations is a properly calculated molecular interaction potential, but it also cannot be ruled out that there are still some difficulties in the process of scattering calculations itself.

4. Conclusions

From the analysis of the obtained experimental recordings, a set of collisional parameters for modeling the absorption coefficient in a mixture of CO with N_2 at pressures ranging within about 10^{-4}–2 atm and temperatures from 238 to 353 K has been obtained, meeting high modern standards in terms of accuracy. Increased reliability of the obtained data is reached by using spectrometers with different principles of measurement and operating pressure ranges. The obtained experimental data on speed-dependent broadening parameters are used for validation of *ab initio* calculation results. The latter were found to be close to the measured ones but their systematic deviation exceeds the estimated uncertainty intervals demonstrating the need to improve the theory.

CRediT authorship contribution statement

D.S. Makarov: Writing – original draft, Visualization, Software, Methodology, Formal analysis. **I.N. Vilkov:** Investigation. **T.A. Galanina:** Supervision, Investigation, Data curation. **E.A. Serov:** Writing – review & editing, Validation, Data curation. **M.A. Koshelev:** Writing – review & editing, Validation, Methodology. **M.Yu. Tretyakov:** Writing – review & editing, Supervision, Funding acquisition.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

The authors express their gratitude to Artem Blokhin for help with data treatment.

The spectra recordings and their analysis were conducted under the RSF project No. 22-17-00041 (https://rscf.ru/en/project/22-17-00041/). The theoretical spectra were produced from *ab initio* data with the support of the Russian State Project FFUF-2024-0016. The experimental part of the work was performed using the research facilities "CKP-7" (USU No 3589084).

Data availability

Data will be made available on request.

References

[1] Jóźwiak H, Thibault F, Cybulski H, Wcisło P. *Ab initio* investigation of the CO–N_2 quantum scattering: The collisional perturbation of the pure rotational R(0) line in CO. J Chem Phys 2021;154(5):054314.

[2] Connor BJ, Radford HE. Pressure broadening of the CO J=1-0 rotational transition by N_2, O_2, and air. J Mol Spectrosc 1986;119(1):229–31.

[3] Colmont JM, Monnanteuil N. Self, nitrogen and oxygen broadening of the 115-GHz line of carbon monoxide. J Quant Spectrosc Radiat Transfer 1986;35(2):81–5.

[4] Nissen N, Doose J, Guarnieri A, Mäder H, Markov VN, Golubyatnikov GY, Leonov I, Shanin VN, Krupnov AF. Foreign gas broadening studies of the J' ← J=1 ← 0 rotational line of CO by frequency and time domain techniques. Z Naturforsch. A 1999;54(3–4):218–24.

[5] Puzzarini C, Dore L, Cazzoli G. A comparison of lineshape models in the analysis of modulated and natural rotational line profiles: Application to the pressure broadening of OCS and CO. J Mol Spectrosc 2002;216(2):428–36.

[6] Paredes-Roibás D, Martínez RZ, Jóźwiak H, Thibault F. Collisional line broadening and mixing in the Raman spectrum of CO perturbed by N_2: Experimental measurements and theoretical calculations. J Quant Spectrosc Radiat Transfer 2021;275:107868.

[7] Serov EA, Stolarczyk N, Makarov DS, Vilkov IN, Golubiatnikov GY, Balashov AA, Koshelev MA, Wcisło P, Thibault F, Tretyakov MY. CO-Ar collisions: *ab initio* model matches experimental spectra at a sub percent level over a wide pressure range. J Quant Spectrosc Radiat Transfer 2021;272:107807.

[8] Tretyakov MY, Serov EA, Makarov DS, Vilkov IN, Golubiatnikov GY, Galanina TA, Koshelev MA, Balashov AA, Simonova AA, Thibault F. Pure rotational R(0) and R(1) lines of CO in Ar baths: experimental broadening, shifting and mixing parameters in a wide pressure range versus *ab initio* calculations. Phys Chem Chem Phys 2023;25:1310–30.

[9] Tretyakov MY, Koshelev MA, Makarov DS, Tonkov MV. Precise measurements of collision parameters of spectral lines with a spectrometer with radioacoustic detection of absorption in the millimeter and submillimeter ranges. Instrum. Exp Tech 2008;51(1):78–88.

[10] Koshelev MA, Leonov II, Serov EA, Chernova AI, Balashov AA, Bubnov GM, Andriyanov AF, Shkaev AP, Parshin VV, Krupnov AF, Tretyakov MY. New frontiers in modern resonator spectroscopy. IEEE Trans Terahertz Sci Technol 2018;8:773–83.

[11] Gordon IE, Rothman LS, Hargreaves RJ, Hashemi R, Karlovets EV, Skinner FM, Conway EK, Hill C, Kochanov RV, Tan Y, Wcisło P, Finenko AA, Nelson K, Bernath PF, Birk M, Boudon V, Campargue A, Chance KV, Coustenis A, Drouin BJ, Flaud JM, Gamache RR, Hodges JT, Jacquemart D, Mlawer EJ, Nikitin AV, Perevalov VI, Rotger M, Tennyson J, Toon GC, Tran H, Tyuterev VG, Adkins EM, Baker A, Barbe A, Canè E, Császár AG, Dudaryonok A, Egorov O, Fleisher AJ, Fleurbaey H, Foltynowicz A, Furtenbacher T, Harrison JJ, Hartmann J-M, Horneman V-M, Huang X, Karman T, Karns J, Kassi S, Kleiner I, Kofman V, Kwabia-Tchana F, Lavrentieva N, Lee T, Long D, Lukashevskaya AA, Lyulin OM, Makhnev VY, Matt W, Massie ST, Melosso M, Mikhailenko SN, Mondelain D, Müller HSP, Naumenko OV, Perrin A, Polyansky OL, Raddaoui E, Raston PL, Reed ZD, Rey M, Richard C, Tóbiás R, Sadiek I, Schwenke DW, Starikova E, Sung K, Tamassia F, Tashkun SA, Vander Auwera J, Vasilenko IA, Vigasin AA, Villanueva GL, Vispoel B, Wagner G, Yachmenev A, Yurchenko SN. The HITRAN2020 molecular spectroscopic database. J Quant Spectrosc Radiat Transfer 2022;277:107949.

[12] Koshelev MA, Markov VN. Broadening of the J=3←2 spectral line of carbon monoxide by pressure of CO, N_2 and O_2. J Quant Spectrosc Radiat Transfer 2009;110(8):526–7.

[13] Predoi-Cross A, Esteki K, Rozario H, Naseri H, Latif S, Thibault F, Malathy Devi V, Smith MAH, Mantz AW. Theoretical and revisited experimentally retrieved He-broadened line parameters of carbon monoxide in the fundamental band. J Quant Spectrosc Radiat Transfer 2016;184:322–40.

[14] Ngo NH, Lisak D, Tran H, Hartmann JM. An isolated line-shape model to go beyond the voigt profile in spectroscopic databases and radiative transfer codes. J Quant Spectrosc Radiat Transfer 2013;129:89–100.

[15] Rosenkranz PW. Shape of the 5 mm oxygen band in the atmosphere. IEEE Trans Antennas and Propagation 1975;23(4):498–506.

[16] Kochanov RV, Gordon IE, Rothman LS, Wcisło P, Hill C, Wilzewski JS. HITRAN application programming interface (HAPI): A comprehensive approach to working with spectroscopic data. J Quant Spectrosc Radiat Transfer 2016;177:15–30.

[17] Koshelev MA, Vilkov IN, Makarov DS, Tretyakov MY, Vispoel B, Gamache RR, Cimini D, Romano F, Rosenkranz PW. Water vapor line profile at 183-GHz: Temperature dependence of broadening, shifting, and speed-dependent shape parameters. J Quant Spectrosc Radiat Transfer 2021;262:107472.

[18] Wcisło P, Thibault F, Zaborowski M, Wójtewicz S, Cygan A, Kowzan G, Masłowski P, Komasa J, Puchalski M, Pachucki K, Ciuryło R, Lisak D. Accurate deuterium spectroscopy for fundamental studies. J Quant Spectrosc Radiat Transfer 2018;213:41.

[19] Martínez RZ, Bermejo D, Thibault F, Wcisło P. Testing the *ab initio* quantum-scattering calculations for the D_2—He benchmark system with stimulated Raman spectroscopy. J Raman Spectrosc 2018;49(8):1339–49.

[20] Thibault F, Martínez RZ, Bermejo D, Wcisło P. Line-shape parameters for the first rotational lines of HD in He. Mol Astrophys 2020;19:100063.

[21] Gancewski M, Jóźwiak H, Quintas-Sánchez E, Dawes R, Thibault F, Wcisło P. Fully quantum calculations of O_2—N_2 scattering using a new potential energy surface: Collisional perturbations of the oxygen 118 GHz fine structure line. J Chem Phys 2021;155(12):124307.

[22] Słowiński M, Jóźwiak H, Gancewski M, Stankiewicz K, Stolarczyk N, Tan Y, Wang J, Liu A-W, Hu S-M, Kassi S, Campargue A, Patkowski K, Żuchowski PS, Ciuryło R, Thibault F, Wcisło P. Collisional line-shape effects in accurate He-perturbed H_2 spectra. J Quant Spectrosc Radiat Transfer 2022;277:107951.