

012 - Central Limit Theorem

EPIB 607

Sahir Rai Bhatnagar
Department of Epidemiology, Biostatistics, and Occupational Health
McGill University

`sahir.bhatnagar@mcgill.ca`

slides compiled on September 26, 2021



Standard deviation and variance of a random variable Y

- $Y \sim \text{unknown_distribution}(\mu, \sigma)$
- Standard Deviation σ , and Variance σ^2 , of a random variable Y with mean μ .

$$\text{Var}[Y] = \sigma^2 = \text{mean of } (Y - \mu)^2$$

$$\text{SD}[Y] = \sigma$$

$$\text{Var}[Y \pm a \text{ constant}] = \text{Var}[Y]$$

$$\text{SD}[Y \pm a \text{ constant}] = \sigma$$

$$\text{Var}[Y \times a \text{ constant}] = \text{constant}^2 \times \text{Var}[Y]$$

$$\text{SD}[Y \times a \text{ constant}] = |\text{constant}| \times \sigma$$

Rules for Variances and SDs of sums and means of n independent random variables

Sums

$$\text{Var}[Y_1 + Y_2 + \cdots + Y_n] = \sigma^2 + \sigma^2 + \cdots + \sigma^2 = n \times \sigma^2$$

$$\text{SD}[Y_1 + Y_2 + \cdots + Y_n] = \sqrt{n} \times \sigma$$

Rules for Variances and SDs of sums and means of n independent random variables

Sums

$$\text{Var}[Y_1 + Y_2 + \cdots + Y_n] = \sigma^2 + \sigma^2 + \cdots + \sigma^2 = n \times \sigma^2$$


$$\text{SD}[Y_1 + Y_2 + \cdots + Y_n] = \sqrt{n} \times \sigma$$

Means

$$\text{Var}\left[\frac{Y_1 + Y_2 + \cdots + Y_n}{n}\right] = \frac{1}{n} \times \sigma^2$$

$$\text{SD}\left[\frac{Y_1 + Y_2 + \cdots + Y_n}{n}\right] = \sqrt{\frac{1}{n}} \times \sigma$$


Age-distribution of the entire population of Dublin¹



¹<http://www.census.nationalarchives.ie/help/about19011911census.html>


Distribution of 10000 Bootstrap samples of size 2

```
## [1] Ages of sampled persons in first 2 samples of size 2
##      [,1] [,2]
## [1,]  24  11
## [2,]  65   4
```




Distribution of 10000 Bootstrap samples of size 4

```
## [1] Ages of sampled persons in first 2 samples of size 4
##      [,1] [,2] [,3] [,4]
## [1,]   60   53   14   50
## [2,]   58   23    2   35
```




Distribution of 10000 Bootstrap samples of size 10


```
## [1] Ages of sampled persons in first 2 samples of size 10
##      [,1] [,2] [,3] [,4] [,5] [,6] [,7] [,8] [,9] [,10]
## [1,]   42  11  16  18   3  48  28   1   7   5
## [2,]   23  20  20  29  38  15  82  68  18  16
```



Distribution of 10000 Bootstrap samples of size 16



Distribution of 10000 Bootstrap samples of size 64



Properties of the sample mean: The Central Limit Theorem (CLT)

The sampling distribution of \bar{Y} is Normal if Y is Normal. What probability distribution does the sample mean follow if Y is not Normal?

Properties of the sample mean: The Central Limit Theorem (CLT)

The sampling distribution of \bar{Y} is Normal if Y is Normal. What probability distribution does the sample mean follow if Y is not Normal?

As sample size increases, the distribution of \bar{Y} becomes closer to a Normal distribution, no matter what the distribution of sampled variable Y !

(This is true as long as the distribution has a finite variance.)

The Central Limit Theorem (CLT)

- The sampling distribution of \bar{y} is, for a large enough n , close to Gaussian in shape no matter what the shape of the distribution of individual Y values.
- This phenomenon is referred to as the CENTRAL LIMIT THEOREM
- The CLT applied also to a sample proportion, slope, correlation, or any other statistic created by aggregation of individual observations

Theorem 1 (Central Limit Theorem).

if $Y \sim ???(\mu_Y, \sigma_Y)$, then

$$\bar{y} \sim \mathcal{N}(\mu_Y, \sigma_Y/\sqrt{n})$$

Standard error (SE) of a sample statistic

- Recall: When we are talking about the variability of a **statistic**, we use the term **standard error** (not standard deviation). The standard error of the sample mean is σ/\sqrt{n} .

Standard error (SE) of a sample statistic

- Recall: When we are talking about the variability of a **statistic**, we use the term **standard error** (not standard deviation). The standard error of the sample mean is σ/\sqrt{n} .





Remark 1 (SE vs. SD).

In quantifying the instability of the sample mean (\bar{y}) statistic, we talk of SE of the mean (SEM)





$SE(\bar{y})$ describes how far \bar{y} could (typically) deviate from μ ;

$SD(y)$ describes how far an individual y (typically) deviates from μ (or from \bar{y}).





CLT in action: Binomial($n = 5, p = 0.8$) distribution




CLT in action: Uniform($a = 0$, $b = 1$) distribution




CLT in action: Exponential($\lambda = 1$) distribution




CLT in action: Depths of the ocean




CLT in action: Depths of the ocean




CLT in action: Depths of the ocean




CLT in action: Depths of the ocean




CLT in action: Depths of the ocean




CLT in action: Depths of the ocean




CLT in action: Depths of the ocean




CLT in action: Depths of the ocean




CLT in action: Depths of the ocean




CLT in action: Depths of the ocean



CLT in action: Depths of the ocean




CLT in action: Depths of the ocean



How long does it take for the CLT to 'kick in'?

- How *fast* or slowly the CLT will **kick in** is a function of how symmetric, or how asymmetric and **CLT-unfriendly**, the distribution of Y (the depths of the ocean) is



Quadruple the work, half the benefit




Figure: When the sample size increases from 4 to 16, the spread of the sampling distribution for the mean is reduced by a half, i.e., the range is cut in half. This is known as the curse of the \sqrt{n}

Session Info

```
R version 4.0.4 (2021-02-15)
Platform: x86_64-pc-linux-gnu (64-bit)
Running under: Pop!_OS 21.04
```

```
Matrix products: default
BLAS:   /usr/lib/x86_64-linux-gnu/openblas-pthread/libblas.so.3
LAPACK: /usr/lib/x86_64-linux-gnu/openblas-pthread/libopenblaspr0.3.13.so
```

attached base packages:

```
[1] tools      stats      graphics  grDevices  utils      datasets  methods
[8] base
```

other attached packages:

```
[1] DT_0.16 mosaic_1.7.0 Matrix_1.3-2 mosaicData_0.20.1
[5] ggformula_0.9.4 ggstance_0.3.4 lattice_0.20-41 kableExtra_1.2.1
[9] socviz_1.2 gapminder_0.3.0 here_0.1 NCStats_0.4.7
[13] FSA_0.8.30 forcats_0.5.1 stringr_1.4.0 dplyr_1.0.7
[17] purrr_0.3.4 readr_1.4.0 tidyr_1.1.3 tibble_3.1.3
[21] ggplot2_3.3.5 tidyverse_1.3.0 knitr_1.33
```

loaded via a namespace (and not attached):

```
[1] fs_1.5.0 lubridate_1.7.9 webshot_0.5.2 httr_1.4.2
[5] rprojroot_2.0.2 backports_1.2.1 utf8_1.2.2 R6_2.5.1
[9] DBI_1.1.1 colorspace_2.0-2 withr_2.4.2 tidyrselect_1.1.1
[13] gridExtra_2.3 leaflet_2.0.3 curl_4.3.2 compiler_4.0.4
[17] cli_3.0.1 rvest_1.0.0 pacman_0.5.1 xml2_1.3.2
[21] ggdendro_0.1.22 mosaicCore_0.8.0 scales_1.1.1 digest_0.6.27
[25] foreign_0.8-81 rmarkdown_2.9.7 rio_0.5.16 pkgconfig_2.0.3
[29] htmltools_0.5.2 highr_0.9 dbplyr_1.4.4 fastmap_1.1.0
[33] htmlwidgets_1.5.3 rlang_0.4.11 readxl_1.3.1 rstudioapi_0.13
[37] farver_2.1.0 generics_0.1.0 jsonlite_1.7.2 crosstalk_1.1.1
[41] zip_2.2.0 car_3.0-9 magrittr_2.0.1 Rcpp_1.0.7
[45] munsell_0.5.0 fansi_0.5.0 abind_1.4-5 lifecycle_1.0.0
[49] stringi_1.7.3 carData_3.0-4 MASS_7.3-53.1 plyr_1.8.6
[53] grid_4.0.4 blob_1.2.1 ggrepel_0.8.2 crayon_1.4.1
[57] cowplot_1.1.0 haven_2.3.1 splines_4.0.4 hms_1.0.0
[61] pillar_1.6.2 reprex_0.3.0 glue_1.4.2 evaluate_0.14
[65] data.table_1.14.0 modelr_0.1.8 vctrs_0.3.8 tweenr_1.0.1
[69] cellranger_1.1.0 gtable_0.3.0 polyclip_1.10-0 assertthat_0.2.1
[73] TeachingDemos_2.12 xfun_0.25 ggforce_0.3.2 openxlsx_4.1.5
[77] RColorMan_2.0.7.2 viridisLite_0.4.0 ellipsis_0.3.2
```