

SAIMEDHA



HYDERABAD-VIJYAWADA-TIRUPATI

ECET-IES-GATE-PSUs

Max Marks:25 **ECET-2024**

Time : 1 hrs

Topic: Matrices

Date:03-01-2024

- A square matrix (a_{ij}) in which $a_{ij} = 0$ for $i \neq j$ and $a_{ij} = k$ (constant) for $i = j$ is
a) Unit matrix b) Scalar matrix c) Null matrix d) Diagonal matrix
- $A_{n \times n}$ and $B_{n \times n}$ are diagonal matrices then $AB = \dots\dots\dots$ matrix
a) square b) diagonal c) scalar d) rectangular
- If $A = [a_{ij}]_{3 \times 3}$ is a square matrix so that $a_{ij} = i^2 - j^2$ then A is a
a) unit matrix b) symmetric matrix c) skew symmetric matrix d) orthogonal matrix
- If $\begin{bmatrix} x+3 & 2y+x \\ z-1 & 4a-z \end{bmatrix} = \begin{bmatrix} 0 & -7 \\ 3 & 2a \end{bmatrix}$, $(x+y+z+a) =$ a)-1 b)0 c)1 d)8
- If $A = \begin{bmatrix} 6 & 10 & 100 \\ 7 & 1 & 0 \\ 0 & 9 & 10 \end{bmatrix}$ then $\text{Tr}(A^T) =$ a)-17 b)17 c)-1/17 d)1/17
- If $A = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}$, then $A^4 = \dots\dots\dots$ a)16A b)32I c)4A d)8A
- If $A^2 = A$, $B^2 = B$, $AB = BA = 0$ then $(A+B)^2 =$
a)A-B b)A+B c) $A^2 - B^2$ d)0
- If $A = \begin{bmatrix} 2 & x-3 & x-2 \\ 3 & -2 & -1 \\ 4 & -1 & -5 \end{bmatrix}$ is a symmetric matrix then $x =$
a)0 b)3 c)6 d)8
- If $A = \begin{bmatrix} a & b & c \\ b & c & a \\ c & a & b \end{bmatrix}$ then cofactor of a_{21} is a) $b^2 - ac$ b) $ac - b^2$ c) $a^2 - bcd$ d) $bc - a^2$
- $\begin{vmatrix} x+y+2z & x & y \\ z & y+z+2x & y \\ z & x & z+x+2y \end{vmatrix} =$
a) $(x+y+z)^3$ b) $2(x+y+z)^3$ c) $x+y+z$ d) $(x+y+z)^2$
- If a,b,c are in A.P., then $\begin{vmatrix} x+1 & x+2 & x+a \\ x+2 & x+3 & x+b \\ x+3 & x+4 & x+c \end{vmatrix}$
a) $\frac{a+b}{2}$ b)ab c)0 d)abc
- $\begin{vmatrix} a^2+2a & 2a+1 & 1 \\ 2a+1 & a+2 & 1 \\ 3 & 3 & 1 \end{vmatrix} =$ a) $(1-a)^3$ b) $(a-1)^2c(a-1)^3$ d) $(a+1)^2$
- If A is $\begin{bmatrix} 8 & -6 & 2 \\ -6 & 7 & -4 \\ 2 & -4 & \lambda \end{bmatrix}$ is a singular matrix then $\lambda =$ a)3 b)4 c)2 d)5
- If $A = \begin{bmatrix} 0 & 1 & 2 \\ 1 & 2 & 3 \\ 3 & 1 & 1 \end{bmatrix}$ then $\text{Adj}(A) =$
a) $\begin{bmatrix} -1 & 8 & -5 \\ 1 & -6 & 3 \\ -1 & 2 & -1 \end{bmatrix}$ b) $\begin{bmatrix} -1 & 1 & -1 \\ 8 & -6 & 2 \\ -5 & 3 & -1 \end{bmatrix}$ c) $\begin{bmatrix} 1 & -1 & 1 \\ 8 & -6 & 2 \\ -5 & 3 & -1 \end{bmatrix}$ d) $\begin{bmatrix} -1 & -8 & 5 \\ -1 & 6 & -3 \\ 1 & -2 & 1 \end{bmatrix}$

15. If $A = \begin{bmatrix} 2 & 2 \\ -3 & 2 \end{bmatrix}$, $B = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$, then $(B^{-1} A^{-1})^{-1} =$

a) $\begin{bmatrix} 2 & -2 \\ 2 & 3 \end{bmatrix}$ b) $\begin{bmatrix} 3 & -2 \\ 2 & 2 \end{bmatrix}$ c) $\frac{1}{10} \begin{bmatrix} 2 & 2 \\ -2 & 3 \end{bmatrix}$ d) $\frac{1}{10} \begin{bmatrix} 3 & 2 \\ -2 & 2 \end{bmatrix}$

16. The number of solutions of the equations $3x-2y=5$, $6x-4y=10$ are ...

a) 0 b) 1 c) 10 d) infinity

17. $(A+AB)^T = XA^T \Rightarrow x =$ a) $B^T b) I + Bc) I + B^T d) B^T \cdot A^T$

18. If $A = \begin{pmatrix} 0 & 0 \\ 1 & 1 \end{pmatrix}$ then the value of $A + A^2 + A^3 + \dots + A^n =$ a) $Ab) nA$ c) $(n+1)A$ d) 0

19. With usual notations in ΔABC $\frac{\sin A}{\sin^2 A} \frac{\sin B}{\sin^2 B} \frac{\sin C}{\sin^2 C} =$

a) $\frac{1}{8R^3} (a-b) (b-c) (c-a)$ b) $8R^3$ c) $(a-b) (b-c) (c-a)$ d) $\frac{1}{8R} (a-b) (b-c) (c-a)$

20. The matrix $\begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}$ has

a) one inverse b) two inverse c) no inverse d) cannot be said

21. If $A = \begin{bmatrix} 9 & 0 & 0 \\ 0 & 10 & 0 \\ 0 & 0 & 8 \end{bmatrix}$ then A^{-1}

a) $\begin{bmatrix} \frac{1}{9} & 0 & 0 \\ 0 & 10 & 0 \\ 0 & 0 & \frac{1}{8} \end{bmatrix}$ b) $\begin{bmatrix} 9 & 0 & 0 \\ 0 & \frac{1}{10} & 0 \\ 0 & 0 & 8 \end{bmatrix}$ c) $\begin{bmatrix} 9 & 0 & 0 \\ 0 & 10 & 0 \\ 0 & 0 & \frac{1}{8} \end{bmatrix}$ d) $\begin{bmatrix} \frac{1}{9} & 0 & 0 \\ 0 & \frac{1}{10} & 0 \\ 0 & 0 & \frac{1}{8} \end{bmatrix}$

22. The inverse of $\begin{bmatrix} 2 & 1 & 2 \\ 1 & 0 & 1 \\ 2 & 2 & 1 \end{bmatrix}$ is

a) $\begin{bmatrix} 4 & -3 & -2 \\ -3 & -2 & 2 \\ -2 & 2 & 1 \end{bmatrix}$ b) $\begin{bmatrix} -2 & 3 & 1 \\ -1 & -2 & 0 \\ 2 & -2 & -1 \end{bmatrix}$ c) $\begin{bmatrix} -2 & 3 & 1 \\ 1 & -2 & 0 \\ 2 & -2 & -1 \end{bmatrix}$ d) $\begin{bmatrix} 2 & 3 & 1 \\ 1 & 2 & 0 \\ 2 & 2 & 1 \end{bmatrix}$

23. If $A = \begin{bmatrix} -1 & -2 & -2 \\ 2 & 1 & -2 \\ 2 & -2 & 1 \end{bmatrix}$ then $\text{Adj } A =$ a) A^T b) $2A^T$ c) $3A^T$ d) $4A^T$

24. $\begin{vmatrix} z+y & x & x \\ y & z+x & y \\ z & z & x+y \end{vmatrix} =$ a) xyz b) $4xyz$ c) $2xyz$ d) $3xyz$

25. $L = \begin{bmatrix} 2 & 3 & 5 \\ 4 & 1 & 2 \\ 1 & 2 & 1 \end{bmatrix} = P + Q$, P is a symmetric matrix, Q is a skew-symmetric matrix then

$P =$ a) $\begin{bmatrix} 2 & 3 & 5 \\ 4 & 1 & 2 \\ 1 & 2 & 1 \end{bmatrix}$ b) $\begin{bmatrix} 2 & -7 & 6 \\ -7 & 1 & 4 \\ 6 & 4 & 1 \end{bmatrix}$ c) $\begin{bmatrix} 2 & 7/2 & 3 \\ 7/2 & 1 & 2 \\ 3 & 2 & 1 \end{bmatrix}$ d) $\begin{bmatrix} 4 & 7 & 6 \\ 7 & 2 & 4 \\ 6 & 4 & 2 \end{bmatrix}$

Key: Matrices

Date: 28-12-23

1-2	2-2	3-3	4-3	5-2	6-4	7-2	8-3	9-2	10-2
11-3	12-3	13-1	14-2	15-1	16-4	17-3	18-2	19-1	20-3
21-4	22-3	23-3	24-2	25-3					