Pursuit, Rendezvous, and Evasion With Constant-Power, Variable-I_{sp} Rockets

Jan Olšina

Independent Researcher, Prague, Czech Republic
email: jan.olsina@gmail.com, ORCID: 0009-0003-9816-3958.

***Abstract*—We study pursuit and evasion by constant-useful-power rockets with continuously variable exhaust speed. Position-only interception is optimized for minimum guaranteed time or minimum worst-case propellant expenditure by a deadline; exact position-and-velocity rendezvous is optimized for minimum guaranteed time. Both capture modes are treated in free space and in a Kepler field. An inverse-mass transformation converts propellant constraints into integral bounds on squared acceleration. We derive the Hamilton–Jacobi–Isaacs equations, feedback selectors, characteristic ordinary differential equations, and global verification theorems. For ideal free-space rendezvous with instantaneous observation of the evader's acceleration, we obtain a complete analytical solution: a square-root resource gap determines a unique cubic time equation, and explicit opposing strategies prove its optimality for arbitrary initial relative position and velocity. When the evader has at least the pursuer's remaining acceleration budget, a constructive survival strategy also attains the least worst-case escape expenditure. A planar Clohessy–Wiltshire rendezvous theorem gives the local Kepler counterpart, with an explicit opposing strategy and a sufficient nonlinear capture tube. These exact guarantees assume that the pursuer observes the evader's current acceleration. Further results include an exact active-interception family, closed-form ballistic benchmarks, mass and exhaust-speed histories, and a rigorous gravity-compensated Kepler capture bound. General orbital values remain nonlinear feedback problems. Strict position-only escape can have an unattained fuel infimum, so attained optima and limiting values are distinguished throughout.**

I. Introduction

The companion article [1] studies a single rocket transferring between specified endpoint states. Its constant-power propulsion law makes acceleration a particularly convenient control variable: propellant expenditure is quadratic in acceleration after transforming mass. In a chase, however, the destination is itself controlled by an opponent. A trajectory optimized against one predicted evader trajectory does not establish interception against a different response.

The appropriate object is therefore a *causal feedback strategy*. A strategy specifies the next control from the observed game state and history; a trajectory is an outcome of two interacting strategies. Dynamic programming and Hamilton–Jacobi–Isaacs (HJI) theory provide the corresponding value equations [2]. Target-hitting and survival games also require careful treatment of the target boundary and possible infinite capture times [3].

Two capture modes are distinguished: interception requires position proximity, whereas rendezvous requires coincident position and velocity. The objectives are:

- A1) free-space interception or exact rendezvous in minimum guaranteed time;
- A2) free-space interception by time T with minimum worst-case pursuer fuel;
- B1) minimum-time interception or exact rendezvous in a Kepler field;
- B2) the deadline-constrained minimum-fuel game in that field.

When the evader can force the relevant survival objective, its secondary objective is to minimize its own worst-case fuel use. For A2 and B2, "win" means surviving through the deadline; for A1 and B1 it means avoiding capture forever. These are different objectives.

Initial positions, velocities, engine parameters, and deadlines are retained as parameters. Free-space moment identities permit explicit solutions in several regimes, including the complete ideal minimum-time rendezvous game under the specified response convention. The Kepler problem retains nonlinear dependence on both orbital states. Strict escape introduces a separate existence question: some admissible instances have no exactly optimal escaping path.

II. Rocket Model and Rules of the Game

A. From Propellant Mass to an Acceleration Budget

Use subscripts P and E for pursuer and evader. Let $P_i > 0$ be useful jet power, m_i instantaneous mass, m_{i0} initial mass, m_{id} dry mass, and $\mathbf{a}_i \in \mathbb{R}^d$ thrust acceleration. We consider $d = 2$ or 3, and occasionally a one-dimensional benchmark. The companion propulsion model is

$$\dot{m}_i = -\frac{m_i^2 \left\|\mathbf{a}_i\right\|^2}{2P_i}, \qquad v_{ei} = \frac{2P_i}{m_i \left\|\mathbf{a}_i\right\|}. \tag{1}$$

The exhaust-speed formula applies on powered arcs. At zero acceleration the engine is off, and exhaust speed need not be assigned. Specific impulse is $I_{sp,i} = v_{ei}/g_0$. Thrust, acceleration, and specific impulse are linked by (1); they cannot be chosen independently at fixed useful power.

Define the initial and remaining acceleration budgets by

$$B_i = 2P_i \left(\frac{1}{m_{id}} - \frac{1}{m_{i0}}\right), \tag{2}$$

$$b_i(t) = 2P_i \left(\frac{1}{m_{id}} - \frac{1}{m_i(t)}\right). \tag{3}$$

Direct differentiation gives the exact reduction

$$\boxed{\dot{b}_i = -\left\|\mathbf{a}_i\right\|^2, \quad b_i(0) = B_i, \quad b_i(t) \geq 0.} \tag{4}$$

Thus admissible ideal controls are measurable, locally square-integrable accelerations satisfying

$$\int_0^t \left\|\mathbf{a}_i(s)\right\|^2 \mathrm{d}s \leq B_i \quad \text{for every pre-termination } t. \tag{5}$$

At $b_i = 0$, only $\mathbf{a}_i = 0$ is admissible. Propellant exhaustion does not remove the spacecraft: it subsequently follows a ballistic trajectory.

The units of b_i are $\mathrm{m}^2\,\mathrm{s}^{-3}$, not joules. We use the term *acceleration budget*, although this is often called an energy-type integral constraint in differential games. Actual mass and propellant expenditure are recovered exactly:

$$m_i(b_i) = \left(\frac{1}{m_{id}} - \frac{b_i}{2P_i}\right)^{-1}, \tag{6}$$

$$\Phi_i(c) = m_{i0} - \left(\frac{1}{m_{i0}} + \frac{c}{2P_i}\right)^{-1}. \tag{7}$$

Here $c = \int \left\|\mathbf{a}_i\right\|^2 \mathrm{d}t$. Since Φ_i is continuous and strictly increasing, minimizing worst-case c is exactly equivalent to minimizing worst-case propellant mass. Powers and masses enter the *ideal uncapped kinematic game* only through B_P, B_E; they are still needed to reconstruct mass and specific impulse.

B. Ideal and Bounded Engines

Unless stated otherwise, the primary model is the ideal uncapped model of the companion article: $\mathbf{a}_i \in \mathbb{R}^d$ while $b_i > 0$. It permits arbitrarily small positive exhaust speed and arbitrarily large acceleration, subject to (5); it also permits unbounded exhaust speed near a vanishing powered acceleration. Impulsive velocity jumps are not included, because an L^2 acceleration has vanishing impulse on intervals whose length tends to zero.

For finite engineering limits, let

$$0 < v_{ei,\min} \leq v_{ei} \leq v_{ei,\max} < \infty. \tag{8}$$

At full power the admissible acceleration set is

$$\mathcal{U}_i(b_i) = \{\mathbf{0}\} \cup \{\mathbf{a} : a_{i,\min}(b_i) \leq \|\mathbf{a}\| \leq a_{i,\max}(b_i)\}, \tag{9}$$

$$a_{i,\min}(b) = \frac{2P_i}{m_i(b)v_{ei,\max}}, \qquad a_{i,\max}(b) = \frac{2P_i}{m_i(b)v_{ei,\min}}.$$

At $b = 0$ this is replaced by $\{\mathbf{0}\}$. A ceiling on exhaust speed alone gives a *minimum nonzero acceleration*, not a maximum acceleration. An artificial bound $\|\mathbf{a}_i\| \leq A_i$ is another useful regularization, but is a different model until a justified limit is taken. Section IX gives selectors for these sets.

C. Conventions Shared With the Transfer Article

For the same physical minimum-time objective, a smooth cost-to-go V uses minimizing dynamic programming here. The companion uses the maximizing Hamiltonian $\mathcal{H}_T = \lambda \cdot f - 1$. The sign and coordinate changes are:

Quantity	Correspondence
Kinematic covectors	$\lambda_x = -V_x$, $\lambda_v = -V_v$
Budget and units	$b = 2P(m_d^{-1} - m^{-1})$, $\mathrm{m}^2/\mathrm{s}^3$
Resource covector	$\lambda_b = -V_b$
Mass covector	$\lambda_m = (2P/m^2)\lambda_b = -(2P/m^2)V_b$
Interior gain	$C_m = \lambda_m m^2 = -2PV_b$, $k_T = P/C_m$
Normalized budget	$\bar{b} = b/J_0$, $J_0 = R_0^2/T_0^3$
Exhaust parameters	$\nu_\pm = v_{e,\pm}/(R_0/T_0)$

The covector formulas compare the same objective in the same units; changing objective units also rescales its covectors. For a differentiable game value the sign reversal exchanges min and max in the entire Isaacs operator. It does not convert the evader into a second independent minimum-time optimizer. The ideal interior constancy of C_m fails on mass-dependent exhaust boundaries, as stated in Section IX and Appendix A of the companion article [1].

D. Motion and Capture

In an inertial frame,

$$\dot{\mathbf{x}}_i = \mathbf{v}_i, \tag{10a}$$

$$\dot{\mathbf{v}}_i = \mathbf{g}(\mathbf{x}_i) + \mathbf{a}_i, \tag{10b}$$

$$\dot{b}_i = -\left\|\mathbf{a}_i\right\|^2, \tag{10c}$$

where

$$\mathbf{g}(\mathbf{x}) = \begin{cases} \mathbf{0}, & \text{A: free space,} \\ -\mu\mathbf{x}/\left\|\mathbf{x}\right\|^3, & \text{B: Kepler field.} \end{cases} \tag{11}$$

Both spacecraft feel the same central body, but generally experience different gravitational accelerations.

Take a prescribed capture radius $R \geq 0$ and define

$$\mathcal{C} = \{z : \|\mathbf{x}_E - \mathbf{x}_P\| \leq R\}, \tag{12}$$

$$\tau = \inf\{t \geq 0 : z(t) \in \mathcal{C}\}, \qquad \inf \varnothing = +\infty. \tag{13}$$

The position-only game stops at first entry into $\mathcal{C}$. A positive R describes interception within a specified range; $R = 0$ describes exact point interception. Its terminal velocity is unrestricted. Exact rendezvous instead uses

$$\mathcal{C}_J = \{z : \mathbf{x}_E = \mathbf{x}_P,\ \mathbf{v}_E = \mathbf{v}_P\}, \tag{14}$$

$$\tau_J = \inf\{t \geq 0 : z(t) \in \mathcal{C}_J\}. \tag{15}$$

The simultaneous equalities are essential: crossing the same point at different velocities does not terminate a rendezvous game. The superscript J denotes its minimum-time value. Rendezvous is treated as a terminal event; attachment and subsequent holding dynamics are not part of the kinematic model. Since $\mathcal{C}_J \subseteq \mathcal{C}$, minimum-time values satisfy $V^J \geq V$ for the same initial state and information convention. Deadline-fuel objectives use position-only capture.

The Kepler field is singular at the central body. Results below apply to noncolliding admissible trajectories. A global numerical problem must prescribe a collision rule or a minimum central distance and the associated individual viability constraints. One consistent choice is to restrict each pilot to trajectories with $\|\mathbf{x}_i\| \geq r_{\min} > 0$, on the set of states from which that individual restriction is viable. Such constraints require their own state-constraint boundary treatment; a collision

is not silently counted as successful evasion. No particular value of $r_{\min}$ is imposed here. Local equations are valid throughout the nonsingular domain, and global certificates must cover the entire region actually used.

E. Information and Objectives

Both pilots know the model, clock, positions, velocities, and remaining budgets exactly. They may change controls at every instant and use the observation history, but cannot anticipate future choices. A pursuer strategy α maps an admissible evader input history to a pursuer input history nonanticipatively; β is the corresponding evader strategy. Admissibility includes the strategy's own budget against *every* opposing input. General value constructions may use an explicitly specified nonanticipative or sampled-feedback formulation; the exact response theorems below specifically use instantaneous acceleration observation and cannot simply be transferred to a delayed sampled implementation [2].

Notation such as $\sup_{\mathbf{a}_E}$ below means the supremum over all admissible opposing input functions, including functions generated by causal state feedback. It does not mean a preannounced maneuver. Two unrestricted nonanticipative response maps should not simply be composed if doing so would create an ill-defined algebraic loop; the verification results use well-posed feedback dynamics.

For minimum time, the pursuer's robust value is

$$V(z) = \inf_{\alpha} \sup_{\mathbf{a}_E} \tau[z; \alpha[\mathbf{a}_E], \mathbf{a}_E]. \tag{16}$$

The dual evader value is $\sup_{\beta} \inf_{\mathbf{a}_P} \tau$. Replacing τ by τ_J defines V^J and its dual. A common value is used only where game determinacy or a verification certificate establishes equality. The explicit free-space strategy results use instantaneous observation of $\mathbf{a}_E$ by the pursuer, and state-and-resource feedback by the evader. This information pattern is used in integral-constraint pursuit games [4]; it permits the response $\mathbf{a}_P = \mathbf{a}_E + \mathbf{a}_0$ without anticipating future input. It does not assert the same exact guarantees for delayed sensing.

For a remaining horizon s, define the extended payoff

$$J_s = \begin{cases} \displaystyle\int_0^{\tau} \|\mathbf{a}_P(t)\|^2 \, \mathrm{d}t, & \tau \le s, \\ +\infty, & \tau > s, \end{cases} \tag{17}$$

and the robust fuel value

$$W(s, z) = \inf_{\alpha} \sup_{\mathbf{a}_E} J_s. \tag{18}$$

The dimensional propellant answer at the initial state is $\Phi_P(W(T, z_0))$ when the value is finite. There is no further preference for earlier capture in this second objective.

The phrase "least fuel" in a game needs an aggregation rule because different responses produce different expenditures. We use worst-case expenditure. If the evader cannot win, it maximizes the pursuer's stated objective: capture time for A1/B1, or required pursuer fuel for A2/B2. If it can win, it instead minimizes its own worst-case fuel while retaining a guaranteed win. This last preference is solved separately in Section VII. The pursuer's guarantee continues to quantify over all admissible responses, regardless of whether those responses are fuel-economical.

III. A Common Isaacs Operator

Let the full state be

$$z = (\mathbf{x}_P, \mathbf{v}_P, \mathbf{x}_E, \mathbf{v}_E, b_P, b_E). \tag{19}$$

For a differentiable function U, define the uncontrolled drift

$$\mathcal{D}_g U = \sum_{i=P,E} \left[\mathbf{v}_i \cdot U_{\mathbf{x}_i} + \mathbf{g}(\mathbf{x}_i) \cdot U_{\mathbf{v}_i} \right]. \tag{20}$$

Subscripts denote partial derivatives. For a running cost $\ell \|\mathbf{a}_P\|^2$, with $\ell \in \{0, 1\}$, the pursuer-minimizing operator is

$$\begin{aligned} \mathcal{H}_\ell[U] = \mathcal{D}_g U & \\ + \inf_{\mathbf{a}_P \in \mathcal{U}_P} & \left[U_{\mathbf{v}_P} \cdot \mathbf{a}_P + (\ell - U_{b_P}) \|\mathbf{a}_P\|^2 \right] \\ + \sup_{\mathbf{a}_E \in \mathcal{U}_E} & \left[U_{\mathbf{v}_E} \cdot \mathbf{a}_E - U_{b_E} \|\mathbf{a}_E\|^2 \right]. \end{aligned} \tag{21}$$

At a fixed state the two controls occur in separate summands. Therefore the pointwise infimum and supremum commute whenever the extrema are finite. This establishes the algebraic Isaacs condition; it does not by itself establish continuity at capture, existence of an optimal strategy, or uniqueness of a singular exit-time boundary problem.

A. Uncapped Interior Selectors

Suppose both budgets are positive and

$$c_P = \ell - U_{b_P} > 0, \qquad c_E = U_{b_E} > 0. \tag{22}$$

Completing squares gives

$$\boxed{\mathbf{a}_P^* = -\frac{U_{\mathbf{v}_P}}{2c_P}, \qquad \mathbf{a}_E^* = \frac{U_{\mathbf{v}_E}}{2c_E},} \tag{23}$$

$$\boxed{\mathcal{H}_\ell[U] = \mathcal{D}_g U - \frac{\|U_{\mathbf{v}_P}\|^2}{4c_P} + \frac{\|U_{\mathbf{v}_E}\|^2}{4c_E}.} \tag{24}$$

For example,

$$\mathbf{h} \cdot \mathbf{a} + c \|\mathbf{a}\|^2 = c \|\mathbf{a} + \mathbf{h}/(2c)\|^2 - \frac{\|\mathbf{h}\|^2}{4c}, \qquad c > 0. \tag{25}$$

Thus the selectors are global pointwise extrema, not merely stationary controls.

Budget monotonicity implies $V_{b_P} \le 0$, $V_{b_E} \ge 0$ wherever these derivatives exist; the same inequalities hold for W in the ideal model. The strict signs needed in (22) can nevertheless fail. If a quadratic coefficient vanishes while the associated linear coefficient is nonzero, the corresponding unbounded extremum is infinite. If both vanish the selector can be nonunique. Never divide by a zero budget derivative. Use (21), a constrained face equation, or a specified bounded-engine model at such points.

At $b_P = 0$ or $b_E = 0$, set the exhausted ship's acceleration to zero and omit its optimization term. This is a lower-dimensional game on a viable budget face, not an absorbing failure boundary.

B. Boundary and Regularity Scope

The value equations below are dynamic-programming characterizations. On smooth regions their classical form gives the feedbacks above. At shocks and competing trajectory branches, the appropriate generalized solutions are viscosity solutions selected by the game's dynamic programming principle, with target and state-constraint boundaries [3]. Smoothness is not presumed throughout the state space.

With compact controls, a nonsingular locally Lipschitz drift, and appropriate boundary/comparison hypotheses, standard HJI theory supplies the value interpretation. These hypotheses must be checked for the selected constrained domain. The ideal L^2 model is unbounded, so compact-control theorems cannot simply be cited as an unconditional existence proof for it. In particular, formulas (23)–(24) are valid on its coercive interior branches, and the verification theorem below applies directly to any admissible solution satisfying its hypotheses. Elsewhere the exact strategy definitions remain authoritative. Simultaneously sending two acceleration bounds to infinity, or a capture radius to zero, requires a separate convergence and attainment argument.

IV. Free-Space Interception and Rendezvous

A. Relative-State Reduction

Set

$$\mathbf{r} = \mathbf{x}_E - \mathbf{x}_P, \qquad \mathbf{w} = \mathbf{v}_E - \mathbf{v}_P. \tag{26}$$

Translations and Galilean boosts do not affect capture in free space. The complete reduced state is

$$z_A = (\mathbf{r}, \mathbf{w}, b_P, b_E), \tag{27}$$

with

$$\dot{\mathbf{r}} = \mathbf{w}, \tag{28a}$$

$$\dot{\mathbf{w}} = \mathbf{a}_E - \mathbf{a}_P, \tag{28b}$$

$$\dot{b}_P = -\left\|\mathbf{a}_P\right\|^2, \qquad \dot{b}_E = -\left\|\mathbf{a}_E\right\|^2. \tag{28c}$$

Consequently $U_{\mathbf{v}_P} = -U_{\mathbf{w}}$ and $U_{\mathbf{v}_E} = U_{\mathbf{w}}$. The operator becomes

$$\begin{aligned}\mathcal{H}^A_\ell[U] = \mathbf{w}\cdot U_{\mathbf{r}} &+ \inf_{\mathbf{a}_P\in\mathcal{U}_P}\left[-U_{\mathbf{w}}\cdot\mathbf{a}_P + (\ell - U_{b_P})\left\|\mathbf{a}_P\right\|^2\right] \\ &+ \sup_{\mathbf{a}_E\in\mathcal{U}_E}\left[U_{\mathbf{w}}\cdot\mathbf{a}_E - U_{b_E}\left\|\mathbf{a}_E\right\|^2\right].\end{aligned} \tag{29}$$

B. A1: Minimum Guaranteed Capture Time

The autonomous exit-time value solves, on its finite continuation region,

$$\boxed{1 + \mathcal{H}^A_0[V_A] = 0, \qquad V_A|_{\mathcal{C}} = 0.} \tag{30}$$

The relevant solution is the capture-time value (16), with budget-face and outer-domain conditions; the target condition alone does not select a solution. On a regular uncapped branch, put $c_P = -V_{A,b_P}$ and $c_E = V_{A,b_E}$. Then

$$1 + \mathbf{w}\cdot V_{A,\mathbf{r}} - \frac{\left\|V_{A,\mathbf{w}}\right\|^2}{4c_P} + \frac{\left\|V_{A,\mathbf{w}}\right\|^2}{4c_E} = 0, \tag{31}$$

and the optimal feedback accelerations are

$$\boxed{\mathbf{a}^*_P = \frac{V_{A,\mathbf{w}}}{2c_P}, \qquad \mathbf{a}^*_E = \frac{V_{A,\mathbf{w}}}{2c_E}.} \tag{32}$$

These are evaluated at the actual joint state throughout the chase. Both accelerations are parallel to the same relative-velocity gradient on this regular branch, but that direction is not generally the current line of sight.

If the verification conditions in Section VI hold, the pursuer catches by $V_A(z_{A0})$ against every admissible evader input, and the evader can prevent capture before that time against every pursuer input. This is a global minimax statement. It neither claims that distance decreases monotonically nor requires such a property. Monotone closing would be an additional constraint: initially $\mathrm{d}\left\|\mathbf{r}\right\|/\mathrm{d}t = \mathbf{r}\cdot\mathbf{w}/\left\|\mathbf{r}\right\|$, which cannot be made immediately negative by an L^2 acceleration if it is initially positive.

C. Minimum-Time Exact Rendezvous

The free-space rendezvous target is $\mathbf{r} = \mathbf{w} = 0$. Its value satisfies

$$\boxed{1 + \mathcal{H}^A_0[V^J_A] = 0, \qquad V^J_A|_{\mathcal{C}_J} = 0.} \tag{33}$$

On a regular uncapped branch this is

$$1 + \mathbf{w}\cdot V^J_{A,\mathbf{r}} - \frac{\left\|V^J_{A,\mathbf{w}}\right\|^2}{4(-V^J_{A,b_P})} + \frac{\left\|V^J_{A,\mathbf{w}}\right\|^2}{4V^J_{A,b_E}} = 0, \tag{34}$$

with feedback

$$\mathbf{a}^*_P = \frac{V^J_{A,\mathbf{w}}}{2(-V^J_{A,b_P})}, \qquad \mathbf{a}^*_E = \frac{V^J_{A,\mathbf{w}}}{2V^J_{A,b_E}}. \tag{35}$$

Unlike the interception value, V^J_A does not vanish at $\mathbf{r} = 0, \mathbf{w} \neq 0$. Theorem 3 applies with $\mathcal{C}_J$ replacing $\mathcal{C}$. For ideal engines, the value and opposing strategies can also be obtained explicitly.

1) The Minimum-Norm Rendezvous Correction: Reset time to zero at the initial state of the calculation. For a prescribed horizon $h > 0$, let $\mathbf{a}_h$ be a correction such that $\dot{\mathbf{w}} = -\mathbf{a}_h$ reaches $\mathbf{r}(h) = \mathbf{w}(h) = 0$. Its moments must satisfy

$$\int_0^h (h-t)\mathbf{a}_h(t)\mathrm{d}t = \mathbf{r}_0 + h\mathbf{w}_0, \quad \int_0^h \mathbf{a}_h(t)\mathrm{d}t = \mathbf{w}_0. \tag{36}$$

The Gram matrix of the functions $h - t$ and 1, and its inverse, are

$$G_h = \begin{pmatrix} h^3/3 & h^2/2 \\ h^2/2 & h \end{pmatrix}, \quad G_h^{-1} = \begin{pmatrix} 12/h^3 & -6/h^2 \\ -6/h^2 & 4/h \end{pmatrix}. \tag{37}$$

Projection onto their span gives the unique least-norm correction

$$\boxed{\mathbf{a}_h(t) = \frac{6\mathbf{r}_0}{h^2} + \frac{4\mathbf{w}_0}{h} - \left(\frac{12\mathbf{r}_0}{h^3} + \frac{6\mathbf{w}_0}{h^2}\right)t.} \tag{38}$$

Write this as $\mathbf{a}_h(t) = \mathbf{A}_h + \mathbf{B}_h t$, thereby defining its constant and linear coefficients. Indeed, any other control with the same moments differs by a function orthogonal to both 1 and t. Its

squared norm is the squared norm of (38) plus a nonnegative orthogonal contribution. Thus

$$\boxed{\begin{aligned} C_J(h) &= \int_0^h \|\mathbf{a}_h\|^2 \,\mathrm{d}t \\ &= \frac{12\,\|\mathbf{r}_0\|^2}{h^3} + \frac{12\mathbf{r}_0\cdot\mathbf{w}_0}{h^2} + \frac{4\,\|\mathbf{w}_0\|^2}{h}. \end{aligned}} \tag{39}$$

This is an auxiliary fixed-time correction cost, used to determine the minimum-time game. It obeys the useful identity

$$C_J'(h) = -\frac{4}{h^4}\,\|3\mathbf{r}_0 + h\mathbf{w}_0\|^2 \le 0. \tag{40}$$

For a nonzero relative phase state, C_J is strictly decreasing: its derivative can vanish at most at an isolated positive time. It decreases from $+\infty$ to zero.

2) A Constructive Escape Lemma:

Proposition 1 (Rendezvous avoidance with resource parity)**.** *In ideal free space, every state outside $\mathcal{C}_J$ with $b_E \ge b_P$ admits a perpetual rendezvous-avoidance feedback. It can be chosen so that its cumulative expenditure never exceeds the pursuer's cumulative expenditure.*

Proof. Write $p = b_P$ and introduce an evader ledger q with $q(0) = p(0)$, even if the actual evader reserve is larger. Both q and the actual reserve decrease at $-\|\mathbf{a}_E\|^2$. Choose $\lambda > 0$ with $\mathbf{r}_0 - \lambda\mathbf{w}_0 \neq 0$ and define

$$\begin{aligned} \mathbf{e} &= \frac{\mathbf{r}_0 - \lambda\mathbf{w}_0}{\|\mathbf{r}_0 - \lambda\mathbf{w}_0\|}, \\ Y(t) &= \mathbf{e}\cdot[\mathbf{r}(t) - (t+\lambda)\mathbf{w}(t)], \\ \mathbf{c}(t) &= -(t+\lambda)\mathbf{e}, \qquad \delta = q - p. \end{aligned} \tag{41}$$

Such a λ always exists outside rendezvous, and $Y(0) > 0$. Since $\dot{Y} = \mathbf{c}\cdot(\mathbf{a}_E - \mathbf{a}_P)$, use

$$\boxed{\mathbf{a}_E = \frac{\delta}{Y}\mathbf{c}(t), \qquad \dot{q} = -\|\mathbf{a}_E\|^2.} \tag{42}$$

This depends on observed resources and state, not on current pursuer acceleration. While $Y > 0$,

$$\begin{aligned} \dot{\delta} &= \|\mathbf{a}_P\|^2 - \|\mathbf{a}_E\|^2, \\ \frac{\mathrm{d}}{\mathrm{d}t}(\delta Y^2) &= Y^2\,\|\mathbf{a}_P - \mathbf{a}_E\|^2 \ge 0. \end{aligned} \tag{43}$$

The scalar equation for δ preserves $\delta \ge 0$. Hence $q \ge p \ge 0$, so the evader never spends more than the pursuer and respects its physical reserve.

The identity also prevents a finite first zero of Y. If δY^2 becomes positive at any preceding time t_1, then $\delta \le q(0)$ gives

$$Y(t)^2 \ge \frac{\delta(t_1)Y(t_1)^2}{q(0)} > 0. \tag{44}$$

If it remains zero, (43) forces $\mathbf{a}_P = \mathbf{a}_E$ almost everywhere; with $\delta(0) = 0$, both controls then vanish and Y stays constant. The zero-budget case is simply coasting. Local well-posedness on $Y > 0$, these bounds, and the finite acceleration budgets permit continuation over every finite interval. Since rendezvous would imply $Y = 0$, it never occurs. □

3) The Exact Value and Opposing Optimal Strategies:

Theorem 1 (Analytical free-space rendezvous game)**.** *For ideal uncapped engines with instantaneous evader-acceleration observation by the pursuer, a state already in $\mathcal{C}_J$ has $V_A^J = 0$. At every other state,*

$$\boxed{V_A^J = \begin{cases} H, & b_P > b_E, \\ +\infty, & b_P \le b_E, \end{cases}} \tag{45}$$

where H is the unique positive solution of

$$\boxed{C_J(H) = (\sqrt{b_P} - \sqrt{b_E})^2.} \tag{46}$$

Equivalently, putting $D_b = \sqrt{b_P} - \sqrt{b_E} > 0$,

$$D_b^2 H^3 - 4\,\|\mathbf{w}_0\|^2 H^2 - 12(\mathbf{r}_0\cdot\mathbf{w}_0)H - 12\,\|\mathbf{r}_0\|^2 = 0. \tag{47}$$

The pursuer attains this time with $\mathbf{a}_P = \mathbf{a}_E + \mathbf{a}_H$. An optimal evader uses $\mathbf{a}_E = k\mathbf{a}_H$ on $[0, H]$, where $k = \sqrt{b_E}/D_b$, until its remaining reserve equals the pursuer's; if equality occurs before H, it switches to (42) *with time reset at that state. If it reaches H without capture or an earlier switch, it subsequently coasts.*

Proof. Existence and uniqueness of H follow from (40). The pursuer's response cancels the evader's acceleration from relative motion and enforces both moment conditions. Minkowski's inequality gives

$$\|\mathbf{a}_P\|_{L^2(0,H)} \le \sqrt{b_E} + \sqrt{C_J(H)} = \sqrt{b_P}. \tag{48}$$

It is admissible against every evader input and guarantees rendezvous by H.

To prove the opposing lower bound, first consider the baseline evader control $k\mathbf{a}_H$. Suppose a pursuer achieves rendezvous at some $t < H$ before the resource-equality switch. Its endpoint moments require the correction $\mathbf{a}_t$. Since the baseline restricted to $[0, t]$ is affine, orthogonal projection yields

$$\int_0^t \|\mathbf{a}_P\|^2\,\mathrm{d}s - \int_0^t \|k\mathbf{a}_H\|^2\,\mathrm{d}s \ge C_J(t) + 2k\int_0^t \mathbf{a}_H\cdot\mathbf{a}_t\mathrm{d}s. \tag{49}$$

Writing $\mathbf{a}_H = \mathbf{A}_H + \mathbf{B}_H s$ and using (36) gives

$$\int_0^t \mathbf{a}_H\cdot\mathbf{a}_t\mathrm{d}s = \mathbf{A}_H\cdot\mathbf{w}_0 - \mathbf{B}_H\cdot\mathbf{r}_0 = C_J(H). \tag{50}$$

Strict decrease of C_J implies that the right side of (49) exceeds

$$(1+2k)C_J(H) = b_P(0) - b_E(0). \tag{51}$$

Such an expenditure difference would require $b_P(t) < b_E(t)$. Therefore rendezvous cannot occur while $b_P(t) > b_E(t)$, or at the first equality before H. At that equality the state is outside $\mathcal{C}_J$, and Proposition 1 supplies perpetual avoidance. The baseline evader fits its budget because $k^2 C_J(H) = b_E(0)$; the switched strategy uses only its remaining reserve. Thus no pursuer can force rendezvous before H.

Against the optimal pursuer response, the two controls are $(1+k)\mathbf{a}_H$ and $k\mathbf{a}_H$. Their remaining-resource difference stays positive for $t < H$ and reaches zero at H, so the evader does not switch early. Their common terminal time is exactly

H. When $b_P \le b_E$ initially, Proposition 1 proves perpetual avoidance immediately. □

The proof is global and uses neither a differentiable value function nor a shooting extremal. No game PDE is needed for this ideal free-space rendezvous value. A real-radical expression follows by defining

$$\begin{aligned} a &= 4\|\mathbf{w}_0\|^2/D_b^2, \quad b = 12\mathbf{r}_0\cdot\mathbf{w}_0/D_b^2,\\ c &= 12\|\mathbf{r}_0\|^2/D_b^2,\\ p_c &= -b - a^2/3, \qquad q_c = -2a^3/27 - ab/3 - c,\\ \Delta_c &= (q_c/2)^2 + (p_c/3)^3. \end{aligned} \tag{52}$$

Then

$$\begin{aligned} H = \frac{a}{3} &+ \operatorname{cbrt}(-q_c/2 + \sqrt{\Delta_c})\\ &+ \operatorname{cbrt}(-q_c/2 - \sqrt{\Delta_c}), \end{aligned} \tag{53}$$

where cbrt denotes the real cube root. Physical values of the coefficients have $\Delta_c \ge 0$: the cubic has a unique positive root, and it is strictly negative for negative arguments, by Cauchy–Schwarz applied to its quadratic terms. Repeated roots occur only in degenerate cases. Monotone scalar root finding avoids cancellation in the radical formula. At $3\mathbf{r}_0 + H\mathbf{w}_0 = 0$, the implicit time derivative is singular, but the root and the strategy construction remain valid. Simple limiting cases are

$$\mathbf{w}_0 = 0: \quad V_A^J = \left(\frac{12\|\mathbf{r}_0\|^2}{D_b^2}\right)^{1/3}, \tag{54}$$

$$\mathbf{r}_0 = 0,\ \mathbf{w}_0 \neq 0: \quad V_A^J = \frac{4\|\mathbf{w}_0\|^2}{D_b^2}. \tag{55}$$

The second case is a rendezvous problem even though position-only capture has already occurred.

4) Least-Fuel Winning Escape and the Saddle Trajectories: For $b_E \ge b_P$ outside $\mathcal{C}_J$, the least worst-case acceleration expenditure for perpetual rendezvous avoidance is exactly

$$\boxed{F_{A,\infty}^J = b_P.} \tag{56}$$

The feedback (42) with initial ledger $q(0) = b_P$ attains an upper bound of b_P. If a winning evader strategy had a strictly smaller uniform total expenditure cap $c < b_P$, the pursuer response in Theorem 1, computed with evader cap c, would rendezvous in finite time against that strategy. This contradiction proves the matching lower bound. Thus the least-fuel winning strategy is attained in this ideal rendezvous game, even though strict position-only escape can exhibit nonattainment. At an arbitrary current state with evader mass m_E, the corresponding mass expenditure is

$$m_E - \left(m_E^{-1} + \frac{b_P}{2P_E}\right)^{-1}. \tag{57}$$

For $b_P > b_E$, define $s = t/H$ and

$$\mathbf{Q}(t) = \tfrac{1}{2}\mathbf{A}_H t^2 + \tfrac{1}{6}\mathbf{B}_H t^3, \tag{58}$$

$$E_H(t) = \|\mathbf{A}_H\|^2 t + (\mathbf{A}_H\cdot\mathbf{B}_H)t^2 + \tfrac{1}{3}\|\mathbf{B}_H\|^2 t^3. \tag{59}$$

Along the opposing optimal controls, the dimensional trajectories are

$$\begin{aligned} \mathbf{x}_P(t) &= \mathbf{x}_{P0} + \mathbf{v}_{P0}t + (1+k)\mathbf{Q}(t),\\ \mathbf{x}_E(t) &= \mathbf{x}_{E0} + \mathbf{v}_{E0}t + k\mathbf{Q}(t),\\ b_P(t) &= b_P(0) - (1+k)^2 E_H(t),\\ b_E(t) &= b_E(0) - k^2 E_H(t). \end{aligned} \tag{60}$$

Velocities are the derivatives of these cubic paths; both remaining budgets vanish at H. Their deterministic relative path is

$$\mathbf{r}_J(t) = \mathbf{r}_0(1 - 3s^2 + 2s^3) + H\mathbf{w}_0(s - 2s^2 + s^3). \tag{61}$$

Under an arbitrary evader response the pursuer's cancellation strategy retains (61), although the individual trajectories and fuel histories change.

The mass and specific impulse along the saddle are also explicit. Put $\gamma_P = 1 + k$ and $\gamma_E = k$. Then

$$m_i(t) = \left[m_{i0}^{-1} + \frac{\gamma_i^2 E_H(t)}{2P_i}\right]^{-1}, \tag{62}$$

$$v_{e,i}(t) = \frac{2P_i}{m_i(t)\gamma_i\|\mathbf{A}_H + \mathbf{B}_H t\|}, \qquad I_{sp,i}(t) = v_{e,i}(t)/g_0. \tag{63}$$

The second formula applies on powered arcs; zero acceleration is coasting. Near a zero of an affine acceleration, the ideal exhaust speed can diverge, so a finite exhaust-speed ceiling changes the solution. For $\mathbf{w}_0 = 0$, the normalized expenditure has the simple form

$$E_H(t) = C_J(H)(3s - 6s^2 + 4s^3), \qquad s = t/H. \tag{64}$$

The correction reverses direction at $s = 1/2$, with final velocity matching supplied by the second half of the maneuver.

D. A2: Minimum Fuel With Capture by a Deadline

Let $s = T - t$ be remaining time. The value satisfies

$$\boxed{\partial_s W_A = \mathcal{H}_1^A[W_A].} \tag{65}$$

Its stopping and terminal data are

$$\begin{aligned} W_A(s,z) &= 0 && \text{for } z \in \mathcal{C},\ s \ge 0,\\ W_A(0,z) &= +\infty && \text{for } z \notin \mathcal{C}. \end{aligned} \tag{66}$$

The infinite condition enforces guaranteed capture rather than imposing a finite penalty for missing the deadline. The equation is solved on the feasible capture domain, or equivalently with a budgeted reachability construction as in Section X.

For $c_P = 1 - W_{A,b_P} > 0$ and $c_E = W_{A,b_E} > 0$,

$$\partial_s W_A = \mathbf{w}\cdot W_{A,\mathbf{r}} - \frac{\|W_{A,\mathbf{w}}\|^2}{4c_P} + \frac{\|W_{A,\mathbf{w}}\|^2}{4c_E}, \tag{67}$$

with

$$\boxed{\mathbf{a}_P^* = \frac{W_{A,\mathbf{w}}}{2c_P}, \qquad \mathbf{a}_E^* = \frac{W_{A,\mathbf{w}}}{2c_E}.} \tag{68}$$

Capture ends fuel accounting at its first occurrence. There is no terminal requirement $\tau = T$: an earlier intercept may use less fuel, for example if the unforced paths already cross. Thus solving a fixed-terminal-time interception problem alone does not solve A2.

E. An Exact Five-State Symmetry Reduction

For isotropic thrust constraints and either capture target, the value is invariant under simultaneous orthogonal transformations of $\mathbf{r}, \mathbf{w}$. For $d = 2$ and 3 it can therefore be written as

$$\begin{aligned} U &= \widehat{U}(\rho, \xi, \omega, b_P, b_E), \\ \rho = \|\mathbf{r}\|^2, \quad \xi &= \mathbf{r} \cdot \mathbf{w}, \quad \omega = \|\mathbf{w}\|^2. \end{aligned} \tag{69}$$

The Gram domain is $\rho \geq 0$, $\omega \geq 0$, $\xi^2 \leq \rho\omega$. The chain rule gives

$$\mathbf{w} \cdot U_{\mathbf{r}} = 2\xi \widehat{U}_\rho + \omega \widehat{U}_\xi, \tag{70}$$

$$U_{\mathbf{w}} = \widehat{U}_\xi \mathbf{r} + 2\widehat{U}_\omega \mathbf{w}, \tag{71}$$

$$\|U_{\mathbf{w}}\|^2 = \rho \widehat{U}_\xi^2 + 4\xi \widehat{U}_\xi \widehat{U}_\omega + 4\omega \widehat{U}_\omega^2. \tag{72}$$

Substituting (70) and (72) into the free-space operators produces a five-state PDE, plus the horizon variable for a deadline objective. The rendezvous target is $\rho = \omega = 0$, which also forces $\xi = 0$. The boundary $\xi^2 = \rho\omega$ represents collinear geometry, not an artificial absorbing boundary; consistency with the original vector game supplies its treatment. This reduction changes neither the fuel constraints nor the adversarial quantifiers.

There is also an exact similarity law. Under positive scale factors α, β,

$$\widetilde{t} = \alpha t, \quad \widetilde{\mathbf{r}} = \beta \mathbf{r}, \quad \widetilde{\mathbf{w}} = \frac{\beta}{\alpha}\mathbf{w}, \quad \widetilde{b}_i = \frac{\beta^2}{\alpha^3} b_i, \tag{73}$$

the ideal dynamics retain their form, with $\widetilde{R} = \beta R$. Both time values scale by α, and a deadline-fuel acceleration expenditure scales by β^2/α^3 when its deadline is scaled by α. This follows by scaling every admissible strategy and its opponent, so it holds for minimax values rather than only for extremal trajectories.

V. Interception and Rendezvous in a Kepler Field

A. Why the Free-Space Relative Game Is Insufficient

Relative acceleration is now

$$\dot{\mathbf{w}} = -\mu \frac{\mathbf{x}_E}{\|\mathbf{x}_E\|^3} + \mu \frac{\mathbf{x}_P}{\|\mathbf{x}_P\|^3} + \mathbf{a}_E - \mathbf{a}_P. \tag{74}$$

The first two terms cannot be inferred from $\mathbf{r}, \mathbf{w}$ alone. Either retain both inertial states or retain one full orbital state in addition to relative position and velocity. The full state has $4d + 2$ components: ten in the plane and fourteen in space. A common rotation is a symmetry; in the planar isotropic problem it can remove one additional state variable. A common translation or velocity boost is not a symmetry of the central field.

B. B1: Minimum Guaranteed Capture Time

Use (20) with $\mathbf{g} = -\mu\mathbf{x}/\|\mathbf{x}\|^3$. The value is characterized by

$$\boxed{1 + \mathcal{H}_0^B[V_B] = 0, \qquad V_B|_{\mathcal{C}} = 0,} \tag{75}$$

with the same capture-time interpretation and the specified gravitational-domain boundaries. On a regular uncapped branch,

$$1 + \mathcal{D}_g V_B - \frac{\|V_{B,\mathbf{v}_P}\|^2}{4(-V_{B,b_P})} + \frac{\|V_{B,\mathbf{v}_E}\|^2}{4V_{B,b_E}} = 0. \tag{76}$$

The feedback accelerations are

$$\boxed{\mathbf{a}_P^* = -\frac{V_{B,\mathbf{v}_P}}{2(-V_{B,b_P})}, \qquad \mathbf{a}_E^* = \frac{V_{B,\mathbf{v}_E}}{2V_{B,b_E}}.} \tag{77}$$

The two velocity gradients are generally different, so the thrust directions need not be parallel. Gravity can reward maneuvers that temporarily increase separation. Neither constant bearing nor a fixed acceleration–braking sequence is a general optimum.

C. Minimum-Time Exact Rendezvous

The Kepler rendezvous value uses the full orbital state and target (14):

$$\boxed{1 + \mathcal{H}_0^B[V_B^J] = 0, \qquad V_B^J|_{\mathcal{C}_J} = 0.} \tag{78}$$

Its regular uncapped form is

$$1 + \mathcal{D}_g V_B^J - \frac{\|V_{B,\mathbf{v}_P}^J\|^2}{4(-V_{B,b_P}^J)} + \frac{\|V_{B,\mathbf{v}_E}^J\|^2}{4V_{B,b_E}^J} = 0, \tag{79}$$

and the feedback accelerations are

$$\mathbf{a}_P^* = -\frac{V_{B,\mathbf{v}_P}^J}{2(-V_{B,b_P}^J)}, \qquad \mathbf{a}_E^* = \frac{V_{B,\mathbf{v}_E}^J}{2V_{B,b_E}^J}. \tag{80}$$

Equations (10) and (80), with fuel-face and central-body boundary conditions, are the closed-loop trajectory equations. Global optimality follows from Theorem 3 with the rendezvous target whenever its hypotheses hold. A velocity mismatch at a position crossing is a continuation state, even if a position-only intercept would already have terminated.

The free-space cubic does not remain exact in a central field: the extra acceleration needed to shape relative motion depends on both evolving positions. In particular, $b_P > b_E$ alone does not provide the free-space rendezvous time in this field. A separate exact limiting result is available when the pursuer is out of fuel. If the initial phase states differ, a coasting evader can never have the same position and velocity as the coasting pursuer at a finite nonsingular time. Otherwise uniqueness of the Kepler flow, integrated backward from that common state, would imply identical initial phase states. Thus $V_B^J = +\infty$ and evader coasting is a zero-fuel optimal perpetual escape strategy whenever $b_P = 0$, the initial phase states differ, and the ballistic trajectories remain admissible. The same argument applies in free space.

1) An Explicit Gravity-Compensated Capture Certificate: In the instantaneous-response model, select a horizon h and the free-space rendezvous correction (38). Use

$$\boxed{\mathbf{a}_P = \mathbf{a}_E + \mathbf{g}(\mathbf{x}_E) - \mathbf{g}(\mathbf{x}_P) + \mathbf{a}_h(t).} \tag{81}$$

Then relative motion is exactly (61), with H replaced by h, against every evader input. The differential gravity has

been compensated, not neglected. The remaining question is a uniform bound on the required pursuer resource.

Suppose the segment joining the two positions stays in a region $\|\mathbf{x}\| \geq r_* > 0$. Since the largest singular value of $D\mathbf{g}$ there is at most

$$K_g = 2\mu/r_*^3, \tag{82}$$

the mean-value integral gives

$$\|\mathbf{g}(\mathbf{x}_E) - \mathbf{g}(\mathbf{x}_P)\| \leq K_g \|\mathbf{r}_J(t)\|. \tag{83}$$

The required polynomial quadrature is explicit:

$$\begin{aligned} I_J(h) &= \int_0^h \|\mathbf{r}_J(t)\|^2 \,\mathrm{d}t \\ &= \frac{13h}{35}\|\mathbf{r}_0\|^2 + \frac{11h^2}{105}\mathbf{r}_0\cdot\mathbf{w}_0 + \frac{h^3}{105}\|\mathbf{w}_0\|^2. \end{aligned} \tag{84}$$

Minkowski's inequality proves the sufficient condition

$$\boxed{\sqrt{b_P} \geq \sqrt{b_E} + \sqrt{C_J(h)} + K_g\sqrt{I_J(h)}.} \tag{85}$$

Under the stated corridor condition, this guarantees exact rendezvous by h, and hence $V_B^J \leq h$. It is a certified upper bound rather than a global optimal-time formula.

The corridor can itself be certified against all evader controls. Choose a fixed unit vector $\mathbf{n}$ and require, for every $0 \leq t \leq h$,

$$\begin{aligned} \mathbf{n}\cdot(\mathbf{x}_{E0} + t\mathbf{v}_{E0}) - \frac{\mu t^2}{2r_*^2} - \sqrt{b_E t^3/3} \\ - \max\{\mathbf{n}\cdot\mathbf{r}_J(t), 0\} > r_*. \end{aligned} \tag{86}$$

Before any first exit from the half-space $\mathbf{n}\cdot\mathbf{x} \geq r_*$, the gravitational displacement along $\mathbf{n}$ is bounded below by $-\mu t^2/(2r_*^2)$ and the evader thrust displacement by $-\sqrt{b_E t^3/3}$. Equation (86) therefore excludes such an exit for either ship, because $\mathbf{x}_P = \mathbf{x}_E - \mathbf{r}_J$. The whole joining segment is then in that convex half-space. This first-exit argument establishes the corridor rather than assuming a favorable realized evader path. If a central exclusion radius is imposed, choose r_* at least as large.

For $\mathbf{w}_0 = 0$ and $L = \|\mathbf{r}_0\|$, the resource inequality simplifies to

$$\sqrt{b_P} - \sqrt{b_E} \geq \frac{\sqrt{12}L}{h^{3/2}} + K_g L\sqrt{13h/35}. \tag{87}$$

The gravity correction grows with the allowed time in this conservative certificate, whereas the free-space correction decreases. The zero-gravity limit recovers the exact free-space time equation.

The same construction certifies point interception using

$$\begin{aligned} \mathbf{a}_h^I(t) &= \frac{3(h-t)}{h^3}(\mathbf{r}_0 + h\mathbf{w}_0), \\ \mathbf{r}_I(t) &= \mathbf{r}_0 + t\mathbf{w}_0 - \frac{3s^2 - s^3}{2}(\mathbf{r}_0 + h\mathbf{w}_0), \quad s = t/h. \end{aligned} \tag{88}$$

Replace C_J by $C_I = 3\|\mathbf{r}_0 + h\mathbf{w}_0\|^2/h^3$, and I_J by

$$I_I = \frac{17h}{35}\|\mathbf{r}_0\|^2 + \frac{6h^2}{35}\mathbf{r}_0\cdot\mathbf{w}_0 + \frac{2h^3}{105}\|\mathbf{w}_0\|^2. \tag{89}$$

Use $\mathbf{r}_I$ in (86). The endpoint is exact position coincidence, so it also suffices for a positive capture radius. These compensation strategies are for ideal engines; finite instantaneous engine limits require checking their full acceleration histories.

D. The Exact Local CW Rendezvous Game

Near a prograde circular reference orbit of mean motion n, the planar Clohessy–Wiltshire (CW) equations retain the linear solar tide and Coriolis terms [5]. In physical units, with rotating state $s = (x, y, \dot{x}, \dot{y})$, write $\dot{s} = A_n s + B\mathbf{a}$, where

$$A_n = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 3n^2 & 0 & 0 & 2n \\ 0 & 0 & -2n & 0 \end{pmatrix}, \qquad B = \begin{pmatrix} 0 & 0 \\ 0 & 0 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}. \tag{90}$$

Let $z = s_E - s_P$, $\Phi(t) = \exp(A_n t)$ and define

$$\begin{aligned} K(t) &= \Phi(-t)B, \qquad M(h) = \int_0^h K(t)K(t)^T \mathrm{d}t, \\ C_{\mathrm{CW}}(h) &= z_0^T M(h)^{-1} z_0, \qquad a_h(t) = K(t)^T M(h)^{-1} z_0. \end{aligned} \tag{91}$$

Here C_{CW} and b_i both have units $\mathrm{m}^2/\mathrm{s}^3$. Controllability makes $M(h)$ positive definite for $h > 0$. For $z_0 \neq 0$, C_{CW} decreases strictly and continuously from infinity to zero; $C'_{\mathrm{CW}}(h) = -|K(h)^T M(h)^{-1} z_0|^2$. Strict decrease follows from controllability on each nonempty interval; the large-time limit follows from the trigonometric-polynomial CW transition matrix.

Theorem 2 (Ideal planar CW rendezvous)**.** *Assume unrestricted square-integrable accelerations, no additional state constraints, and instantaneous observation of the evader's current acceleration by the pursuer. For $z_0 \neq 0$ and $b_P > b_E$, the guaranteed rendezvous time is the unique $H > 0$ satisfying*

$$\boxed{C_{\mathrm{CW}}(H) = (\sqrt{b_P} - \sqrt{b_E})^2.} \tag{92}$$

It is infinite when $b_P \leq b_E$, and zero at $z_0 = 0$.

Proof. For the upper bound, use $\mathbf{a}_P = \mathbf{a}_E + a_H$. Then $z(t) = \Phi(t)[z_0 - \int_0^t K(s)a_H(s)\mathrm{d}s]$, so $z(H) = 0$. Minkowski gives $\|\mathbf{a}_P\|_2 \leq \sqrt{b_E} + \sqrt{C_{\mathrm{CW}}(H)} = \sqrt{b_P}$ against every allowed evader.

For the opposing bound put $k = \sqrt{b_E}/(\sqrt{b_P} - \sqrt{b_E})$. The evader starts with $\mathbf{a}_E = ka_H$ until resource parity. Any rendezvous at $t < H$ requires a pursuer correction with moments $\int_0^t K(s)(\mathbf{a}_P - \mathbf{a}_E)\mathrm{d}s = z_0$. Minimum-norm orthogonality gives

$$\begin{aligned} \int_0^t (|\mathbf{a}_P|^2 - |\mathbf{a}_E|^2)\mathrm{d}s &\geq C_{\mathrm{CW}}(t) + 2kC_{\mathrm{CW}}(H) \\ &> b_P(0) - b_E(0). \end{aligned} \tag{93}$$

Thus rendezvous is impossible before or at first resource parity. The planned evader burn spends at most $b_E(0)$ by H.

At parity, reset time at the nonzero state z_* and put $l = z_*/|z_*|$. Introduce a virtual evader reserve $q(0) = b_P(0)$, $\dot{q} = -|\mathbf{a}_E|^2$. Define

$$\begin{aligned} Y = l^T\Phi(-t)z, \quad c = B^T\Phi(-t)^T l, \quad \delta = q - b_P, \\ \mathbf{a}_E = (\delta/Y)c. \end{aligned} \tag{94}$$

Initially $Y > 0$, $\delta = 0$. While $Y > 0$, the feedback preserves $\delta \geq 0$ and

$$(\delta Y^2)' = Y^2 |\mathbf{a}_P - \mathbf{a}_E|^2 \geq 0. \tag{95}$$

If the product becomes positive, $\delta \leq q(0)$ prevents Y from vanishing at finite time. If it stays zero, both controls vanish almost everywhere and Y stays constant. The virtual budget bounds actual expenditure. The same strategy starts immediately when $b_E \geq b_P$; if $b_P = 0$ it is simply coasting. This proves the lower bound. □

The result concerns full-state rendezvous. In the dimensional limit $n \to 0$, A_n becomes the free-space double integrator and

$$C_{\mathrm{CW}}(h) \longrightarrow \frac{12|\mathbf{r}_0|^2}{h^3} + \frac{12\mathbf{r}_0 \cdot \mathbf{w}_0}{h^2} + \frac{4|\mathbf{w}_0|^2}{h} = C_J(h). \tag{96}$$

Hence the value equation reduces to the free-space cubic in Theorem 1. Taking this physical limit precedes any normalization that sets $n = 1$.

1) A Sufficient Nonlinear Extension and Explicit Tube: Let the reference radius be R_c, $n^2 = \mu/R_c^3$, and use normalized position q, time $\tau = nt$, acceleration $u = \mathbf{a}/(R_c n^2)$ and budget $\bar{b} = b/(R_c^2 n^3)$. In these units the exact rotating Kepler field is

$$q'' + 2Jq' = \mathrm{diag}(3,0)q + u + N(q), \quad J = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \tag{97}$$

where

$$N(q) = -\frac{e_1 + q}{|e_1 + q|^3} + e_1 - \mathrm{diag}(2,-1)q. \tag{98}$$

On $|q| \leq \epsilon < 1$, Taylor's formula and $\|D^2 g(x)[a,b]\| \leq 6|a||b|/|x|^4$ give

$$|N| \leq N_\epsilon = \frac{3\epsilon^2}{(1-\epsilon)^4}, \qquad \mathrm{Lip}(N) \leq L_\epsilon = \frac{6\epsilon}{(1-\epsilon)^4}. \tag{99}$$

Compute the preceding CW objects with $n = 1$, normalized states and horizon h. The response $u_P = u_E + a_h + N(q_E) - N(q_P)$ makes relative motion exactly the CW correction path z_h. If both ships stay in the ball, Minkowski yields the sufficient resource condition

$$\begin{aligned} \sqrt{\bar{b}_P} \geq \sqrt{\bar{b}_E} + \sqrt{C_{\mathrm{CW}}(h)} \\ + L_\epsilon \left(\int_0^h |\Pi z_h|^2 \mathrm{d}\tau \right)^{1/2}, \end{aligned} \tag{100}$$

Here $\Pi = (I, 0)$. The ball can be certified rather than assumed. Define

$$\begin{aligned} g_h &= h + h^2 + 2h^3/3, \\ G_1 &= h^2/2 + h^3/3 + h^4/6, \\ G_2 &= h^3/3 + h^4/2 + 7h^5/15 + 2h^6/9 + 4h^7/63, \\ F_h(s) &= |q| + (3h^2/2 + h^3)|q_x| + g_h|v|, \\ D_h &= F_h(z_0) + \sqrt{C_{\mathrm{CW}}(h) G_2}, \\ E_h &= F_h(s_{E0}) + \sqrt{\bar{b}_E G_2}. \end{aligned} \tag{101}$$

TABLE I
PURSUER VALUE EQUATIONS AND CAPTURE MODES

Case	Value equation	Initial answer
A1	$1 + \mathcal{H}_0^A[V_A] = 0$	$V_A(z_{A0})$
A1, rendezvous	$1 + \mathcal{H}_0^A[V_A^J] = 0$	(45)–(47)
A2	$(W_A)_s = \mathcal{H}_1^A[W_A]$	$\Phi_P(W_A(T, z_{A0}))$
B1	$1 + \mathcal{H}_0^B[V_B] = 0$	$V_B(z_0)$
B1, rendezvous	$1 + \mathcal{H}_0^B[V_B^J] = 0$	$V_B^J(z_0)$
B2	$(W_B)_s = \mathcal{H}_1^B[W_B]$	$\Phi_P(W_B(T, z_0))$

The explicit sufficient test is

$$\begin{aligned} E_h + D_h + N_\epsilon G_1 < \epsilon, \\ \sqrt{\bar{b}_P} \geq \sqrt{\bar{b}_E} + \sqrt{C_{\mathrm{CW}}(h)} + L_\epsilon \sqrt{h}\, D_h. \end{aligned} \tag{102}$$

Indeed, the CW matrix exponential obeys $\|\Pi\Phi(\tau)B\| \leq \tau + \tau^2 + 2\tau^3/3$ and $|\Pi\Phi(\tau)s| \leq F_h(s)$ for $0 \leq \tau \leq h$. Variation of constants and Cauchy–Schwarz bound the evader displacement by $E_h + N_\epsilon G_1$ and the relative path by D_h. The first inequality excludes a first exit of either ship; the second supplies its uniform fuel bound. Impose $R_c(1-\epsilon) \geq r_{\min}$ for a physical exclusion radius. Failure of this sufficient test is inconclusive.

This local certificate incorporates the linear tide exactly and compensates only the nonlinear remainder, whose Lipschitz bound is $O(\epsilon)$. It can therefore be sharper locally than compensating the entire field with K_g. It does not uniformly dominate the inertial half-space certificate above, which remains useful at wide separations. These are ideal-engine results: a finite-band guarantee additionally requires admissibility of the feedback against every allowed opponent. Delayed acceleration observation invalidates the cancellation used by this proof, so the exact value and stated capture time are no longer guaranteed.

E. B2: Minimum Fuel With Capture by a Deadline

The fuel value solves

$$\boxed{\partial_s W_B = \mathcal{H}_1^B[W_B],} \tag{103}$$

with the stopping and terminal conditions (66), and the Kepler-domain state constraints. Its regular uncapped form is

$$\partial_s W_B = \mathcal{D}_g W_B - \frac{\|W_{B,\mathbf{v}_P}\|^2}{4(1 - W_{B,b_P})} + \frac{\|W_{B,\mathbf{v}_E}\|^2}{4W_{B,b_E}}, \tag{104}$$

with

$$\boxed{\mathbf{a}_P^* = -\frac{W_{B,\mathbf{v}_P}}{2(1 - W_{B,b_P})}, \qquad \mathbf{a}_E^* = \frac{W_{B,\mathbf{v}_E}}{2W_{B,b_E}}.} \tag{105}$$

Equations (10) and (105), together with $\dot{s} = -1$, reconstruct the optimal feedback outcomes once the value has been obtained. The worst-case propellant infimum is $\Phi_P(W_B(T, z_0))$; it is attained when an optimal admissible selector exists. If $W_B = +\infty$, no admissible pursuer strategy meets the specified robust deadline.

VI. Global Optimality Verification

The distinction between an extremal and a sufficient certificate is essential. The following results prove optimality of *feedback strategies against arbitrary opposing controls*. They do not assume the opponent follows its extremal trajectory. In the minimum-time theorem, write $\mathcal{C}$ and τ for either the interception target and its hitting time or the exact rendezvous target $\mathcal{C}_J$ and τ_J.

Theorem 3 (Minimum-time verification)**.** *Let Ω be a region outside $\mathcal{C}$ on which a finite nonnegative function V is continuously differentiable, satisfies $1 + \mathcal{H}_0[V] = 0$, and extends continuously to $V = 0$ on the capture boundary. Suppose $V > 0$ outside capture and pointwise minimizing and maximizing selectors exist. Assume the selected feedbacks generate well-posed, admissible trajectories, and the differential inequalities below remain valid along all opposing responses up to capture. In particular, the pursuer selector cannot exit the certificate domain into an uncertified region. Then the pursuer guarantees $\tau \le V(z_0)$ and the evader guarantees $\tau \ge V(z_0)$. If both selectors are played in a well-posed joint trajectory, its capture time is exactly $V(z_0)$.*

Proof. For the pursuer selector and any admissible evader input, separability and pointwise optimality imply

$$1 + DV \cdot f(z, \mathbf{a}_P^*, \mathbf{a}_E) \le 0. \tag{106}$$

Hence $V(z(t)) \le V(z_0) - t$ before capture. Continuation beyond $V(z_0)$ would contradict nonnegativity; continuation exactly to that time outside the target contradicts $V > 0$. Thus capture occurs by $V(z_0)$. The stated domain and boundary assumptions exclude loss of the inequality before that conclusion.

Conversely, for the evader selector and every pursuer input,

$$1 + DV \cdot f(z, \mathbf{a}_P, \mathbf{a}_E^*) \ge 0. \tag{107}$$

At any finite capture time this gives $0 \ge V(z_0) - \tau$, or $\tau \ge V(z_0)$. If capture never occurs, the same lower bound holds trivially. Both inequalities together prove the saddle value and global minimax optimality. □

Theorem 4 (Deadline-fuel verification)**.** *Let $W(s, z)$ be a finite continuously differentiable solution of $W_s = \mathcal{H}_1[W]$ on a continuation domain with $W = 0$ at capture. Suppose the pursuer selector is admissible and guarantees capture by s_0 against every opponent while remaining in that domain. Suppose a maximizing evader selector is also admissible and the inequalities below are valid until capture. Then the pursuer guarantees expenditure at most $W(s_0, z_0)$, and the evader forces expenditure at least $W(s_0, z_0)$ or failure to meet the deadline. Thus W is the global minimax value there.*

Proof. Along a physical trajectory, $\dot{s} = -1$ and

$$\frac{\mathrm{d}}{\mathrm{d}t} W(s(t), z(t)) = -W_s + DW \cdot f. \tag{108}$$

For the pursuer selector and arbitrary evader controls,

$$\|\mathbf{a}_P^*\|^2 + \frac{\mathrm{d}W}{\mathrm{d}t} \le 0. \tag{109}$$

Integrating to its guaranteed capture gives

$$\int_0^\tau \|\mathbf{a}_P^*\|^2 \, \mathrm{d}t \le W(s_0, z_0). \tag{110}$$

For the evader selector and arbitrary pursuer controls the inequality reverses. If the pursuer captures by the deadline, integration gives expenditure at least $W(s_0, z_0)$. Otherwise $J_{s_0} = +\infty$. These are the required opposing global bounds. □

The capture-feasibility hypothesis in Theorem 4 is substantive. A smooth solution of an interior fuel PDE and its pointwise selectors do not prove that capture will occur. Feasibility must come from the correct terminal/viability construction or a separate reachability certificate. This prevents a spurious low-fuel strategy from winning on paper by never intercepting.

For nonsmooth value functions, dynamic programming replaces ordinary derivatives by the appropriate viscosity and sub-/superoptimality statements. Compatible feedback approximations can provide ε-optimal guarantees under the relevant comparison and strategy-existence hypotheses. The existence of a value, the existence of ε-optimal strategies, and the existence of an exactly optimal classical feedback are separate claims. No unconditional existence of the last object is asserted here.

VII. The Evader's Least-Fuel Winning Strategy

The survival construction below applies to either capture mode by using its own forbidden target and first hitting time. For a deadline-fuel interception game, winning means survival through the deadline. For either minimum-time game, an evader win means perpetual avoidance. In free-space rendezvous, (45) and (56) already give the complete winning set and its attained least-fuel escape cost. In the Kepler rendezvous game, the same reversed-role operator below defines the least-fuel perpetual-survival problem with forbidden set $\mathcal{C}_J$.

A. Finite-Deadline Survival

For a remaining horizon s, let

$$\mathfrak{E}_s(z) = \{\beta : \tau[z; \mathbf{a}_P, \beta[\mathbf{a}_P]] > s \text{ for every admissible } \mathbf{a}_P\}. \tag{111}$$

Strict inequality is necessary because capture at the deadline is a pursuer win. Define

$$F(s, z) = \inf_{\beta \in \mathfrak{E}_s(z)} \sup_{\mathbf{a}_P} \int_0^s \|\mathbf{a}_E(t)\|^2 \, \mathrm{d}t. \tag{112}$$

Set $F = +\infty$ when $\mathfrak{E}_s$ is empty. The actual evader's least worst-case propellant value is $\Phi_E(F(s, z))$ where finite. Exact attainment requires an additional check.

In this auxiliary game the evader minimizes and the pursuer maximizes. A pursuer able to force capture gives the evader infinite loss, whatever the pursuer's own fuel cost. The operator is

$$\begin{aligned} \mathcal{H}_E[F] = \mathcal{D}_g F &+ \sup_{\mathbf{a}_P \in \mathcal{U}_P} [F_{\mathbf{v}_P} \cdot \mathbf{a}_P - F_{b_P} \|\mathbf{a}_P\|^2] \\ &+ \inf_{\mathbf{a}_E \in \mathcal{U}_E} [F_{\mathbf{v}_E} \cdot \mathbf{a}_E + (1 - F_{b_E}) \|\mathbf{a}_E\|^2]. \end{aligned} \tag{113}$$

The finite-horizon equation and data are

$$F_s = \mathcal{H}_E[F], \tag{114}$$

$$F(0,z) = 0 \qquad (z \notin \mathcal{C}),$$

$$F(s,z) = +\infty \qquad (z \in \mathcal{C},\ s \geq 0). \tag{115}$$

They are understood on the strict survival domain, with the strategy definition (112) resolving singular boundaries.

For a regular ideal branch, $F_{b_P} > 0$ and $1 - F_{b_E} > 0$ give

$$\mathcal{H}_E[F] = \mathcal{D}_g F + \frac{\|F_{\mathbf{v}_P}\|^2}{4F_{b_P}} - \frac{\|F_{\mathbf{v}_E}\|^2}{4(1-F_{b_E})}, \tag{116}$$

$$\mathbf{a}_P^\dagger = \frac{F_{\mathbf{v}_P}}{2F_{b_P}}, \qquad \mathbf{a}_E^\dagger = -\frac{F_{\mathbf{v}_E}}{2(1-F_{b_E})}. \tag{117}$$

Here $F_{b_P} \geq 0$ and $F_{b_E} \leq 0$ express that a better-equipped pursuer makes survival more expensive, whereas additional evader reserve cannot make it harder. For free space, substitute $F_{\mathbf{v}_P} = -F_{\mathbf{w}}$ and $F_{\mathbf{v}_E} = F_{\mathbf{w}}$; the same five-state reduction applies.

Proposition 2 (Verification of least-fuel deadline survival)**.** *Suppose the minimizing selector in* (113) *stays strictly outside* $\mathcal{C}$ *through* s_0 *against every opposing input and remains admissible. Suppose a smooth certificate* F *satisfies* (114) *with terminal value zero along those surviving trajectories. If a maximizing selector provides the reverse inequality for every winning evader strategy, then* (117)*, or its constrained counterpart, attains* (112)*.*

Proof. For the minimizing evader selector,

$$\left\|\mathbf{a}_E^\dagger\right\|^2 + \frac{\mathrm{d}}{\mathrm{d}t}F(s(t),z(t)) \leq 0 \tag{118}$$

against every pursuer input. Integration through s_0 bounds its cost above by $F(s_0,z_0)$. The maximizing pursuer selector gives the reverse inequality against every surviving evader strategy. Non-survival is assigned infinite cost. The opposing bounds establish optimality among guaranteed winning strategies. □

This auxiliary selector need not maximize W or V. Once the evader can win, its own fuel preference has replaced delay maximization. Simply adding a small fuel penalty to the pursuer's payoff does not in general reproduce this lexicographic rule.

B. Perpetual Escape

For A1/B1, define the stronger class

$$\mathfrak{E}_\infty(z) = \{\beta : \tau = +\infty \text{ for every } \mathbf{a}_P\}, \tag{119}$$

and its cost

$$F_\infty(z) = \inf_{\beta \in \mathfrak{E}_\infty(z)} \sup_{\mathbf{a}_P} \int_0^\infty \|\mathbf{a}_E(t)\|^2 \,\mathrm{d}t. \tag{120}$$

An empty strategy class again gives $+\infty$. On a regular region a stationary candidate satisfies

$$\mathcal{H}_E[F_\infty] = 0. \tag{121}$$

This equation requires a perpetual-survival viability condition and a condition at infinity. A sufficient verification framework requires the selected evader to remain safe forever and, for the upper bound, $F_\infty \geq 0$; for a matching lower bound it also requires the residual certificate along the maximizing response to every competing finite-cost winning strategy to tend to zero. Integrating the preceding differential inequalities then proves equality of total infinite-horizon costs. Without such conditions a stationary solution can select a trajectory that is eventually captured.

In particular, $V(z) = +\infty$ means that the pursuer has no finite *uniform* bound on capture time. It does not alone prove the existence of one evader strategy that avoids capture forever. Nor does

$$\text{for every } s \text{ there exists } \beta_s \in \mathfrak{E}_s(z) \tag{122}$$

automatically imply $\mathfrak{E}_\infty(z) \neq \varnothing$. The infinite-horizon winning set must be established through the actual viability game. Under suitable compactness and closed-safety hypotheses, viability constructions can justify horizon limits; strict avoidance of a closed capture target needs additional care. We do not identify these logically different conditions without proof.

C. An Explicit Nonattainment Counterexample

Nonattainment already occurs with unrestricted planar or spatial steering. Let the exhausted pursuer be at rest at the origin, and let the evader start from $(-L, R)$ with velocity $(u, 0)$, where $L, u > 0$. The unused third coordinate can be set to zero. Coasting grazes the closed capture ball at time $t_* = L/u$ and therefore loses. Fix $0 < h < t_*$ and apply lateral acceleration $(0, \varepsilon)$ for $0 \leq t \leq h$, then coast. The second coordinate is

$$x_{E,2}(t) = R + \begin{cases} \varepsilon t^2/2, & 0 \leq t \leq h, \\ \varepsilon(ht - h^2/2), & t \geq h. \end{cases} \tag{123}$$

For every $\varepsilon > 0$ it exceeds R at every positive time, so the evader escapes forever. The total acceleration expenditure is $\varepsilon^2 h$, tending to zero. Zero expenditure forces zero acceleration almost everywhere and hence the losing coasting path. Thus the escape-cost infimum is zero and is not attained, for every $R \geq 0$, in both $d = 2$ and $d = 3$. The same argument applies to deadline survival for $T \geq t_*$. Any positive available evader budget permits a sufficiently small ε.

This fully multidimensional example suffices to disprove universal existence of a least-fuel winning trajectory in the stated problem. With on/off engines and a fixed admissible positive lateral acceleration, arbitrarily short initial pulses give the same zero-cost limit, so nonattainment is not solely an artifact of an unbounded exhaust-speed ceiling. We next give a one-dimensional example with a positive, explicitly calculable infimum; its restriction to one-dimensional steering is intentional.

Consider a one-dimensional free-space game. The pursuer has exhausted its fuel and is stationary at $x_P = 0$. Let $x_E(0) = R + L$, $\dot{x}_E(0) = -u$, with $L, u > 0$, and define clearance $y = x_E - R$. Capture occurs as soon as $y \leq 0$. Assume the evader's available acceleration budget exceeds

$$C_* = \frac{4u^3}{9L}. \tag{124}$$

This is an admissible boundary-state instance of the two-rocket model, and works also for a positive capture radius.

Theorem 5 (Winning escape need not have a least-fuel trajectory)**.** *In this instance perpetual escape is possible. Its acceleration-expenditure infimum is C_*, but every escaping trajectory spends strictly more than C_*. Hence no least-propellant escaping trajectory exists. The same nonattainment holds for deadline survival whenever $T \geq 3L/u$.*

Proof. Let $t_* = 3L/u$. For any trajectory surviving through t_*,

$$0 < y(t_*) = L - ut_* + \int_0^{t_*} (t_* - t)a_E(t)\mathrm{d}t \tag{125}$$
$$= -2L + \int_0^{t_*} (t_* - t)a_E(t)\mathrm{d}t.$$

Cauchy–Schwarz therefore gives

$$2L < \left(\frac{t_*^3}{3}\right)^{1/2} \left(\int_0^{t_*} a_E(t)^2\mathrm{d}t\right)^{1/2}, \tag{126}$$

so the expenditure is strictly greater than

$$\frac{12L^2}{t_*^3} = \frac{4u^3}{9L} = C_*. \tag{127}$$

For any $0 < \delta < L$, set $t_\delta = 3(L-\delta)/u$ and use

$$y_\delta(t) = \delta + (L-\delta)(1 - t/t_\delta)^3, \quad 0 \leq t \leq t_\delta, \tag{128}$$

$$a_{E,\delta}(t) = \frac{2u}{t_\delta}(1 - t/t_\delta). \tag{129}$$

Thereafter coast at rest with $y_\delta = \delta$. This trajectory remains strictly outside capture forever and spends

$$C_\delta = \frac{4u^2}{3t_\delta} = \frac{4u^3}{9(L-\delta)} \longrightarrow C_* \quad \text{as } \delta \downarrow 0. \tag{130}$$

For sufficiently small positive δ, it fits any budget $B_E > C_*$. Equality $\delta = 0$ reaches the capture boundary at t_* and loses. Thus the infimum is sharp and unattained. The argument through t_* also applies to every deadline $T \geq t_*$. Strict monotonicity of (7) transfers the conclusion to propellant mass. □

For completeness, on $y > 0$ and relative velocity $w < 0$, the smooth expression

$$F_*(y, w) = \frac{4(-w)^3}{9y} \tag{131}$$

satisfies the formal stationary interior equation

$$w(F_*)_y - \frac{(F_*)_w^2}{4} = 0. \tag{132}$$

Its minimizing selector is $a_E = -(F_*)_w/2 = 2w^2/(3y)$. Starting from $(L, -u)$, that selector is precisely the limiting $\delta = 0$ profile, which is captured. This is a concrete demonstration that the stationary PDE alone does not supply the missing safety and attainment conditions.

The family (128) is an explicit ε-optimal solution of the stated preference. If a minimum safety clearance is prescribed as a closed constraint, the corresponding constrained problem is different and may have an attained minimum. One must not silently replace strict escape by that modified objective.

VIII. Ordinary Differential Equations for Trajectory Reconstruction

The PDEs determine globally meaningful feedback values. Their characteristics provide ordinary differential equations suitable for indirect numerical methods and for reconstructing smooth saddle trajectories. Satisfying these ODEs alone gives candidates, just as PMP shooting in the companion transfer problem gives candidates unless a sufficiency argument is supplied [6], [1].

Introduce position costates $\boldsymbol{\lambda}_{x_i}$, velocity costates $\boldsymbol{\lambda}_{v_i}$, and budget costates η_i. We use the dynamic-programming *minimization* convention; its costates have the opposite sign to a single-player maximization convention with Hamiltonian $\lambda \cdot f - L$ used in the companion article. The unoptimized characteristic Hamiltonian is

$$K = \sigma + \ell_P \left\|\mathbf{a}_P\right\|^2 + \ell_E \left\|\mathbf{a}_E\right\|^2 + \sum_{i=P,E} \left[\boldsymbol{\lambda}_{x_i} \cdot \mathbf{v}_i + \boldsymbol{\lambda}_{v_i} \cdot (\mathbf{g}(\mathbf{x}_i) + \mathbf{a}_i) - \eta_i \left\|\mathbf{a}_i\right\|^2\right]. \tag{133}$$

Use $(\sigma, \ell_P, \ell_E) = (1,0,0)$ for capture time, $(0,1,0)$ for pursuer fuel, and $(0,0,1)$ for evader fuel. The first two optimize by pursuer-min/evader-max; the last reverses those roles.

On uncapped interior arcs the complete state–costate system is

$$\dot{\mathbf{x}}_i = \mathbf{v}_i, \tag{134a}$$
$$\dot{\mathbf{v}}_i = \mathbf{g}(\mathbf{x}_i) + \mathbf{a}_i^*, \tag{134b}$$
$$\dot{b}_i = -\left\|\mathbf{a}_i^*\right\|^2, \tag{134c}$$
$$\dot{\boldsymbol{\lambda}}_{x_i} = -D\mathbf{g}(\mathbf{x}_i)^\top \boldsymbol{\lambda}_{v_i}, \tag{134d}$$
$$\dot{\boldsymbol{\lambda}}_{v_i} = -\boldsymbol{\lambda}_{x_i}, \tag{134e}$$
$$\dot{\eta}_i = 0. \tag{134f}$$

For the pursuer objectives,

$$\mathbf{a}_P^* = -\frac{\boldsymbol{\lambda}_{v_P}}{2(\ell_P - \eta_P)}, \qquad \mathbf{a}_E^* = \frac{\boldsymbol{\lambda}_{v_E}}{2\eta_E}, \tag{135}$$

on the branches $\ell_P - \eta_P > 0$, $\eta_E > 0$. For the escape-cost game,

$$\mathbf{a}_P^\dagger = \frac{\boldsymbol{\lambda}_{v_P}}{2\eta_P}, \qquad \mathbf{a}_E^\dagger = -\frac{\boldsymbol{\lambda}_{v_E}}{2(1-\eta_E)}, \tag{136}$$

on its corresponding branches. If necessary add $\dot{s} = -1$ and a quadrature for the selected running cost. Then recover $m_i, v_{ei}, I_{sp,i}$ using (6) and (1).

A. Free-Space Characteristics

In the relative formulation let $\mathbf{p}_r, \mathbf{p}_w$ be the kinematic costates. Then

$$\dot{\mathbf{p}}_r = 0, \qquad \dot{\mathbf{p}}_w = -\mathbf{p}_r, \qquad \dot{\eta}_P = \dot{\eta}_E = 0. \tag{137}$$

On regular arcs, $\mathbf{p}_w = \mathbf{A} - \mathbf{B}t$ and

$$\mathbf{a}_P^* = \frac{\mathbf{p}_w}{2(\ell_P - \eta_P)}, \qquad \mathbf{a}_E^* = \frac{\mathbf{p}_w}{2\eta_E}. \tag{138}$$

Hence both accelerations are affine, velocities are quadratic, positions are cubic, and budget depletion is cubic on such

an arc. This extends the affine-acceleration structure in the companion article, but only to a *smooth interior saddle arc*. Exhaustion, finite engine bounds, capture boundaries, and changes of active value-function branch can interrupt it. A precomputed pair of cubic trajectories is not a feedback guarantee against deviations.

B. Kepler Characteristics

For $r_i = \|\mathbf{x}_i\|$,

$$D\mathbf{g}(\mathbf{x}_i) = -\mu\left(\frac{I}{r_i^3} - \frac{3\mathbf{x}_i\mathbf{x}_i^\top}{r_i^5}\right), \tag{139}$$

so the costate equations become

$$\dot{\boldsymbol{\lambda}}_{x_i} = \mu\left(\frac{I}{r_i^3} - \frac{3\mathbf{x}_i\mathbf{x}_i^\top}{r_i^5}\right)\boldsymbol{\lambda}_{v_i}, \tag{140}$$

$$\dot{\boldsymbol{\lambda}}_{v_i} = -\boldsymbol{\lambda}_{x_i}. \tag{141}$$

Equivalently,

$$\ddot{\boldsymbol{\lambda}}_{v_i} = D\mathbf{g}(\mathbf{x}_i)^\top\boldsymbol{\lambda}_{v_i}. \tag{142}$$

These Cartesian equations avoid polar-coordinate singularities at a vanishing tangential velocity and apply equally to planar and spatial motion away from the central body.

There are useful rotational first integrals on smooth uncapped characteristic arcs:

$$\mathbf{J}_i = \mathbf{x}_i \times \boldsymbol{\lambda}_{x_i} + \mathbf{v}_i \times \boldsymbol{\lambda}_{v_i} = \text{constant}. \tag{143}$$

Differentiation cancels the velocity terms, the central-gravity terms cancel by the displayed form of $D\mathbf{g}$, and $\mathbf{a}_i \times \boldsymbol{\lambda}_{v_i} = 0$ for the regular selectors. In planar motion the cross product is scalar. These are costate integrals, not the physical orbital angular momenta of the powered ships. Simultaneous rotational invariance of a differentiable game value gives $\mathbf{J}_P + \mathbf{J}_E = 0$ along its characteristics. They provide algebraic checks for orbital trajectory reconstruction.

C. Terminal Conditions and Their Limits

For a regular capture endpoint on $\|\mathbf{x}_E - \mathbf{x}_P\| = R > 0$, let $\mathbf{n} = (\mathbf{x}_E - \mathbf{x}_P)/R$. With no terminal state cost and no other active terminal constraints, transversality gives

$$\begin{aligned} \boldsymbol{\lambda}_{x_P}(\tau) &= -\nu\mathbf{n}, & \boldsymbol{\lambda}_{x_E}(\tau) &= \nu\mathbf{n}, \\ \boldsymbol{\lambda}_{v_P}(\tau) &= 0, & \boldsymbol{\lambda}_{v_E}(\tau) &= 0. \end{aligned} \tag{144}$$

For $R = 0$, use a vector multiplier for $\mathbf{x}_E - \mathbf{x}_P = 0$ instead. Budget endpoint constraints and central-distance constraints introduce their own multipliers; inactive free budget endpoints give $\eta_i(\tau) = 0$. These statements concern regular endpoint branches, not singular value gradients at a target.

For exact rendezvous, both endpoint differences are constrained. With vector multipliers $\boldsymbol{\nu}$ and $\boldsymbol{\zeta}$, the corresponding conditions are

$$\begin{aligned} \boldsymbol{\lambda}_{x_P}(\tau_J) &= -\boldsymbol{\nu}, & \boldsymbol{\lambda}_{x_E}(\tau_J) &= \boldsymbol{\nu}, \\ \boldsymbol{\lambda}_{v_P}(\tau_J) &= -\boldsymbol{\zeta}, & \boldsymbol{\lambda}_{v_E}(\tau_J) &= \boldsymbol{\zeta}. \end{aligned} \tag{145}$$

The common terminal position and velocity are free, but their differences are zero. Setting both terminal velocity costates to zero would instead solve the position-only problem. The same rendezvous transversality applies in free space and in the Kepler field, subject to any active state constraints.

For free terminal time in the time game, $K = 0$ along a regular autonomous saddle extremal. For fuel minimization, an interior capture time $0 < \tau < T$ also has the free-time transversality condition; an active deadline has a time-constraint multiplier and does not permit imposing that condition unaltered. One must allow capture before T.

Notice that zero terminal velocity costates and free terminal budgets can make the strictly coercive formulas degenerate. In particular, one cannot insist that an entirely interior uncapped arc with constant nonzero budget costates extends smoothly to every unconstrained terminal state. Budget-active or singular terminal arcs are often essential. At budget exhaustion and other state constraints, use constrained optimal-control conditions and the reduced face dynamics. Ordinary shooting across such boundaries without the correct multipliers is not justified.

IX. Finite Engine Bounds and Switching

For a compact admissible radius set

$$S_i(b_i) = \{0\} \cup [a_{i,\min}(b_i), a_{i,\max}(b_i)], \tag{146}$$

pointwise vector optimization reduces to a scalar problem. For

$$Q(\mathbf{a}) = \mathbf{h}\cdot\mathbf{a} + c\|\mathbf{a}\|^2, \tag{147}$$

the minimizing powered direction is $-\mathbf{h}/\|\mathbf{h}\|$ and the maximizing direction is $+\mathbf{h}/\|\mathbf{h}\|$. Thus

$$\min_{\mathbf{a}\in\mathcal{U}_i} Q = \min_{r\in S_i}[-\|\mathbf{h}\| r + cr^2], \tag{148}$$

$$\max_{\mathbf{a}\in\mathcal{U}_i} Q = \max_{r\in S_i}[+\|\mathbf{h}\| r + cr^2]. \tag{149}$$

At $\mathbf{h} = 0$ direction is immaterial. For the minimizing quadratic with $c > 0$, compare zero with the interval-clipped radius $\|\mathbf{h}\|/(2c)$. For the maximizing quadratic with $c < 0$, compare zero with the interval-clipped radius $-\|\mathbf{h}\|/(2c)$. In all other sign cases, comparing the interval endpoints and zero is sufficient. These rules solve the pointwise optimization globally, including coast and boundary arcs.

Insert the resulting extrema in (21) or (113). For a mass-independent ball bound $\|\mathbf{a}\| \le A$, a coercive selector is simply radially clipped at A. An exhaust-speed ceiling alone retains the gap between coast and minimum powered acceleration described in the companion article.

With mass-dependent control boundaries, (134) must be replaced by the canonical equations of the *optimized constrained Hamiltonian*. In particular, $\dot{\eta}_i = -\partial K^*/\partial b_i$ can be nonzero on exhaust-boundary arcs. The constant-budget-costate result is for unconstrained interior arcs, not a universal first integral of the bounded game. This agrees with the finite-exhaust appendix of the companion article [1]: on a fixed-exhaust burn $\dot{\lambda}_m = (2P/V)|\lambda_v|/m^2$, and $C_m = \lambda_m m^2$ varies. Its proved free-space switching sequence is ceiling-only; adding a lower exhaust limit introduces maximum-acceleration arcs whose complete topology is left open there.

Rapid switching or relaxed controls may improve existence properties, but relaxation must retain the pair $(\mathbf{a}, -\|\mathbf{a}\|^2)$

when taking convex combinations. Replacing an averaged acceleration by its square would generally give the wrong propellant flow. If a relaxed optimum exists, additional work is needed to show exact realization by an ordinary control rather than approximation through fast switching.

X. Numerically Specified Global Solution

A. Exact Winning Sets and Resource Thresholds

Define the robust capture set at horizon s by

$$\mathcal{K}_s = \{z : \exists \alpha \ \forall \mathbf{a}_E, \tau \le s\}. \tag{150}$$

Then, directly from (16),

$$\boxed{V(z) = \inf\{s \ge 0 : z \in \mathcal{K}_s\}.} \tag{151}$$

The infimum need not itself be a feasible horizon. Thus $V(z) < T$ implies feasibility by T, whereas the equality $V(z) = T$ requires attainment or a direct membership test in $\mathcal{K}_T$.

For rendezvous, define $\mathcal{K}_s^J$ by replacing τ with τ_J in (150); then $V^J = \inf\{s : z \in \mathcal{K}_s^J\}$. Theorem 1 proves attainment of every finite ideal free-space rendezvous value. General orbital or bounded-engine rendezvous needs the target-specific reachability construction.

For the ideal model, or bounds independent of the artificial fuel ledger, A2/B2 can be calculated without storing an infinite terminal fuel cost. Keep the actual initial state and impose an additional pathwise pursuer expenditure cap c. Let $\mathcal{K}_T(c)$ be its robust capture set. Then

$$\boxed{W(T, z) = \inf\{c \in [0, b_P] : z \in \mathcal{K}_T(c)\}.} \tag{152}$$

An empty set gives $+\infty$. Proof: a strategy guarantees a worst-case expenditure at most c if and only if it guarantees capture with that additional cap. Taking the infimum gives (152). For an ideal engine, the artificial cap can replace the pursuer budget in the kinematic reachability calculation. For exhaust-speed limits that depend on actual mass, retain the true b_P and add a separate ledger q with

$$\dot{q} = -\left\|\mathbf{a}_P\right\|^2, \qquad q(0) = c, \qquad q \ge 0. \tag{153}$$

Changing a fuel allowance must not inadvertently change the spacecraft's initial mass and thrust bounds.

The evader cost has the analogous exact construction: minimize a pathwise expenditure cap over strategies that survive to T, or survive forever. For perpetual escape the quantifiers are $\exists \beta \, \forall \mathbf{a}_P \, \forall t \ge 0$, with one common strategy. Threshold values give infima. Theorem 5 proves that the boundary cap itself may fail to permit a winning strategy.

B. A Continuous Reachability PDE

For a positive-radius target, let $\varphi(z) = \|\mathbf{r}\| - R$; it may be smoothly truncated away from the target on a numerical domain. A useful finite-horizon game payoff is the smallest clearance attained on a path:

$$L(s, z) = \inf_{\alpha} \sup_{\mathbf{a}_E} \min_{0 \le t \le s} \varphi(z(t)). \tag{154}$$

With budgets included in the state, its dynamic-programming obstacle equation is

$$\boxed{\max\{L - \varphi, \ L_s - \mathcal{H}_0[L]\} = 0, \qquad L(0, z) = \varphi(z).} \tag{155}$$

The obstacle records capture at *any* earlier time, rather than only endpoint proximity. Continuous reachability and reach-avoid HJI formulations are developed in [7], [8]. State and resource constraints must be imposed consistently when using this equation.

Under a value/strategy existence theory, $L < 0$ supplies a robust capture certificate with interior penetration, and $L > 0$ supplies the opposite strict-margin certificate through the dual game. At $L = 0$, exact capture requires attention to attainment and target regularity: an infimum of miss distances equal to zero is not automatically an attained interception. The exact set (150) remains the definition. For $R = 0$, $\varphi \ge 0$, so there is no negative interior level; use the exact target game or a justified target-limit construction. Simply taking a small numerical miss distance as a proof of point capture is invalid.

Similarly, closed safety sets with a prescribed margin can be propagated by viability methods. Their relation to the original strict escape condition, especially on an infinite horizon, must be verified rather than assumed.

For exact rendezvous, one possible terminal function is

$$\varphi_J(z) = \max\left\{\frac{\|\mathbf{x}_E - \mathbf{x}_P\|}{L_*}, \frac{\|\mathbf{v}_E - \mathbf{v}_P\|}{V_*}\right\}, \tag{156}$$

where $L_*, V_* > 0$ only set units. Its zero set is $\mathcal{C}_J$. Equation (155) then describes the smallest joint mismatch, but there is no negative interior level. A zero infimum or two small residuals alone do not prove attained exact rendezvous. Positive tolerances in both position and velocity define a different, thickened target; convergence to the exact target requires a separate argument.

C. Differential Equations and Boundary Data to Supply

A numerical mathematical specification is as follows:

1) Choose P_i, m_{i0}, m_{id}, the initial positions and velocities, R, and, when relevant, T and μ. Specify ideal or bounded engines, observation convention, and the central-body boundary rule.
2) Compute B_i by (2). Use the relative state (28) in free space, optionally reduced by (69); use (10) in the Kepler field.
3) Obtain the winning domain with (150) and an HJI/viability construction such as (155). Treat $b_i = 0$ by the exact ballistic face game. Account for outer-domain boundaries rather than setting arbitrary terminal values there.
4) For minimum-time interception, use (30) or (75), or find the threshold in (151). For ideal free-space rendezvous, use the cubic (47) and its explicit strategies. For Kepler rendezvous, use (78) and $\mathcal{K}_s^J$. For A2/B2, use (65) or (103), or search the monotone fuel threshold (152).
5) On an evader-winning state, solve (114) or the perpetual-survival problem (120). Check whether the resulting

infimum is attained; otherwise construct a winning ε-optimal strategy.

6) Reconstruct trajectories using the value-gradient feedback and state ODEs, or use (134) as characteristic equations with proper switching and terminal conditions. Evaluate controls from the actual state when the opponent deviates.

This specifies the mathematical inputs to a numerical method without implementing one. A convergent monotone, consistent and stable HJI discretization needs a comparison principle for the chosen boundary problem; convergence results for pursuit–evasion discretizations require corresponding regularity assumptions [9]. A computed grid function is not by itself an exact global proof. Certified upper and lower value bounds or verified differential inequalities, together with capture feasibility, provide a stronger certificate. Shooting only one state–costate boundary-value problem cannot supply the global adversarial guarantee.

D. Numerical Limits That Must Be Kept Separate

Increasing only the pursuer's admissible control set can improve its capture value; increasing only the evader's can worsen it. Enlarging both sets simultaneously need not produce a monotone sequence of game values. Therefore "solve with larger and larger common acceleration bounds" is not, by itself, a proof of the ideal unbounded game.

Likewise, a finite-horizon safety calculation cannot establish perpetual escape without a valid limiting strategy, and an ε-capture policy need not attain exact point capture. These are mathematical limitations of limit interchange, not missing ODE terms. They are especially relevant at resource exhaustion, grazing capture, and the strict-survival boundary demonstrated above.

XI. Analytic Free-Space Checks and Useful Bounds

A. An Exhausted, Ballistic Evader

Suppose $b_E = 0$, so the evader cannot thrust. For a candidate interception time $h > 0$, define

$$\mathbf{q}(h) = \mathbf{r}_0 + \mathbf{w}_0 h. \tag{157}$$

$$\mathbf{d}(h) = \begin{cases} (1 - R/\left\|\mathbf{q}(h)\right\|)\mathbf{q}(h), & \left\|\mathbf{q}(h)\right\| > R, \\ \mathbf{0}, & \left\|\mathbf{q}(h)\right\| \le R. \end{cases} \tag{158}$$

The exact minimum acceleration expenditure to be in the capture ball at time h is

$$\boxed{c(h) = \frac{3\left\|\mathbf{d}(h)\right\|^2}{h^3}.} \tag{159}$$

Indeed, the pursuer must supply a weighted acceleration integral whose norm is at least $\|\mathbf{d}(h)\|$, and

$$\left\| \int_0^h (h-t)\mathbf{a}_P(t)\mathrm{d}t \right\|^2 \le \frac{h^3}{3} \int_0^h \left\|\mathbf{a}_P(t)\right\|^2 \mathrm{d}t. \tag{160}$$

Equality is attained by

$$\boxed{\mathbf{a}_P(t) = \frac{3(h-t)}{h^3}\mathbf{d}(h), \quad 0 \le t \le h.} \tag{161}$$

Consequently, in this special case,

$$V_A = \inf\{h > 0 : c(h) \le b_P\}, \tag{162}$$

$$W_A(T) = \inf_{0<h\le T} c(h). \tag{163}$$

The second equality requires that this infimum be feasible within b_P. Otherwise $W_A = +\infty$. An initially captured state has value zero. If a constructed endpoint path captures earlier, terminate it then. Conversely every first-capture path obeys the same lower bound at its own capture time. These observations justify the first-hit interpretation of (162)–(163), not just the endpoint problem.

For point interception ($R = 0$), put $R_2 = \|\mathbf{r}_0\|^2$, $Q = \mathbf{r}_0 \cdot \mathbf{w}_0$, and $W_2 = \|\mathbf{w}_0\|^2$. Then

$$\begin{aligned} c(h) &= 3(R_2 + 2Qh + W_2h^2)/h^3, \\ c'(h) &= -\frac{3}{h^4}(3R_2 + 4Qh + W_2h^2). \end{aligned} \tag{164}$$

For an initially separated state, the earliest feasible time is the smallest positive root of

$$b_P h^3 - 3W_2h^2 - 6Qh - 3R_2 = 0, \tag{165}$$

if such a root exists. When $b_P > 0$ at least one exists. For the deadline-fuel value it is sufficient to compare $c(T)$ with $c(h_-)$ and $c(h_+)$ for the real candidates

$$h_\pm = \frac{-2Q \pm \sqrt{4Q^2 - 3R_2W_2}}{W_2} \tag{166}$$

that lie in $(0, T]$. If $W_2 = 0$, only T is needed. The minimum is the exact fuel value when it is at most b_P; otherwise capture is infeasible. This finite comparison also covers a zero-cost ballistic collision. Initial approach can make $c(h)$ nonmonotone, so simply choosing the latest time can be suboptimal.

For zero relative velocity and initial excess separation $D = \|\mathbf{r}_0\| - R > 0$,

$$\boxed{V_A = (3D^2/b_P)^{1/3}, \qquad W_A(T) = 3D^2/T^3,} \tag{167}$$

provided the relevant budget is available. In dimensional propulsion parameters the time is

$$V_A = \left[\frac{3D^2}{2P_P(m_{Pd}^{-1} - m_{P0}^{-1})}\right]^{1/3}. \tag{168}$$

The companion rest-to-rest transfer has expenditure $12D^2/T^3$ instead of $3D^2/T^3$. The factor of four arises because interception here does not require matching terminal velocity.

B. An Exact Interception Game With Two Powered Ships

Theorem 6 (Equal-velocity interception with a resource margin)**.** *Suppose $\mathbf{w}_0 = 0$, $D = \|\mathbf{r}_0\| - R > 0$, $b_P > b_E$, and $b_P \ge 4b_E$. Assume ideal uncapped controls and instantaneous*

observation of the evader's current acceleration by the pursuer. Then,

$$\boxed{V_A = H_I = \left[\frac{3D^2}{(\sqrt{b_P}-\sqrt{b_E})^2}\right]^{1/3}}. \tag{169}$$

Put $k = \sqrt{b_E}/(\sqrt{b_P}-\sqrt{b_E}) \le 1$, $\mathbf{n} = \mathbf{r}_0/\|\mathbf{r}_0\|$, *and*

$$\mathbf{a}_0(t) = \frac{3D}{H_I^2}(1 - t/H_I)\mathbf{n}. \tag{170}$$

The response $\mathbf{a}_P = \mathbf{a}_E + \mathbf{a}_0$ *guarantees capture by* H_I. *The evader control* $\mathbf{a}_E = k\mathbf{a}_0$ *on* $[0, H_I]$, *followed by coasting if needed, prevents capture before* H_I. *Their saddle controls are* $(1+k)\mathbf{a}_0$ *and* $k\mathbf{a}_0$.

Proof. The correction has squared norm $C = 3D^2/H_I^3$ and reduces the relative separation to R at H_I. Minkowski's inequality proves the upper bound since $(1+k)^2 C = b_P$.

For the lower bound fix the stated evader control. At $t = sH_I$, $0 < s < 1$, its powered displacement is $kDf(s)\mathbf{n}$, where $f(s) = (3s^2 - s^3)/2$. To enter the radius-R ball at that time, any pursuer must spend at least

$$C\, s^{-3}[1 + kf(s)]^2. \tag{171}$$

This is larger than $b_P = (1+k)^2 C$, because

$$1 + kf(s) - (1+k)s^{3/2} = (1-k)(1-s^{3/2}) + k[1 + f(s) - 2s^{3/2}] > 0. \tag{172}$$

Indeed $f(s) \ge s^3$ and $1 + s^3 - 2s^{3/2} = (1 - s^{3/2})^2 > 0$. The evader uses exactly b_E by H_I and is admissible. These opposing inequalities prove global optimality. □

Here $b_P \ge 4b_E$ is a sufficient condition for this particular explicit interception saddle. No claim of necessity is made. With $R = 0$, comparison with (54) gives

$$\frac{V_A^J}{V_A} = 4^{1/3} \qquad (\mathbf{w}_0 = 0,\ b_P \ge 4b_E,\ b_P > b_E). \tag{173}$$

The same resource gap produces a longer boarding time because the relative velocity must also vanish. For the interception saddle, replace E_H in (62) by

$$E_I(t) = C(3s - 3s^2 + s^3), \qquad s = t/H_I, \tag{174}$$

and use $\gamma_i \|\mathbf{a}_0(t)\|$ as the acceleration in the exhaust formula. Thus positions, velocities, masses, and specific impulses are all elementary functions of time.

C. A Certified Zero-Fuel Evader Win by a Deadline

Even if the evader has fuel available, it can elect to coast. Compute (159) for that coasting motion. If

$$c(h) > b_P \qquad \text{for every } 0 < h \le T, \tag{175}$$

then no pursuer control can intercept the coasting evader by T. Thus coasting is a guaranteed evader win with exactly zero fuel and is globally fuel-optimal. Conversely, if some $h \le T$ has $c(h) \le b_P$, the coasting strategy is not a guaranteed win, although a powered evasion strategy may still be one. This is a directly usable analytical certificate rather than an engine-superiority heuristic.

D. A Free-Space Capture Bound Under Ideal Nonanticipative Response

The following sufficient bound makes its information assumption explicit. Suppose the nonanticipative pursuer may use the evader's current acceleration, equivalently its almost-everywhere reconstruction from exact past velocity observations. For a prescribed time h, let

$$\mathbf{a}_0(t) = \frac{3(h-t)}{h^3}(\mathbf{r}_0 + \mathbf{w}_0 h). \tag{176}$$

The nonanticipative response

$$\mathbf{a}_P(t) = \mathbf{a}_E(t) + \mathbf{a}_0(t) \tag{177}$$

forces $\mathbf{r}(h) = 0$ for every admissible evader input. Minkowski's inequality gives

$$\left(\int_0^h \|\mathbf{a}_P\|^2 \,\mathrm{d}t\right)^{1/2} \le \sqrt{b_E} + \frac{\sqrt{3}\,\|\mathbf{r}_0 + \mathbf{w}_0 h\|}{h^{3/2}}. \tag{178}$$

Therefore a sufficient capture-by-h condition is

$$\boxed{\sqrt{b_P} \ge \sqrt{b_E} + \frac{\sqrt{3}\,\|\mathbf{r}_0 + \mathbf{w}_0 h\|}{h^{3/2}}.} \tag{179}$$

It respects the budget pathwise; if capture occurs earlier, the response is stopped. Since the second term tends to zero as $h \to \infty$, $b_P > b_E$ suffices for some finite guaranteed interception time in this ideal response model. If $\mathbf{w}_0 = 0$, the resulting upper bound is

$$V_A \le \left[\frac{3\,\|\mathbf{r}_0\|^2}{(\sqrt{b_P}-\sqrt{b_E})^2}\right]^{1/3}. \tag{180}$$

Theorem 6 proves equality in the stated equal-velocity resource-margin regime and replaces the distance by its excess over R. Outside that regime, (179) is only a sufficient bound. The strategy uses the stated exact nonanticipative information convention; delayed measurements or restricted acceleration sensing require a different guarantee. It also need not satisfy finite instantaneous acceleration or exhaust-speed bounds. In a Kepler field, the additional unequal-gravity cost must be included, as in Section V-C.

E. Exact Prescribed-Time Endpoint Expenditure

Under instantaneous observation of the evader's current acceleration, an auxiliary endpoint game gives a useful explicit fuel guarantee. Prescribe $h > 0$ and continue the dynamics to h even if the target was met earlier. Require either exact position coincidence or exact rendezvous at h. Let $C(h)$ be, respectively,

$$C_I(h) = \frac{3\,\|\mathbf{r}_0 + h\mathbf{w}_0\|^2}{h^3} \quad \text{or} \quad C_J(h). \tag{181}$$

With an evader expenditure cap b_E, the minimum worst-case pursuer expenditure is

$$\boxed{M(h) = \left(\sqrt{b_E} + \sqrt{C(h)}\right)^2.} \tag{182}$$

This statement presumes the pursuer has at least that budget; otherwise the robust endpoint requirement is infeasible.

For the upper bound, use the minimum-norm correction $\mathbf{a}_h$ and $\mathbf{a}_P = \mathbf{a}_E + \mathbf{a}_h$. For the lower bound when $C(h) > 0$, let $\mathbf{a}_E = \sqrt{b_E/C(h)}\,\mathbf{a}_h$. The endpoint moments force the orthogonal projection of every feasible pursuer control onto the moment subspace to be $(1 + \sqrt{b_E/C(h)})\mathbf{a}_h$. Its squared norm is (182). When $C(h) = 0$, choose any evader control of squared norm b_E in that subspace; matching its moments forces at least the same expenditure. The case $b_E = 0$ follows directly. This proves both bounds against arbitrary opposing controls.

Consequently, the first-interception deadline value satisfies

$$W_A(T) \le \inf_{0<h\le T} \left(\sqrt{b_E} + \sqrt{C_I(h)}\right)^2 \tag{183}$$

whenever a minimizing endpoint candidate fits the pursuer's reserve. For an initially separated point target, its candidates are exactly T and the admissible times (166), because the displayed objective increases with C_I. Positive interception radius can only improve this upper bound. First-hit stopping can reduce expenditure, so the endpoint equality must not be asserted as an equality for the general deadline game.

F. A Numerical Ballistic Check and the First-Hit Distinction

Take a one-dimensional point target with $r_0 = 1000$ m, $w_0 = -10$ m/s, and $b_E = 0$. The exact endpoint requirement costs

$$c(h) = \frac{3(1000 - 10h)^2}{h^3} \quad [\mathrm{m}^2/\mathrm{s}^3]. \tag{184}$$

At $h = 100$ s, the two coasting paths meet and $c(100) = 0$. Thus $W_A(200) = 0$: the first-hit game terminates at 100 s with zero fuel. If instead coincidence is imposed at exactly 200 s and motion continues through earlier meetings, $c(200) = 0.375$ and the endpoint minimax is $M(200) = (\sqrt{0} + \sqrt{0.375})^2 = 0.375$. For $P_P = 1$ MW and $m_{P0} = 1000$ kg, this expenditure corresponds to 0.187465 kg of propellant by (7); coasting still costs zero.

For comparison, with the same initial state but an active evader allowed $b_E = 0.04$, the fixed-time endpoint minimax becomes $M(200) = (0.2 + \sqrt{0.375})^2 = 0.659949$ in the same units, provided the pursuer has that reserve. Against this active evader zero-cost first-hit capture no longer follows; the distinction is between the first meeting and an enforced endpoint at the deadline.

XII. Conclusion

The variable-specific-impulse chase is a differential game in kinematic state and two remaining acceleration budgets. Free-space and Kepler dynamics enter the same separable Isaacs construction. Minimum guaranteed time uses a unit running cost and an absorbing capture target; minimum fuel by a deadline uses squared pursuer acceleration, zero cost at first capture, and infeasibility outside capture at the deadline. The evader's preference for least-fuel winning escape requires a separate survival game with the optimization roles reversed.

Exact position-and-velocity rendezvous has a complete analytical solution in ideal free space under instantaneous response: the pursuer wins precisely when its remaining acceleration budget exceeds the evader's, and its optimal time is the unique positive root of an explicit cubic. A resource-parity escape feedback and a moment-projection lower bound prove optimality of both strategies. The least worst-case fuel cost of winning perpetual rendezvous avoidance is attained and equals the pursuer's remaining acceleration budget in the transformed units. Explicit active-interception and ballistic families, together with prescribed-time endpoint minimax formulas, supply further closed-form results.

For general orbital interception and rendezvous, the HJI boundary problems, feedback selectors, and state–costate ODEs specify the numerical mathematical solution. The planar CW game gives an exact local linear value and an opposing parity strategy. Its explicit nonlinear tube and the inertial half-space bound provide complementary sufficient Kepler capture certificates. The verification theorems establish global optimality when feasibility, admissibility, regularity, and boundary conditions are certified; shooting extremals alone do not. Strict position-only escape can have an unattained fuel infimum, in which case winning sets, infimal costs, and winning ε-optimal strategies are the appropriate solution objects.

AI-assistance acknowledgment

AI assistance was used for mathematical discussion, derivations, code development and editing.

References

[1] J. Olšina, "Time-optimal heliocentric transfers with a constant-power, variable-i_{sp} engine," 2025. Code and data: https://github.com/irigi/VariableISPRocketTrajectories.

[2] L. C. Evans and P. E. Souganidis, "Differential games and representation formulas for solutions of Hamilton–Jacobi–Isaacs equations," *Indiana University Mathematics Journal*, vol. 33, no. 5, pp. 773–797, 1984.

[3] P. Soravia, "Pursuit–evasion problems and viscosity solutions of Isaacs equations," *SIAM Journal on Control and Optimization*, vol. 31, no. 3, pp. 604–623, 1993.

[4] G. I. Ibragimov, A. A. Azamov, and M. Khakestari, "Solution of a linear pursuit-evasion game with integral constraints," *ANZIAM Journal*, vol. 52, pp. E59–E75, 2011.

[5] W. H. Clohessy and R. S. Wiltshire, "Terminal guidance system for satellite rendezvous," *Journal of the Aerospace Sciences*, vol. 27, no. 9, pp. 653–658, 1960.

[6] L. S. Pontryagin, V. G. Boltyanskii, R. V. Gamkrelidze, and E. F. Mishchenko, *The Mathematical Theory of Optimal Processes*. Oxford, UK: Pergamon Press, 1962.

[7] I. M. Mitchell, A. M. Bayen, and C. J. Tomlin, "A time-dependent Hamilton–Jacobi formulation of reachable sets for continuous dynamic games," *IEEE Transactions on Automatic Control*, vol. 50, no. 7, pp. 947–957, 2005.

[8] K. Margellos and J. Lygeros, "Hamilton–Jacobi formulation for reach-avoid differential games." arXiv:0911.4625, 2009.

[9] P. Soravia, "Estimates of convergence of fully discrete schemes for the Isaacs equation of pursuit-evasion differential games via maximum principle," *SIAM Journal on Control and Optimization*, vol. 36, no. 1, pp. 1–11, 1998.