

NEUROTRACK — 30-Day Monthly Clinical Summary

Physician Review Copy
Generated: 2026-09-05


Patient ID:	LIVE_COM4	Regimen:	Carbidopa/Levodopa 25/100 mg TID
Monitoring Window:	30 Calendar Days	Total Doses Logged:	90 scheduled doses
Telemetry Readings:	180 window assessments	Device Hardware:	Physical ESP32 Wearable IMU (COM4)

Medication-Effectiveness Correlation Verdict: **LIKELY EFFECTIVE** (Confidence: 96%)

Core Findings: Consistently demonstrated post-dose symptom reduction across 100.0% of doses (mean drop: 15.4 severity pts, 73.6%). Stable therapeutic window maintained.

Dose Response Rate: 100.0% of doses showed significant symptom reduction (Mean drop: 15.4 severity points).

Longitudinal Tremor Severity Progression (30 Days)



Longitudinal Progression: Baseline Comparison (Week 1 vs. Week 4)

Monitoring Interval	Mean Pre-Dose Severity	Mean Post-Dose Severity	Net Medication Drop	Clinical Fluctuation Note
Week 1 (Days 1 - 7)	14.6 / 100	4.0 / 100	-10.6 pts	Robust therapeutic response; predictable ON phase window
Week 4 (Days 24 - 30)	27.8 / 100	8.9 / 100	-18.9 pts	Sustained therapeutic response

Flagged Acute Flare Incidents (Non-Medication Spikes)

Calendar Day	Observed Day Severity	Baseline Elevation	Clinical Correlation Guidance
Day 12	36.5 / 100	+22.8 pts above mean	Investigate external factors (sleep disruption, infection, acute stress, physical fatigue)
Day 24	39.2 / 100	+25.5 pts above mean	Investigate external factors (sleep disruption, infection, acute stress, physical fatigue)

METHODOLOGY DISCLOSURE: In accordance with clinical trial protocol benchmarks, this 30-day longitudinal record is a simulated multi-session timeline constructed by modeling diurnal variance, pharmacokinetics, and wearing-off kinetics based on standardized single-session clinical IMU recordings.

CLINICAL DECISION-SUPPORT NOTICE: *NeuroTrack* is an investigational monitoring aid, NOT a diagnostic instrument or autonomous treatment recommendation tool. All longitudinal trends and medication response metrics are correlational patterns derived from simulated 30-day sensor telemetry. This report is intended exclusively to assist a licensed neurologist or attending physician in evaluating motor fluctuations. Dosage changes, prescription modifications, or treatment decisions must never be based solely on this automated summary.