

Ordinary differential equations

Martin van Sint Annaland, Ivo Roghair

m.v.sintannaland@tue.nl

Chemical Process Intensification,
Eindhoven University of Technology

Part I

Explicit and Implicit methods

Today's outline

- 1 Introduction
- 2 Euler's method
 - Forward Euler
- 3 Rates of convergence
- 4 Runge-Kutta methods
 - RK2 methods
 - RK4 method
- 5 Step size control
- 6 Solving ODEs in Matlab

Overview

Ordinary differential equations

An equation containing a function of one independent variable and its derivatives, in contrast to a *partial differential equation*, which contains derivatives with respect to more independent variables.

Main question

How to solve

$$\frac{dy}{dx} = f(y(x), x) \quad \text{with} \quad y(x=0) = y_0$$

accurately and efficiently?

What is an ODE?

- Algebraic equation:

$$f(y(x), x) = 0 \quad \text{e.g. } -\ln(K_{eq}) = (1 - \zeta)$$

- First order ODE:

$$f\left(\frac{dy}{dx}(x), y(x), x\right) = 0 \quad \text{e.g. } \frac{dc}{dt} = -kc^n$$

- Second order ODE:

$$f\left(\frac{d^2y}{dx^2}(x), \frac{dy}{dx}(x), y(x), x\right) = 0 \quad \text{e.g. } \mathcal{D}\frac{d^2c}{dx^2} = -\frac{kc}{1 + Kc}$$

About second order ODEs

Very often a second order ODE can be rewritten into a system of first order ODEs (whether it is handy depends on the boundary conditions!)

More general

Consider the second order ODE:

$$\frac{d^2 y}{dx^2} + q(x) \frac{dy}{dx} = r(x)$$

Now define and solve using z as a new variable:

$$\frac{dy}{dx} = z(x)$$


$$\frac{dz}{dx} = r(x) - q(x)z(x)$$

Importance of boundary conditions

The nature of boundary conditions determines the appropriate numerical method. Classification into 2 main categories:


- *Initial value problems (IVP)*

We know the values of all y_i at some starting position x_s , and it is desired to find the values of y_i at some final point x_f .



- *Boundary value problems (BVP)*

Boundary conditions are specified at more than one x . Typically, some of the BC are specified at x_s and the remainder at x_f .



Overview

Initial value problems:

- Explicit methods
 - First order: forward Euler
 - Second order: improved Euler (RK2)
 - Fourth order: Runge-Kutta 4 (RK4)
 - Step size control
- Implicit methods
 - First order: backward Euler
 - Second order: midpoint rule

Boundary value problems

- Shooting method

Today's outline

- ① Introduction
- ② Euler's method
Forward Euler
- ③ Rates of convergence
- ④ Runge-Kutta methods
RK2 methods
RK4 method
- ⑤ Step size control
- ⑥ Solving ODEs in Matlab

Euler's method

Consider the following single initial value problem:

$$\frac{dc}{dt} = f(c(t), t) \quad \text{with} \quad c(t=0) = c_0 \quad (\text{initial value problem})$$

Easiest solution algorithm: Euler's method, derived here via Taylor series expansion:

$$c(t_0 + \Delta t) \approx c(t_0) + \left. \frac{dc}{dt} \right|_{t_0} \Delta t + \frac{1}{2} \left. \frac{d^2c}{dt^2} \right|_{t_0} (\Delta t)^2 + \mathcal{O}(\Delta t^3)$$


Neglect terms with higher order than two: $\left. \frac{dc}{dt} \right|_{t_0} = \frac{c(t_0 + \Delta t) - c(t_0)}{\Delta t}$

Substitution:


$$\frac{c(t_0 + \Delta t) - c(t_0)}{\Delta t} = f(c_0, t_0) \Rightarrow c(t_0 + \Delta t) = c(t_0) + \Delta t f(c_0, t_0)$$

Euler's method: graphical example

$$\frac{c(t_0 + \Delta t) - c(t_0)}{\Delta t} = f(c_0, t_0) \Rightarrow c(t_0 + \Delta t) = c(t_0) + \Delta t f(c_0, t_0)$$



Euler's method - solution method



Start with $t = t_0$, $c = c_0$, then calculate at discrete points in time:
 $c(t_1 = t_0 + \Delta t) = c(t_0) + \Delta t f(c_0, t_0)$.

Pseudo-code Euler's method: $\frac{dy}{dx} = f(x, y)$ and $y(x_0) = y_0$.

- 1 Initialize variables, functions; set $h = \frac{x_1 - x_0}{N}$
- 2 Set $x = x_0$, $y = y_0$
- 3 While $x < x_{\text{end}}$ do

$y_{i+1} = y_i + h f(x_i, y_i)$


Euler's method - example

First order reaction in a batch reactor:

$$\frac{dc}{dt} = -kc \quad \text{with} \quad c(t=0) = 1 \text{ mol m}^{-3}, \quad k = 1 \text{ s}^{-1}, \quad t_{\text{end}} = 2 \text{ s}$$

Time [s]	Concentration [mol m ⁻³]
$t_0 = 0$	$c_0 = 1.00$
$t_1 = t_0 + \Delta t$	$c_1 = c_0 + \Delta t \cdot (-kc_0)$
$= 0 + 0.1 = 0.1$	$= 1 + 0.1 \cdot (-1 \cdot 1) = 0.9$
$t_2 = t_1 + \Delta t$	$c_2 = c_1 + \Delta t \cdot (-kc_1)$
$= 0.1 + 0.1 = 0.2$	$= 0.9 + 0.1 \cdot (-1 \cdot 0.9) = 0.81$
$t_3 = t_2 + \Delta t$	$c_3 = c_2 + \Delta t \cdot (-kc_2)$
$= 0.2 + 0.1 = 0.3$	$= 0.81 + 0.1 \cdot (-1 \cdot 0.81) = 0.729$
...	...
$t_{i+1} = t_i + \Delta t$	$c_{i+1} = c_i + \Delta t \cdot (-kc_i)$
...	...
$t_{20} = 2.0$	$c_{20} = c_{19} + \Delta t \cdot (-kc_{19}) = 0.121577$



Euler's method - example



Problems with Euler's method

The question is: What step size, or how many steps to use?

- 1 *Accuracy* \Rightarrow need information on numerical error!
- 2 *Stability* \Rightarrow need information on stability limits!




Accuracy

Comparison with analytical solution for $k = 1 \text{ s}^{-1}$:

$$c(t) = c_0 \exp(-kt) \Rightarrow \zeta = 1 - \exp(-kt) \Rightarrow \zeta_{\text{analytical}} = 0.864665$$

N	ζ	$\frac{\zeta_{\text{numerical}} - \zeta_{\text{analytical}}}{\zeta_{\text{analytical}}}$
20	0.878423	0.015912
40	0.871488	0.007891
80	0.868062	0.003929
160	0.866360	0.001961
320	0.865511	0.000979



Accuracy

For Euler's method: Error halves when the number of grid points is doubled, i.e. error is proportional to Δt : first order method.

Error estimate:

$$\left. \frac{dx}{dt} \right|_{t_0} = \frac{x(t_0 + \Delta t) - x(t_0)}{\Delta t} + \frac{1}{2} \left. \frac{d^2x}{dt^2} \right|_{t_0} (\Delta t) + \mathcal{O}(\Delta t)^2$$

$$\frac{x(t_0 + \Delta t) - x(t_0)}{\Delta t} = f(x_0, t_0) - \frac{1}{2} \left. \frac{d^2x}{dt^2} \right|_{t_0} (\Delta t) + \mathcal{O}(\Delta t)^2$$

Today's outline

- ① Introduction
- ② Euler's method
Forward Euler
- ③ Rates of convergence
- ④ Runge-Kutta methods
RK2 methods
RK4 method
- ⑤ Step size control
- ⑥ Solving ODEs in Matlab

Errors and convergence rate

L_2 norm (Euclidean norm)

$$\|\mathbf{v}\|_2 = \sqrt{v_1^2 + v_2^2 + \dots + v_n^2} = \sqrt{\sum_{i=1}^n v_i^2}$$

L_∞ norm (maximum norm)

$$\|\mathbf{v}\|_\infty = \max(|v_1|, \dots, |v_n|)$$

Absolute difference

$$\epsilon_{\text{abs}} = \left\| \mathbf{y}_{\text{numerical}} - \mathbf{y}_{\text{analytical}} \right\|_{2,\infty}$$

Relative difference

$$\epsilon_{\text{rel}} = \frac{\left\| \mathbf{y}_{\text{numerical}} - \mathbf{y}_{\text{analytical}} \right\|_{2,\infty}}{\left\| \mathbf{y}_{\text{analytical}} \right\|_{2,\infty}}$$

Errors and convergence rate

Convergence rate (or: order of convergence) r

$$\epsilon = \lim_{\Delta x \rightarrow 0} c(\Delta x)^r$$

- A first order method reduces the error by a factor 2 when increasing the number of steps by a factor 2
- A second order method reduces the error by a factor 4 when increasing the number of steps by a factor 2

Computing the rate of convergence

When the analytical solution is available, choose ① or ② for a particular number of grid points N :

- ① Compute the relative or absolute error vector $\bar{\epsilon}$. Take the norm to compute a single error value ϵ following:
 - Based on L_1 -norm: $\epsilon = \frac{\|\bar{\epsilon}\|_1}{N}$
 - Based on L_2 -norm: $\epsilon = \frac{\|\bar{\epsilon}\|_2}{\sqrt{N}}$
 - Based on L_∞ -norm: $\epsilon = \|\bar{\epsilon}\|_\infty$
- ② Compute the relative or absolute error at a single indicative points (e.g. middle of domain, outlet).

Compare to calculations with different number of steps: $\epsilon_1 = c(\Delta x_1)^r$ and $\epsilon_2 = c(\Delta x_2)^r$ and solve for r :

$$\frac{\epsilon_2}{\epsilon_1} = \frac{c(\Delta x_2)^r}{c(\Delta x_1)^r} = \left(\frac{\Delta x_2}{\Delta x_1} \right)^r \Rightarrow \log \left(\frac{\epsilon_2}{\epsilon_1} \right) = \log \left(\frac{\Delta x_2}{\Delta x_1} \right)^r$$

$$\Rightarrow r = \frac{\log \left(\frac{\epsilon_2}{\epsilon_1} \right)}{\log \left(\frac{\Delta x_2}{\Delta x_1} \right)} = \frac{\log \left(\frac{\epsilon_2}{\epsilon_1} \right)}{\log \left(\frac{N_1}{N_2} \right)} \quad \text{in the limit of } \Delta x \rightarrow 0 \quad \text{or} \quad N \rightarrow \infty$$

Computing the rate of convergence

When the analytical solution is not available:

- ① Compute the solution with $N + 1$, N , $N - 1$ and $N - 2$ grid points
- ② Select a single indicative grid point (e.g. middle of domain, outlet) that lies at exactly the same position in each computation
- ③ Use the solution c at this grid point for various grid sizes to compute:

$$r = \frac{\log \frac{c_{N+1} - c_N}{c_N - c_{N-1}}}{\log \frac{c_N - c_{N-1}}{c_{N-1} - c_{N-2}}}$$

- ④ Alternative for simulations with $2N$, N and $\frac{N}{2}$ grid points:

$$r = \frac{\log \left| \frac{c_{2N} - c_N}{c_N - c_{\frac{N}{2}}} \right|}{\log \left| \frac{2N}{N} \right|}$$

Example: Euler's method — order of convergence

N	ζ	$\frac{\zeta_{\text{numerical}} - \zeta_{\text{analytical}}}{\zeta_{\text{analytical}}}$	$r = \frac{\log\left(\frac{\epsilon_i}{\epsilon_{i-1}}\right)}{\log\left(\frac{N_{i-1}}{N_i}\right)}$
20	0.878423	0.015912	—
40	0.871488	0.007891	1.011832
80	0.868062	0.003929	1.005969
160	0.866360	0.001961	1.002996
320	0.865511	0.000979	1.001500

⇒ Euler's method is a first order method (as we already knew from the truncation error analysis)

Wouldn't it be great to have a method that can give the answer using much less steps? ⇒ Higher order methods

Today's outline


- ① Introduction
- ② Euler's method
Forward Euler
- ③ Rates of convergence
- ④ Runge-Kutta methods
RK2 methods
RK4 method
- ⑤ Step size control
- ⑥ Solving ODEs in Matlab

Runge-Kutta methods


Propagate a solution by combining the information of several Euler-style steps (each involving one function evaluation) to match a Taylor series expansion up to some higher order.

Euler: $y_{i+1} = y_i + hf(x_i, y_i)$ with $h = \Delta x$, i.e.
slope = $k_1 = f(x_i, y_i)$.

Euler's method



RK2 method



Classical second order Runge-Kutta (RK2) method

This method is also called Heun's method, or improved Euler method:

- 1 Approximate the slope at x_i : $k_1 = f(x_i, y_i)$
- 2 Approximate the slope at x_{i+1} : $k_2 = f(x_{i+1}, y_{i+1})$ where we use Euler's method to approximate $y_{i+1} = y_i + hf(x_i, y_i) = y_i + hk_1$
- 3 Perform an Euler step with the average of the slopes:
$$y_{i+1} = y_i + h\frac{1}{2}(k_1 + k_2)$$

In pseudocode:

```
x = x0, y = y0
while x < xend do
  xi+1 = xi + h
  k1 = f(xi, yi)
  k2 = f(xi + h, yi + hk1)
  yi+1 = yi + h $\frac{1}{2}$ (k1 + k2)
end while
```

Runge-Kutta methods — derivation

$$\frac{dy}{dx} = f(x, y(x))$$

Using Taylor series expansion: $y_{i+1} = y_i + h \left. \frac{dy}{dx} \right|_i + \frac{h^2}{2} \left. \frac{d^2y}{dx^2} \right|_i + \mathcal{O}(h^3)$

$$\left. \frac{dy}{dx} \right|_i = f(x_i, y_i) \equiv f_i$$

$$\left. \frac{d^2y}{dx^2} \right|_i = \left. \frac{d}{dx} f(x, y(x)) \right|_i = \left. \frac{\partial f}{\partial x} \right|_i + \left. \frac{\partial f}{\partial y} \right|_i \left. \frac{dy}{dx} \right|_i = \left. \frac{\partial f}{\partial x} \right|_i + \left. \frac{\partial f}{\partial y} \right|_i f_i \quad (\text{chain rule})$$

Substitution gives:

$$y_{i+1} = y_i + hf_i + \frac{h^2}{2} \left(\left. \frac{\partial f}{\partial x} \right|_i + \left. \frac{\partial f}{\partial y} \right|_i f_i \right) + \mathcal{O}(h^3)$$

$$y_{i+1} = y_i + \frac{h}{2} f_i + \frac{h}{2} \left(f_i + h \left. \frac{\partial f}{\partial x} \right|_i + hf_i \left. \frac{\partial f}{\partial y} \right|_i \right) + \mathcal{O}(h^3)$$

Runge-Kutta methods — derivation

Note multivariate Taylor expansion:

$$\begin{aligned} f(x_i + h, y_i + k) &= f_i + h \left. \frac{\partial f}{\partial x} \right|_i + h \left. \frac{\partial f}{\partial y} \right|_i + \mathcal{O}(h^2) \\ \Rightarrow \frac{h}{2} \left(f_i + h \left. \frac{\partial f}{\partial x} \right|_i + h f_i \left. \frac{\partial f}{\partial y} \right|_i \right) &= \frac{h}{2} f(x_i + h, y_i + h f_i) + \mathcal{O}(h^3) \end{aligned}$$

Concluding:

$$y_{i+1} = y_i + \frac{h}{2} f_i + \frac{h}{2} f(x_i + h, y_i + h f_i) + \mathcal{O}(h^3)$$

Rewriting:

$$\begin{aligned} k_1 &= f(x_i, y_i) \\ k_2 &= f(x_i + h, y_i + h k_1) \\ \Rightarrow y_{i+1} &= y_i + \frac{h}{2} (k_1 + k_2) \end{aligned}$$

Runge-Kutta methods — derivation

Generalization: $y_{i+1} = y_i + h(b_1 k_1 + b_2 k_2) + \mathcal{O}(h^3)$

with $k_1 = f_i$, $k_2 = f(x_i + c_2 h, y_i + a_{2,1} h k_1)$

(Note that classical RK2: $b_1 = b_2 = \frac{1}{2}$ and $c_2 = a_{2,1} = 1$.)

Bivariate Taylor expansion:

$$f(x_i + c_2 h, y_i + a_{2,1} h k_1) = f_i + c_2 h \left. \frac{\partial f}{\partial x} \right|_i + a_{2,1} h k_1 \left. \frac{\partial f}{\partial y} \right|_i + \mathcal{O}(h^2)$$

$$y_{i+1} = y_i + h(b_1 k_1 + b_2 k_2) + \mathcal{O}(h^3)$$

$$= y_i + h[b_1 f_i + b_2 f(x_i + c_2 h, y_i + a_{2,1} h k_1)] + \mathcal{O}(h^3)$$

$$= y_i + h \left[b_1 f_i + b_2 \left\{ f_i + c_2 h \left. \frac{\partial f}{\partial x} \right|_i + a_{2,1} h k_1 \left. \frac{\partial f}{\partial y} \right|_i + \mathcal{O}(h^2) \right\} \right] + \mathcal{O}(h^3)$$

$$= y_i + h(b_1 + b_2) f_i + h^2 b_2 \left(c_2 \left. \frac{\partial f}{\partial x} \right|_i + a_{2,1} f_i \left. \frac{\partial f}{\partial y} \right|_i \right) + \mathcal{O}(h^3)$$

Comparison with Taylor:

$$y_{i+1} = y_i + h f_i + \frac{h^2}{2} \left(\left. \frac{\partial f}{\partial x} \right|_i + \left. \frac{\partial f}{\partial y} \right|_i f_i \right) + \mathcal{O}(h^3)$$

Runge-Kutta methods — derivation

$$y_{i+1} = y_i + h(b_1 + b_2)f_i + h^2 b_2 \left(c_2 \frac{\partial f}{\partial x} \Big|_i + a_{2,1} f_i \frac{\partial f}{\partial y} \Big|_i \right) + \mathcal{O}(h^3)$$

$$y_{i+1} = y_i + hf_i + \frac{h^2}{2} \left(\frac{\partial f}{\partial x} \Big|_i + \frac{\partial f}{\partial y} \Big|_i f_i \right) + \mathcal{O}(h^3)$$

\Rightarrow 3 eqns and 4 unknowns \Rightarrow multiple possibilities!

- 1 Classical RK2:

$$b_1 = b_2 = \frac{1}{2} \text{ and } c_2 = a_{2,1} = 1$$

- 2 Midpoint rule (modified Euler):

$$b_1 = 0, b_2 = 1, c_2 = a_{2,1} = \frac{1}{2}$$

Second order Runge-Kutta methods

Classical RK2 method
(= Heun's method,
improved Euler method)

$$k_1 = f_i$$

$$k_2 = f(x_i + h, y_i + hk_1)$$



$$y_{i+1} = y_i + \frac{1}{2}h(k_1 + k_2)$$

Explicit midpoint rule
(modified Euler method)

$$k_1 = f_i$$

$$k_2 = f(x_i + \frac{1}{2}h, y_i + \frac{1}{2}hk_1)$$


$$y_{i+1} = y_i + hk_2$$



Second order Runge-Kutta method — Example

First order reaction in a batch reactor: $\frac{dc}{dt} = -kc$ with
 $c(t = 0) = 1 \text{ mol m}^{-3}$, $k = 1 \text{ s}^{-1}$, $t_{\text{end}} = 2 \text{ s}$.

Time [s]	C [mol m ⁻³]	$k_1 = hf(x_i, y_i)$	$k_2 = hf(x_i + \frac{1}{2}h, y_n + \frac{1}{2}k_1)$
0	1.00	$0.1 \cdot (-1 \cdot 1) = -0.1$	$0.1 \cdot (-1 \cdot (1 - 0.5 \cdot 0.1)) = -0.095$
0.1	$1 - 0.095 = 0.905$	$0.1 \cdot (-1 \cdot 0.905) = -0.0905$	$0.1 \cdot (-1 \cdot (0.905 - 0.5 \cdot 0.0905)) = -0.085975$
...
2	0.1358225	-0.0135822	-0.0129031




RK2 method — order of convergence

N	ζ	$\frac{\zeta_{\text{numerical}} - \zeta_{\text{analytical}}}{\zeta_{\text{analytical}}}$	$r = \frac{\log\left(\frac{\epsilon_i}{\epsilon_{i-1}}\right)}{\log\left(\frac{N_{i-1}}{N_i}\right)}$
20	0.864178	5.634×10^{-4}	—
40	0.864548	1.355×10^{-4}	2.056
80	0.864636	3.323×10^{-5}	2.028
160	0.864658	8.229×10^{-6}	2.014
320	0.864663	2.048×10^{-6}	2.007

⇒ RK2 is a second order method. Doubling the number of cells reduces the error by a factor 4!

Can we do even better?

RK4 method (classical fourth order Runge-Kutta method)



$$k_1 = f(x_i, y_i)$$

$$k_2 = f(x_i + \frac{1}{2}h, y_i + \frac{1}{2}hk_1)$$

$$k_3 = f(x_i + \frac{1}{2}h, y_i + \frac{1}{2}hk_2)$$

$$k_4 = f(x_i + h, y_i + hk_3)$$

$$y_{i+1} = y_i + h \left(\frac{1}{6}k_1 + \frac{1}{3}(k_2 + k_3) + \frac{1}{6}k_4 \right)$$

RK4 method — order of convergence

N	ζ	$\frac{\zeta_{\text{numerical}} - \zeta_{\text{analytical}}}{\zeta_{\text{analytical}}}$	$r = \frac{\log\left(\frac{\epsilon_j}{\epsilon_{j-1}}\right)}{\log\left(\frac{N_{j-1}}{N_j}\right)}$
20	0.864664472	2.836×10^{-7}	—
40	0.864664702	1.700×10^{-8}	4.060
80	0.864664716	1.040×10^{-9}	4.030
160	0.864664717	6.435×10^{-11}	4.015
320	0.864664717	4.001×10^{-12}	4.007

⇒ RK4 is a fourth order method: Doubling the number of cells reduces the error by a factor 16!

Can we do even better?

Today's outline

- ① Introduction
- ② Euler's method
Forward Euler
- ③ Rates of convergence
- ④ Runge-Kutta methods
RK2 methods
RK4 method
- ⑤ Step size control**
- ⑥ Solving ODEs in Matlab


Adaptive step size control

The step size (be it either position, time or both (PDEs)) cannot be decreased indefinitely to favour a higher accuracy, since each additional grid point causes additional computation time. It may be wise to adapt the step size according to the computation requirements.

Globally two different approaches can be used:

- 1 Step doubling: compare solutions when taking one full step or two consecutive halve steps
- 2 Embedded methods: Compare solutions when using two approximations of different order

Adaptive step size control: step doubling



- RK4 with one large step of h : $y_{i+1} = y_1 + ch^5 + \mathcal{O}(h^6)$
- RK4 with two steps of $\frac{1}{2}h$: $y_{i+1} = y_2 + 2c(\frac{1}{2}h)^5 + \mathcal{O}(h^6)$

Adaptive step size control: step doubling

- Estimation of truncation error by comparing y_1 and y_2 :

$$\Delta = y_2 - y_1$$

- If Δ too large, reduce step size for accuracy
- If Δ too small, increase step size for efficiency.

- Ignoring higher order terms and solving for c :

$$\Delta = \frac{15}{16}ch^5 \Rightarrow ch^5 = \frac{16}{15}\Delta \Rightarrow y_{i+1} = y_2 + \frac{\Delta}{15} + \mathcal{O}(h^6)$$

(local Richardson extrapolation)

Note that when we specify a tolerance tol , we can estimate the maximum allowable step size as: $h_{\text{new}} = \alpha h_{\text{old}} \left| \frac{tol}{\Delta} \right|^{\frac{1}{5}}$ with α a safety factor (typically $\alpha = 0.9$).

Adaptive step size control: embedded methods

Use a special fourth and a fifth order Runge Kutta method to approximate y_{i+1}

- The fourth order method is special because we want to use the same positions for the evaluation for computational efficiency.
- RK45 is the preferred method (minimum number of function evaluations) (this is built in Matlab as `ode45`).

Today's outline

- ① Introduction
- ② Euler's method
Forward Euler
- ③ Rates of convergence
- ④ Runge-Kutta methods
RK2 methods
RK4 method
- ⑤ Step size control
- ⑥ Solving ODEs in Matlab

Solving ODEs in Matlab

Matlab provides convenient procedures to solve (systems of) ODEs automatically.

The procedure is as follows:

- 1 Create a function that specifies the ODE(s). Specifically, this function returns the $\frac{dy}{dx}$ value (vector).
- 2 Initialise solver variables and settings (e.g. step size, initial conditions, tolerance), in a separate script
- 3 Call the ODE solver function, using a *function handle* to the ODE function described in point 1.
 - The ODE solver will return the vector for the independent variable, and a solution vector (matrix for systems of ODEs).

Solving ODEs in Matlab: example 1

We solve the system: $\frac{dx}{dt} = -k_1x + k_2, k_1 = 0.2, k_2 = 2.5$

- Create an anonymous function handle:


```
>> myEqn = @(t,x) (-0.2*x + 2.5)
```

- Solve with a call to

`ode45`(function_handle, timespan, initial_condition):

```
>> ode45(myEqn, [0 40], 0);
```

- By omitting the output of this function, the graph is automatically drawn.



Solving ODEs in Matlab: example 2

We solve the system:

$$\frac{dx}{dt} = \begin{cases} -\frac{k_1}{x^2} & t \leq 10 \\ \frac{k_2}{x} - \frac{k_1}{x^2} & t > 10 \end{cases} \quad \text{with } k_1 = 0.5, k_2 = 1, x(0) = 2$$

Create an ODE function

```
function [dxdt] = myEqnFunction(t,x)
k1 = 0.5;
k2 = 1;
dxdt = (t>10)*k2/x - k1/x^2;
```


Create a solution script

```
x_init = 2;           % Initial condition
tspan = [0 20];      % Time span
options = odeset('RelTol',1e-8,'AbsTol',1e-8);
[t,x] = ode45(@myEqnFunction,tspan,x_init,options);
```

Solving ODEs in Matlab: example 2

Plot the solution:

```
plot(t,x,'r-x')
```



Note the refinement in regions where large changes occur.

Solving ODEs in Matlab: example

A few notes on working with `ode45` and other solvers. If we want to give additional arguments (e.g. `k1` and `k2`) to our ODE function, we can list them in the function line:

```
function [dxdt] = myEqn(t,x,k1,k2)
```

The additional arguments can now be set in the solver script by *adding them after the options*:

```
[t,x] = ode45(@myEqn,tspan,x_0,options,k1,k2);
```

- Of course, in the solver script, the variables do not have to be called `k1` and `k2`:

```
[t,x] = ode45(@myODE,tspan,x_0,options,q,u);
```

- These variables may be of any type (vectors, matrix, struct). Especially a struct is useful to carry many values in 1 variable.

Solving systems of ODEs in Matlab: example

You have noticed that the step size in t varied. This is because we have given just the begin and end times of our time span:

```
tspan = [0 10];
```

You can also solve at specific steps, by supplying all steps explicitly, e.g.:

```
tspan = linspace(0,10,101);
```

This example provides 101 explicit time steps between 0 and 10 seconds.

Note that the results are interpolated to these data points afterwards; you do not influence the efficiency and accuracy of the solver algorithm this way!