

CHE 201 Fall 2019 Midterm

Abdulla Amer

TOTAL POINTS

71 / 100

QUESTION 1

20 pts

1.1 a 6 / 10

- 0 pts Correct
- 2 pts first one is wrong
- ✓ - 2 pts second one is wrong
- 2 pts third one is wrong
- ✓ - 2 pts fourth one is wrong
- 2 pts fifth one is wrong

1.2 b 3 / 5

- 0 pts Correct
- ✓ - 2 pts gage pressure is not converted to absolute pressure
- 2 pts T is not converted to K
- 5 pts Not done or completely wrong
- 3 pts only pressure is calculated

1.3 c 0 / 5

- 0 pts Correct
- ✓ - 5 pts incorrect

QUESTION 2

2 50 pts

2.1 a 3 / 3

- ✓ - 0 pts Correct

2.2 b 8 / 8

- ✓ - 0 pts Correct
- 1 pts PV state 2 is wrong
- 1 pts PV state 3 is wrong
- 1 pts TV state 2 is wrong
- 1 pts TV state 3 is wrong
- 8 pts incorrect

- 1 pts PV state 1 is wrong
- 1 pts TV state 1 is wrong
- 4 pts PV graph is wrong

2.3 c 5 / 5

- ✓ - 0 pts Correct
- 1 pts work expansion compression
- 2 pts KE and PE is zero
- 2 pts closed system is not listed

2.4 d 2 / 2

- ✓ - 0 pts Correct
- 1 pts ΔE equation is not written
- 1 pts not simplified

2.5 e 12 / 15

- 0 pts Correct
- 3 pts calculation error
- ✓ - 3 pts v_3 is wrong
- 3 pts v_2 is wrong
- 15 pts incorrect
- 3 pts P is wrong
- 3 pts unit conversion
- 3 pts W_{12} is zero
- 2 pts not multiplied with mass

2.6 f 12 / 15

- 0 pts Correct
- ✓ - 3 pts u_3 is wrong
- 3 pts u_1 is wrong
- 2 pts not multiplied with mass
- 2 pts x value is wrong
- 2 pts calculation error
- 15 pts incorrect
- 3 pts q not calculated
- 3 pts not clear calculation

2.7 g 1 / 2

- 0 pts Correct

✓ - 0.5 pts $M3v$ not calculated /wrong

✓ - 0.5 pts $m2v$ not calculated/ wrong

- 1 pts $m1v$ not calculated / wrong

✓ - 10 pts incorrect or not done

- 2 pts calculation error

QUESTION 3

3 30 pts

3.1 a 2 / 2

✓ - 0 pts Correct

- 2 pts not done

- 1 pts boundary not labeled

3.2 b 2 / 2

✓ - 0 pts Correct

- 0.5 pts Delta KE and PE is zero not listed

- 1 pts ideal gas is not listed

- 1 pts closed system is not listed

3.3 C 2 / 3

- 0 pts Correct

✓ - 1 pts Z is not listed or wrong

- 3 pts not done

- 2 pts delta E equation is not listed

- 1 pts not simplified

3.4 d 3 / 3

✓ - 0 pts Correct

- 3 pts wrong

- 1 pts slope needs to be negative

- 1 pts Not curved, plotted as linear

3.5 e 10 / 10

✓ - 0 pts Correct

- 10 pts not correct or not done

- 6 pts error with integration

- 4 pts not clear explanation

- 2 pts calculation error

3.6 f 0 / 10

- 0 pts Correct

NAME: Abdullah Amer

Please fill in answers in the space supplied and show all your work to receive full credit.
Useful Tables at end of Exam.

1. a) (10 pts) State whether the following statements are **True or False**

F An ideal gas at a given state expands to a fixed final volume at constant pressure or at constant temperature. Work is greater in constant temperature case.

T A two-phase liquid-vapor mixture with equal volumes of saturated vapor and saturated liquid has a quality of 0.5.

F As pressure decreases from the critical pressure, the values of the specific volume of saturated vapor and saturated liquid approach each other.

T If a system undergoes a polytropic process, the system must consist of an ideal gas.

T It takes more energy to vaporize 1kg of saturated liquid water at 100°C than it would at 120°C.

b) (5 pts) The gage pressure of an automobile tire is measured to be 210kPa before a trip and 220kPa after the trip at a location where the atmospheric pressure is 95kPa. Assuming the volume of the tire remains constant and the air temperature before the trip is 25°C, determine the air temperature in the tire after the trip. (Air is ideal gas)

$$\begin{aligned}
 P_1 &= 210 \text{ kPa} \\
 P_2 &= 220 \text{ kPa} \\
 P_{\text{atm}} &= 95 \text{ kPa} \\
 T_1 &= 25^\circ\text{C}
 \end{aligned}
 \quad
 \frac{T_2}{T_1} = \frac{P_2}{P_1}
 \quad
 T_2 = \frac{P_2}{P_1} T_1 = \frac{220 \text{ kPa}}{210 \text{ kPa}} (273 + 25) = 281.5 \text{ K}$$

c) (5pts) The specific volume of saturated liquid water at 0.01°C is $1.002 \times 10^{-3} \text{ m}^3/\text{kg}$. What would be the specific volume of saturated solid at the same temperature, greater or less than saturated liquid value? Why?

less, because as temperature increases molecules move at faster rates, and the nature of solid molecules are more joined than liquid so they move at slower rates therefore occupying less volume.

NAME: _____

CHE 201 MidTerm Exam 1




October 11th, 2019

2. (50 pts) 2kg of liquid-vapor mixture of water contained in a piston-cylinder assembly undergoes two processes in series from an initial state where the temperature is 120°C.

- **Process 1-2** System is heated at constant volume to a saturated vapor state at 200°C.
- **Process 2-3** System is compressed at constant pressure until it reaches to 150°C.

Kinetic and potential effects can be ignored.

- (3 pts) Draw a schematic of the system and label the system boundary.
- (8 pts) Carefully sketch the process on a:
 - P-v diagram (Pressure-specific volume)
 - T-v diagram (Temperature-specific volume)
- (5 pts) State the engineering model
- (2 pts) Write the simplified energy balance
- (15 pts) Calculate the work transferred in kJ. Is work done on the system or by the system?
- (15 pts) Calculate the heat transferred in kJ. Is heat transferred to the system or from the system?
- (2 pts) Calculate the mass of vapor in each state in kg.



- c) Model
- closed system
 - ΔKE and $\Delta PE = 0$
 - work expansion / contraction
 - Heat gain / loss

d) $\Delta E = Q - W$

$$\Delta KE + \Delta PE + \Delta U = Q - W \rightarrow \Delta U = Q - W$$

NAME: _____

CHE 201 MidTerm Exam 1

October 11th, 2019

$$\textcircled{c}) \quad w_{1-2} = 0$$

$$w_{2-3} = \int_{v_2}^{v_1} P \, dv$$

$$N_1 = N_2 = N_3 = 0.1274 \, \text{m}^3/\text{kg}$$

$$N_g = N_F - x(N_g - N_F)$$

$$x = 0.14$$

$$N_3 = 0.056 \, \text{m}^3/\text{kg}$$

$$w_{2-3} = 15.54 \, \text{bar} \cdot \frac{10^5 \, \text{Pa}}{1 \, \text{bar}} \left(0.056 \, \text{m}^3/\text{kg} - 0.1274 \, \text{m}^3/\text{kg} \right)$$

$$= -110.96 \, \text{kJ/kg} \times 2 \, \text{kg} = -221.91 \, \text{kJ}$$

$$\textcircled{f}) \quad Q = \Delta U + w$$

$$Q = (U_3 - U_1) + w$$

$$U_F = U_F + x(U_g - U_F) = 787.112 \, \text{kJ/kg}$$

$$U_3 = U_F + x(U_g - U_F) = 901.57 \, \text{kJ/kg}$$

$$Q = m(U_3 - U_1) + w$$

$$= 2(901.57 \, \text{kJ/kg} - 787.112 \, \text{kJ/kg}) - (221.91 \, \text{kJ}) = 7.01 \, \text{kJ}$$

$$\textcircled{g}) \quad x = \frac{m_{\text{vap}}}{m}$$

$$m_{\text{vap}} = m x = 0.14(2 \, \text{kg}) = 0.28 \, \text{kg}$$

NAME: _____

CHE 201 MidTerm Exam 1

October 11th, 2019

3. (30 points) A piston cylinder device initially contains 0.4 m^3 of air at 100 kPa and 240°C . The air is now compressed to 0.1 m^3 in such a way that the temperature inside the cylinder remains constant. The air is modeled as an ideal gas and kinetic and potential energy effects are negligible.

- (2 pts) Draw a schematic of the system and label the system boundary.
- (2 pts) State the engineering model
- (3 pts) Write the simplified energy balance and compressibility factor (Z).
- (3 pts) Draw P-v diagram (Pressure-specific volume)
- (10 pts) Determine the work (kJ)
- (10 pts) Determine the heat (kJ)



$$V_1 = 0.4 \text{ m}^3 \quad V_2 = 0.1 \text{ m}^3$$

$$P_1 = 100 \text{ kPa}$$

$$T_1 = 240^\circ\text{C} + 273 = T_2$$

$$P_2 = \frac{RT_2}{V_2}$$

$$= \frac{8.314 \text{ kJ/kmol} \cdot \left(\frac{1 \text{ kmol}}{28.97 \text{ kg}} \right) (513 \text{ K})}{0.1 \text{ m}^3} \left(\frac{1 \text{ kPa}}{1 \text{ kJ}} \right)$$

$$= 1472240.9 \text{ Pa} \times \frac{1 \text{ kPa}}{10^3 \text{ Pa}} = 1472.24 \text{ kPa}$$


b) Model:-

- closed system
- ideal gas
- $\Delta KE = \Delta PE = 0$

$$c) \Delta E = Q - W$$

$$\Delta PE + \Delta KE + \Delta U = \cancel{Q} - W$$

$$\Delta U = W$$



NAME: _____

CHE 201 MidTerm Exam 1

October 11th, 2019

$$e) \quad W = P_1 V_1 \ln \frac{V_2}{V_1}$$

$$= (100 \text{ kPa}) (0.4 \text{ m}^3) \ln \left(\frac{0.1}{0.4} \right)$$

$$= -55.45 \text{ kJ}$$

$$f) \quad Q = 0$$

since temperature is constant

