

MATH676 — Tropical Geometry

Based on the lectures by Renzo Cavalieri

Notes written by Ignacio Rojas

Fall 2023

Please note that these notes were not provided or endorsed by the lecturer and have been significantly altered after the class. They may not accurately reflect the content covered in class and any errors are solely my responsibility.

This is a topics course on this stuff

Requirements

Knowledge on stuff

TO DO:

- ◇ Write info on course description and requirements.
- ◇ Polish info from day 1
- ◇ Polish last part of day 2
- ◇ Continue adding info from Renzo's digital notes

Contents

Contents	2
1 Combinatorial Shadow of Algebraic Geometry	3
1.1 Day 1 20230821	3
1.2 Day 2 20230823	4
1.3 Interim	8
1.4 Day 3 20230825	10
Index	11
Bibliography	13

Chapter 1

Combinatorial Shadow of Algebraic Geometry

1.1 Day 1 | 20230821

Think of an algorithm where the input is an algebraic variety and the output is a combinatorial object, a piecewise linear object.

Example 1.1.1. Consider as an input a line in the plane. Say $V(x + y - 1)$, then an output would be a tropical line. If we remain in the plane and consider a higher degree polynomial, say an elliptic curve, as an output we obtain a tropical cubic.

Leaving the plane behind and thinking of abstract nodal curves, we can think of a sphere attached to a torus which is attached to a genus 2 torus, then the corresponding object is what we call the dual graph.

Right now we do not know the specific algorithm, but we can observe that the outputs are *more simple* than the inputs. So the important question is:

What algebraic information does the simplified object remember? How do we extract the information the object remembers? And once we know how to work with this objects, can we return to algebraic geometry from any kind of these objects?

Observe that the number of ends which go to infinity corresponds with the degree.

1.2 Day 2 | 20230823

Algebraic Geometry on \mathbb{T}

Let us talk about ways to get into tropical geometry. We will first define the tropical semifield which the base set over which we will do algebraic geometry.

Definition 1.2.1. The tropical semifield is the set $(\mathbb{R} \cup \{-\infty\})$ equipped with tropical addition and multiplication:

$$\begin{cases} x \oplus y = \max(x, y) \\ x \odot y = x + y \end{cases}$$

With this set we can make multivariable polynomials


$$p(\underline{x}) : (\mathbb{R} \cup \{-\infty\})^n \rightarrow \mathbb{R} \cup \{-\infty\}$$

which gives rise to their *tropicalization*, a piecewise linear function $\text{Trop}(p) : \mathbb{R}^n \rightarrow \mathbb{R}$.

Example 1.2.2. Consider the polynomial

$$p(x, y) = x \oplus y \oplus 0,$$

its tropicalization is $\text{Trop}(p)(x, y) = \max(x, y, 0)$ which indeed is a piecewise linear function from \mathbb{R}^2 to \mathbb{R} . Observe that the surface is not smooth where the planes meet,



this is what we will call the *corner locus* or *tropical hypersurface*.

Definition 1.2.3. The tropical hypersurface $V(\text{Trop}(p))$ is the codimension 1 locus in \mathbb{R}^n where the function is non-linear (corner locus).

Example 1.2.4. If we consider higher degree tropical polynomials, they will become linear in the usual sense. Consider

$$p(x) = 3x^2 = 3 \odot x \odot x = 3 + x + x = 3 + 2x$$

which is indeed linear.

Valued fields

Definition 1.2.5. The field of Puiseux series or rational functions over \mathbb{C} is $\mathbb{C}(t)$ where the elements are of the form

$$f(t) = \sum_{i=k_0}^{\infty} a_i t^{i/n}.$$

The lower bound k_0 could be negative and the exponents, are rational with bounded denominators.

Consider the valuation

$$\text{val}_0 : \mathbb{C}(t) \rightarrow \mathbb{R} \cup \{\infty\}, \begin{cases} 0 \mapsto \infty \\ f \mapsto \text{order of vanishing at } 0. \end{cases}$$

This order of vanishing is the value α such that f/t^α approaches a finite non-zero value. The corresponding coefficient in the series expansion of f for this value is called the valuation coefficient.

Example 1.2.6. What happens to the order of vanishing when you add two functions? Consider $f = t^2$, $g = t^3$, then $f + g = t^2 + t^3$ which has order of vanishing 2. Observe that $2 = \min(2, 3)$.

In general what happens is that

$$\text{val}_0(f_1 + f_2) \geq \min(\text{val}_0 f_1, \text{val}_0 f_2), \quad \text{and} \quad \text{val}_0(f_1 f_2) = \text{val}_0(f_1) + \text{val}_0(f_2).$$

We can do algebraic geometry over this field! Let K be the field of rational functions, if $p(\underline{x}) \in K[\underline{x}]$ then we consider the algebraic variety

$$X = V(p) = \{ \mathbf{x} : p(\mathbf{x}) = 0 \} \subseteq K^n.$$

Taking the image through the n -fold valuation, we will obtain a set in $(\mathbb{R} \cup \{\infty\})^n$. The tropicalization of X is the image via this map: and here $\text{Trop}(V(p))$ is the tropical hypersurface for p .

$$\begin{array}{ccc}
 \text{val}_0 : K^n & \longrightarrow & (\mathbb{R} \cup \{\infty\})^n \\
 \cup & & \cup \\
 V(p) & \xrightarrow{\quad} & \overline{\text{val}_0(V(p))} \\
 \parallel & & \parallel \\
 \{\mathbf{x} : p(\mathbf{x}) = 0\} & & \text{Trop}(V(p))
 \end{array}$$

Example 1.2.7. Consider the polynomial in $K[x, y]$

$$p(x, y) = tx + y + t^2,$$

then the variety is $X = \{(x, y) : tx + y + t^2 = 0\}$ which we can solve to $y = -tx - t^2$.

If we choose $x = 0$ then y becomes $-t^2$. Now we take the valuation of $(0, -t^2)$ and so $(\infty, 2) \in \text{Trop}(X)$.

Amoebas

Let us return to the usual stage and consider $p \in \mathbb{C}[x]$ which defines an algebraic variety $X = V(p) \subseteq \mathbb{C}^n$. Now consider the map which sends every coordinate's modulus to its logarithm in base t :

$$\mathbb{C}^n \rightarrow (\mathbb{R} \cup \{-\infty\})^n, \quad (z_1, \dots, z_n) \rightarrow (\log_t |z_1|, \dots, \log_t |z_n|).$$

The image of X under this map, $\log_t(X)$, is the t -amoeba of X . If we take the limit as $t \rightarrow \infty$ then we get the *spine* of the amoeba.

Example 1.2.8. When $p(x, y) = x + y - 1$ then we can describe $V(p)$ via the parametrization $(x, 1 - x)$. So the corresponding t -amoeba in the real case is


$$\{(\log_t |x|, \log_t |1 - x|) : x \in \mathbb{R}\}$$

and we ordinarily take the limit, we see that the functions converge to zero point-by-point. But the set is actually approaching the spine!


Observe that the spine approaches the tropical hypersurface associated to p . In other words we have that the tropical hypersurface is $\lim_{t \rightarrow \infty} \log_t(V(p))$.

Degenerations

We may parametrize any algebraic variety with a time variable, then converting the information to a graph, edges code the information about how fast the node forms related to the length.



Consider a family of of what, what is this family of?! Stuff? Curve in $P^1 \times P^1$ which eventually becomes P^2 ?



It is too early to understand this point of view. We will set everything up to get to it.


In general, the big idea will be to explore and understand these perspectives in the case of plane curves. We want to show how they are equivalent and then recover classical algebraic geometry results in terms of tropical geometry.

1.3 Interim

Tropical Arithmetics

Minimizing Tolls

Consider a set of cities connected by a network of toll-ways: If we only care about



minimizing toll expenses, what would be the cheapest way to go from one given city to another? Let us record the information as an incidence matrix.

$$M = \begin{pmatrix} 0 & \infty & 2 \\ x & 0 & y \\ \infty & 1 & 0 \end{pmatrix}$$

In this matrix, the rows determine the outbound city, while the columns are the destination. Each entry records the cost of a toll and tolls are considered to be infinite when the road does not exist. We can also think of M as recording the cheapest toll to go from one city to another with at most one move.

But if we wanted to find the cheapest way from one city to another in **two** moves, we could use M^2 with standard matrix multiplication. However we must redefine our basic operations as follows:

$$+ = \min, \quad \cdot = +$$

$$\begin{aligned}
 \begin{pmatrix} 0 & \infty & 2 \\ x & 0 & y \\ \infty & 1 & 0 \end{pmatrix}^2 &= \begin{pmatrix} \min \begin{pmatrix} 0+0 \\ \infty+x \\ 2+\infty \end{pmatrix} & \min \begin{pmatrix} 0+\infty \\ \infty+0 \\ 2+1 \end{pmatrix} & \min \begin{pmatrix} 0+2 \\ \infty+y \\ 2+0 \end{pmatrix} \\ \min \begin{pmatrix} x+0 \\ 0+x \\ y+\infty \end{pmatrix} & \min \begin{pmatrix} x+\infty \\ 0+0 \\ y+1 \end{pmatrix} & \min \begin{pmatrix} x+2 \\ 0+y \\ y+0 \end{pmatrix} \\ \min \begin{pmatrix} \infty+0 \\ 1+x \\ 0+\infty \end{pmatrix} & \min \begin{pmatrix} \infty+\infty \\ 1+0 \\ 0+1 \end{pmatrix} & \min \begin{pmatrix} \infty+2 \\ 1+y \\ 0+0 \end{pmatrix} \end{pmatrix} \\
 &= \begin{pmatrix} 0 & 3 & 2 \\ x & \min(0, y+1) & \min(x+2, y) \\ 1+x & 1 & \min(0, 1+y) \end{pmatrix}.
 \end{aligned}$$


Observe that $1 + y$ can be the minimum in the diagonal when we allow *negative tolls*.

Remark 1.3.1. If we disallow negative tolls, the products M^n eventually stabilize to a matrix whose entries record the cheapest way to get from one city to another in n steps.

This gives us an intuition that minimization problems correspond to linear algebra problems over $(\mathbb{T}, +, \cdot)$ which is precisely $(\mathbb{R} \cup \{\infty\}, \min, +)$.

Forgetting phases

Recall that any complex number can be written as $z = re^{i\theta}$ where $r \geq 0$. Consider the map $T_t : \mathbb{C} \rightarrow \{-\infty\} \cup \mathbb{R}$, $z \mapsto \log_t(r)$. This map is surjective, and this we can see by



checking it is right-invertible. Observe that:

$$\begin{cases} T_t^{-1}(x) = \{ t^x e^{i\theta} \} \subseteq \mathbb{C}, & \text{for } x \in \mathbb{R}, \\ T_t^{-1}(-\infty) = 0. \end{cases}$$

With this in hand, we wish to define an exotic addition and multiplication on $\{-\infty\} \cup \mathbb{R}$ using T_t . We will dequantize!

1. COMBINATORIAL SHADOW OF ALGEBRAIC GEOMETRY

We begin with **hyper-addition**, the output will be a subset of $\{-\infty\}$ so it's not a binary operation by itself.

$$x \diamondsuit_t y := T(T_t^{-1}(x) + T_t^{-1}(y)) = [\log_t(|t^x - t^y|), \log_t(t^x + t^y)].$$

This is an interval in $\{-\infty\} \cup \mathbb{R}$, in order to make \diamondsuit_t into an operation we take a limit:

$$\begin{array}{ccc} x \diamondsuit_t y & \xrightarrow{\lim_{t \rightarrow \infty}} & x \diamondsuit y = \lim_{t \rightarrow \infty} x \diamondsuit_t y \\ \downarrow \max & & \downarrow \max \\ x +_t y & \xrightarrow{\lim_{t \rightarrow \infty}} & x + y = \max(x, y) \end{array}$$

1.4 Day 3 | 20230825

Index

dual graph, 3

Puiseux series, 5

tropical hypersurface, 4

tropical semifield, 4

Bibliography

- [1] D. Maclagan and B. Sturmfels. *Introduction to Tropical Geometry*. Graduate Studies in Mathematics. American Mathematical Society, 2021.