

Exercise 1. Consider a region R bounded by the curves

$$y=x, \quad y=-x, \quad \text{and} \quad x=1,$$

additionally the region has density $\rho(y)=e^y$. Now suppose we rotate the region about the axis $y=-4$. Do the following:

- i) Draw the region in question.
- ii) Draw the solid of revolution obtained after rotation.
- iii) Which method should we use to find the volume of this shape?
- iv) Find the bounds of the region. Label them either as $a \leq x \leq b$ or $c \leq y \leq d$.
- v) Find the **GREATER** and **LOWER** curves by writing their equations.
- vi) Find the parameters (*either R, r or r, h*) used to build your area function.
- vii) With the previous information, write out the integral which represents the mass of the solid obtained.

i) See diagram.

ii) See diagram.

iii) As we have a rotation about a $y=b$ line, we can either use shells in y or rings in x . The density is in terms of y so we should use shells in y .

iv) The regions starts at $y=-1$ and ends at $y=1$. So $-1 \leq y \leq 1$.

v) We must divide the region into two pieces:

$$\begin{cases} \text{GREATER: } x=1 \\ \text{LOWER1: } x=-y \text{ for } -1 \leq y \leq 0 \end{cases} \quad \text{and} \quad \begin{cases} \text{GREATER: } x=1 \\ \text{LOWER2: } x=y \text{ for } 0 \leq y \leq 1 \end{cases}$$

vi) As we are using shells we must find r, h . These are:

$$\begin{cases} h_1 = \text{GREATER} - \text{LOWER1} = [1 - (-y)] & \text{for } -1 \leq y \leq 0 \\ h_2 = \text{GREATER} - \text{LOWER2} = [1 - (y)] & \text{for } 0 \leq y \leq 1 \\ r = \text{dist}(\text{axis}, \text{coordinate}) = y - (-4) \end{cases}$$

vii) Adding this facts together we get

$$m = \int_{-1}^0 2\pi(y+4)(1+y)e^y dy + \int_0^1 2\pi(y+4)(1-y)e^y dy.$$

Exercise 2. Consider the tank formed after rotating the curve $y=x^3$ with $0 \leq x \leq 1$ about the axis $x=0$. Suppose tank is filled with *radioactive waste* with density $\rho(y)=100+25y^2$. Do the following:

