

Exercise 1. Define a *line* in \mathbb{P}^2 to be a closed subset of the form $L = \{ [x : y : z] : ax + by + cz = 0 \}$ for some constants $a, b, c \in \mathbb{C}$, not all zero.

- i) If $(a, b, c) = (1, 0, 0)$, we saw in class that $\mathbb{P}^2 \setminus L = \{ [x : y : z] : x \neq 0 \} = U_x$ could be identified with \mathbb{C}^2 .

Similarly, show that for any line L there is a bijection $\mathbb{P}^2 \setminus L \simeq \mathbb{C}^2$.

- ii) Prove that any two distinct lines L_1 and L_2 intersect in a single point.
 iii) Prove that there is a unique line L through any two distinct points in \mathbb{P}^2 .

Answer

- i) constants
 ii) Let $L_1, L_2 \subseteq \mathbb{P}^2$ be two distinct lines with direction (a, b, c) and (d, e, f) . Since they are distinct this means that $\nexists \lambda((d, e, f) = \lambda(a, b, c))$. A point $[x : y : z]$ in the intersection of L_1 and L_2 must satisfy the system of equations

$$\begin{cases} ax + by + cz = 0, \\ dx + ey + fz = 0. \end{cases}$$

Solutions to this system of equations are parametrized in terms of z in the following manner

$$[x : y : z] = \left[\frac{bf - ce}{ae - bd}z : \frac{cd - af}{ae - bd}z : z \right],$$

and ordinarily this would give us an infinite number of solutions. However in \mathbb{P}^2 this corresponds to the point $[bf - ce : cd - af : ae - bd]$.

- iii) Let us now consider two points $[x : y : z], [u : v : w] \in \mathbb{P}^2$ which are distinct. Once again, consider a system of equations

$$\begin{cases} ax + by + cz = 0, \\ au + bv + cw = 0. \end{cases}$$

There is an infinite number of solutions to this system for $(a, b, c) \in \mathbb{C}^3$.aaaaaaa

Exercise 2. Consider the sequence $(p_n)_{n \in \mathbb{N}} \subseteq \mathbb{C}^3$ with $p_n = (n^3, 2n^2, 3n^3)$. Identifying \mathbb{C}^3 with $\{x_0 \neq 0\} \subseteq \mathbb{P}^3$, what is the limit of p_n as $n \rightarrow \infty$?

Answer

We can identify p_n with the sequence $\tilde{p}_n = [n^3 : 2n^2 : 3n^3 : 1]$. Now for $n \neq 0$ it holds that

$$\tilde{p}_n = \left[1 : \frac{2}{n} : 3 : \frac{1}{n^3} \right] \xrightarrow{n \rightarrow \infty} [1 : 0 : 3 : 0].$$

This coincides with the limit of p_n in the usual sense which is ∞ and $[1 : 0 : 3 : 0]$ is a point at infinity.

Exercise 3. In \mathbb{A}^2 , let $V = \mathbb{V}(x)$, $W = \mathbb{V}(x - 1)$ and $Z = \mathbb{V}(y - x^2)$. Let \overline{V} , \overline{W} and \overline{Z} denote their respective *projective closures* in \mathbb{P}^2 . Find the points in the intersections $\overline{V} \cap \overline{W}$, $\overline{V} \cap \overline{Z}$ and $\overline{W} \cap \overline{Z}$.

Answer

First, let us parametrize the varieties in question as points of \mathbb{A}^2 :

$$\begin{cases} \mathbb{V}(x) = \{x = 0\} = \{(0, t) : t \in \mathbb{C}\}, \\ \mathbb{V}(x - 1) = \{x = 1\} = \{(1, t) : t \in \mathbb{C}\}, \\ \mathbb{V}(y - x^2) = \{y = x^2\} = \{(t, t^2) : t \in \mathbb{C}\}. \end{cases}$$

For each one of those sets, their projective closure corresponds to the embedding of the points inside \mathbb{P}^2 along with their limit points. In the case of V we have

$$\overline{V} = \{[0 : t : 1] : t \in \mathbb{C}\} \cup \{\text{limit points}\} = \{[0 : t : 1] : t \in \mathbb{C}\} \cup \{[0 : 1 : 0]\}.$$