

Modern Physics Letters A
© World Scientific Publishing Company

Holographic dark energy models in $f(Q, T)$ gravity

Xuwei Zhang*

*University Department, University Name, Address
City, State ZIP/Zone, Country[†]
author@emailaddress*

Second Author

*Group, Laboratory, Address
City, State ZIP/Zone, Country*

Received (Day Month Year)

Revised (Day Month Year)

The abstract should summarize the context, content and conclusions of the paper in less than 200 words. It should not contain any references or displayed equations. Typeset the abstract in 8 pt Times Roman with baselineskip of 10 pt, making an indentation of 1.5 pica on the left and right margins.

Keywords: Keyword1; keyword2; keyword3.

PACS Nos.: include PACS Nos.

1. Introduction

Over the past decades, a series of discoveries in cosmology have profoundly changed our understanding of the universe. In 1998, the accelerated expansion of the universe was discovered through the study of Type Ia supernovae^(?,?). This fact had been later confirmed by many other cosmological observations, such as the measurement of temperature anisotropy and polarization in the cosmic microwave background (CMB) radiation^(?,?); Baryon acoustic oscillations (Baryon Acoustic Oscillations, BAO) peak length scale^(?,?); the study of the large-scale structure (LSS) of the universe^(?,?) and use Cosmic Chronometers to direct measurement of Hubble parameter^(?,?). These observations suggest the existence of a mysterious energy in our universe who has high negative pressure and increasing density.

*Typeset names in 8 pt Times Roman. Use the footnote to indicate the present or permanent address of the author.

[†]State completely without abbreviations, the affiliation and mailing address, including country and e-mail address. Typeset in 8 pt Times Italic.

2. $f(Q, T)$ gravity theory and Holographic dark energy

In Weyl-Cartan geometry the connection can be decomposed into three parts: the Christoffel symbol $\hat{\Gamma}^\alpha_{\mu\nu}$, the contortion tensor $K^\alpha_{\mu\nu}$ and the disformation tensor $L^\alpha_{\mu\nu}$, so that the general affine connection can be expressed as

$$\Gamma^\alpha_{\mu\nu} = \hat{\Gamma}^\alpha_{\mu\nu} + K^\alpha_{\mu\nu} + L^\alpha_{\mu\nu} \quad (1)$$

The first term $\hat{\Gamma}^\alpha_{\mu\nu}$ is the Levi-civita connection of metric $g_{\mu\nu}$, given by

$$\hat{\Gamma}^\alpha_{\mu\nu} = \frac{1}{2}g^{\alpha\beta}(\partial_\mu g_{\beta\nu} + \partial_\nu g_{\beta\mu} - \partial_\beta g_{\mu\nu}) \quad (2)$$

The second term $K^\alpha_{\mu\nu}$ is the contortion tensor

$$K^\alpha_{\mu\nu} = \frac{1}{2}T^\alpha_{\mu\nu} + T_{(\mu}{}^\alpha{}_{\nu)} \quad (3)$$

The Last term is distortion tensor $L^\alpha_{\mu\nu}$, given by

$$L^\alpha_{\mu\nu} = \frac{1}{2}Q^\alpha_{\mu\nu} - Q_{(\mu}{}^\alpha{}_{\nu)} \quad (4)$$

Enclose $Q^\alpha_{\mu\nu}$ with nonmetricity tensor

$$Q_{\rho\mu\nu} \equiv \nabla_\rho g_{\mu\nu} = \partial_\rho g_{\mu\nu} - \Gamma^\beta_{\rho\mu} g_{\beta\nu} - \Gamma^\beta_{\rho\nu} g_{\mu\beta} , \quad (5)$$

$$T^\lambda_{\mu\nu} \equiv \Gamma^\lambda_{\mu\nu} - \Gamma^\lambda_{\nu\mu} , \quad (6)$$

$$R^\sigma_{\rho\mu\nu} \equiv \partial_\mu \Gamma^\sigma_{\nu\rho} - \partial_\nu \Gamma^\sigma_{\mu\rho} + \Gamma^\alpha_{\nu\rho} \Gamma^\sigma_{\mu\alpha} - \Gamma^\alpha_{\mu\rho} \Gamma^\sigma_{\nu\alpha} , \quad (7)$$

In weyl geometry, affine connection is not compatible with the metric tensor as

$$Q_{\alpha\mu\nu} = \nabla_\alpha g_{\mu\nu} = -w_\alpha g_{\mu\nu} \quad (8)$$

In gravity, we consider the action is

$$S = \int \left(\frac{1}{2}f(Q, T) + \mathcal{L}_m \right) \sqrt{-g} d^4x \quad (9)$$

where f is an arbitrary function of the non-metricity and T is the trace of the matter-energy-momentum tensor, \mathcal{L}_m is known as matter Lagrangian and $g = \det(g_{\mu\nu})$ denotes determinant of metric tensor.

Vary the action we can get

$$\delta S = \int \left(\frac{1}{2}\delta[f(Q, T)\sqrt{-g}] + \delta(\mathcal{L}_m\sqrt{-g}) \right) d^4x \quad (10)$$

$$= \int \frac{1}{2} \left(-\frac{1}{2}f g_{\mu\nu} \sqrt{-g} \delta g^{\mu\nu} + f_Q \sqrt{-g} \delta Q + f_T \sqrt{-g} \delta T - \frac{1}{2}T_{\mu\nu} \sqrt{-g} \delta g^{\mu\nu} \right) d^4x \quad (11)$$

Field equation is

$$-\frac{2}{\sqrt{-g}}\nabla_\alpha(f_Q\sqrt{-g}P^\alpha_{\mu\nu}) - \frac{1}{2}f g_{\mu\nu} + f_T(T_{\mu\nu} + \Theta_{\mu\nu}) - f_Q(P_{\mu\alpha\beta}Q^\nu{}_{\alpha\beta} - 2Q^{\alpha\beta}{}_{\mu}P_{\alpha\beta\nu}) = T_{\mu\nu} \quad (12)$$

In FLRW metric, given by

$$ds^2 = -dt^2 + a^2(t)\delta_{ij}dx^i dx^j \quad (13)$$

then we can get Friedmann equations

$$\rho = \frac{f}{2} - 6f_Q H^2 - \frac{2f_T}{1+f_T}(\dot{f}_Q H + f_Q \dot{H}) \quad (14)$$

$$p = -\frac{f}{2} + 6f_Q H^2 + 2(\dot{f}_Q H + f_Q \dot{H}) \quad (15)$$

EoS parameter

$$w = \frac{p}{\rho} = -1 + \frac{4f_Q H + f_Q \dot{H}}{(1+f_T)(f - 12f_Q H^2) - 4f_T(\dot{f}_Q H + f_Q \dot{H})} \quad (16)$$

In our universe $w < -1/3$, where ρ and p denote total fluid energy density and pressure of the universe, which $\rho = \rho_m + \rho_{de}$, $p = p_m + p_{de}$.

Effective EoS parameter denote geometry qualities.

$$\rho_{\text{eff}} = 3H^2 = \frac{f}{4f_Q} - \frac{1}{2f_Q}[(1+f_T)\rho + f_T p] \quad (17)$$

$$-p_{\text{eff}} = 2\dot{H} + 3H^2 = \frac{f}{4f_Q} - \frac{2\dot{f}_Q H}{f_Q} + \frac{1}{2f_Q}[(1+f_T)\rho + (2+f_T)p] \quad (18)$$

EoS parameter for equivalent dark energy

$$w_{\text{eff}} = \frac{p_{\text{eff}}}{\rho_{\text{eff}}} = -\frac{f - 8\dot{f}_Q H + 2[(1+f_T)\rho + (2+f_T)p]}{f - 2[(1+f_T)\rho + f_T p]} \quad (19)$$

Combine, we can get the evolution equation of Hubble parameter H Deceleration parameter

$$q = -\frac{\ddot{a}a}{\dot{a}^2} = \frac{1}{2}(1+3w) = \frac{1}{2}\left(1+3\frac{p_{\text{eff}}}{\rho_{\text{eff}}}\right) = -1 + \frac{3(4\dot{f}_Q H - f + p)}{f - [(1+f_T)\rho + f_T p]} \quad (20)$$

We assume that

$$f(Q, T) = \alpha Q^n + \beta T \quad (21)$$

where $Q = 6H^2$, $T = -\rho + 3p$, so that we can derive $f_Q = \alpha n Q^{n-1} = \alpha n 6^{n-1} H^{2n-2}$, $f_T = \beta$, $\dot{f}_Q = 2\alpha n(n-1)6^{n-1} H^{2n-3} \dot{H}$

3. Cosmic solutions

Hybrid Expansion Law (HEL)

$$H(z) = \frac{H_0}{\sqrt{2}} \sqrt{1 + (1+z)^{2m}} \quad (22)$$

$$\dot{H}(z) = \frac{H_0}{\sqrt{2}} \frac{2m(1+z)^{2m-1}}{\sqrt{1 + (1+z)^{2m}}} \quad (23)$$

4 *Authors' names*

use deceleration parameter to simplify calculation

$$\dot{H} = -H^2(1+q) \quad (24)$$

get

$$\rho = \frac{f}{2} - 2f_Q(-H^2(1+q) + 3H^2 + \frac{f_T}{1+f_T} \frac{\dot{f}_Q}{f_Q} H) \quad (25)$$

$$p = -\frac{f}{2} + 2f_Q(-H^2(1+q) + 3H^2 + \frac{\dot{f}_Q}{f_Q} H) \quad (26)$$

Holographic dark energy with Hubble cutoff is

$$\rho_{de} = 3c^2 H^2 \quad (27)$$

$$p_{de} = \frac{\alpha (6H(z)^2)^{n-1} \left(2n(-2\beta + (3\beta + 2)n - 1)\dot{H}(z) + 3(\beta + 1)(2n - 1)H(z)^2 \right)}{(\beta + 1)(2\beta + 1)} \quad (28)$$

dark energy EoS parameter

$$w_{de} = \frac{p_{de}}{\rho_{de}} = \frac{\alpha (6H(z)^2)^{n-1} \left(2n(-2\beta + (3\beta + 2)n - 1)\dot{H}(z) + 3(\beta + 1)(2n - 1)H(z)^2 \right)}{3c^2 H^2 (\beta + 1)(2\beta + 1)} \quad (29)$$

where $\dot{H}(z) = \frac{d}{dt} H(z) = -\frac{dH(z)}{dz} H(z)(1+z)$

stability parameter of in this situation can be described as

$$c_s^2 = \frac{dp_{de}}{d\rho_{de}} = \frac{\alpha(2n(-2\beta + (3\beta + 2)n - 1))6^{n-1}((2n - 2)H^{2n-3}\dot{H}^2 + \ddot{H}H^{2n-2}) + \alpha 3(\beta + 1)(2n - 1)6^{n-1}2nH^{2n}}{6c^2 H \dot{H} (\beta + 1)(2\beta + 1)} \quad (30)$$

4. Tsallis entropy dark energy in different IR cutoff

Tsallis proposed a modified black hole entropy

$$S_\delta = \gamma A^\delta \quad (31)$$

Tsallis holographic dark energy density

$$\rho_{de} = BH^{4-2\delta} \quad (32)$$

5. Age of the universe

6. Observational data and constraint

7. Conclusion

Appendix A. Appendix information