




EleutherAI Math Reading Group

Differential Geometry I: Manifolds



Motivation

Manifold hypothesis







Latent coordinates

e.g. Autoencoders,
GANs, ...

↳ Space that we
can move around
in


"locally linear"

Implicit vs Explicit equations




$$t \mapsto (r \cos t, r \sin t)$$

Coordinate Systems



Cartesian coordinates




Polar coordinates

$$\begin{cases} r = \sqrt{x^2 + y^2} \\ \theta = \arctan\left(\frac{y}{x}\right) \quad [\text{atan2}(y, x)] \end{cases}$$




θ undefined at $(0, 0)$



Most coordinate systems are
only defined locally

Definition: An n -dimensional manifold M is a collection of points together with a collection of coordinate systems $\{\psi_i : M_i \rightarrow U_i\}_{i \in I}$ [Atlas] [charts]


- ① Each coordinate system ψ_i is a bijective map between $U_i \subset \mathbb{R}^n$ and $M_i \subset M$



- ② The map $\psi_i \circ \psi_j^{-1}$ is differentiable on $\psi_j(M_i \cap M_j)$

- ③ Every point $p \in M$ is in some M_i

$$\simeq \bigcup_{i \in I} M_i = M$$



Example

Circle $S^1 = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 = 1\}$

- $\Psi_1: M_1 \rightarrow (-1, 1)$ $\Psi_3: M_3 \rightarrow (-1, 1)$


$$(x, y) \mapsto x$$

$$(x, y) \mapsto y$$

- $\Psi_2: M_2 \rightarrow (-1, 1)$ $\Psi_4: M_4 \rightarrow (-1, 1)$

$$(x, y) \mapsto x$$

$$(x, y) \mapsto y$$




- $\phi_1: N_1 \rightarrow (-\pi, \pi)$

$$(x, y) \mapsto \text{atan}(\text{y}, \text{x})$$

- $\phi_2: N_2 \rightarrow (-\pi, \pi)$

$$(x, y) \mapsto \text{atan}(\text{y}, -\text{x})$$




Manifolds are topological spaces

- A neighbourhood of $p_0 \in M$ is a subset $\mathcal{N} \subset M$ s.t

$\forall p \in \mathcal{N}, \|\psi_i^{-1}(p) - \psi_i^{-1}(p_0)\| < \varepsilon \Rightarrow p \in \mathcal{N}$
for some $\varepsilon > 0$

[NB: ε depends on the chart,
but the property of being a
neighborhood does not]



- A subset of M is open if it contains a neighbourhood of each of its points

→ "open sets" define a topology on M

Product Manifold

Let M, N be (m/n) -manifolds, then

$M \times N = \{(p, q) \mid p \in M, q \in N\}$ is a manifold with
the coordinate system $(x^1, \dots, x^m, y^1, \dots, y^n)$


$$\text{with } (x^1, \dots, x^m)(p) = \psi(p) \quad p \in M$$

$$(y^1, \dots, y^n)(q) = \phi(q) \quad q \in N$$

Example:

$$S^1 \times S^1 \cong T^2$$

$$\begin{array}{c} \text{Blue circle} \\ \times \\ \text{Orange circle} \end{array} \cong$$



Example : Nelson's car

Configuration Space of a car


$$M = \mathbb{R}^2 \times T^2$$



(x, y)
position

(θ, ϕ)

heading \rightarrow "absolute
heading"



More generally, any kind of industrial robot



Tangent vectors

$$\varphi: [a, b] \rightarrow M$$

differentiable curve $\varphi(t)$ on a manifold M

+ coordinate system (x^i) around $\varphi(t)$


$$\rightarrow (\xi^i) = (\dot{x}^i(t)) = \left(\frac{d}{dt} x^i \circ \varphi \right)(t) \quad \text{Coordinate tangent vector}$$

change of coordinates: $\tilde{x} = f(x(p))$

$$\tilde{\xi}^j = \left(\frac{\partial \tilde{x}^j}{\partial x^i} \right)_p \xi^i \quad \star$$

↳ Jacobian

A tangent vector v at p is any object represented by some (ξ^i) which transforms according to \star



Tangent Space



The set of all vectors around φ is a vector space

$$T_\varphi(M)$$

[\approx infinitesimal displacements
around φ]

Given a coordinate system (x^i) around φ , we can
construct a basis of $T_\varphi M$

$$e_k = \left(\frac{\partial}{\partial x^k} \right)_\varphi$$




Example : Polar coordinates

$$(x, y) = (r \cos \theta, r \sin \theta)$$

$$r = \sqrt{x^2 + y^2}$$

$$\theta \Rightarrow \operatorname{atan}(y, x) = \operatorname{atan}\left(\frac{y}{x}\right)$$



$$\frac{\partial}{\partial r} = \frac{\partial x}{\partial r} \frac{\partial}{\partial x} + \frac{\partial y}{\partial r} \frac{\partial}{\partial y} = \cos \theta \frac{\partial}{\partial x} + \sin \theta \frac{\partial}{\partial y}$$

$$\frac{\partial}{\partial \theta} = \frac{\partial x}{\partial \theta} \frac{\partial}{\partial x} + \frac{\partial y}{\partial \theta} \frac{\partial}{\partial y} = -r \sin \theta \frac{\partial}{\partial x} + r \cos \theta \frac{\partial}{\partial y}$$

$$\sim \begin{bmatrix} \frac{\partial}{\partial r} \\ \frac{\partial}{\partial \theta} \end{bmatrix} = \begin{bmatrix} \cos \theta & \sin \theta \\ -r \sin \theta & r \cos \theta \end{bmatrix} \begin{bmatrix} \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \end{bmatrix}$$

Differentials

Let $f: M \rightarrow N$ be a differentiable map between manifolds

$p \in M$, (x^i) a coordinate system around p
 (y^j) coordinates around $f(p)$


f induces a map $T_p M \rightarrow T_{f(p)} N$
linear

$$\xi^i \frac{\partial}{\partial x^i} \mapsto \eta^j \frac{\partial}{\partial y^j}$$

with

$$\eta^j = \left(\frac{\partial y^j}{\partial x^i} \right) \xi^i = \underline{d f_p \xi^i}$$

Moreover, this map is independent
of the coordinate systems



↳ differential / Jacobian of f
at p

Covectors & 1-forms

Let $f: M \rightarrow \mathbb{R}$ be a differentiable map, then

$df_p: T_p M \rightarrow \mathbb{R}$ is a linear form on $T_p M$

i.e. $df_p \in T_p M^*$ [cotangent space]

In particular, the coordinate differentials dx^i at p satisfy $dx^i \frac{\partial}{\partial x^j} = f_j^i \Rightarrow \underline{(dx^i)}$ is the dual basis of $\underline{\frac{\partial}{\partial x^i}}$

Change of coordinates: $w = \eta_i dx^i$


$$\tilde{\eta}_i = \left(\frac{\partial x^i}{\partial \tilde{x}^j} \right)_p \eta_j \quad [\text{constant}]$$

inverse Jacobian

Riemannian metrics

Since manifolds are usually "embedded" in Euclidean Space

We can inherit a metric on the manifold



- length of a curve $\rho: [a, b] \rightarrow \mathbb{R}^n$:
$$\int_a^b \sqrt{\sum_{i=1}^n \left(\frac{dx^i}{dt} \right)^2} dt$$


"length of tangent at curve"
- Intuition: take the shortest paths between points on the manifold

Examples and notations : \mathbb{R}^3

$$ds = \sqrt{dx^2 + dy^2 + dz^2}$$

length element

$$ds^2 = dx^2 + dy^2 + dz^2$$



cylindrical
coordinates

$$ds^2 = dr^2 + r^2 d\theta^2 + dz^2$$

spherical
coordinates

$$ds^2 = d\ell^2 + \ell^2 \sin^2 \phi d\theta^2 + d\phi^2$$

Sphere :

$$ds^2 = R^2 \sin^2 \phi d\theta^2 + d\phi^2$$

Riemann metric

A Riemann metric on M is a map $g: M \rightarrow ((T_p M)^2 \rightarrow \mathbb{R})$

which associates to each point of M a non-degenerate symmetric bilinear form on $T_p M$

$$\rightarrow ds^2 = g_{ij} dx^i dx^j \quad [\text{quadratic form}]$$

$$g_{ij} = g\left(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial x^j}\right)$$

\rightarrow Can compute norms of tangent vectors!

[also inner-products]


N.B. This is a covariant tensor

$$\xi^i \frac{\partial}{\partial x^i} \cdot \eta^j \frac{\partial}{\partial x^j} = g_{ij} \xi^i \eta^j = \langle \xi, \eta \rangle$$

Distance between two points

let M be a manifold, $p, q \in M$

- a path between p and q is a curve $\gamma: [0, 1] \rightarrow M$
- M is path connected if there is a path between any two points $\gamma(0) = p$, $\gamma(1) = q$
- length of a path $\gamma: \int_0^1 L(\gamma, \dot{\gamma}) dt$
- where $L(\gamma, \dot{\gamma}) = \sqrt{g_{ij} \frac{dx^i}{dt} \frac{dx^j}{dt}}$
- $\gamma = (x^1, \dots, x^n)$
- Distance between p and q = $\inf_{\gamma} \int_0^1 L(\gamma, \dot{\gamma}) dt$



Tangent vectors

$$\varphi: [a, b] \rightarrow M$$

differentiable curve $\varphi(t)$ on a manifold M

+ coordinate system (x^i) around $\varphi(t)$


$$\rightarrow (\xi^i) = (\dot{x}^i(t)) = \left(\frac{d}{dt} x^i \circ \varphi \right)(t) \quad \text{Coordinate tangent vector}$$

change of coordinates: $\tilde{x} = f(x(p))$

$$\tilde{\xi}^j = \left(\frac{\partial \tilde{x}^j}{\partial x^i} \right)_p \xi^i \quad \star$$

↳ Jacobian

A tangent vector v at p is any object represented by some (ξ^i) which transforms according to \star



Tangent Space



The set of all vectors around φ is a vector space

$$T_\varphi(M)$$

[\approx infinitesimal displacements
around φ]

Given a coordinate system (x^i) around φ , we can
construct a basis of $T_\varphi M$

$$e_k = \left(\frac{\partial}{\partial x^k} \right)_\varphi$$




Example : Polar coordinates

$$(x, y) = (r \cos \theta, r \sin \theta)$$

$$r = \sqrt{x^2 + y^2}$$

$$\theta \Rightarrow \operatorname{atan}(y, x) = \operatorname{atan}\left(\frac{y}{x}\right)$$



$$\frac{\partial}{\partial r} = \frac{\partial x}{\partial r} \frac{\partial}{\partial x} + \frac{\partial y}{\partial r} \frac{\partial}{\partial y} = \cos \theta \frac{\partial}{\partial x} + \sin \theta \frac{\partial}{\partial y}$$

$$\frac{\partial}{\partial \theta} = \frac{\partial x}{\partial \theta} \frac{\partial}{\partial x} + \frac{\partial y}{\partial \theta} \frac{\partial}{\partial y} = -r \sin \theta \frac{\partial}{\partial x} + r \cos \theta \frac{\partial}{\partial y}$$

$$\sim \begin{bmatrix} \frac{\partial}{\partial r} \\ \frac{\partial}{\partial \theta} \end{bmatrix} = \begin{bmatrix} \cos \theta & \sin \theta \\ -r \sin \theta & r \cos \theta \end{bmatrix} \begin{bmatrix} \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \end{bmatrix}$$

Differentials

Let $f: M \rightarrow N$ be a differentiable map between manifolds

$p \in M$, (x^i) a coordinate system around p
 (y^j) coordinates around $f(p)$


f induces a map $T_p M \rightarrow T_{f(p)} N$
linear

$$\xi^i \frac{\partial}{\partial x^i} \mapsto \eta^j \frac{\partial}{\partial y^j}$$

with

$$\eta^j = \left(\frac{\partial y^j}{\partial x^i} \right) \xi^i = \underline{d f_p \xi^i}$$

Moreover, this map is independent
of the coordinate systems



↳ differential / Jacobian of f
at p

Covectors & 1-forms

Let $f: M \rightarrow \mathbb{R}$ be a differentiable map, then

$df_p: T_p M \rightarrow \mathbb{R}$ is a linear form on $T_p M$

i.e. $df_p \in T_p M^*$ [cotangent space]

In particular, the coordinate differentials dx^i at p satisfy $dx^i \frac{\partial}{\partial x^j} = f_j^i \Rightarrow \underline{(dx^i)}$ is the dual basis of $\underline{\frac{\partial}{\partial x^i}}$

Change of coordinates: $w = \eta_i dx^i$


$$\tilde{\eta}_i = \left(\frac{\partial x^i}{\partial \tilde{x}^j} \right)_p \eta_j \quad [\text{constant}]$$

inverse Jacobian

Riemannian metrics

Since manifolds are usually "embedded" in Euclidean Space

We can inherit a metric on the manifold



- length of a curve $\rho: [a, b] \rightarrow \mathbb{R}^n$:
$$\int_a^b \sqrt{\sum_{i=1}^n \left(\frac{dx^i}{dt} \right)^2} dt$$


"length of tangent at curve"
- Intuition: take the shortest paths between points on the manifold

Examples and notations : \mathbb{R}^3

$$ds = \sqrt{dx^2 + dy^2 + dz^2}$$

length element

$$ds^2 = dx^2 + dy^2 + dz^2$$



cylindrical
coordinates

$$ds^2 = dr^2 + r^2 d\theta^2 + dz^2$$

spherical
coordinates

$$ds^2 = d\ell^2 + \ell^2 \sin^2 \phi d\theta^2 + d\phi^2$$

Sphere :

$$ds^2 = R^2 \sin^2 \phi d\theta^2 + d\phi^2$$

Riemann metric

A Riemann metric on M is a map $g: M \rightarrow ((T_p M)^2 \rightarrow \mathbb{R})$

which associates to each point of M a non-degenerate symmetric bilinear form on $T_p M$

$$\rightarrow ds^2 = g_{ij} dx^i dx^j \quad [\text{quadratic form}]$$

$$g_{ij} = g\left(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial x^j}\right)$$

\rightarrow Can compute norms of tangent vectors!

[also inner-products]


N.B. This is a covariant tensor

$$\xi^i \frac{\partial}{\partial x^i} \cdot \eta^j \frac{\partial}{\partial x^j} = g_{ij} \xi^i \eta^j = \langle \xi, \eta \rangle$$

Distance between two points



let M be a manifold, $p, q \in M$

- a path between p and q is a curve $\gamma: [0, 1] \rightarrow M$
- M is path connected if there is a path between any two points $\gamma(0) = p$, $\gamma(1) = q$
- length of a path $\gamma: \int_0^1 L(\gamma, \dot{\gamma}) dt$
- where $L(\gamma, \dot{\gamma}) = \sqrt{g_{ij} \frac{dx^i}{dt} \frac{dx^j}{dt}}$
- $\gamma = (x^1, \dots, x^n)$
- Distance between p and q = $\inf_{\gamma} \int_0^1 L(\gamma, \dot{\gamma}) dt$




EleutherAI Math Reading Group

Differential Geometry I: Manifolds



Motivation

Manifold hypothesis







Latent coordinates

e.g. Autoencoders,
GANs, ...

↳ Space that we
can move around
in


"locally linear"

Implicit vs Explicit equations




$$t \mapsto (r \cos t, r \sin t)$$

Coordinate Systems



Cartesian coordinates




Polar coordinates

$$\begin{cases} r = \sqrt{x^2 + y^2} \\ \theta = \arctan\left(\frac{y}{x}\right) \quad [\text{atan2}(y, x)] \end{cases}$$




θ undefined at $(0, 0)$



Most coordinate systems are
only defined locally

Definition: An n -dimensional manifold M is a collection of points together with a collection of coordinate systems $\{\psi_i : M_i \rightarrow U_i\}_{i \in I}$ [Atlas] [charts]


- ① Each coordinate system ψ_i is a bijective map between $U_i \subset \mathbb{R}^n$ and $M_i \subset M$



- ② The map $\psi_i \circ \psi_j^{-1}$ is differentiable on $\psi_j(M_i \cap M_j)$

- ③ Every point $p \in M$ is in some M_i

$$\simeq \bigcup_{i \in I} M_i = M$$



Example

Circle $S^1 = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 = 1\}$

- $\Psi_1: M_1 \rightarrow (-1, 1)$ $\Psi_3: M_3 \rightarrow (-1, 1)$


$$(x, y) \mapsto x$$

$$(x, y) \mapsto y$$

- $\Psi_2: M_2 \rightarrow (-1, 1)$ $\Psi_4: M_4 \rightarrow (-1, 1)$

$$(x, y) \mapsto x$$

$$(x, y) \mapsto y$$




- $\phi_1: N_1 \rightarrow (-\pi, \pi)$

$$(x, y) \mapsto \text{atan}(\text{y}, \text{x})$$

- $\phi_2: N_2 \rightarrow (-\pi, \pi)$

$$(x, y) \mapsto \text{atan}(\text{y}, -\text{x})$$




Manifolds are topological spaces

- A neighbourhood of $p_0 \in M$ is a subset $N \subset M$ s.t

$\forall p \in N, \|\psi_i^{-1}(p) - \psi_i^{-1}(p_0)\| < \varepsilon \Rightarrow p \in N$
for some $\varepsilon > 0$

[NB: ε depends on the chart,
but the property of being a
neighborhood does not]



- A subset of M is open if it contains a neighbourhood of each of its points

→ "open sets" define a topology on M

Product Manifold

Let M, N be (m/n) -manifolds, then

$M \times N = \{(p, q) \mid p \in M, q \in N\}$ is a manifold with
the coordinate system $(x^1, \dots, x^m, y^1, \dots, y^n)$


$$\text{with } (x^1, \dots, x^m)(p) = \psi(p) \quad p \in M$$

$$(y^1, \dots, y^n)(q) = \phi(q) \quad q \in N$$

Example:

$$S^1 \times S^1 \cong T^2$$

$$\begin{array}{c} \text{Blue circle} \\ \times \\ \text{Orange circle} \end{array} \cong$$



Example : Nelson's car

Configuration Space of a car


$$M = \mathbb{R}^2 \times T^2$$



(x, y)
position

(θ, ϕ)

heading \rightarrow "absolute
heading"



More generally, any kind of industrial robot

