

Computational analysis of tongue reconstruction surgery: the impact of free flap site and volume on the tongue anatomy and biomechanics

Amir Reza Isazadeh^{1*}, Lindsey Westover², Hadi Seikaly^{3,4} and Daniel Aalto^{1,4}

1 Department of Communication Sciences and Disorders, Faculty of Rehabilitation Medicine, University of Alberta, Edmonton, AB, Canada

2 Department of Mechanical Engineering & Department of Biomedical Engineering, Faculty of Engineering, University of Alberta, Edmonton, AB, Canada

3 Division of Otolaryngology, Department of Surgery, Faculty of Medicine and Dentistry, University of Alberta, Edmonton, AB, Canada

4 Institute for Reconstructive Sciences in Medicine (iRSM), Misericordia Community Hospital, Edmonton, AB, Canada

* Corresponding author, Email: isazadeh@ualberta.ca

Abstract

Tongue reconstruction requires a series of decisions—including donor site and flap volume—tailored to patient needs to restore anatomy and preserve speech and swallowing. The impact of these interdependent choices is difficult to evaluate, as clinical outcomes depend on case-specific factors. This study presents an automated framework that simulates free flap tongue reconstruction and quantifies the anatomical and biomechanical effects of key surgical decisions. The framework leverages biomechanically optimized flap design to generate a multi-component virtual flap, which is then computationally sutured to the resection site, and its long-term tissue atrophy is simulated to predict the final neotongue state. Using this platform, we conducted a 120-simulation factorial study to systematically quantify how key surgical decisions, namely donor site selection (Radial Forearm vs. Anterolateral Thigh) and flap overbulking, influence the final anatomical and biomechanical outcomes. Four clinically

plausible tongue reconstruction scenarios, six levels of flap stiffness, and five levels of flap overbulking were simulated. The results reveal a fundamental, physics-based trade-off: while increasing flap overbulking was the dominant factor in restoring preoperative anatomy ($p < 0.001$), it came at the cost of a predictable increase in biomechanical strain imposed on the native tongue ($p < 0.001$). Furthermore, stiffer flaps induced significantly higher baseline strain ($p < 0.001$), providing a quantitative measure for tissue tethering. The anatomical benefit of overbulking was significantly modulated by tissue properties, proving more effective with more pliable forearm flaps ($p < 0.001$). These findings provide a biomechanical rationale for clinically observed functional trade-offs and support practices like the "beavertail" modification of the forearm flap, which adds volume while preserving pliability. This work presents an open-source robust computational testbed for systematically evaluating interdependent surgical variables. It provides quantitative, physics-based insights into the clinical trade-offs of surgical planning. Ultimately, the framework's automation and scalability offer a pathway toward personalized, simulation-informed surgical planning for tongue reconstruction.

Keywords: Patient-specific surgical simulation, Free flap tongue reconstruction, Finite Element Analysis in surgery, Virtual Surgical Planning, Biomechanical modeling

1 Introduction

Surgical removal of the cancerous defect from the tongue, known as glossectomy, is a primary treatment for tongue cancer. The standard of care to address this is microvascular free flap reconstruction, where tissue is transplanted from a donor site to rebuild the tongue. However, this procedure can severely compromise vital functions such as speech and swallowing [1]. A successful tongue reconstruction is a complex process that relies on a series of interdependent surgical decisions tailored to the individual. The surgeon's strategy is critically influenced by the defect's specific characteristics, including its size, volume, location, and geometry of resection. Key surgical considerations involve donor site selection and free flap design, both its planar geometry and the volume, to restore the tongue's intricate three-dimensional anatomy [2,3]. While recent research has begun to address the challenge of

patient-specific free flap design in terms of its planar geometry via template-based [3–5] or algorithmic approaches [6, 7], other critical decisions that significantly impact surgical outcomes remain largely guided by surgeon experience.

The preservation of residual tongue mobility is the foremost determinant of functional recovery after reconstruction—mobility for speech and swallowing precedes bulk. Consistent with this priority, the Radial Forearm Free Flap (RFFF) is often preferred for defects where pliability and fine shape conformity are paramount: it is thin, pliable, and readily contours to the mobile anterior tongue, supporting articulation [8–10]. At the same time, the Anterolateral Thigh (ALT) flap remains a strong option when substantial bulk is needed, particularly for base-of-tongue defects where palate contact is essential for swallowing, or when forearm donor-site morbidity to wrist function should be avoided, as many patients would prioritize hand function and cosmesis [8, 11–13]. This choice entails a familiar trade-off: the greater stiffness of a bulky ALT flap can tether the residual mobile tongue and impair speech articulation, whereas the more compliant RFFF mitigates tethering but may provide insufficient long-term volume for optimal swallowing [9, 12]. Potential donor-site morbidity further complicates decision-making. This decision is therefore not merely about defect size but is a complex, experience-based judgment on the long-term anatomical and functional consequences. Several studies have investigated the impact of donor site choice on the function and quality of life outcomes [8, 12, 13]. Nevertheless, assessing the impact of donor site selection on the biomechanics and anatomy of the tongue brings granular insights.

A universal challenge in reconstructive surgery is the significant volume loss, or atrophy, that free flaps undergo in the months following transplantation, primarily due to denervation, muscle disuse, and fat resorption. To mitigate this, surgeons intentionally harvest a flap larger than the resection defect, a practice known as “overbulking.” The extent of overbulking is mostly based on the surgeon’s expertise because evidence-based recommendations are limited to a handful of studies and depend on individual circumstances [14–19]. The clinical consequences of miscalculation are severe. Insufficient overbulking can lead to a neotongue with inadequate volume, which severely impairs speech intelligibility and swallowing. Conversely, excessive overbulking can produce a bulky and immobile tongue, leading to difficulties with mastication,

articulation, and even potential airway obstruction [20].

The interdependent choices of donor-site selection and flap overbulking are highly patient-specific, often necessitating intraoperative, experience-based judgment. The choice of donor site impacts not only the provided volume but also the extent of mechanical property mismatch between the donor tissue and the native tongue. Likewise, the decision on the extent of flap overbulking must account for patient-specific variables such as patient anatomy and the extent of the resection, BMI, age, postoperative radiation therapy (RT), and individual healing responses and expected tissue atrophy [17,18]. These decisions are not made in isolation, which creates a complex, case-specific decision matrix.

Designing a prospective randomized clinical trial to systematically isolate these variables presents ethical and methodological challenges [21,22]. Ethically, random assignment of patients to surgical arms that may be considered suboptimal for their specific defect is difficult to justify. Methodologically, the integrity of such a trial would be challenged by numerous confounding variables. Patient-specific factors like tumor extent and location, age, overall health, and healing rates, combined with treatment-related factors like flap design, donor site selection, and radiation dosage, introduce significant variability that can obscure the true effect of the surgical decisions. While controlling for this array of confounders is theoretically possible, it would require a large, expensive, and logistically complex multi-center study, making a traditional clinical investigation a difficult first step.

Biomechanical modeling offers a viable and ethical alternative to investigate these variables and gain insights into their interactions. Buchaillard and colleagues [23,24], for instance, utilized a generic Finite Element tongue model to simulate hemiglossectomy and floor-of-mouth reconstruction by modifying stiffness in affected areas. The approach to modeling surgical removal and free flap reconstruction, though simplified, provided insight into the functional outcome aligned with the clinical understanding. Building on their generic tongue model, several studies introduced multibody dynamics, rigid-soft tissue interactions, advanced hyperelastic and visco-hyperelastic models, and various muscle modeling techniques, all serving to improve the fidelity of the simulation of the complex biomechanics of the oropharynx as accurately as possible [25–28]. Further, to streamline the patient-specific model generation, Harandi et al. [29,30]

developed a method for patient-specific oropharyngeal modeling based on patient MRI data. They used elastic registration to adapt generic tongue and other intraoral models to individual anatomy. Kappert et al. [31] advanced this approach by developing an atlas model that incorporated detailed muscle fiber orientations from Diffusion-weighted MRI (DW-MRI). While effective for assessing patient-related nuances and how they impact function, refinements are needed to implement these techniques in patient-specific surgical intervention analysis.

To our knowledge, a limited number of studies have explored the simulation of surgical interventions on the tongue. Foundational work simulated the functional consequences of glossectomy by altering the material properties of the affected regions to represent the reconstructed tissue [23]. This approach of modifying regional soft tissue properties has since been adapted to investigate other clinical phenomena, such as the biomechanical impact of radiation-induced fibrosis [32]. To advance the simulation of surgical procedures beyond solely property modification, Kappert et al. [33] developed a framework for simulating small resections that are treated with primary closure. Their method replicated the surgical act of suturing by computationally pulling the resection edges together, generating a new finite element model from the deformed geometry, and increasing the stiffness in the scar region.

This study builds on these simulation works by developing an end-to-end patient-specific simulation framework. Such a high-fidelity simulation models the free flap as an independent, multi-component body that is virtually sutured to the resected tongue model. The simulation computationally replicates the surgical workflow, including the attachment and tensioning of sutures to reconstruct the tongue. Furthermore, it incorporates clinically-reported biological phenomena, such as tissue atrophy, to estimate the final, post-remodeling anatomical and biomechanical state.

The proposed robust, automated workflow serves as a computational testbed for surgical planning and systematic studies. Accordingly, we conducted a computational factorial experiment to decouple and quantify the effects of two critical surgical decisions: donor-site selection and flap overbulking. By systematically varying the flap's biomechanical properties and its initial subcutaneous volume, this research can isolate the independent and interactive effects of these variables on the final anatomical and biomechanical outcomes. Such computational modeling allows for the simulation of

numerous scenarios and provides quantitative physics-based insight that is otherwise
difficult to obtain from clinical studies alone.

2 Method

2.1 Patient-Specific Tongue Model

This study is built upon a high-resolution, patient-specific 3D model of the tongue, previously derived from an MRI scan of a healthy male participant in his 40s [7]. Data acquisition was approved by the Health Research Ethics Board of Alberta (Protocol HREBA.CC-17-0597), with written informed consent obtained from the participant upon recruitment on 30/07/2019. The tongue model establishes the baseline preoperative anatomy for all simulations herein. To ensure consistency across representative cases, we adopted the same four clinically plausible tongue reconstruction scenarios and used "resection models" identical to those in our prior work. For each case, the "resected tongue model" was generated via a Boolean subtraction of the corresponding resection model from the baseline tongue anatomy. The resulting volumetric mesh serves as the recipient site for the virtual free flap reconstruction, and the resection model provides the geometric basis for subsequent flap design (Fig 1).

To establish a stable anatomical reference frame and define physiological boundary conditions, a rigid jaw model was also created. This was accomplished by first extracting the mandibular surface from the patient's MRI data. Subsequently, a high-resolution generic jaw model was registered to this surface using an elastic registration algorithm guided by manually placed anatomical landmarks. The resulting registered jaw model serves as a rigid, stationary foundation for the simulation, and defines the anatomical locations of the tongue's muscular attachments.

2.2 Biomechanically Optimized Flap Design

As an input for the surgical simulation, a patient-specific three-dimensional free flap model is generated, which is composed of skin and adipose layers (Fig 2B-D). The process begins by employing the hiFEM algorithm, as detailed in previous work [7], to produce a biomechanically optimized planar skin template for each resection case

(Fig 2C). A key challenge, however, is the unknown geometric model of the subcutaneous adipose tissue.

Mechanically, adipose tissue exhibits a very low shear modulus, allowing it to undergo large, quasi-fluidic deformations with relatively low elastic energy storage [34]. Its primary contribution to the flap's integrity is volumetric bulk rather than a predefined geometric shape. This principle is clinically leveraged in techniques like "beavertail" augmentation, where supplementary adipose tissue is malleably shaped to increase flap volume [9]. Consequently, the critical modeling requirement is to define an initial adipose geometry that has the necessary volume and maintains geometric congruency with the planar skin template.

To achieve this, a reproducible initial 3D shape for the adipose layer is generated by computationally "unfolding" the original 3D resection volume (Fig 2B). This transformation is guided by the deterministic 3D-to-2D nodal mapping generated by the hiFEM solution. This step ensures the resulting adipose layer is perfectly congruent with the planar design from the hiFEM algorithm, and creates a geometrically consistent foundation for assembling the composite flap model. The multilayer flap model is discretized with high-quality volumetric tetrahedral mesh using fTetWild [35], which guarantees congruent meshing and a seamless interface between the skin and fat domains.

2.3 Simulation of Tongue Reconstruction Surgery

The surgical procedure of tongue reconstruction was computationally simulated to establish the postoperative neotongue configuration (Fig 3i–iii). This involves inseting the multi-component flap onto the resected tongue and applying sutures to close the defect. First, a series of suture points is established along the flap's perimeter, spaced at clinically-relevant 5 mm intervals [36]. To establish a corresponding set of attachment points on the resected tongue model, the simulation leverages the deterministic 3D-to-2D nodal mapping generated by the hiFEM solution. This critical step ensures that the boundary deformation is guided by the optimized geometry derived from the flap design stage.

A multi-step numerical approach was developed to simulate the application of

sutures and ensure a realistic distribution of forces. To mimic how a physical suture engages a small area of tissue and avoid artificial stress concentrations, the force at each suture point was applied to a small cluster of nodes within a 2 mm radius rather than to a single node. The force is distributed to the cluster nodes via linear-spring elements that connect all nodes in a cluster to a central point. The corresponding central points on the flap and the resected tongue are then drawn together using time-activated, one-dimensional tension elements that shorten to a near-zero length over the simulation step. This sequence computationally replicates the surgical action of tightening and securing sutures.

Following the virtual suturing, a subsequent step allows the system to equilibrate for five minutes. This simulates the immediate viscoelastic stress relaxation that occurs in soft tissues after being stretched and sutured. By modeling this phenomenon, internal stresses are allowed to redistribute throughout the neotongue. The final, equilibrated geometry after this relaxation represents the immediate, postoperative "Day 0" anatomical state (Fig 3iii), which provides a stable configuration for the subsequent simulation of biological remodeling.

2.4 Simulation of postoperative flap atrophy

A critical challenge in free flap tongue reconstruction is the significant, time-dependent volume loss that occurs postoperatively. Clinical studies have reported a wide range of atrophy, with long-term volume reduction ranging from approximately 20% to nearly 60% [14,17,18,37]. The final flap volume is heavily influenced by factors such as tissue composition, patient age, BMI, and most critically, the administration of adjuvant radiation therapy. For instance, at one year post-surgery, ALT flaps in patients receiving radiotherapy exhibited an average volume loss of 44.2%, compared to just 19.8% in non-irradiated patients [14]. Similarly, radial forearm free flaps (RFFF) have shown a mean volume loss of 42.7% at ten months post-radiotherapy [17]. Given that radiotherapy is a common component of treatment for advanced cancers, and considering this body of evidence, a total volume loss of 40% was selected for this study as a clinically representative value to model the extent of flap atrophy over a one-year period. While this value serves as a clinically representative average for this systematic

study, its application to an individual patient would require personalization based on known risk factors for atrophy, such as the planned radiotherapy dose, patient age, and BMI.

To simulate this phenomenon, the model is evolved from its "Day 0" state (Fig 3iii) to an estimated one-year postoperative state (Fig 3iv). This remodeling was implemented by applying a time-dependent, isotropic negative kinematic growth to the finite elements of the flap's subcutaneous tissue. Consistent with clinical observations, atrophy was confined to the flap; the native tongue was assumed to maintain its volume. The simulation progresses over the defined one-year timeframe, with appropriate time-scaling to ensure computational tractability. The final, equilibrated geometry represents the estimated one-year anatomical and biomechanical state of the neotongue, and this 3D model serves as the primary output for the subsequent quantitative analysis.

2.5 Boundary Conditions and Contact Mechanics

A stable and physiologically realistic reference frame was established by defining the anatomical boundary conditions. The jaw model was treated as a rigid body and was fully constrained in all degrees of freedom. The tongue's physiological connection to this frame was modeled by fixing the nodes of the tongue mesh at its anatomical attachment points to the mandible and hyoid bone, which include the genioglossus, mylohyoid, and hyoglossus muscle origins as illustrated in Fig 2F and G. This configuration ensures that all simulated deformations originate from a stable anatomical base. As the objective was to estimate the final, stable anatomy at rest, muscle activation was not included in the model; however, gravity was applied as a global body force.

The physical interface between the flap and native tongue was modeled using a robust contact formulation. A penalty-based contact algorithm was enforced between the surfaces of the flap and the resected tongue to prevent interpenetration. To computationally model the desired surgical outcome of complete tissue apposition, a "sticky" contact formulation was also specified at the interface of the flap and the resected tongue surface. The simulation results confirmed that this approach prevented the formation of gaps or voids, and yielded a seamless integration at the tongue-flap interface. Additionally, a penalty contact was defined between the tongue and the jaw

model to prevent interpenetration.

227

2.6 Constitutive Models

228

All soft tissues (tongue, skin, and subcutaneous adipose tissue) were modeled as nearly incompressible, non-linear hyperelastic materials. Native tongue adopted Yeoh hyperelastic model, while donor flap skin and adipose tissue was modeled with Ogden hyperelastic model, both following the FElbio definition as below respectively [38].

229

230

231

232

$$\Psi_{\text{Yeoh}} = \sum_{i=1}^2 c_i (\tilde{I}_1 - 3)^i + \frac{k}{2} (\ln J)^2 \quad (1)$$

$$\Psi_{\text{Ogden}} = \sum_{i=1}^2 \frac{c_i}{m_i^2} \left(\tilde{\lambda}_1^{m_i} + \tilde{\lambda}_2^{m_i} + \tilde{\lambda}_3^{m_i} - 3 \right) + \frac{k}{2} (\ln J)^2 \quad (2)$$

Where Ψ represents the strain energy density function. In both models, the volumetric response is governed by the material's bulk modulus, k , and the volume ratio, J , which is the determinant of the deformation gradient tensor. The Yeoh model's isochoric behavior is a function of the first deviatoric strain invariant, $\tilde{I}_1$, with c_i being the material constants. In the Ogden formulation, the isochoric response is described in terms of the deviatoric principal stretches, $\tilde{\lambda}_i$, where c_i and m_i are material parameters.

233

234

235

236

237

238

To simulate the immediate postoperative behavior, the hyperelastic models were augmented with uncoupled viscoelastic properties to capture stress relaxation. The deviatoric part of the second Piola-Kirchhoff stress, $\mathbf{S}$, is calculated using the following integral:

239

240

241

242

$$\mathbf{S}(t) = \int_{-\infty}^t G(t-s) \frac{d\mathbf{S}^e}{ds} ds, \quad G(t) = 1 + \sum_{i=1}^N \gamma_i \exp\left(-\frac{t}{\tau_i}\right) \quad (3)$$

Where the current stress is determined by the entire history of the elastic deviatoric stress, $\mathbf{S}^e$, derived from the hyperelastic potential. The material's time-dependent behavior is characterized by the relaxation function, $G(t)$, which is represented by a Prony series. In this series, γ_i and τ_i are the relaxation constants and relaxation times for each of the N terms.

243

244

245

246

247

2.7 Soft Tissue Parameterization

To investigate the impact of donor site selection, material parameters for the Forearm skin and the Thigh skin were derived from a thorough literature review of experimental data reported in the literature (Fig 4). These experimental curves reveal significant variability in tissue stiffness, both within and between donor sites. To account for this clinically observed range, a bounding approach was implemented. To account for the observed biomechanical variability, three representative uniaxial stress-stretch curves were defined for each donor site, corresponding to three stiffness levels (Fig 4).

To incorporate the observed range of tissue properties into the simulation, a two-term Ogden material model was calibrated to each of the three stiffness levels for both the Forearm and Thigh skin. This yielded six distinct sets of material parameters that encapsulate the clinically relevant range of soft tissue properties (three for each donor site). This methodology enables the factorial study to simulate reconstructions with three levels of tissue stiffness and analyze how tissue biomechanics influence surgical outcomes.

Table 1 summarizes the calibrated hyperelastic parameters for the Forearm and Thigh skin across the three stiffness levels, as well as the constitutive parameters for the other soft tissues used in the simulation. All tissues were assigned viscoelastic properties using the Prony series (Eq 3) to model stress relaxation.

2.8 Simulation Stability and Robustness

To ensure the robustness and automation required for the 120-simulation factorial study, a strategy was implemented using the open-source finite element solver FEBio, which is specifically designed for biomechanical applications [38]. The significant non-linearities inherent to the simulation, primarily arising from non-linear materials and contacts, were managed using an implicit dynamic solver with a robust automatic time-stepping algorithm. The highly non-linear system of equations was solved using a quasi-Newton (BFGS) method with a non-symmetric stiffness matrix.

The stability of the simulations was further enhanced through careful model setup. Contact mechanics, a common source of numerical instability, were configured by tuning the penalty parameters to ensure minimal surface penetration without inducing

Table 1. Material parameters used in the finite element simulations.

Soft Tissue	Model	Parameters (Values)	Source
Native Tongue	Yeoh (Hyperelastic)	$(c_1, c_2, k) = (1.037kPa, 0.486kPa, 100kPa)$	[24]
Native Tongue	Prony (Viscoelastic)	$(\tau_1, \tau_2, \tau_3, \tau_4) = (0.05s, 1.0s, 20s, 400s);$ $(\gamma_1, \gamma_2, \gamma_3, \gamma_4) = (39.0, 4.526, 4.895, 3.211)$	[39] [†]
Subcutaneous Adipose	Ogden (Hyperelastic)	$(c_1, m_1, k) = (9.0kPa, 6.0kPa, 100kPa)$	[40]
Subcutaneous Adipose	Prony (Viscoelastic)	$(\tau_1, \tau_2, \tau_3) = (0.1s, 1.0s, 10.0s);$ $(\gamma_1, \gamma_2, \gamma_3) = (4.428, 8.692, 0.738)$	[41] [†]
Forearm Skin (Stiffness Levels)	Ogden (Hyperelastic)	Level 1: $(c_1, c_2, m_1, m_2, k) = (41.5kPa, 1.39Pa, 1.223, 41.672, 200kPa);$ Level 2: $(c_1, c_2, m_1, m_2, k) = (61.3kPa, 4.38Pa, 7.059, 39.062, 200kPa);$ Level 3: $(c_1, c_2, m_1, m_2, k) = (102.1kPa, 5.20Pa, 7.660, 38.691, 200kPa)$	‡
Thigh Skin (Stiffness Levels)	Ogden (Hyperelastic)	Level 1: $(c_1, c_2, m_1, m_2, k) = (90.0kPa, -10.0Pa, 13.431, 31.000, 200kPa);$ Level 2: $(c_1, c_2, m_1, m_2, k) = (180.0kPa, -60.0Pa, 14.531, 36.333, 200kPa);$ Level 3: $(c_1, c_2, m_1, m_2, k) = (360.0kPa, -120.0Pa, 13.839, 35.274, 200kPa)$	‡
Skin	Prony (Viscoelastic)	$(\tau_1, \tau_2, \tau_3, \tau_4) = (0.3791s, 4.1987s, 6.9961s, 180.02s);$ $(\gamma_1, \gamma_2, \gamma_3, \gamma_4) = (3.649, 0.452, 0.553, 0.627)$	[42] [†]

Notes: [†] The referenced studies define the relaxation function differently from Eq 3. The reported viscoelastic parameters have been mathematically converted to be consistent with the formulation used in this study. Relaxation coefficients (γ_i) are dimensionless. [‡] Material profiles for the three stiffness levels of Forearm skin were calibrated using experimental data from several published studies [43–46]. Similarly, the profiles for Thigh skin were calibrated based on data reported by [47].

ill-conditioning of the stiffness matrix. To improve the numerical performance for nearly
incompressible materials, uncoupled formulations were used for the hyperelastic and
viscoelastic constitutive models. A critical component of the strategy was the meshing
approach. Given the complex, organic geometries of the tongue and flap, tetrahedral
elements were chosen over hexahedral elements due to the availability of automated and
robust meshing algorithms, such as the state-of-the-art mesher fTetWild [35]. To
mitigate the risk of volumetric and bending locking with linear elements, higher-order
quadratic tetrahedral elements (tet10) were used. These elements have been shown to
reliably capture complex deformations and contact mechanics without the locking
artifacts, while providing solution accuracy that is comparable or superior to hexahedral
elements with the same number of degrees of freedom [48].

Finally, a mesh convergence analysis was conducted to ensure that the simulation
outcomes were independent of the mesh discretization. Key simulations were repeated
with progressively refined mesh densities, and the results were evaluated using key
outcome metrics: maximum displacement and maximum von Mises equivalent strain.
The analysis demonstrated that the solution converged as the mesh was refined. The
mesh density selected for the full factorial study was chosen from the converged region
to maintain computational tractability. The detailed results of this convergence analysis
are presented in the Supporting information.

2.9 Outcome Metrics

To quantitatively evaluate the results of the simulations, two primary metrics were
established to assess the final state of the neotongue. These metrics focus on the
anatomical fidelity and the biomechanical state of the reconstructed tongue, as
visualized in Fig 3v–vi.

The first metric, anatomical state, quantifies how closely the final, post-remodeling
neotongue morphology restores the patient’s original, preoperative anatomy. The
preoperative native tongue 3D model serves as the ground truth for this comparison.
Since both the simulated and original tongue models are anatomically fixed to the same
rigid jaw model, they share a common coordinate system, allowing for direct spatial
comparison without the need for further registration. The discrepancy between the two

surfaces is quantified by the maximum surface deviation, calculated as the maximum Hausdorff distance. To ensure this metric is robust against numerical artifacts or isolated mesh irregularities, the 99th percentile of the Hausdorff distances is reported, providing a stable measure of the largest regional anatomical deviation.

The second metric, imposed tongue deformation, evaluates the biomechanical strain imparted to the native tongue tissue as a result of the reconstruction and subsequent remodeling. To provide a robust and physically meaningful measure of deformation, the von Mises equivalent strain (ε_{eq}) was selected, hereafter referred to as Effective Strain. This scalar invariant distills the complex, three-dimensional strain state into a single value, permitting a direct and objective comparison across all simulated scenarios. The equivalent strain was calculated from the Lagrange strain tensor, $\mathbf{E}$, as follows:

$$\varepsilon_{eq} = \sqrt{\frac{2}{3} \mathbf{dev} \mathbf{E} : \mathbf{dev} \mathbf{E}} \quad (4)$$

Where $\mathbf{dev} \mathbf{E} = \mathbf{E} - \frac{1}{3} \text{tr}(\mathbf{E}) \mathbf{I}$ is the deviatoric part of the strain tensor, $:$ denotes the double contraction, $\text{tr}(\mathbf{E})$ is the trace of $\mathbf{E}$, and $\mathbf{I}$ is the identity tensor. This formulation is particularly advantageous as it is based on the deviatoric part of the strain tensor, which mathematically isolates the distortional component of deformation (i.e., shear and tensile strain) from the hydrostatic (volumetric) component. This allows the metric to specifically quantify the mechanically significant distortion that can lead to tissue tethering and functional impairment, while remaining insensitive to the uniform volumetric shrinkage associated with the simulated flap atrophy.

2.10 The Factorial Study Design

A computational factorial experiment was designed to systematically investigate the anatomical and biomechanical impact of key surgical decisions. This approach allows for a controlled analysis of how different factors independently and interactively influence the outcome metrics. The study examined three primary independent variables:

- Patient-Specific Defect Geometry: Four distinct, clinically-plausible glossectomy scenarios were used (Fig 1, Cases A–D). These cases represent several distinct anatomical defects.

- Donor Site Biomechanics: The mechanical properties of the flap’s skin layer were varied to simulate tissue harvested from two common donor sites: the radial forearm and the thigh. To account for natural patient-to-patient variability in tissue stiffness, three distinct stiffness profiles, as detailed in section 2.7, were defined for each site.
- Flap Overbulking: Five percentage levels of adipose bulk increase were simulated: 0%, 22.2%, 44.4%, 66.7%, and 88.9%. Overbulking was achieved by scaling the flap’s subcutaneous adipose component in the thickness direction (Fig 2D). This approach reflects the clinical practice of elevating a thicker subcutaneous fat layer (for thigh flaps) or harvesting extra beavertail fat tissue (for forearm flaps). This method isolates the effect of flap’s bulk from the flap’s skin paddle size. The selected linear levels span a clinically relevant range, which includes the calculated volume needed to perfectly match the resection size after 40% atrophy (the 66.7% level).

A full factorial design was implemented by combining every level of each factor. This comprehensive approach ensures that all possible interactions between the variables could be studied. The total number of unique simulations conducted was the product of the levels for each factor: (4 defect geometries) $\times$ (6 skin stiffness levels) $\times$ (5 overbulking levels), resulting in 120 distinct simulation setups.

2.11 Statistical Analysis

To quantify the influence of the experimental factors on the simulated outcomes, Ordinary Least Squares (OLS) linear regression was employed for each of the two primary metrics. The independent variables from the factorial design were encoded for the regression models. *Case* was treated as a four-level categorical variable representing the defect geometries, with Case A serving as the reference. *Overbulking* was included as a continuous numerical variable. To decouple the effect of donor site biomechanics, two predictors were created: *is_thigh*, a binary variable (0 = Forearm, 1 = Thigh), and *stiffness_level*, an ordinal variable coded as -1, 0, +1 for the three stiffness levels for each of the donor sites. This encoding treats stiffness as a numerical factor and assumes a linear effect for changes in stiffness within a given donor site.

A hierarchical approach was used to build and select the most appropriate model. An initial model included only the main effects of each predictor ($Outcome \approx Case + Overbulking + is_thigh + stiffness_level$). A second, more complex model was then fitted, containing all main effects plus all two-way scientifically relevant interaction terms. The final model for interpretation was selected by comparing these nested models using an analysis of variance (ANOVA) F-test. This test determined if the interaction terms provided a statistically significant improvement in explaining outcome variance, with the adjusted R-squared also being evaluated to ensure that added complexity was justified by increased explanatory power.

Following selection, the final model for each outcome was subjected to diagnostic checks to validate the underlying assumptions of OLS regression. The normality of the model's residuals was assessed through visual inspection of quantile-quantile (Q-Q) plots. Multicollinearity was assessed using the Variance Inflation Factor (VIF), ensuring that all VIF scores were below the common threshold of 10 to confirm the stability of the model's coefficients. All statistical analyses were conducted using the *statsmodels* open-source library [49].

2.12 Data and Code Availability

The simulation framework is made openly available [50] under MIT license, without restriction, enabling full reproducibility of all 120 simulations and direct application to new patient cases. Patient-specific baseline tongue MRI data cannot be shared due to privacy restrictions and ethics board requirements; however, all inputs required to reproduce the simulation outcomes and statistical results are included in this repository (hosted on GitHub at <https://github.com/is-Amir/VSP-BioSim> and archived with a citable DOI: 10.5281/zenodo.17438091).

This open-source release provides the research community with immediate access to a robust, scalable platform for patient-specific tongue reconstruction simulation and empowers researchers to adapt this methodology to other soft-tissue reconstruction and virtual surgical planning applications.

3 Results

The computational framework demonstrated high stability and robustness, successfully completing all 120 unique simulations in the full factorial study without numerical failures. The simulations provided quantitative outcomes for the final anatomy and biomechanics of the neotongue across the full spectrum of surgical variables tested.

The overall trends are visualized in Fig 5. A clear trade-off emerges between the two primary outcomes: increasing the level of subcutaneous tissue overbulking consistently increased the maximum Effective Strain imposed on the native tongue (top row), while simultaneously decreasing the maximum surface deviation from the original anatomy (middle row). The final, post-remodeling tongue volume was a direct and linear function of the initial overbulking percentage (bottom row). The 66.7% overbulking level consistently resulted in a final flap volume that closely compensated for the simulated 40% atrophy. At this level, the final tongue volume closely matched the preoperative native tongue volume, resulting in a percentage change close to zero.

To quantify the factors influencing the first dependent variable (maximum surface deviation) a hierarchical linear regression analysis was performed. A formal model comparison using an ANOVA F-test revealed that the model including two-way interactions provided a significantly better fit to the data than a main effects model ($F = 26.12$, $p < 0.001$). The interaction model explained a substantial portion of the variance ($\text{Adj. } R^2 = 0.993$) and was therefore selected for interpretation.

The regression results, detailed in Table 2, identify *overbulking* as the most dominant factor for restoring the original tongue anatomy. The significant negative coefficient (-3.420 , $p < 0.001$) indicates that increasing the flap volume consistently reduces the maximum surface deviation. However, the beneficial effect of overbulking is significantly modulated by donor site selection (coeff = 0.494 , $p < 0.001$), stiffness level (coeff = 0.160 , $p < 0.001$), and defect geometry (all $p < 0.001$).

A separate regression analysis was conducted for the second dependent variable: Maximum Effective Strain. The model comparison showed that adding interaction terms did not provide any significant improvement over the main effects model ($F = 0.069$, $p = 0.999$). The main effects model was therefore selected as the superior representation of the dependent variable, explaining 86.3% of the variance in maximum effective strain.

Table 2. OLS regression results for maximum surface deviation (interaction model). Dependent variable: maximum surface deviation (mm). Reference: *Case A*; *is_thigh* = 0 (Forearm).

Predictor	Coefficient	Std. Error	t-statistic	p-value
Intercept	8.000	0.036	220.77	< 0.001
<i>Case B</i> vs. <i>A</i>	1.101	0.046	24.01	< 0.001
<i>Case C</i> vs. <i>A</i>	1.594	0.046	34.77	< 0.001
<i>Case D</i> vs. <i>A</i>	-0.578	0.046	-12.61	< 0.001
<i>overbulking</i>	-3.420	0.067	-51.38	< 0.001
<i>is_thigh</i>	-0.232	0.032	-7.15	< 0.001
<i>stiffness_level</i>	-0.035	0.023	-1.54	0.125
<i>overbulking</i> × <i>Case B</i>	0.482	0.084	5.72	< 0.001
<i>overbulking</i> × <i>Case C</i>	0.335	0.084	3.98	< 0.001
<i>overbulking</i> × <i>Case D</i>	0.625	0.084	7.42	< 0.001
<i>overbulking</i> × <i>is_thigh</i>	0.494	0.060	8.30	< 0.001
<i>overbulking</i> × <i>stiffness_level</i>	0.160	0.036	4.39	< 0.001
<i>is_thigh</i> × <i>stiffness_level</i>	-0.065	0.023	-2.82	0.006

The results for the main effect model, presented in Table 3, describe a simple, additive system. All main effects—*overbulking*, *is_thigh*, and *stiffness_level*—were highly significant positive predictors of strain. Increasing flap volume, using tissue from the Thigh, and increasing skin stiffness each independently and consistently increased the maximum Effective Strain experienced by the native tongue tissue. The defect geometry (*Case*) also had a significant effect, with Cases B and C (larger resections) resulting in higher strain than the reference Case A, while Case D was not significantly different. The lack of significant interactions implies that the strain-inducing effect of *overbulking* was consistent and not modulated by other independent variables.

Table 3. OLS regression results for maximum Effective Strain (main effects model). Dependent variable: maximum Effective Strain. Reference: *Case A*; *is_thigh* = 0 (Forearm).

Predictor	Coefficient	Std. Error	t-statistic	p-value
Intercept	0.226	0.005	49.27	< 0.001
<i>Case B</i> vs. <i>A</i>	0.051	0.005	10.37	< 0.001
<i>Case C</i> vs. <i>A</i>	0.052	0.005	10.67	< 0.001
<i>Case D</i> vs. <i>A</i>	0.004	0.005	0.81	0.422
<i>overbulking</i>	0.093	0.006	16.75	< 0.001
<i>is_thigh</i>	0.043	0.003	12.32	< 0.001
<i>stiffness_level</i>	0.023	0.002	10.74	< 0.001

4 Discussion

The fundamental goal of tongue reconstruction is twofold: to restore the anatomical volume required for effective tongue-palate contact during swallowing and speech articulation [9,11], and to preserve the dynamic mobility of the residual tongue essential for chewing, food propulsion, and clear speech [2,8,9]. This study provides a mechanistic and quantitative explanation for the clinical challenges inherent in balancing these objectives. Our simulations reveal a fundamental, physics-based trade-off: surgical decisions aimed at restoring neotongue’s bulk through flap overbulking invariably impose significant and predictable strain on the remaining native tongue tissue. This finding suggests that the ”ideal” reconstruction is one that optimally balances volume restoration against the biomechanical cost.

These insights were enabled by the primary methodological achievement of this work: the development of a novel, end-to-end computational pipeline for surgical simulation. This framework uniquely integrates patient-specific model generation, biomechanically-optimized flap design, the simulation of tongue reconstruction, and the modeling of biological atrophy. Its power as a robust and automated virtual testbed was exemplified through the successful execution of a 120-simulation factorial study, which enabled such a systematic investigation.

The selection of a donor site is a foundational surgical decision that imposes a distinct and quantifiable biomechanical cost on the reconstructed tongue. Our regression analysis (Table 3) reveals a clear, quantitative relationship: the stiffer the flap tissue, the greater the biomechanical strain imposed on the residual native tongue. Specifically, the model demonstrated that Thigh flaps induced a significantly greater mechanical strain on the surrounding native tongue than their Forearm counterparts. This biomechanical penalty was further amplified by increasing stiffness levels within each donor site. This finding provides the first physics-based, quantitative measure of the clinical concept of ”tethering”; the mechanical anchoring of the mobile tongue remnant by a stiff, non-compliant flap.

This quantified tethering effect provides a direct mechanical rationale for the functional trade-offs reported extensively in the clinical literature. Studies consistently associate the thin, pliable Radial Forearm Free Flap (RFFF) with superior speech

articulation, a function dependent on preserving tongue mobility [8,9,12]. Our results support this clinical observation, demonstrating that the lower material stiffness of the Forearm flap results in the lowest imposition of strain, thereby creating a more favorable biomechanical environment for the complex movements of speech. In contrast, for extensive resections creating large volumetric deficits, such as total and subtotal glossectomies and those involving the tongue base, the ALT flap is the preferred reconstructive choice to restore the required bulk [8,13]. Our simulations support the rationale for this choice but also reveal its inherent biomechanical cost: it simultaneously imposes a greater mechanical strain, which can compromise the mobility required for vital functions.

The clinical practice of overbulking is a direct response to the challenge of postoperative atrophy, yet it forces surgeons into a delicate balancing act to avoid creating a neotongue that is too bulky and immobile [20]. The computational framework provides a clear explanation for this surgical dilemma by quantifying its consequences. The results reveal a fundamental conflict: increasing flap volume is, by a significant margin, the most powerful surgical parameter for reducing maximum surface deviation, which is a direct proxy for restoring the preoperative anatomy (Table 2, $\text{coeff} = -3.420$, $p < 0.001$). However, this anatomical gain comes at a direct and unavoidable biomechanical cost, similar to the findings regarding stiffness levels. As shown in Table 3, every increase in overbulking imposes a linear and predictable increase in the strain imparted to the residual native tongue ($\text{coeff} = 0.093$, $p < 0.001$). This finding quantitatively defines the trade-off between form and function. The "ideal" reconstruction is therefore not simply a matter of perfectly matching the original volume, but rather a patient-specific compromise that optimizes anatomical fidelity against the predictable biomechanical strain that may tether the tongue and impair function; what we know as "too bulky and immobile".

While the biomechanical cost of overbulking is straightforward and additive, its anatomical benefit is highly conditional, modulated by both the properties of the flap and the geometry of the defect. The regression analysis for anatomical deviation (Table 2) is characterized by significant interaction terms, showing that the effectiveness of adding volume is not constant. The most critical interaction is with the donor site (*overbulking* x *is_thigh*, $\text{coeff} = 0.494$, $p < 0.001$). The positive coefficient reveals that

the anatomical benefit of adding volume is significantly less pronounced for the stiffer Thigh flaps compared to the more pliable Forearm flaps. This finding introduces a key nuance: while a Thigh flap shows a minor anatomical advantage at baseline (*is_thigh*, $\text{coeff} = -0.232$, $p < 0.001$), the Forearm flap's final anatomy is more responsive to the surgeon's corrective overbulking. We interpret this as a difference in "tunability"; the compliant Forearm flap is more malleable, allowing the added subcutaneous volume to be effectively shaped by the surrounding native tongue, leading to better anatomical conformity. This effect is reinforced by the *overbulking* $\times$ *stiffness_level* interaction ($\text{coeff} = 0.160$, $p < 0.001$), which reveals that for any given donor site, the anatomical return on investment from overbulking is greatest when using the most pliable tissue available. This provides a direct, physics-based rationale for the clinical preference for the beavertail modification of the RFFF, which increases the subcutaneous bulk while maintaining the flap's overall pliability [9, 51, 52]. Furthermore, the significant *overbulking* $\times$ *Case* interactions highlight that the benefit of adding volume is not universal; it is dictated by the unique geometrical constraints of the resection site. Taken together, the model demonstrates that while overbulking is universally beneficial for restoring anatomy, its clinical effectiveness is highly variable and depends on the flap's inherent biomechanics and the specific demands of the defect.

Together, the surgical decisions of donor site selection and flap overbulking are not independent variables but a coupled system with compounding effects. The choice of donor site establishes a baseline biomechanical cost in the form of tissue strains and tensions. Moreover, it dictates the therapeutic effectiveness of overbulking; a pliable Forearm flap imposes less initial strain and allows its final anatomy to be more effectively 'tuned' with added volume. Conversely, a stiffer Thigh flap introduces a higher baseline strain and shows diminished anatomical returns from the same corrective overbulking. Therefore, an optimal reconstructive strategy must consider these decisions in unison, recognizing that the inherent properties of the chosen flap define the potential and the limitations of subsequent efforts to restore anatomical form. This interdependence underscores the complexity of surgical planning and highlights the need for patient-specific models such as the one presented herein to navigate this complex decision-making process.

The presented framework introduces a novel approach for simulating the complex

procedure of free flap reconstruction. This is achieved by modeling the flap as an independent, multi-component body that is virtually sutured to the resection site. This approach fundamentally advances two established paradigms in surgical simulation. First, it moves beyond the "property modification" approach, where the outcome of a reconstruction is simulated by altering the material properties of elements within a single, continuous mesh. Foundational work by Buchaillard et al. [23], later adopted by van Alphen et al. [53] and more recently Al-Zanoon et al. [32], effectively used this simplification to explore the functional consequences of flap stiffness mismatch or fibrotic tissue. Our framework, in contrast, models the flap as a distinct biomechanical entity. This allows explicit capture of the physics of the flap-tongue interface, the induced deformations by suture tensioning, and the final postoperative equilibrium state. In essence, it simulates the surgical process, not just its endpoint material state. Second, our work extends the "primary closure" simulation paradigm, pioneered by Kappert et al. [33]. Their method represented a major advance by computationally replicating the surgical act of suturing, successfully modeling the deformation caused by pulling native tissue edges together to close a small defect. Our framework adapts this concept to the more complex and common scenario of microvascular free flap reconstruction.

Another key innovation of this pipeline is the simulation of postoperative tissue atrophy, which enables the model to evolve from its immediate "Day 0" state to a predicted one-year post-remodeling anatomy; the state most relevant to long-term patient function. By modeling this crucial remodeling phase, the framework provides a biomechanical basis to investigate the clinical practice of flap overbulking. The focus on long-term biological change complements prior simulation efforts that have also investigated post-treatment tissue alteration. Al-Zanoon et al. [32], for instance, modeled the development of radiation-induced fibrosis by increasing the stiffness of affected tissue elements. Our approach is complementary, modeling atrophy via a prescribed volumetric change, another critical remodeling phenomenon that dictates the final tongue morphology. Furthermore, while Wang et al. [54] specifically studied the functional consequences of base-of-tongue volume loss on swallowing, their work focused on the hydrodynamics of bolus flow driven by a prescribed boundary motion. Our framework, in contrast, simulates the solid mechanics of how the reconstructed neotongue itself deforms and settles into a new passive equilibrium as the flap tissue

atrophies. This provides direct insight into the final resting anatomy and the resulting strains imposed on the native tongue—a different set of biomechanical outcomes.

The successful execution of a 120-simulation factorial study demonstrates the framework’s core design principles: automation and robustness. This scalability marks a methodological shift. We present a high-throughput virtual testbed that contrasts with a significant body of research focused on building baseline models of the highest fidelity. Such landmark efforts include the comprehensive FRANK model, which integrates numerous head and neck components [25–27], and patient-specific models derived from DW-MRI that incorporate detailed muscle fiber architecture [31]. Our framework, however, makes a deliberate, pragmatic trade-off. Since our primary goal is to predict the final, passive anatomy at rest after biological remodeling, we do not model the active muscle architecture, which is critical for functional simulation but not for this specific question. Instead, the framework employs well-established isotropic hyperelastic models [28] rather than more complex anisotropic formulations, which have proven exceptionally difficult to parameterize from in vivo data [31]. This trade-off prioritizes simulation scalability over baseline complexity. The focus is therefore not on perfecting the preoperative model but on robustly simulating the postoperative consequences that arise from interdependent surgical decisions. Ultimately, this capability distinguishes our approach from prior tongue surgery simulations, which have necessarily focused on single-case analyses or a small number of scenarios [23, 33]. It is this scalability that opens the door to systematic, multi-parameter investigations of surgical variables.

This degree of automation and robustness is what transforms the framework from a research platform into a viable tool for patient-specific preoperative planning. The laborious, manual, and expert-driven process of creating high-fidelity, patient-specific biomechanical models has traditionally been a major bottleneck, limiting their scalability for large-scale or clinical applications [27, 29–31]. In contrast, our framework is designed for clinical tractability. The pipeline requires only two triangulated meshes—the baseline tongue and the planned resection—which directly align with the modern VSP workflows where a surgeon delineates the resection on medical images. From this point, the entire workflow—from biomechanically optimized flap design to volumetric FEM generation and final post-atrophy analysis—is fully automated. The clinical necessity for such a tool is underscored by the results of the factorial study itself.

The significant *overbulking* $\times$ *Case* interaction demonstrates that the optimal surgical strategy is dictated by the unique geometric constraints of the patient's specific defect. The streamlined and robust nature of the pipeline makes it computationally feasible to conduct a tailored series of simulations for an individual patient to optimize a critical variable like the overbulking percentage. This offers a direct, physics-based method to navigate the complex, interdependent surgical decision, and move toward data-driven, individualized surgical planning.

The current framework, while robust, relies on a simplified representation of the tongue's complex biomechanics and postoperative healing. Future work should incorporate muscle fiber orientation data, potentially from DT-MRI, to capture the tongue's anisotropic properties. A key advancement will be the integration of active muscle contraction models, enabling the simulation of dynamic functions like speech and swallowing. Furthermore, the biological remodeling simulation can be enhanced by implementing patient-specific atrophy models that factor in variables like radiation dose, flap composition, and patient-specific factors like age, sex, and BMI. Such studies could also benefit from modeling the formation of stiffer, contracted scar tissue at the incision interfaces. Finally, refining the constitutive parameters through in vivo techniques, such as MR Elastography for the tongue [55] and preoperative mechanical testing of donor skin [43], will increase the model's predictive accuracy.

To translate this framework into a clinically viable tool, several validation and generalization steps are necessary. A prospective clinical study is needed to validate the simulated anatomical outcomes against long-term postoperative imaging from a cohort of patients. The model's generalizability must also be established by applying it to a diverse range of patient anatomies and tumor-specific resection geometries. The robust pipeline is also well-suited for large-scale in silico studies to systematically investigate surgical variables, such as comparing the biomechanically optimized flaps used herein against conventional flap designs sourced from surgeons. Ultimately, this framework has the potential to serve as a research platform for systematically investigating surgical variables and as a tool for conducting patient-specific preoperative simulations to compare optimal reconstructive strategies.

5 Conclusion

This study's primary contribution is the development and application of a robust and automated simulation pipeline capable of modeling key aspects of tongue reconstruction surgery and its long-term anatomical remodeling. This framework provides a novel platform to systematically investigate interdependent surgical variables that are otherwise challenging to assess. Our findings provide a quantitative explanation for the delicate balance surgeons must strike between restoring volume and preserving mobility. The simulations confirm that surgical decisions to increase bulk invariably impart a biomechanical cost in the form of tissue strain, which can lead to the tethering of the mobile tongue. We demonstrated that the choice of donor site and the degree of overbulking are a coupled system. The selection of a stiffer donor site, like the thigh, not only imposes a higher baseline mechanical strain but also limits the anatomical effectiveness of subsequent overbulking. This supports the clinical rationale for using pliable flaps, such as the beavertail-modified forearm flap, to achieve anatomical conformity with a lower biomechanical penalty. Crucially, the significant interaction between surgical strategy and the patient's unique defect geometry confirms that a universal approach is suboptimal. The results highlight the potential for this framework to be used for patient-specific preoperative planning, allowing surgeons to computationally explore different scenarios and gain insight into the specific case. Methodological refinements should include anisotropic material properties and active muscle modeling, and the framework's predictive accuracy must ultimately be validated through prospective clinical studies. This work lays the foundation for a powerful research platform and, following clinical validation, a potential decision-support tool to help surgeons optimize and individualize reconstructive strategies.

Funding

This work was supported by the Alberta Cancer Foundation (grant number 27 601). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

1. Lam L, Samman N. Speech and Swallowing Following Tongue Cancer Surgery and Free Flap Reconstruction - A Systematic Review. *Oral Oncol.* 2013 Jun;49(6):507-24. doi:10.1016/j.oraloncology.2013.03.001.
2. Urken ML. Multidisciplinary Head and Neck Reconstruction: A Defect-Oriented Approach. Lippincott Williams & Wilkins; 2012.
3. Baskin RM, Seikaly H, Sawhney R, Danan D, Burt M, Idris S, et al. Tongue Reconstruction: Rebuilding Mobile Three-Dimensional Structures from Immobile Two-Dimensional Substrates, a Fresh Cadaver Study. *Head Neck-J Sci Spec Head Neck.* 2019 Oct;41(10):3693-9. doi:10.1002/hed.25889.
4. Liu MD, Liu DZ, Al-Aroomi MA, Xiong JJ, Liu S, Sun CF, et al. Reconstruction of Large Defects of Anterior Floor of Mouth with Free Flaps Using a Novel Individualized Flap Design Method. *J Cranio-MaxilloFac Surg.* 2023 Sep;51(9):574-9. doi:10.1016/j.jcms.2023.08.006.
5. Liu MD, Xue XM, Al-Aroomi MA, Xiong JJ, Liu S, Sun CF, et al. A Novel Flap Design Technique for Subtotal Tongue Reconstruction with an “Individualized and Convenient Tongue Model”. *Oral Oncol.* 2023 Oct;145:106531. doi:10.1016/j.oraloncology.2023.106531.
6. Xu J, Lai F, Liu Y, Tan Z, Zheng C, Wang J, et al. Novel Computer-Aided Reconstruction of Soft Tissue Defects Following Resection of Oral and Oropharyngeal Squamous Cell Carcinoma. *World J Surg Oncol.* 2022 Jun;20(1):196. doi:10.1186/s12957-022-02654-7.
7. Isazadeh AR, Zenke J, Westover L, Seikaly H, Aalto D. Patient-Specific Virtual Surgical Planning for Tongue Reconstruction: Evaluating Hyperelastic Inverse FEM with Four Simulated Tongue Cancer Cases. *Phys Med Biol.* 2025 Jul;70:145017. doi:10.1088/1361-6560/adebd8.
8. Marra C, Pinto V, Benanti E, De Maria F, Pinelli M, Spaggiari A, et al. Radial Forearm Flap versus Antero-Lateral Thigh Flap in Oral Soft Tissue Reconstruction: Update and Statistical Analysis on Our 77 Consecutive Cases for

- an Objective Selection Criteria. *Acta Biomed.* 2023 Oct;94(5):e2023252.
doi:10.23750/abm.v94i5.15174.
9. Seikaly H, Rieger J, O'Connell D, Ansari K, AlQahtani K, Harris J. Beavertail Modification of the Radial Forearm Free Flap in Base of Tongue Reconstruction: Technique and Functional Outcomes. *Head Neck.* 2009 Feb;31(2):213-9.
doi:10.1002/hed.20953.
 10. Seikaly H, Rieger J, Wolfaardt J, Moysa G, Harris J, Jha N. Functional Outcomes After Primary Oropharyngeal Cancer Resection and Reconstruction With the Radial Forearm Free Flap. *Laryngoscope.* 2003 May;113(5):897-904.
doi:10.1097/00005537-200305000-00023.
 11. Wang X, Yan G, Zhang G, Li J, Liu J, Zhang Y. Functional Tongue Reconstruction with the Anterolateral Thigh Flap. *World J Surg Onc.* 2013 Dec;11(1):303. doi:10.1186/1477-7819-11-303.
 12. Sepúlveda L, Peralta J G, Ampuero E V, Fernández A F. Comparison of Functional Outcomes of Radial Forearm and Anterolateral Thigh Free Flaps after Hemiglossectomy in Patients with Tongue Cancer: A Systematic Review and Meta-Analysis. *Rev Otorrinolaringol Cir Cabeza Cuello.* 2025 Mar;85(1):95-104.
doi:10.4067/s0718-48162025000100549.
 13. Cai YC, Li C, Zeng Df, Zhou YQ, Sun RH, Shui CY, et al. Comparative Analysis of Radial Forearm Free Flap and Anterolateral Thigh Flap in Tongue Reconstruction after Radical Resection of Tongue Cancer. *ORL-J Oto-Rhino-Laryngol Head Neck Surg.* 2019 Nov;81(5-6):252-64.
doi:10.1159/000502151.
 14. Tarsitano A, Battaglia S, Cipriani R, Marchetti C. Microvascular Reconstruction of the Tongue Using a Free Anterolateral Thigh Flap: Three-dimensional Evaluation of Volume Loss after Radiotherapy. *J Cranio-MaxilloFac Surg.* 2016 Sep;44(9):1287-91. doi:10.1016/j.jcms.2016.04.031.
 15. Vincent A, Kohlert S, Lee T, Inman J, Ducic Y. Free-Flap Reconstruction of the Tongue. *Seminars in Plastic Surgery.* 2019 Feb;33(01):038-45.
doi:10.1055/s-0039-1677789.

16. Kim MS, Oh KH, Cho JG, Baek SK, Woo JS, Jung KY, et al. Assessment of Chronological Volume Changes in Radial Forearm Free Flaps for Tongue Cancer. *ORL*. 2020;82(1):40-6. doi:10.1159/000504196.
17. Razavi CR, Hostetter J, Shukla A, Cheng Z, Aygun N, Boahene K, et al. Predictors of Free Flap Volume Loss in Nonosseous Reconstruction of Head and Neck Oncologic Defects. *ENT-Ear Nose Throat J*. 2022 Jan;101(1):48-53. doi:10.1177/0145561320938903.
18. Gewirtz JI, Zhao S, Brock G, Luttrull MD, Sethuraman S, Kang SY, et al. A Pilot Study: Free Flap Atrophy in Tongue Reconstruction Using 3D Volumetric Analysis. *Ann Otol Rhinol Laryngol*. 2024 Mar;133(3):253-60. doi:10.1177/00034894231204720.
19. Zirk M, Niknazemi M, Riekert M, Kreppel M, Linz C, Lentzen MP. Comparative Analysis of Volumetric Changes between Resection Volume of Oral Tongue Cancer and Post Operative Volume of Radial Forearm Flaps. *Clin Oral Invest*. 2024 Aug;28(9):498. doi:10.1007/s00784-024-05885-y.
20. Bashorun OH, Johnson RM, Johnson EA. Flap Debulking and Secondary Revisions in Head and Neck Reconstruction: A Systematic Review with Clinical Applications. *Semin Plast Surg*. 2023 Feb;37(01):073-82. doi:10.1055/s-0042-1760444.
21. Reynolds T. Why Randomized Surgical Oncology Trials Are So Scarce. *JNCI Journal of the National Cancer Institute*. 1999 Jul;91(14):1182-3. doi:10.1093/jnci/91.14.1182.
22. Bielecki K. Are Randomized Clinical Trials Necessary in Cancer Surgery? *Nowotwory Journal of Oncology*. 2017 Jun;66(6):494-8. doi:10.5603/NJO.2016.0088.
23. Buchaillard S, Brix M, Perrier P, Payan Y. Simulations of the Consequences of Tongue Surgery on Tongue Mobility: Implications for Speech Production in Post-Surgery Conditions. *Int J Med Robot Comput Assist Surg*. 2007 Sep;3(3):252-61. doi:10.1002/rcs.142.

24. Buchaillard S, Perrier P, Payan Y. A Biomechanical Model of Cardinal Vowel Production: Muscle Activations and the Impact of Gravity on Tongue Positioning. *J Acoust Soc Am*. 2009 Oct;126(4):2033-51. doi:10.1121/1.3204306.
25. Stavness I, Lloyd JE, Payan Y, Fels S. Coupled Hard-Soft Tissue Simulation with Contact and Constraints Applied to Jaw-Tongue-Hyoid Dynamics. *Numer Methods Biomed Eng*. 2011 Mar;27(3):367-90. doi:10.1002/cnm.1423.
26. Stavness I, Nazari MA, Flynn C, Perrier P, Payan Y, Lloyd JE, et al. Coupled Biomechanical Modeling of the Face, Jaw, Skull, Tongue, and Hyoid Bone. In: Magnenat-Thalmann N, Ratib O, Choi HF, editors. *3D Multiscale Physiological Human*. London: Springer; 2014. p. 253-74. doi:10.1007/978-1-4471-6275-9_11.
27. Anderson P, Fels S, Harandi NM, Ho A, Moisik S, Sánchez CA, et al. FRANK: A Hybrid 3D Biomechanical Model of the Head and Neck. In: Payan Y, Ohayon J, editors. *Biomechanics of Living Organs*. vol. 1 of *Translational Epigenetics*. Oxford: Academic Press; 2017. p. 413-47. doi:10.1016/B978-0-12-804009-6.00020-1.
28. Hermant N, Perrier P, Payan Y. Human Tongue Biomechanical Modeling. In: Payan Y, Ohayon J, editors. *Biomechanics of Living Organs*. vol. 1 of *Translational Epigenetics*. Oxford: Academic Press; 2017. p. 395-411. doi:10.1016/B978-0-12-804009-6.00019-5.
29. Harandi N, Stavness I, Woo J, Stone M, Abugharbieh R, Fels S. Subject-Specific Biomechanical Modelling of the Oropharynx: Towards Speech Production. *Computer Methods in Biomechanics and Biomedical Engineering: Imaging & Visualization*. 2017 Nov;5(6):416-26. doi:10.1080/21681163.2015.1033756.
30. Harandi N, Woo J, Stone M, Abugharbieh R, Fels S. Variability in Muscle Activation of Simple Speech Motions: A Biomechanical Modeling Approach. *The Journal of the Acoustical Society of America*. 2017 Apr;141(4):2579-90. doi:10.1121/1.4978420.
31. Kappert KDR, Voskuilen L, Smeele LE, Balm AJM, Jasperse B, Nederveen AJ, et al. Personalized Biomechanical Tongue Models Based on Diffusion-Weighted

- MRI and Validated Using Optical Tracking of Range of Motion. *Biomech Model Mechanobiol.* 2021 Jun;20(3):1101-13. doi:10.1007/s10237-021-01435-7.
32. Al-Zanoon N, Cummine J, Jeffery CC, Westover L, Aalto D. The Effect of Simulated Radiation Induced Fibrosis on Tongue Protrusion. *Biomech Model Mechanobiol.* 2024 Jun;23:1649-60. doi:10.1007/s10237-024-01860-4.
 33. Kappert KDR, van Alphen MJA, van Dijk S, Smeele LE, Balm AJM, van der Heijden F. An Interactive Surgical Simulation Tool to Assess the Consequences of a Partial Glossectomy on a Biomechanical Model of the Tongue. *Comput Methods Biomech Biomed Eng.* 2019 Jun;22(8):827-39. doi:10.1080/10255842.2019.1599362.
 34. Sun Z, Gepner BD, Cottler PS, Lee SH, Kerrigan JR. In Vitro Mechanical Characterization and Modeling of Subcutaneous Adipose Tissue: A Comprehensive Review. *Journal of Biomechanical Engineering.* 2021 Jul;143(7):070803. doi:10.1115/1.4050286.
 35. Hu Y, Schneider T, Wang B, Zorin D, Panozzo D. Fast Tetrahedral Meshing in the Wild. *ACM Trans Graph.* 2020 Aug;39(4). doi:10.1145/3386569.3392385.
 36. Jeong WS, Oh TS. Oral and Oropharyngeal Reconstruction with a Free Flap. *Arch Craniofac Surg.* 2016;17(2):45. doi:10.7181/acfs.2016.17.2.45.
 37. Sakakibara A, Kusumoto J, Sakakibara S, Hasegawa T, Akashi M, Minamikawa T, et al. Long-Term Effects on Volume Change in Musculocutaneous Flaps after Head and Neck Reconstruction. *J Reconstr Microsurg.* 2019 May;35(4):235-43. doi:10.1055/s-0038-1672134.
 38. Maas SA, Ellis BJ, Ateshian GA, Weiss JA. FEBio: Finite Elements for Biomechanics. *Journal of Biomechanical Engineering.* 2012 Jan;134(1):011005. doi:10.1115/1.4005694.
 39. Vaidya AJ, Wheatley BB. An Experimental and Computational Investigation of the Effects of Volumetric Boundary Conditions on the Compressive Mechanics of Passive Skeletal Muscle. *Journal of the Mechanical Behavior of Biomedical Materials.* 2020 Feb;102:103526. doi:10.1016/j.jmbbm.2019.103526.

40. Sun Z, Lee SH, Gepner BD, Rigby J, Hallman JJ, Kerrigan JR. Comparison of Porcine and Human Adipose Tissue Loading Responses under Dynamic Compression and Shear: A Pilot Study. *J Mech Behav Biomed Mater*. 2021 Jan;113:104112. doi:10.1016/j.jmbbm.2020.104112.
41. Sun Z, Gepner BD, Lee SH, Rigby J, Cottler PS, Hallman JJ, et al. Multidirectional Mechanical Properties and Constitutive Modeling of Human Adipose Tissue under Dynamic Loading. *Acta Biomaterialia*. 2021 Jul;129:188-98. doi:10.1016/j.actbio.2021.05.021.
42. Then C, Stassen B, Depta K, Silber G. New Methodology for Mechanical Characterization of Human Superficial Facial Tissue Anisotropic Behaviour in Vivo. *Journal of the Mechanical Behavior of Biomedical Materials*. 2017 Jul;71:68-79. doi:10.1016/j.jmbbm.2017.02.022.
43. Flynn C, Taberner A, Nielsen P. Modeling the Mechanical Response of In Vivo Human Skin Under a Rich Set of Deformations. *Ann Biomed Eng*. 2011 Jul;39(7):1935-46. doi:10.1007/s10439-011-0292-7.
44. Griffin MF, Leung BC, Premakumar Y, Szarko M, Butler PE. Comparison of the Mechanical Properties of Different Skin Sites for Auricular and Nasal Reconstruction. *Journal of Otolaryngology - Head & Neck Surgery*. 2017 Jan;46(1):33. doi:10.1186/s40463-017-0210-6.
45. Ammam I, Guillermin A, Ouillon L, Vargiolu R, Perrot JL, Zahouani H. Coupling Tensile Test with LC-OCT and Ultrasound Imaging: Investigation of the Skin Sublayers Mechanical Behaviour. *R Soc Open Sci*. 2024 Jun;11(6):231712. doi:10.1098/rsos.231712.
46. Jacquet E, Chambert J, Pauchot J, Sandoz P. Intra- and Inter-individual Variability in the Mechanical Properties of the Human Skin from in Vivo Measurements on 20 Volunteers. *Skin Research and Technology*. 2017 Nov;23(4):491-9. doi:10.1111/srt.12361.
47. Zhou M, González PJ, Van Haasterecht L, Soyly A, Mihailovski M, Van Zuijlen P, et al. Uniaxial Mechanical Stretch Properties Correlated with Three-Dimensional

- Microstructure of Human Dermal Skin. *Biomech Model Mechanobiol.* 2024 Jun;23(3):911-25. doi:10.1007/s10237-023-01813-3.
48. Maas SA, Ellis BJ, Rawlins DS, Weiss JA. Finite Element Simulation of Articular Contact Mechanics with Quadratic Tetrahedral Elements. *J Biomech.* 2016 Mar;49(5):659-67. doi:10.1016/j.jbiomech.2016.01.024.
49. Seabold S, Perktold J. Statsmodels: Econometric and Statistical Modeling with Python. In: 9th Python in Science Conference; 2010. .
50. Isazadeh AR. VSP-BioSim: Patient-Specific Virtual Surgical Planning via Biomechanical Simulations. <https://github.com/is-Amir/VSP-BioSim>; 2025. doi:10.5281/zenodo.17438091.
51. Dziegielewski PT, Rieger J, Shama MA, O'Connell DA, Harris JR, Seikaly H. Beavertail Modification of the Radial Forearm Free Flap in Total Oral Glossectomy Reconstruction: Technique and Functional Outcomes. *Oral Oncol.* 2019 Sep;96:71-6. doi:10.1016/j.oraloncology.2019.07.004.
52. Hanubal KS, Reschly WJ, Conrad D, Festa BM, Weiss JP, Shama M, et al. The Beavertail Modified Radial Forearm Free Flap: Retrospective Review of a Versatile Technique to Increase Flap Bulk in the Head and Neck. *Microsurgery.* 2023 Nov;43(8):767-74. doi:10.1002/micr.31028.
53. van Alphen MJA, Kreeft AM, van der Heijden F, Smeele LE, Balm AJM. Towards Virtual Surgery in Oral Cancer to Predict Postoperative Oral Functions Preoperatively. *Br J Oral Maxillofac Surg.* 2013 Dec;51(8):747-51. doi:10.1016/j.bjoms.2013.06.012.
54. Wang J, Ho AK, Papadopoulos-Nydam G, Rieger J, Inamoto Y, Fels S, et al. Simulated Volume Loss in the Base of Tongue in a Virtual Swallowing Model. *Comp Meth Biomech Biomed Eng.* 2019;7(4):389-95. doi:10.1080/21681163.2017.1382392.
55. Cheng S, Gandevia SC, Green M, Sinkus R, Bilston LE. Viscoelastic Properties of the Tongue and Soft Palate Using MR Elastography. *Journal of Biomechanics.* 2011 Feb;44(3):450-4. doi:10.1016/j.jbiomech.2010.09.027.




Fig 1. Patient-Specific Tongue Model, Resection Scenarios, and Flap Designs. Overview of the four simulated tongue cancer cases along with their biomechanically-optimized flap designs. The texture mapping is generated with hiFEM and maps each point in the resection model to its corresponding point in the planar flap design.




Fig 2. Workflow for free flap model generation and surgical simulation pre-processing. Panel (A) Shows the two primary inputs: the "Resected Tongue Model" (red) and "Resection model" (textured). First, the hiFEM algorithm creates an optimized 2D planar skin template for the flap (C). This template's deterministic 3D-to-2D mapping then guides the computational "unfolding" of the resection model to generate a multi-component free flap model (B). To simulate the clinical practice of "overbulking," the flap's volume is increased by scaling the thickness of its yellow adipose component (D). Next, the multi-component flap is positioned at the resection site (E), and one-dimensional tension elements (blue lines) are defined to connect corresponding suture points on the flap and the resected tongue. Anatomical boundary conditions are established by fixing nodes (yellow dots) that represent the tongue's muscular attachments to the mandible and hyoid bone (F, G);




Fig 3. Surgical simulation pipeline: reconstruction to long-term remodeling. (i) Initial simulation setup, (ii) The suture model brings the flaps proximal to the resected tongue model and simulates suturing. (iii) After the 5-minute idling for stresses to relax with the viscoelastic model present, this is "Day 0" immediately after reconstruction. (iv) shows the tongue model after one-year of post-surgery atrophy modeling. At the very end, the effective strain (v) and Hausdorff distance (vi) are calculated and used for the statistical analysis.




Fig 4. Literature review of experimental data on the material parameters for the Forearm skin (i) and the Thigh skin (ii) represented as the stress-stretch response to a uniaxial tensile test.






Fig 5. Visualization of the sensitivity of dependant variables to different skin material properties and different levels of overbulking: overbulking increases Effective Strain (top) while decreasing anatomical deviation (middle); final tongue volume varies linearly with initial overbulking (bottom).

Supporting information

Mesh convergence analysis. We assessed mesh-independence for 12 randomly-selected representative parameter combinations (S1 Fig). Maximum equivalent strain (Eq 3) versus total mesh nodes plateaus between $\sim 24k$ and $\sim 31k$ nodes; the change across the last two refinements is $< 5\%$ for all series. Maximum displacement showed the same plateau (S2 Fig). Consistent with Simulation Stability and Robustness, we therefore used $\sim 24k$ nodes for the full study as mesh-independent within tolerance.



S1 Fig. Mesh convergence analysis—Maximum equivalent strain. Maximum equivalent strain of the neotongue at the final post-remodeling equilibrium versus total mesh nodes. Each curve is one of 12 representative parameter combinations and is labeled with a three-letter code in the form [Case][Donor/Stiffness][Overbulking], where Cases A–D are the four defect geometries; Donor/Stiffness A–C = Forearm (soft→stiff), D–F = Thigh (soft→stiff); Overbulking A–E = 0%, 22.2%, 44.4%, 66.7%, 88.9%.



S2 Fig. Mesh convergence analysis—Maximum displacement. Maximum nodal displacement of the neotongue at the final post-remodeling equilibrium (mm) versus total mesh nodes for 12 representative parameter combinations (same [Case][Donor/Stiffness][Overbulking] codes as in S1 Fig). Two panels are shown for better readability; horizontal axis is shared.