Notes for Operations Research & More

Lan Peng, PhD Student

Department of Industrial and Systems Engineering University at Buffalo, SUNY lanpeng@buffalo.edu

July 24, 2019

July 24, 2019

Contents

I	Preliminary Topics	7			
1	Review of Linear Algebra	9			
2	Convex Sets				
3	Convex Functions and Generalizations	13			
II	Linear Programming	15			
1	Formulation	17			
4	4.1 Typical Problems	17			
	4.2 Formulation Skills	17			
	4.2.1 Absolute Value	17			
	4.2.2 A Minimax Objective	17			
	4.2.3 A fractional Objective	18			
	4.2.4 A range Constraint	18			
5	Simplex Method	19			
	5.1 Basic Feasible Solutions and Extreme Points	19			
	5.2 Simplex Method	19			
	5.2.1 Simplex Method Algorithm	19			
	5.2.2 Simplex Method Tableau	19			
	5.2.3 Simplex Method as a Search Algorithm	19			
	5.3 Revised Simplex Method	19			
	5.4 Simplex Method with Bounded Variables	19			
	5.5 Artificial Variable	19			
	5.5.1 Two-Phase Method	19			
	5.5.2 Big-M Method	19			
	5.5.3 Single Artificial Variable Technique	19			
	5.6 Degeneracy and Cycling	19			
	5.6.1 Degeneracy	19			
	5.6.2 Cycling	19 19			
6	Duality Theory and Sensitivity Analysis	21			
	Decomposition Principle	23			
8	Ellipsoid Algorithm	2 5			
9	Projective Algorithm	27			
10	Interior-Point Algorithm	29			

4 CONTENTS

III Graph and Network Theory	31
11 Paths, Trees, and Cycles	33
12 Shortest-Path Problem	35
13 Minimum Spanning Tree Problem	37
14 Maximum Flow Problem	39
15 Minimum Cost Flow Problem	41
16 Assignment and Matching Problem	43
17 Graph Algorithms	45
18 Polygon Triangulation 18.1 Types of Polygons	47 47 48 48 48
IV Integer and Combinatorial Programming	49
19 Formulation 19.1 Typical Problems 19.2 Integer Programming Formulation Skills 19.2.1 A Variable Taking Discontinuous Values 19.2.2 Fixed Costs 19.2.3 Either-or Constraints 19.2.4 Conditional Constraints 19.2.5 Special Ordered Sets 19.2.6 Piecewise Linear Formulations 19.2.7 Conditional Binary Variables 19.2.8 Elimination of Products of Variables	51 51 51 51 51 51 52 52 53 53
20 Branch and Bound	55
21 Branch and Cut	57
22 Packing and Matching	59
23 Traveling Salesman Problem	61
24 Knapsack Problem	63
V Nonlinear Programming	65
25 KKT Optimality Conditions	67
26 Lagrangian Duality	69
27 Unconstrained Optimization	71
28 Penalty and Barrier Functions	73

CONTENTS 5

VI	Algorithms and Computational Complexity	75
29	Computational Complexity	77
	Sorting 30.1 Elementary Sorting Algorithms 30.2 Heap-sort 30.3 Quick-sort 30.4 Sorting in Linear Time 30.5 Medians and Order Statistics	79 79 79 79 79 79
	Data Structures 31.1 Elementary Data Structures 31.2 Hash Tables 31.3 Binary Search Trees 31.4 Red-Black Trees 31.5 B-Trees 31.6 Fibonacci Heaps 31.7 van Emde Boas Trees	81 81 81 81 81 81
	Design and Analysis Techniques 32.1 Dynamic Programming 32.2 Greedy Algorithms 32.3 Amortized Analysis 32.4 Multi-threaded Algorithms 32.5 Matrix Operations 32.6 Polynomials and the FFT 32.7 Number-Theoretic Algorithms 32.8 String Matching 32.9 Computational Geometry 32.10NP-Completeness 32.11Approximation Algorithms	83 83 83 83 83 83 83 83 83
VI	I Heuristics ans Meta-heuristics	85
VI	III Game Theory	87
;	Games with Ordinal Payoffs 33.1 Ordinal Games in Strategic Form	89 89 89
•	Games with Cardinal Payoffs 34.1 Expected Utility Theory	91919191
•	Knowledge, Common Knowledge, Beliefs 35.1 Common Knowledge	93 93 93 93
•	Refinements of Subgame-perfect Equilibrium 36.1 Weak Sequential Equilibrium	95 95 95 95

6 CONTENTS

37	Incomplete Information 37.1 Static Games	. 97
IX	A Probability, Stochastic Processes and Markov Chains	99
38	Probability	101
39	Random Variables 39.1 Relationship between Some Random Variables	. 105
40	Limit Theorems	107
41	The Bernoulli and Poisson Process	109
42	Discrete-Time Markov Chains	111
43	Continuous-Time Markov Chains	113
\mathbf{X}	Queuing Theory	115
44	Queuing Model	117
45	Birth-and-Death Queuing Models	119
46	Multidimensional Birth-and-Death Queuing Models	121
47	Phase-Type Queue	123
48	Bulk Queue	125
49	Imbedded-Markov-Chain Queuing Models	127
50	Queuing Network	129
X	I Inventory Theory	131
X	II Reliability Theory	133
X	III Statistic	135
\mathbf{X}	IV Simulation	137

Part I Preliminary Topics

Review of Linear Algebra

Convex Sets

Convex Functions and Generalizations

Part II Linear Programming

Formulation

4.1 Typical Problems

4.2 Formulation Skills

4.2.1 Absolute Value

Description: Consider the following model statement:

$$\begin{aligned} & \text{min} & & \sum_{j \in J} c_j |x_j|, & & c_j > 0 \\ & \text{s.t.} & & \sum_{j \in J} a_{ij} x_j \gtrsim b_i, & \forall i \in I \\ & & x_j & \text{unrestricted}, & \forall j \in J \end{aligned}$$

Modeling:

$$\min \sum_{j \in J} c_j(x_j^+ + x_j^-), \quad c_j > 0$$
s.t.
$$\sum_{j \in J} a_{ij}(x_j^+ - x_j^-) \gtrsim b_i, \quad \forall i \in I$$

$$x_j^+, x_j^- \ge 0, \quad \forall j \in J$$

4.2.2 A Minimax Objective

Description: Consider the following model statement:

$$\min \quad \max_{k \in K} \sum_{j \in J} c_{kj} x_j$$
s.t.
$$\sum_{j \in J} a_{ij} x_j \gtrsim b_i, \quad \forall i \in I$$

$$x_j \ge 0, \quad \forall j \in J$$

Modeling:

$$\begin{aligned} & \text{min} & z \\ & \text{s.t.} & \sum_{j \in J} a_{ij} x_j \gtrapprox b_i, & \forall i \in I \\ & \sum_{j \in J} c_{kj} x_j \le z, & \forall k \in K \\ & x_j \ge 0, & \forall j \in J \end{aligned}$$

4.2.3 A fractional Objective

Description: Consider the following model statement:

$$\min \quad \frac{\sum_{j \in J} c_j x_j + \alpha}{\sum_{j \in J} d_j x_j + \beta}$$
s.t.
$$\sum_{j \in J} a_{ij} x_j \gtrsim b_i, \quad \forall i \in I$$

$$x_j \ge 0, \quad \forall j \in J$$

Modeling:

$$\min \quad \sum_{j \in J} c_j x_j t + \alpha t$$

$$\text{s.t.} \quad \sum_{j \in J} a_{ij} x_j \gtrsim b_i, \quad \forall i \in I$$

$$\sum_{j \in J} d_j x_j t + \beta t = 1$$

$$t > 0$$

$$x_j \ge 0, \quad \forall j \in J$$

$$(t = \frac{1}{\sum_{j \in J} d_j x_j + \beta})$$

4.2.4 A range Constraint

Description: Consider the following model statement:

$$\min \quad \sum_{j \in J} c_j x_j$$
s.t.
$$d_i \leq \sum_{j \in J} a_{ij} x_j \leq e_i, \quad \forall i \in I$$

$$x_j \geq 0, \quad \forall j \in J$$

Modeling:

$$\begin{aligned} & \text{min} & & \sum_{j \in J} c_j x_j, \quad c_j > 0 \\ & \text{s.t.} & & u_i + \sum_{j \in J} a_{ij} x_j = e_i, \quad \forall i \in I \\ & & x_j \geq 0, \quad \forall j \in J \\ & & & 0 \leq u_i \leq e_i - d_i, \quad \forall i \in I \end{aligned}$$

Simplex Method

~ 1	D .	T3 *1 1	α 1 α	1 13 4	D • 4
5.1	Basic	reasible	Solutions	and Extrem	ne Points

- 5.2 Simplex Method
- 5.2.1 Simplex Method Algorithm
- 5.2.2 Simplex Method Tableau
- 5.2.3 Simplex Method as a Search Algorithm
- 5.3 Revised Simplex Method
- 5.4 Simplex Method with Bounded Variables
- 5.5 Artificial Variable
- 5.5.1 Two-Phase Method
- 5.5.2 Big-M Method
- 5.5.3 Single Artificial Variable Technique
- 5.6 Degeneracy and Cycling
- 5.6.1 Degeneracy
- 5.6.2 Cycling
- 5.6.3 Cycling Prevention Rules

Lexicographic Rule

Bland's Rule

Successive Ratio Rule

Duality Theory and Sensitivity Analysis

Decomposition Principle

Ellipsoid Algorithm

Projective Algorithm

Interior-Point Algorithm

Part III Graph and Network Theory

Paths, Trees, and Cycles

Shortest-Path Problem

Minimum Spanning Tree Problem

Maximum Flow Problem

Minimum Cost Flow Problem


Assignment and Matching Problem

Graph Algorithms

Polygon Triangulation

18.1 Types of Polygons

Def: A **simple polygon** is a closed polygonal curve without self-intersection.




Simple Polygon Non-simple Polygon

Polygons are basic building blocks in most geometric applications. It can model arbitrarily complex shapes, and apply simple algorithms and algebraic representation/manipulation.

18.2 Triangulation

Def: Triangulation is to partition polygon P into non-overlapping triangles using diagonals only. It reduces complex shapes to collection of simpler shapes. Every simple n-gon admits a triangulation which has n-2 triangles.



Triangulation

Theorem: Every polygon has a triangulation

Lemma: Every polygon with more than three vertices has a diagonal.

Proof: (by Meisters, 1975) Let P be a polygon with more than three vertices. Every vertex of a P is either *convex* or *concave*. W.L.O.G. (any polygon must has convex corner) Assume p is a convex vertex. Denote the neighbors of

p as q and r. If $q\bar{r}$ is a diagonal, done, and we call $\triangle pqr$ is an ear. If $\triangle pqr$ is not an ear, it means at least one vertex is inside $\triangle pqr$, assume among those vertexes inside $\triangle pqr$, s is a vertex closest to p, then $p\bar{s}$ is a diagonal.

18.3 Art Gallery Theorem

Problem: The floor plan of an art gallery modeled as a simple polygon with n vertices, there are guards which is stationed at fixed positions with 360 degree vision but cannot see through the walls. How many guards does the art gallery need for the security? (Fun fact: This problem was posted to Vasek Chvatal by Victor Klee in 1973)

Theorem: Every *n*-gon can be guarded with $\lfloor \frac{n}{3} \rfloor$ vertex guards

Lemma: Triangulation graph can be 3-colored.

Proof:

- P plus triangulation is a planar graph
- 3-coloring means there exist a 3-partition for vertices that no edge or diagonal has both endpoints within the same set of vertices.
- Proof by Induction:
 - Remove an ear (there will always exist ear)
 - Inductively 3-color the rest
 - Put ear back, coloring new vertex with the label not used by the boundary diagonal.

18.4 Triangulation Algorithms

18.5 Shortest Path

Part IV

Integer and Combinatorial Programming

Formulation

19.1 Typical Problems

19.2 Integer Programming Formulation Skills

19.2.1 A Variable Taking Discontinuous Values

Description: In algebraic notation:

$$x = 0$$
, or $l \le x \le u$

Modeling:

$$x \le uy$$
$$x \ge ly$$
$$y \in \{0, 1\}$$

where

$$y = \begin{cases} 0, & \text{if } x = 0\\ 1, & \text{if } l \le x \le u \end{cases}$$

19.2.2 Fixed Costs

Description: In algebraic notation:

$$C(x) = \begin{cases} 0 & \text{for } x = 0\\ k + cx & \text{for } x > 0 \end{cases}$$

Modeling:

$$C^*(x, y) = ky + cx$$
$$x \le My$$
$$x \ge 0$$
$$y \in \{0, 1\}$$

where

$$y = \begin{cases} 0, & \text{if } x = 0 \\ 1, & \text{if } x \ge 0 \end{cases}$$

19.2.3 Either-or Constraints

Description: In algebraic notation:

$$\sum_{j \in J} a_{1j} x_j \le b_1 \text{ or } \sum_{j \in J} a_{2j} x_j \le b_2$$

Modeling:

$$\sum_{j \in J} a_{1j} x_j \le b_1 + M_1 y$$

$$\sum_{j \in J} a_{2j} x_j \le b_2 + M_1 (1 - y)$$

$$y \in \{0, 1\}$$

where

$$y = \begin{cases} 0, & \text{if } \sum_{j \in J} a_{1j} x_j \le b_1 \\ 1, & \text{if } \sum_{j \in J} a_{2j} x_j \le b_2 \end{cases}$$

Notice that the sign before M is determined by the inequality \geq or \leq , if it is " \geq ", use "-", if it " \leq ", use "+".

19.2.4 Conditional Constraints

Description: If constraint A is satisfied, then constraint B must also be satisfied

If
$$\sum_{j \in J} a_{1j} x_j \le b_1$$
 then $\sum_{j \in J} a_{2j} x_j \le b_2$

The key part is to find the opposite of the first condition. We are using $A \Rightarrow B \Leftrightarrow \neg B \Rightarrow \neg A$ Therefore it is equivalent to

$$\sum_{j \in J} a_{1j} x_j > b_1 \text{ or } \sum_{j \in J} a_{2j} x_j \le b_2$$

Furthermore, it is equivalent to

$$\sum_{j \in J} a_{1j} x_j \ge b_1 + \epsilon \text{ or } \sum_{j \in J} a_{2j} x_j \le b_2$$

Where ϵ is a very small positive number.

Modeling:

$$\sum_{j \in J} a_{1j} x_j \ge b_1 + \epsilon - M_2 y$$

$$\sum_{j \in J} a_{2j} x_j \le b_2 + M_2 (1 - y)$$

$$y \in \{0, 1\}$$

19.2.5 Special Ordered Sets

SOS1 Description Out of a set of yes-no decisions, at most one decision variable can be yes.

$$x_1 = 1, x_2 = x_3 = \dots = x_n = 0$$
or
 $x_2 = 1, x_1 = x_3 = \dots = x_n = 0$
or ...

Modeling:

$$\sum_{i} x_i = 1, \quad i \in N$$

SOS2 Description 1 Out of a set of binary variables, at most two variables can be nonzero. In addition, the two variables must be adjacent to each other in a fixed order list.

Modeling: If $x_1, x_2, ..., x_n$ is a SOS2, then

$$\sum_{i=1}^{n} x_i \le 2$$

$$x_i + x_j \le 1, \forall i \in \{1, 2, ..., n\}, j \in \{i + 2, i + 3, ..., n\}$$

$$x_i \in \{0, 1\}$$

SOS2 Description 2 There is another type of definition, that is out of a set of nonnegative variables not binary here, at most two variables can be nonzero. In addition, the two variables must be adjacent to each other in a fixed order list. All variables summing to 1.

This definition of SOS2 is used in the following section Piecewise Linear Formulations

19.2.6 Piecewise Linear Formulations

Description: The objective function is a sequence of line segments, e.g. y = f(x), consists k - 1 linear segments going through k given points $(x_1, y_1), (x_2, y_2), ..., (x_k, y_k)$. Denote

$$d_i = \begin{cases} 1, & x \in (x_i, x_{i+1}) \\ 0, & \text{otherwise} \end{cases}$$

Then the objective function is

$$\sum_{i \in \{1,2,\dots,k-1\}} y = d_i f_i(x)$$

Modeling: Given that objective function as a piecewise linear formulation, we can have these constraints

$$\sum_{i \in \{1,2,\dots,k-1\}} d_i = 1$$

$$d_i \in \{0,1\}, i \in \{1,2,\dots,k-1\}$$

$$x = \sum_{i \in \{1,2,\dots,k\}} w_i x_i$$

$$y = \sum_{i \in \{1,2,\dots,k\}} w_i y_i$$

$$w_1 \le d_1$$

$$w_i \le d_{i-1} + di, i \in \{2,3,\dots,k-1\}$$

$$w_k \le d_{k-1}$$

In this case, $w_i \in SOS2$ (second definition)

19.2.7 Conditional Binary Variables

Description: Choose at most n binary variable to be 1 out of $x_1, x_2, ... x_m, m \ge n$. If n = 1 then it is SOS1.

Modeling:

$$\sum_{k \in \{1, 2, \dots, m\}} x_k \le n$$

Description: Choose exactly n binary variable to be 1 out of $x_1, x_2, ... x_m, m \ge n$ **Modeling:**

$$\sum_{k \in \{1,2,\dots,m\}} x_k = n$$

Description: Choose x_j only if $x_k = 1$ Modeling:

$$x_j = x_k$$

Description: "and" condition, iff $x_1, x_2, ..., x_m = 1$ then y = 1 Modeling:

$$y \le x_i, i \in \{1, 2, ..., m\}$$

 $y \ge \sum_{i \in \{1, 2, ..., m\}} x_i - (m - 1)$

19.2.8 Elimination of Products of Variables

Description: For variables x_1 and x_2 ,

$$y = x_1 x_2$$

Modeling: If x_1, x_2 are binary, it is the same as "and" condition of binary variables. If x_1 is binary, while x_2 is continuous and $0 \le x_2 \le u$, then

$$y \le ux_1$$

$$y \le x_2$$

$$y \ge x_2 - u(1 - x_1)$$

$$y \ge 0$$

If both x_1 and x_2 are continuous, it is non-linear, we can use Piecewise linear formulation to simulate.

Branch and Bound

Branch and Cut

Packing and Matching

Traveling Salesman Problem

Knapsack Problem

${\bf Part~V}$ ${\bf Nonlinear~Programming}$

KKT Optimality Conditions

Lagrangian Duality

Unconstrained Optimization

Penalty and Barrier Functions

Part VI Algorithms and Computational Complexity

Computational Complexity

Sorting

- 30.1 Elementary Sorting Algorithms
- 30.2 Heap-sort
- 30.3 Quick-sort
- 30.4 Sorting in Linear Time
- 30.5 Medians and Order Statistics

Data Structures

- 31.1 Elementary Data Structures
- 31.2 Hash Tables
- 31.3 Binary Search Trees
- 31.4 Red-Black Trees
- 31.5 B-Trees
- 31.6 Fibonacci Heaps
- 31.7 van Emde Boas Trees

Design and Analysis Techniques

32.1	Dynamic Programming
32.2	Greedy Algorithms
32.3	Amortized Analysis
32.4	Multi-threaded Algorithms
32.5	Matrix Operations
32.6	Polynomials and the FFT
32.7	Number-Theoretic Algorithms
32.8	String Matching
32.9	Computational Geometry
32.10	NP-Completeness
32.11	Approximation Algorithms

Part VII Heuristics ans Meta-heuristics

Part VIII Game Theory

Games with Ordinal Payoffs

- 33.1 Ordinal Games in Strategic Form
- 33.2 Perfect-information Games
- 33.3 General Dynamic Games

Games with Cardinal Payoffs

- 34.1 Expected Utility Theory
- 34.2 Strategic-form Games
- 34.3 Extensive-form Games

Knowledge, Common Knowledge, Beliefs

- 35.1 Common Knowledge
- 35.2 Adding Beliefs to Knowledge
- 35.3 Rationality

Refinements of Subgame-perfect Equilibrium

- 36.1 Weak Sequential Equilibrium
- 36.2 Sequential Equilibrium
- 36.3 Perfect Bayesian Equilibrium

Incomplete Information

- 37.1 Static Games
- 37.2 Dynamic Games
- 37.3 The Type-Space Approach

Part IX

Probability, Stochastic Processes and Markov Chains

Probability

Random Variables

39.1 Relationship between Some Random Variables




Figure 39.1: Relationship between Some Random Variables

9.2 Discrete Random Variables

Table 39.1: Discrete Random Variables

MGF	$M(t) = \frac{e^{at} - e^{(b+1)t}}{(b-a+1)(1-e^t)}$ $t \in \mathbb{R}$	$M(t) = 1 - p + pe^t$ $t \in \mathbb{R}$	$M(t) = (1 - p + pe^t)^n$ $t \in \mathbb{R}$	$M(t) = e^{\mu(e^t - 1)}$ $t \in \mathbb{R}$	$M(t) = \frac{p}{1 - (1 - p)e^t}$ $t < -\ln(1 - p)$	$M(t) = \left(\frac{p}{1 - (1 - p)e^t}\right)^n$ $t < -\ln(1 - p)$
Variance	$D[X] = \frac{(b-a+1)^2 - 1}{12}$	D[X] = p(1-p)	D[X] = np(1-p)	$D[X] = \mu$	$D[X] = \frac{1-p}{p^2}$	$D[X] = \frac{n(1-p)}{p^2}$
Expectation	$E[X] = \frac{b - a}{2}$	E[X] = p	E[X] = np	$E[X] = \mu$	$E[X] = \frac{1-p}{p}$	$E[X] = \frac{n(1-p)}{p}$
CDF	$F(x) = \frac{x - a + 1}{b - a + 1}$ $x = a, a + 1,, b$	$F(x) = \begin{cases} 0, & x < 0 \\ 1 - p, & 0 \le x \le 1 \\ 1, & x > 1 \end{cases}$	$F(x) = \sum_{k=0}^{x} \binom{n}{k} p^{k} (1-p)^{n-k}$ $x = 0, 1,, n$	$f(x) = \frac{\Gamma(x+1,\mu)}{\Gamma(x+1)}$ x = 0, 1,, n,	$F(x) = 1 - (1 - p)^{x+1}$ x = 0, 1,, n,	$F(x) = 1 - I_p(k+1, n)$ x = 0, 1, 2,, n,
PMF	$f(x) = \frac{1}{b-a+1}$ x = a, a+1,, b	$f(x) = p^{x} (1 - p)^{1 - x}$ $x \in \{0, 1\}$	$f(x) = \binom{n}{x} p^x (1-p)^{n-x}$ $x = 0, 1,, n$	$f(x) = \frac{\mu^x e^{\mu}}{x!}$ $x = 0, 1, \dots, n, \dots$	$f(x) = p(1-p)^x$ x = 0, 1,, n,	$f(x) = \binom{n-1+x}{x} p^n (1-p)^x$ $x = 0, 1, 2,, n,$
Distribution	Discrete Uniform (a, b)	$\mathrm{Bernoulli}(p)$	$\operatorname{Binomial}(n,p)$	$\mathrm{Poisson}(\mu)$	$\operatorname{Geometric}(p)$	$\operatorname{Pascal}(n,p)$

39.3 Continuous Random Variables

Table 39.2: Continuous Random Variables

MGF	$M(t) = \begin{cases} 1, & t = 0\\ \frac{e^{bt} - e^{at}}{t(b - a)}, & t \neq 0 \end{cases}$	$e^{\frac{t(t\sigma^2+2\mu)}{2}}$ $t \in \mathbb{R}$	$rac{1}{1-rac{t}{\lambda}}$	$rac{1}{(1-rac{1}{\zeta})^n}$ $t<\lambda$
Variance	$D[X] = \frac{(b-a)^2}{12}$	$D[X] = \sigma^2$	$D[X] = \frac{1}{\lambda^2}$	$D[X] = \frac{n}{\lambda^2}$
Expectation	$E[X] = \frac{b - a}{2}$	$E[X] = \mu$	$E[X] = \frac{1}{\lambda}$	$E[X] = \frac{n}{\lambda}$
CDF	$F(x) = \frac{x - a}{b - a}$ $x = [a, b]$	$F(x) = \int_{-\infty}^{x} \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$ $x \in \mathbb{R}$	$F(x) = 1 - e^{\lambda x}$ $x > 0$	$F(x) = 1 - \sum_{i=0}^{n-1} \frac{\lambda^n x^n e^{-\lambda x}}{n!}$ $x > 0$
PDF	$f(x) = \frac{1}{b-a}$ $x = [a, b]$	$f(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$ $x \in \mathbb{R}$	$f(x) = \lambda e^{-\lambda x}$ $x > 0$	$f(x) = \frac{\lambda^n x^{n-1} e^{-\lambda x}}{(n-1)!}$ $x > 0$
Distribution	$\operatorname{Uniform}(a,b)$	Normal (μ, σ) $f(x) = \frac{1}{\sqrt{2\pi\sigma}}e^{-\frac{1}{x}}$	$\operatorname{Exponential}\lambda)$	$\operatorname{Erlang}(n,\lambda)$

Limit Theorems

The Bernoulli and Poisson Process

Discrete-Time Markov Chains

Continuous-Time Markov Chains

$\begin{array}{c} {\rm Part} \ {\rm X} \\ \\ {\rm Queuing} \ {\rm Theory} \end{array}$

Queuing Model

Birth-and-Death Queuing Models

Multidimensional Birth-and-Death Queuing Models

Phase-Type Queue

Bulk Queue

Imbedded-Markov-Chain Queuing Models

Queuing Network

Part XI Inventory Theory

Part XII Reliability Theory

Part XIII

Statistic

Part XIV Simulation