



Alcohol dehydration

A Level



Part A Preparation of ethene

In a preparation of ethene, ethanol is added a drop at a time to a heated reagent **Y**. The impure ethene is washed by being bubbled through a solution **Z** and then collected. What are reagent **Y** and solution **Z** likely to be?

	reagent Y	solution Z
A	acidified $\text{K}_2\text{Cr}_2\text{O}_7$	dilute NaOH
B	concentrated H_2SO_4	dilute H_2SO_4
C	concentrated H_2SO_4	dilute NaOH
D	ethanolic NaOH	concentrated H_2SO_4
E	ethanolic NaOH	dilute NaOH

- ☐ **A**
- ☐ **B**
- ☐ **C**
- ☐ **D**
- ☐ **E**

Part B Dehydration of propan-1-ol

Propan-1-ol, $\text{C}_3\text{H}_7\text{OH}$, is dehydrated by passing its vapour over hot aluminium oxide to give a hydrocarbon.

Which structural formula represents the product obtained when the hydrocarbon reacts with bromine?




Figure 1: Possible structures after dehydration and bromination of propan-1-ol

- ☐ **A**
- ☐ **B**
- ☐ **C**
- ☐ **D**

Part A adapted with permission from UCLES, A-Level Chemistry, June 1990, Paper 1, Question 27;

Part B adapted with permission from UCLES, A-Level Chemistry, June 1996, Paper 3, Question 21



Alcohol reactions

The compound C_3H_8O has two isomers that are alcohols. These isomers can undergo a series of reactions with the reagents shown giving organic products.

Deduce the identity of each of the organic products **A** to **D**. Use the [structure editor](#) to generate SMILES strings as your answers.




Figure 1: Reactions of alcohols

In the editor, after drawing your structure, click on the round, yellow smiley face to generate a SMILES string. Copy the SMILES string and paste it in the answer box.

[Using the structure editor](#)

Part A **A**

A is:

Part B B

B is:

Part C C

C is:

Part D D

D is:

Adapted with permission from UCLES, A-Level Modular Sciences, November 1996, Chains and Rings, Question 3

All materials on this site are licensed under the [**Creative Commons license**](#), unless stated otherwise.




Aldehyde and alcohol reactions

A Level



Part A Bioluminescence

The production of light by animals and plants is known as bioluminescence. It sometimes involves the following reaction:



What type of reaction is this?

- ☐ Substitution
- ☐ Elimination
- ☐ Addition
- ☐ Reduction
- ☐ Oxidation

Part B Butan-2-ol with potassium dichromate(VI)

Which of the following are produced when an aqueous solution of butan-2-ol is refluxed with potassium dichromate(VI) in dilute sulfuric acid?

1 butanal

2 butanoic acid

3 butanone

- ☐ 1 and 2 only are correct
- ☐ 1, 2 and 3 are correct
- ☐ 1 only is correct
- ☐ 2 and 3 only are correct
- ☐ 3 only is correct

All materials on this site are licensed under the [**Creative Commons license**](#), unless stated otherwise.

Butanol dehydration

A Level



The four different isomers of butanol (**1** - **4**) can be dehydrated to give four isomers of butene (**A** - **D**)









**1****2****3****4****A****B****C****D**

Figure 1: Isomers of butanol and butene

Part A A

Which isomer(s) of butanol could give rise to butene **A**?

- ☐ 2 only
 - ☐ 3 only
 - ☐ 1 only
 - ☐ 3 and 4 only
 - ☐ 1 and 4 only
 - ☐ 1 and 2 only
 - ☐ 2 and 3 only
 - ☐ 4 only
-

Part B B

Which isomer(s) of butanol could give rise to butene **B**?

- ☐ 3 only
 - ☐ 1 only
 - ☐ 4 only
 - ☐ 3 and 4 only
 - ☐ 2 and 3 only
 - ☐ 1 and 4 only
 - ☐ 1 and 2 only
 - ☐ 2 only
-

Part C C

Which isomer(s) of butanol could give rise to butene **C**?

- ☐ 2 only
 - ☐ 3 only
 - ☐ 1 and 2 only
 - ☐ 2 and 3 only
 - ☐ 1 only
 - ☐ 4 only
 - ☐ 3 and 4 only
 - ☐ 1 and 4 only
-

Part D D

Which isomer(s) of butanol could give rise to butene **D**?

- ☐ 4 only
 - ☐ 3 only
 - ☐ 3 and 4 only
 - ☐ 1 only
 - ☐ 2 only
 - ☐ 1 and 4 only
 - ☐ 2 and 3 only
 - ☐ 1 and 2 only
-

Part E Reaction type

What type of reaction is this?



Butanol oxidation

A Level




Dilute acidified sodium dichromate(VI) is used to distinguish between primary, secondary and tertiary alcohols. Draw full structural formulae of the final organic products (if any) when the following alcohols are treated with this reagent.

Use the [structure editor](#) to generate a SMILES string.


In the editor, after drawing your structure, click on the round, yellow smiley face to generate a SMILES string. Copy the SMILES string and paste it in the answer box.

[Using the structure editor](#)


Part A $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$



Part B $\text{CH}_3\text{CH}_2\text{CH}(\text{OH})\text{CH}_3$



Part C $(\text{CH}_3)_3\text{COH}$



Adapted with permission from UCLES, A-Level Chemistry, June 1990, Paper 2, Question 3

Ketones with HCN steps

A Level



Part A Steps in mechanism

Hydrogen cyanide HCN adds to propanone most readily at a pH value between 9 and 10. Which of the following are likely steps in the reaction at this pH range?






Figure 1: Possible steps in reaction of hydrogen cyanide with propanone at pH 9 - 10

- ☐ 1, 2 and 3 are correct
- ☐ 2 and 3 only are correct
- ☐ 3 only is correct
- ☐ 1 and 2 only are correct
- ☐ 1 only is correct

Part B Rate at different pH values

Hydrogen cyanide adds to ketones. The table shows the relative rates of reaction in aqueous solutions under different conditions.

condition	relative rate
neutral solution	slow
acidified solution	virtually zero
alkaline solution	very rapid

Which of the following is likely to be involved in the rate-determining step of the reaction?

- ☐ H^+
- ☐ H_2O
- ☐ CN^-
- ☐ HCN

Part A adapted with permission from UCLES, A-Level Chemistry, June 1996, Paper 3, Question 37;

Part B adapted with permission from UCLES, A-Level Chemistry, June 1989, Paper 3, Question 37

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.



Physics. *You work it out.*

[Home](#)

Unknown page

Page not found

We're sorry, page not found: /questions/ketones_with_kcn_mechanism

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.



Nucleophilic addition mechanism

A Level



Aldehydes and ketones typically react by nucleophilic addition reactions.

Part A HCN with ketones first step

What is the sequence of curly arrows denoting movement of electrons in the first step of the reaction between propanone and HCN catalysed by KCN?




Figure 1: First step of HCN with propanone catalysed by KCN

For example in the reaction below, if you think the mechanism is as shown, your answer would be **cdef**.




Figure 2: Sequence **cdef**

Part B HCN with ketones second step

What is the sequence of curly arrows denoting possible movement of electrons in the second step of the reaction between propanone and HCN catalysed by KCN that would regenerate the catalyst?




Figure 3: Second step of HCN with propanone catalysed by KCN



Reactions of C₄H₁₀O

A Level



Part A Elimination

A compound C₄H₁₀O reacts with sodium, is not affected by warm acidified potassium dichromate(VI) solution, and eliminates water when warmed with concentrated sulfuric acid. What could the compound be?

- ☐ CH₃CH₂CH₂OCH₃
- ☐ CH₃CH₂CH(OH)CH₃
- ☐ (CH₃)₃COH
- ☐ CH₃CH₂CH₂CH₂OH
- ☐ CH₃CH₂OCH₂CH₃

Part B Oxidation

A compound **X**, C₄H₁₀O, gives the compound **Y**, C₄H₈O, on oxidation. **Y** does **not** give a silver mirror on the addition of Tollens' reagent. Which of the following could **X** be?

- ☐ CH₃CH₂CH₂CH₂OH
- ☐ CH₃CH₂OCH₂CH₃
- ☐ CH₃CH₂CH₂OCH₃
- ☐ (CH₃)₃COH
- ☐ CH₃CH₂CH(OH)CH₃

Part A adapted with permission from OCSEB, A-Level Chemistry, June 1995, Paper 1, Question 28;

Part B created for isaacphysics.org by R. Less



Tollens' reagent

A Level



When propanal reacts with Tollens' reagent, what are the principal inorganic and organic products?

- ☐ AgNO_3 and $\text{CH}_3\text{CH}_2\text{COOH}$
- ☐ Ag and $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$
- ☐ Ag and $\text{CH}_3\text{CH}_2\text{COOH}$
- ☐ Ag_2O and $\text{CH}_3\text{CH}_2\text{COOH}$

Adapted with permission from UCLES, A-Level Chemistry, June 1994, Paper 4, Question 24

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.