



Block on a Rough Table

A Level



This problem involves friction, which is not covered in some Physics A Levels. For more information please check with your teacher.

A block is being pulled along a rough table at constant speed by a horizontal force of magnitude F . The block has mass $m = 2.0 \text{ kg}$ and the coefficient of dynamic friction between the block and the table is $\mu = 0.50$.

(Take $g = 9.81 \text{ m s}^{-2}$) .

Find F .



Essential GCSE Physics 7.4

Year 9 GCSE



The number of widgets made in a factory each week is proportional to the number of workers and the number of hours each worker works. When the factory employed 25 staff, each working 35 hours / week, 65 400 widgets were made each week.

Part A 40 staff for 30 hours/week

How many widgets would be made each week if 40 staff worked for 30 hours per week?

Part B 130000 widgets

If we need 130 000 widgets made each week, and the staff will work 42 hours / week, how many workers are needed?



Physics. *You work it out.*

[Home](#) [Physics](#) [Skills](#) [Relationships](#) [Essential Pre-Uni Physics A10.1](#)

Essential Pre-Uni Physics A10.1



If $V \propto I$ and $I = 0.35 \text{ A}$ when $V = 9.6 \text{ V}$, what will V be when $I = 0.90 \text{ A}$?

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Physics](#) [Skills](#) [Relationships](#) [Essential Pre-Uni Physics A10.3](#)

Essential Pre-Uni Physics A10.3



If $g \propto 1/r^2$ and $g = 9.8 \text{ N kg}^{-1}$ when $r = 6400 \text{ km}$, what will g be when $r = 15000 \text{ km}$?

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Physics](#) [Skills](#) [Relationships](#) [Essential Pre-Uni Physics A10.4](#)

Essential Pre-Uni Physics A10.4



If $E \propto x^2$ and $E = 2.5 \text{ J}$ when $x = 1.5 \text{ cm}$, what will x be when $E = 6.0 \text{ J}$?

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.



Essential Pre-Uni Physics A10.9

GCSE A Level



If $R = \rho L / A$, and $R = 5.0 \, \Omega$ when $L = 65 \, \text{m}$ and $A = 2.5 \, \text{mm}^2$, what will R be when $L = 120 \, \text{m}$ and $A = 1.5 \, \text{mm}^2$ if ρ doesn't change?



Physics. *You work it out.*

[Home](#) [Physics](#) [Skills](#) [Relationships](#) [Essential Pre-Uni Physics A9.1](#)

Essential Pre-Uni Physics A9.1



Give the factor by which the quantity changes to two significant figures. So if it doubles, your answer is 2.0, and if it halves your answer is 0.50.

By what factor does $E = mv^2/2$ change if v doubles?

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.



Essential Pre-Uni Physics A9.10

GCSE A Level



Give the percentage change (to two significant figures) in the quantity. Use "+" or "-" to indicate increase or decrease (so a 3.0 % decrease would be given as -3.0%).

Part A Percentage change in E

In $E = VIt$, what is the percentage change in E if V increases by 1.0 %, I decreases by 2.0 % and t increases by 3.0 %?

Part B Percentage change in R

In $R = \rho L / A$, what is the percentage change in R if L increases by 7.0 % and A increases by 3.0 %?



Essential Pre-Uni Physics A9.2



Give the factor by which the quantity changes to two significant figures. So if it doubles, your answer is 2.0, and if it halves your answer is 0.50.

Part A r is multiplied by 3.3

By what factor does $V = -GM/r$ change if r is multiplied by 3.3?

Part B r is multiplied by 0.64

By what factor does $g = GM/r^2$ change if r is multiplied by 0.64?



Essential Pre-Uni Physics A9.4

GCSE A Level



Give the factor by which the quantity changes to two significant figures. So if it doubles, your answer is 2.0, and if it halves your answer is 0.50.

By what factor does d need to change in $I = P/(4\pi d^2)$ if P were multiplied by 5.2 and I were to not change?
