



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Two Particles on a Pulley](#)

Two Particles on a Pulley

A Level




Figure 1: Particles A and B attached to either end of a string.

Particles A and B are attached to the ends of a light inextensible string. The string passes over a smooth fixed pulley. The particles are released from rest, with the string taut, and A and B at the same height above a horizontal floor. In the subsequent motion A descends with acceleration 1.4 m s^{-2} and strikes the floor 0.80 s after being released. It is given that B never reaches the pulley.

Part A Distance to the floor

Calculate the distance A moves before it reaches the floor. Give your answer to 2 significant figures.

Part B Further B rises

Calculate how much further B rises after A hits the floor. Give your answer to 2 significant figures.

Part C Total time of rising

Calculate the total length of time during which B is rising. Give your answer to 2 significant figures.

Part D Motion of B

Sketch the (t, v) graph for the motion of B from the instant it is released from rest until it reaches a position of instantaneous rest.

To see an example of the sketch, answer the following question:

What is the maximum speed of B ? Give your answer to 3 significant figures.

Part E Mass of A

Before A strikes the floor the tension in the string is 5.88 N . Calculate the mass of A . Give your answer to 2 significant figures.

Part F Mass of B

Calculate the mass of B . Give your answer to 2 significant figures.

Part G Pulley's support 1

The pulley has mass 0.50 kg and is held in a fixed position by a light vertical chain.

Calculate the tension in the chain immediately **before** A strikes the floor. Give your answer to 3 significant figures.

Part H Pulley's support 2

Calculate the tension in the light vertical chain immediately **after** A strikes the floor. Give your answer to 2 significant figures.

Used with permission from UCLES, A Level Maths, January 2008, OCR M1, Question 7

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Kinematics & Calculus](#)

Kinematics & Calculus

A Level




Figure 1: Velocity-time graph of the motion of two particles A and B along the same straight line.

The diagram shows the (t, v) graphs for two particles A and B which move on the same straight line. The units of v and t are m s^{-1} and s respectively. Both particles are at the point S on the line when $t = 0$. The particle A is initially at rest, and moves with acceleration $0.18t \text{ m s}^{-2}$ until the two particles collide when $t = 16 \text{ s}$. The initial velocity of B is $U \text{ m s}^{-1}$ and B has variable acceleration for the first five seconds of its motion. For the next ten seconds of its motion B has a constant velocity of 9 m s^{-1} ; finally B moves with constant deceleration for one second before it collides with A .

Part A t for same velocity

Calculate the value of t at which the two particles have the same velocity.

Part B Calculate U

For $0 \leq t \leq 5$ the distance of B from S is $(Ut + 0.08t^3)$ m.

Calculate U .

Part C Distance from S

Calculate how far B is from S when $t = 5$ s.

Part D v_B when $t = 16$ s

Calculate the velocity of B when $t = 16$ s.

Used with permission from UCLES, A Level Maths, June 2016, OCR M1, Question 7

Gameboard:

STEM SMART Single Maths 16 - Mechanics Revision

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Particle on a Surface](#)

Particle on a Surface

A Level




Figure 1: Two forces of magnitude 6 N and 10 N acting on a particle P .

Two forces of magnitudes 6 N and 10 N separated by an angle of 110° act on a particle P , which rests on a horizontal surface.

Part A Resultant magnitude

Find the magnitude of the resultant of the 6 N and 10 N forces. Give your answer to 3 significant figures.

Part B Resultant angle

Calculate the angle between the resultant force and the 10 N force. Give your answer to 3 significant figures.

Part C Surface force

The two forces act in the same vertical plane. The particle P has weight 20 N and rests in equilibrium on the surface. If you have not studied resolving forces then please visit the [learning_hexagon](#) on the gameboard.

Given that the surface is smooth, find the magnitude of the force exerted on P by the surface. Give your answer to 3 significant figures.

Part D Angle to 10 N force

Given that the surface is smooth, find the angle between the surface and the 10 N force. Give your answer to 3 significant figures.

Used with permission from UCLES, A Level Maths, June 2015, OCR M1, Question 4

Gameboard:

STEM SMART Single Maths 16 - Mechanics Revision

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Pulley with Three Masses](#)

Pulley with Three Masses

A Level




Figure 1: Particles P , Q , and R attached along a string that passes over a pulley.

Particles P and Q , of masses m kg and 0.05 kg respectively, are attached to the ends of a light inextensible string which passes over a smooth pulley. Q is attached to a particle R of mass 0.45 kg by a light inextensible string. The strings are taut, and the portions of the strings not in contact with the pulley are vertical. P is in contact with a horizontal surface when the particles are released from rest. The tension in the string QR is 2.52 N during the descent of R .

Part A Light string

The string is described as light. Explain how this modelling assumption affects calculations when **treating the whole system as a single particle**.

Easier question?

The string is described as light. Explain how this modelling assumption affects calculations when **finding the tension in a vertical string**.

Easier question?

Part B Acceleration of R

Find the acceleration of R during its descent. Give your answer to 2 significant figures.

Part C Tension in PQ

By considering the motion of Q , calculate the tension in the string PQ during the descent of R . Give your answer to 2 significant figures.

Part D Finding m

Find the value of m . Give your answer to 2 significant figures.

Part E Max height of P

R strikes the surface 0.5 s after release and does not rebound. During their subsequent motion, P does not reach the pulley and Q does not reach the surface.

Calculate the greatest height of P above the surface. Give your answer to 2 significant figures.

Adapted with permission from UCLES, A Level Maths, January 2012, OCR M1, Question 7

Gameboard:

STEM SMART Single Maths 16 - Mechanics Revision

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Friction](#)

Friction

A Level



Part A Force diagram 1




Figure 1: Block B connected by a string that passes over a pulley to a particle P .

A block B of mass 0.4 kg and a particle P of mass 0.3 kg are connected by a light inextensible string. The string passes over a smooth pulley at the edge of a rough horizontal table. B is in contact with the table and the part of the string between B and the pulley is horizontal. P hangs freely below the pulley.

The system is in equilibrium with the string taut and P on the point of moving downwards.

Draw a diagram to show the forces associated with the particles and the strings.

Easier question?

Part B Frictional force for B

Find the frictional force between B and the table. Give your answer to 2 significant figures.

Part C Force diagram 2

A horizontal force of magnitude X N, acting directly away from the pulley, is now applied to B . The system is again in equilibrium with the string taut, and with P now on the point of moving upwards.

Draw a diagram to show the forces associated with the particles and the strings. You may assume that the magnitude of the frictional force is unchanged.

Easier question?

Part D Value of X

Find the value of X . Give your answer to 2 significant figures.

Part E Deceleration in curling

In the sport of curling, a heavy stone is projected across a horizontal ice surface. One player projects a stone of weight 180 N, which moves 36 m in a straight line and comes to rest 24 s after the instant of the projection. The only horizontal force acting on the stone after its projection is a constant frictional force between the stone and the ice.

Calculate the deceleration of the stone.

Part F Magnitude of friction

Find the magnitude of the frictional force acting on the stone. Give your answer to 3 significant figures.

Used with permission from UCLES, A Level Maths, January 2007, OCR M1 & January 2012, OCR M1

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.