

# 1. Motivation

## 1.1. Plane curves

- Curves mainly parametrised:  $\alpha : I \rightarrow \mathbb{R}^2$ ,  $I \subset \mathbb{R}$  interval, with a direction.
- **Four vertex theorem**: every closed plane curve has at least 4 vertices.

## 1.2. Surfaces

- Surfaces are 2-dimensional subsets of  $\mathbb{R}^3$ .

# 2. Regular curves in $\mathbb{R}^n$

## 2.1. Regular curves, length and tangent vectors

- Let  $I$  be open interval, then  $\underline{\alpha} : I \rightarrow \mathbb{R}^n$  is **parametrised curve**.
- $\underline{\alpha}$  is **smooth** if  $\underline{\alpha}(u) = (\alpha_1(u), \dots, \alpha_n(u))$  where all  $\alpha_i : I \rightarrow \mathbb{R}$  are smooth maps.
- Image  $\underline{\alpha}(I) \subset \mathbb{R}^n$  is the **trace**.
- **Tangent vector of  $\alpha$  at  $u$**  is

$$\underline{\alpha}'(u) = (\alpha'_1(u), \dots, \alpha'_n(u))$$

- $\underline{\alpha}$  is **regular** if  $\forall u \in I, \underline{\alpha}'(u) \neq 0$ .  $\underline{\alpha}$  is **singular at  $u$**  if  $\underline{\alpha}'(u) = 0$ .
- If  $\underline{\alpha}$  is regular, **unit tangent vector of  $\alpha$  at  $u$**  is

$$\underline{t}(u) = \underline{\alpha}' \frac{u}{\|\underline{\alpha}'(u)\|}$$

- If  $\forall u \in I, \|\underline{\alpha}'(u)\| = 1$  then  $\underline{\alpha}$  is a **unit speed curve**. If  $\forall u \in I, \|\underline{\alpha}'(u)\| = c$ ,  $\underline{\alpha}$  is **constant speed curve**.
- **Example**: unit circle  $\underline{\alpha}(u) = (\cos u, \sin u)$  is regular:  $\alpha'(u) = (-\sin u, \cos u) \neq 0$ . It is unit speed as  $\|\alpha'(u)\| = 1$ .
- **Example**: helix  $\underline{\alpha}(u) = (\cos u, \sin u, u)$ ,  $\alpha'(u) = (-\sin u, \cos u, 1)$ ,  $\|\alpha'(u)\| = \sqrt{2}$  so constant speed.
- **Example**: cusp  $\underline{\alpha}(u) = (u^3, u^2)$ ,  $\alpha'(u) = (3u^2, 2u)$  so  $\alpha'(u) = 0 \iff u = 0$  so  $\underline{\alpha}$  singular at 0.
- **Example**: node  $\underline{\alpha}(u) = (u^3 - u, u^2 - 1)$ . So  $\alpha(-1) = \alpha(1) = (0, 0)$  so it has a self-intersection at the origin.  $\alpha'(u) = (3u^2 - 1, 2u)$  so is regular.
- **Definition**: let  $\underline{\alpha} : I \rightarrow \mathbb{R}^n$ ,  $[a, b] \subset I$ .  $\underline{\alpha}$  is **rectifiable** on  $[a, b]$  if

$$L(\underline{\alpha}|_{[a,b]}) := \sup \left\{ \sum_{i=0}^{n-1} \|\underline{\alpha}(u_{i+1}) - \underline{\alpha}(u_i)\| : n \in \mathbb{N}, a = u_0 < \dots < u_m = b \right\}$$

is finite. Then  $L(\underline{\alpha}|_{[a,b]})$  is the **(arc) length** of  $\underline{\alpha} : [a, b] \rightarrow \mathbb{R}^n$ .

- **Proposition**: let  $\underline{\alpha} : I \rightarrow \mathbb{R}^n$  smooth,  $[a, b] \subset I$ . Then

$$L(\underline{\alpha}|_{[a,b]}) = \int_a^b \|\underline{\alpha}'(u)\| du$$

## 2.2. Reparametrisation

- **Definition:** let  $\underline{\alpha} : I \rightarrow \mathbb{R}^n$  be smooth regular curve. A **parameter change** for  $\alpha$  is a smooth map  $h : J \rightarrow I$ ,  $J \subset \mathbb{R}$  is open interval, where
  - $\forall t \in J, h'(t) \neq 0$
  - $h(J) = I$ .

$\tilde{\alpha} = \underline{\alpha} \circ h : J \rightarrow \mathbb{R}^n$  is a **reparametrisation** of  $\underline{\alpha}$ . If  $h' > 0$ ,  $h$  is **orientation preserving**, otherwise it is **orientation reversing**.

- **Proposition:** let  $\underline{\alpha} : I \rightarrow \mathbb{R}^n$  be smooth, regular curve,  $u_0 \in I$ ,  $\ell : I \rightarrow \mathbb{R}$  defined by

$$\ell(u) = \int_{u_0}^u \|\underline{\alpha}'(t)\| dt$$

Let  $J = \ell(I)$ . Then  $\ell$  is strictly monotone increasing and  $\tilde{\alpha} = \underline{\alpha} \circ \ell^{-1} : J \rightarrow \mathbb{R}^n$  is unit speed.

- **Proposition:** let  $\underline{\alpha} : I \rightarrow \mathbb{R}^n$  be smooth regular curve and  $\tilde{\alpha} := \underline{\alpha} \circ h : J \rightarrow \mathbb{R}^n$  be reparametrisation of  $\underline{\alpha}$  with parameter change  $h : J \rightarrow I$ . Let  $[a, b] \subset I$  and  $[c, d] = h^{-1}([a, b])$ . Then

$$L(\underline{\alpha}|_{[a,b]}) = L(\tilde{\alpha}|_{[c,d]})$$

i.e. length is independent of parametrisation.

### 3. Plane curves

#### 3.1. Unit normal vectors and curvature

- **Definition:** let  $\alpha : I \rightarrow \mathbb{R}^2$  be smooth regular plane curve. The **unit normal vector** of  $\alpha$  at  $u$  is

$$\underline{n}_{\alpha}(u) = \underline{t}(u) \begin{bmatrix} \cos(\pi/2) & \sin(\pi/2) \\ -\sin(\pi/2) & \cos(\pi/2) \end{bmatrix} = (-t_2(u), t_1(u))$$

- **Lemma:** let  $\alpha : I \rightarrow \mathbb{R}^2$  be smooth unit speed plane curve. Then  $\underline{t}'(s) = \alpha''(s)$  is parallel to  $\underline{n}(s)$ .
- **Definition: (signed) curvature**  $\kappa(s)$  of unit speed plane curve  $\alpha : I \rightarrow \mathbb{R}^2$  at  $s \in I$  is defined by

$$\underline{t}'(s) = \kappa(s)\underline{n}(s)$$

Note that we can compute  $\kappa(s)$  by

$$\underline{t}'(s) \cdot \underline{n}(s) = \kappa(s)\underline{n}(s) \cdot \underline{n}(s) = \kappa(s)\|\underline{n}(s)\|^2 = \kappa(s)$$

- **Rule for sign of curvature:** if curve turns clockwise, curvature is negative, if curve turns anti-clockwise, its curvature is positive.
- **Proposition:** let  $\alpha : I \rightarrow \mathbb{R}^2$  be any smooth regular plane curve,  $\alpha(u) = (x(u), y(u))$ . Then its curvative is

$$\kappa(u) = \frac{x'(u)y''(u) - x''(u)y'(u)}{\left((x'(u))^2 + (y'(u))^2\right)^{3/2}}$$