



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Calculus](#) [Differentiation](#) [Powers Using Chain Rule 2](#)

# Powers Using Chain Rule 2

A Level Further A

P P P P P P

## Part A First and second derivatives of $u = 2(3 - 2v)^{\frac{3}{2}}$

Find  $\frac{du}{dv}$  when  $u = 2(3 - 2v)^{\frac{3}{2}}$ .

The following symbols may be useful:  $v$

Find  $\frac{d^2u}{dv^2}$  when  $u = 2(3 - 2v)^{\frac{3}{2}}$ .

The following symbols may be useful:  $v$

## Part B Differentiate $p = \frac{1}{\sqrt{3a+8}}$

Find  $\frac{dp}{da}$  if  $p = \frac{1}{\sqrt{3a+8}}$ .

The following symbols may be useful:  $a$

Created for isaacphysics.org by Julia Riley

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.



# Chain Rule 2

A Level

Further A

P

P

P

P

P

P

**Part A**   Differentiate  $E = B \sin^2(\omega t)$ .

Find  $\frac{dE}{dt}$  if  $E = B \sin^2(\omega t)$ , where  $B$  and  $\omega$  are constants.

The following symbols may be useful: B, E, omega, t

**Part B**   Differentiate  $y = e^{-\frac{x^2}{2\sigma^2}}$

Find  $\frac{dy}{dx}$  if  $y = e^{-\frac{x^2}{2\sigma^2}}$ , where  $\sigma$  is a constant.

The following symbols may be useful: e, sigma, x

Created for isaacphysics.org by Julia Riley

Gameboard:

**STEM SMART Double Maths 11 - The Chain & Product Rules**

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Calculus](#) [Differentiation](#) [Differentiating Exponentials 3](#)

# Differentiating Exponentials 3

A Level Further A  
P P P P P P

## Part A Tangent to $y = e^{2x} - e^{-2x}$

Find the equation of the tangent to the curve  $y = e^{2x} - e^{-2x}$  at the point  $x = \frac{1}{2}$ .

The following symbols may be useful:  $e$ ,  $x$ ,  $y$

## Part B Stationary point of $u = 2e^{3v} - 3v$

Find the coordinates and nature of the stationary point of the function  $u = 2e^{3v} - 3v$ .

Find the  $v$  coordinate of the stationary point.

The following symbols may be useful:  $v$

Find the  $u$  coordinate of the stationary point.

The following symbols may be useful:  $u$

Determine the nature of the stationary point.

- ☐ Minimum
- ☐ Maximum

Gameboard:

**STEM SMART Double Maths 11 - The Chain & Product Rules**

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Calculus](#) [Differentiation](#) [Differentiating Natural Logs](#)

# Differentiating Natural Logs

A Level Further A  
P P P P P P

## Part A Differentiate $u = \ln(2v + 3)$

Find  $\frac{du}{dv}$  if  $u = \ln(2v + 3)$ .

The following symbols may be useful:  $v$

## Part B Stationary point of $p = 2 \ln(2q) - 3q$

Find the coordinates and nature of the stationary point of the function  $p = 2 \ln(2q) - 3q$ .

Find the  $q$  coordinate of the stationary point.

The following symbols may be useful:  $q$

Find the  $p$  coordinate of the stationary point.

The following symbols may be useful:  $p$

Determine the nature of the stationary point.

☐ Maximum

☐ Minimum

Gameboard:

**STEM SMART Double Maths 11 - The Chain & Product Rules**

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Calculus](#) [Differentiation](#) [Differentiating Trig Functions 3](#)

# Differentiating Trig Functions 3

A Level Further A  


## Part A Velocity and acceleration if $x = A \cos(\omega t + \phi)$

The displacement  $x$  of an oscillating particle at time  $t$  is given by  $x = A \cos(\omega t + \phi)$  where  $A$ ,  $\omega$  and  $\phi$  are constants; find expressions for the velocity (the rate of change of displacement) and acceleration (the rate of change of velocity) of the particle.

Find an expression for the velocity (the rate of change of displacement) of the particle.

The following symbols may be useful:  $A$ ,  $a$ ,  $\omega$ ,  $\phi$ ,  $t$ ,  $v$

Find an expression for the acceleration (the rate of change of velocity) of the particle.

The following symbols may be useful:  $A$ ,  $a$ ,  $\omega$ ,  $\phi$ ,  $t$ ,  $v$

## Part B Stationary points of the function $y = \cos(\omega t) + \sin(\omega t)$

Consider the function  $y = \cos(\omega t) + \sin(\omega t)$ , where  $\omega$  is a positive constant.

Find the stationary points of the function in the range  $0 < t < \frac{2\pi}{\omega}$ . How many are there? The stationary point with the lowest value of  $t$  is at  $(t_1, y_1)$  and the stationary point with the second lowest value of  $t$  is at  $(t_2, y_2)$ . Find the values of  $t$  and  $y$  at  $(t_1, y_1)$  and  $(t_2, y_2)$ .

How many stationary points are there?

- ☐ 3
- ☐ 4
- ☐ 1
- ☐ 2
- ☐ 0

Find  $t_1$ , the  $t$  coordinate of the stationary point with the lowest value of  $t$  in the range  $0 < t < \frac{2\pi}{\omega}$ .

The following symbols may be useful:  $\omega$ ,  $\pi$ ,  $t_1$ ,  $y_1$

Find  $t_2$ , the  $t$  coordinate of the stationary point with the second lowest value of  $t$  in the range  $0 < t < \frac{2\pi}{\omega}$ .

The following symbols may be useful:  $\omega$ ,  $\pi$ ,  $t_2$ ,  $y_2$

Created for isaacphysics.org by Julia Riley

Gameboard:

**STEM SMART Double Maths 11 - The Chain & Product Rules**

All materials on this site are licensed under the [Creative Commons license](https://creativecommons.org/licenses/by/4.0/), unless stated otherwise.





# Product Rule 2

A Level

Further A

P

P

P

P

P

P

**Part A**   Differentiate  $y = at^2 e^{\beta t}$

Find  $\frac{dy}{dt}$  where  $y = at^2 e^{\beta t}$ , given that  $a$  and  $\beta$  are constants.

The following symbols may be useful: a, beta, e, t

**Part B**   Differentiate  $\tan \theta$

Find the derivative w.r.t.  $\theta$  of  $\tan \theta$  by writing it as  $\frac{\sin \theta}{\cos \theta}$ .

The following symbols may be useful: cos(), cosec(), cot(), sec(), sin(), tan(), theta

Created for isaacphysics.org by Julia Riley

Gameboard:

**STEM SMART Double Maths 11 - The Chain & Product Rules**

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Differentiation: Products 4ii](#)

# Differentiation: Products 4ii



Differentiate with respect to  $x$ , simplifying your answers where possible.

## Part A $\sin x \tan x$

Differentiate  $\sin x \tan x$ .

The following symbols may be useful:  $x$

## Part B $x^2(x+1)^6$

Differentiate  $x^2(x+1)^6$ .

The following symbols may be useful:  $x$

Used with permission from UCLES A-level Maths papers, 2003-2017.

Gameboard:

**STEM SMART Double Maths 11 - The Chain & Product Rules**

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Differentiation: Quotients 2ii](#)

# Differentiation: Quotients 2ii

A Level



Differentiate with respect to  $x$ , simplifying your answers where possible.

**Part A**  $\frac{\ln x}{x}$

$$y = \frac{\ln x}{x}$$

The following symbols may be useful:  $\text{Derivative}(y, x)$ ,  $\ln()$ ,  $\log()$ ,  $x$ ,  $y$

**Part B**  $\frac{x^2}{\ln x}$

$$y = \frac{x^2}{\ln x}$$

The following symbols may be useful:  $\text{Derivative}(y, x)$ ,  $\ln()$ ,  $\log()$ ,  $x$ ,  $y$

**Part C** Stationary point of  $y = \frac{x^2}{\ln x}$

Determine the exact  $x$ -coordinate of the stationary point of the curve  $y = \frac{x^2}{\ln x}$ .

The following symbols may be useful:  $e$ ,  $\ln()$ ,  $x$

Used with permission from UCLES A-level Maths papers, 2003-2017.

Gameboard:

**STEM SMART Double Maths 11 - The Chain & Product Rules**

All materials on this site are licensed under the [\*\*Creative Commons license\*\*](#), unless stated otherwise.



Physics. *You work it out.*

[Home](#)   [Gameboard](#)   Maths   Differentiation: Chain Rule 5ii

# Differentiation: Chain Rule 5ii



Leaking oil is forming a circular patch on the surface of the sea. The area of the patch is increasing at a rate of 250 square metres per hour. Find the rate at which the radius of the patch is increasing at the instant the area of the patch is 1900 square metres. Give your answer in metres per hour correct to two significant figures.

Used with permission from UCLES A-level Maths papers, 2003-2017.

Gameboard:

**STEM SMART Double Maths 11 - The Chain & Product Rules**

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.




Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Differentiation: Chain Rule 5i](#)

# Differentiation: Chain Rule 5i

A Level



**Figure 1:** A diagram of a container in the form of a right circular cone.

**Figure 1** shows a container in the form of a right circular cone. The angle between the axis and the slant height is  $\alpha$ , where  $\alpha = \tan^{-1}\left(\frac{1}{2}\right)$ . Initially the container is empty, and then liquid is added at the rate of  $14 \text{ cm}^3 \text{ min}^{-1}$ . The depth of liquid in the container at time  $t$  minutes is  $x \text{ cm}$ .

## Part A Volume in Terms of Depth

Find an expression for the volume,  $V \text{ cm}^3$ , of liquid in the container when the depth is  $x \text{ cm}$  in terms of  $x$ .

[The volume of a cone is  $\frac{1}{3}\pi r^2 h$ .]

The following symbols may be useful:  $v$ ,  $\pi$ ,  $x$

## Part B Rate of Change

Find the rate at which the depth of the liquid is increasing at the instant when the depth is 8 cm. Give your answer in  $\text{cm min}^{-1}$  correct to two significant figures.

---

Used with permission from UCLES A-level Maths papers, 2003-2017.

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.