# Special Relativity:

DR LISA JARDINE-WRIGHT


Cavendish Laboratory, Cambridge

#### Copyright Cavendish Laboratory.

For student personal use only. Not for distribution or publication without written permission from the author.

#### Frames of Reference:

# Galilean Coordinate Transformations



- S' moves with velocity v along the x-direction in frame S.
- If the origins of the two frames coincide at  $t=t^{\prime}=0$ , the transformation of an event in S to S' is

$$x' = x - vt$$

$$y' = y$$

$$z' = z$$

$$t' = t$$

• Implicit here the assumption that there is a universal 'absolute' time, running at the same rate in all inertial frames.

### Inertial Frames of Reference

- A frame in which Newton's first law of motion is valid.
- What is Newton's first law?

A body remains in a state of rest or uniform motion unless acted upon by a resultant force.


 You are in a car and cannot see outside but you can see a parcel in the boot of the car that is stationary according to you.

What do you deduce about the motion of the car?

What do you deduce (and why) if the parcel accelerates to the back of the car?

A <u>non-inertial</u> frame is a frame which is accelerating.

### Frames of Reference



Platform view

### Frames of Reference




Platform view

# **Enter Einstein**

Galileo's relativity is not the whole story!

it is actually an approximation to the real truth that works for us in everyday life.

Proposed that there is a single relativity principle for both dynamics and EM but it is NOT Galilean.



# Special Theory of Relativity: Historical Context

#### The "Aether"

 Mid to late 19<sup>th</sup> century the wave theory of light was generally accepted and naturally explained by


Reflection, refraction, interference, diffraction and polarisation.


- It was thought that, like all other known waves, light must travel in a medium – called the luminiferous aether.
- Maxwell's equations predicted electromagnetic waves light travelling with a speed  $c = \sqrt{\frac{1}{\epsilon_0 \mu_0}} \approx 3 \times 10^8 \text{ m s}^{-1}$

# **Aether Experiments**

- Two experiments to deduce the properties of the aether were in direct contradiction to each other
  - 1. "stellar aberration" → evidence for a stationary aether absolute frame.
  - 2. "Michelson-Morley experiment"

# Michaelson-Morley





## **Aether Experiments**

- Two experiments to deduce the properties of the aether were in direct contradiction to each other
  - 1. "stellar aberration"  $\rightarrow$  evidence for a stationary aether absolute frame.
  - 2. "Michelson-Morley experiment" → the aether was at rest w.r.t. the Earth i.e. dragged along by the Earth no absolute frame, no aether?

#### • Contradictions:


- a) Newton's Laws do not need absolute space they apply in all inertial frames.
- b) Maxwell's equations are not Galilean invariant because the speed of light is the same in all frames.

# Einstein's Solution:

There is a single relativity principle for both dynamics and EM but it is NOT Galilean.

#### The Postulates:

- 1. The laws of Physics are the same in all inertial frames;
- 2. The speed of light in a vacuum is the same in all inertial frames.




Train frame, S' with the light clock

Time for A to B to C 
$$\Delta \mathsf{t}_{AC}' = \frac{2h}{c}$$

#### **Physics**

$$Speed = \frac{distance}{time}$$




Train frame, S' with the light clock

Time for A to B to C 
$$\Delta t_{AC}' = \frac{2h}{c}$$








Platform frame, S Standing watching the light clock pass by on the train


#### **Physics**


 $Speed = \frac{distance}{time}$ 




Train frame, S' with the light clock

Time for A to B to C
$$\Delta t'_{AC} = \frac{2h}{C}$$







#### Platform frame, S

$$(c^2 - v^2)\Delta t_{AC}^2 = 4h^2$$

#### **Physics**

$$Speed = \frac{distance}{time}$$

Time for A to B to C
$$\Delta t'_{AC} = \frac{2h}{c}$$

### So what?

What if 
$$v = \frac{4}{5}c$$
?

Distance pulse travels  $\,c\Delta t_{AC}\,$ 

$$(c\Delta t_{AC})^2 = (v\Delta t_{AC})^2 + 4h^2$$

$$(c^2 - v^2)\Delta t_{AC}^2 = 4h^2$$

$$(c^2 - v^2)\Delta t_{AC}^2 = c^2 \Delta t_{AC}'^2$$

$$\left(1 - \frac{v^2}{c^2}\right) \Delta t_{AC}^2 = \Delta t_{AC}^{\prime 2}$$

$$\sqrt{\left(1 - \frac{v^2}{c^2}\right)} \Delta t_{AC} = \Delta t'_{AC}$$

# 30 000 m Secondary cosmic rays $\pi$ + 20 000 m μ- $\pi^-$ Concorde 10000 m **Everest** Image courtesy of CERN

# Does this physics matter?

- Measuring muons on Earth presented a dilemma.
- Their half-life is just 2.2  $\mu$ s.
- How far can they travel in that time?
- The speed of the muon means that we need to take SR into account.

$$\sqrt{1 - \frac{v^2}{c^2}} \Delta t = \Delta t'$$
Earth Muon

# **Staying Younger for Longer?**




$$\sqrt{\left(1 - \frac{v^2}{c^2}\right)\Delta t} = \Delta t'$$

- One twin, Bob, stays on Earth.
- Alice, takes her rocket into space travelling at 4/5 of the speed of light.
- After 5 years have passed for Bob, how many have passed for Alice?


Questions 23 & 24.

Not to scale!

# **Length Contraction**



### **Lorentz Transformations**



$$r = x - vt$$

• Ruler of length  $r_0$  in frame S' has length r in S such that

$$r_0 = \gamma r$$

$$x' = \gamma r$$

$$x' = \gamma (x - \nu t)$$

And

$$x = \gamma(x' + \nu t')$$

• To find times:

$$x = \gamma [\gamma(x - vt) + vt']$$
$$t' = \gamma(t - vx/c^2)$$

### **Lorentz Transformations**

$$\Delta t' = \gamma \left( \Delta t - \frac{v \Delta x}{c^2} \right)$$

$$\Delta x' = \gamma (\Delta x - v \Delta t)$$

$$\Delta y' = \Delta y$$

$$\Delta z' = \Delta z$$

$$\Delta t = \gamma \left( \Delta t' + \frac{v \Delta x'}{c^2} \right)$$

$$\Delta x = \gamma (\Delta x' + \nu \Delta t')$$

$$\Delta y = \Delta y'$$

$$\Delta z = \Delta z'$$

With

$$\gamma = \frac{1}{\sqrt{1 - v^2/c^2}}$$