

Physics. *You work it out.*[Home](#) [Gameboard](#) [Maths](#) [2D Vectors & NII 1ii](#)

2D Vectors & NII 1ii

A Level




Figure 1: Two horizontal forces P and Q acting at the origin of a point O as viewed from above.

Two horizontal forces P and Q act at the origin O of rectangular coordinates Oxy . The components of P in the x - and y -directions are 14 N and 5 N respectively. The components of Q in the x and y directions are -9 N and 7 N respectively as shown in **Figure 1**

Part A Resultant force 1

Write down the x -component of the resultant of P and Q .

Write down the y -component of the resultant of P and Q .

Part B Resultant force 2

Find the magnitude of this resultant of resultant force.

Find the angle the resultant makes with the positive x -axis to 3 significant figures.

Part C Acceleration

The two forces P and Q act on a particles of mass 0.2 kg . Express the acceleration of the particle using ijk notation..

The following symbols may be useful: a , i , j , k

Used with permission from UCLES, A Level, June 2007, OCR M1, Question 1

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [2D Vectors & NII 2ii](#)

2D Vectors & NII 2ii

A Level





Figure 1: Diagram showing three forces acting on a particle.

A particle rests on a smooth horizontal surface. Three horizontal forces of magnitudes 2.5 N, F N and 2.4 N act on the particle on bearings θ° , 180° and 270° respectively. The particle is in equilibrium.

Part A Vector notation

The 2.5 N force may be written in the form $(p\mathbf{i} + q\mathbf{j})\text{N}$. Write down the value of p .

Part B Finding F

Hence, find F .

Part C Finding θ

Find θ to 3 significant figures.

Part D Acceleration

The 2.4 N force suddenly ceases to act on the particle, which has mass 0.2 kg .

Find the magnitude of the acceleration of the particle.

Find the direction of the acceleration of the particle in the form of a bearing.

Used with permission from UCLES, A Level, June 2014, OCR M1, Question 2

Gameboard:

STEM SMART Single Maths 21 - Forces and Motion

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) Maths Friction 2i

Friction 2i

A Level



A particle of mass 2 kg is propelled in a straight line across a rough surface with an initial velocity $\underline{u} = 12\underline{i} + 9\underline{j} \text{ m s}^{-1}$. It comes to rest in 3 seconds.

Assuming that the frictional force is constant throughout the motion and no other forces are acting, what is the magnitude of the frictional force?

Created for isaacphysics.org by Sally Waugh

Gameboard:

STEM SMART Single Maths 21 - Forces and Motion

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Friction 1ii](#)

Friction 1ii

A Level



A block of mass 4 kg is in equilibrium on a rough surface under the influence of a horizontal force, X , of 11.8 N .

Part A Diagram

Draw a diagram to show all the forces on the block.

More practice questions?

Part B Ratio of forces

Find the ratio of the frictional force, F , to the normal reaction, R . Give your answer to one decimal place.

Part C Finding M

A different block, of mass M , is placed on the same surface. A force of 16.8 N is applied in place of force X . Again the system is in equilibrium.

Assuming the same ratio applies to the frictional and normal forces, find the value of M to 3 significant figures.

Gameboard:

STEM SMART Single Maths 21 - Forces and Motion

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.




Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Newton's Laws 2i](#)

Newton's Laws 2i

A Level



A trailer of mass 500 kg is attached to a car of mass 1250 kg by a light rigid horizontal tow-bar. The car and trailer are travelling along a horizontal straight road. The resistance to motion of the trailer is 400 N and the resistance to motion of the car is 900 N. Find both the tension in the tow-bar and the driving force of the car in each of the following cases.

Part A Driving force 1

The car and trailer are travelling at constant speed.

What is the driving force?

Part B Tension 1

What is the tension in the tow-bar?

Part C Driving force 2

The car and trailer have acceleration 0.6 m s^{-2} .

What is the driving force of the car? Give your answer to 4 significant figures.

Part D Tension 2

What is the tension in the tow-bar?

Used with permission from UCLES, A Level Maths, January 2009, OCR M1, Question 2

Gameboard:

STEM SMART Single Maths 21 - Forces and Motion

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Normal Reaction 2i](#)

Normal Reaction 2i

A Level
P P P




Figure 1: Diagram showing a block of mass M kg on a platform which is raised by a hoist.

A box of mass M kg is loaded onto a platform which can be raised by a hoist. The platform, which consists of 2 vertical supports and a horizontal plate, has a total mass of 30 kg. Before the hoist is attached, the box and platform are on horizontal ground and the thrust in each support is 350 N.

Part A Mass of box

Find the mass of the box to 3 significant figures.

Part B Normal reaction 1

The platform is now connected to the hoist cable, raised slowly by a short distance so that it leaves the ground, and brought to a halt.

Find the normal reaction on the box in this stationary position to 3 significant figures.

Part C Tension

Find the tension in the hoist cable.

Part D Normal reaction 2

The hoist suddenly jerks the platform upwards, with an initial acceleration of 1.5 m s^{-2} .

Find the normal reaction on the box to 3 significant figures.

Created for isaacphysics.org by Sally Waugh

Gameboard:

STEM SMART Single Maths 21 - Forces and Motion

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Analysing Systems and Forces 2i](#)

Analysing Systems and Forces 2i

A Level




Figure 1: Man preparing to lower a box down a cliff.

A man is preparing to lower a box down a cliff. He sets up a fixed pulley with a rope as shown in **Figure 1**. He pulls on the rope with a force $-(100\hat{i} + 75\hat{j})\text{ N}$ and the box settles into a stationary position with the rope between the box and pulley vertical.

Part A Force in rope

The force **on the man's hand from the rope** can be written as

$$\underline{\boldsymbol{F}} = \begin{pmatrix} F_1 \\ F_2 \end{pmatrix}$$

Find F_1 .

Find F_2 .

Part B Magnitude of tension

What is the magnitude of the tension in the rope?

Part C Force diagrams

Draw a labelled diagram showing the forces acting in the rope at the pulley.

Easier question?

Draw a labelled diagram showing the forces on the box.

Easier question?

Part D Assumptions

In order to model the system mathematically, it is necessary to make assumptions. Give one assumption you need to make about the pulley and two assumptions about the rope.

Easier question?

Part E Mass of box

Find the mass of the box to 3 significant figures.

Gameboard:

STEM SMART Single Maths 21 - Forces and Motion

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Analysing Systems and Forces 1i](#)

Analysing Systems and Forces 1i

A Level



A small, smooth pulley is suspended from a fixed point by a light chain. A light inextensible string passes over the pulley. Particles P and Q , of masses 0.30 kg and m respectively, are attached to the opposite ends of the string. The particles are released from rest at a height of 0.20 m above horizontal ground with the string taut; the portions of the string not in contact with the pulley are vertical. P strikes the ground with speed 1.4 m s^{-1} . Subsequently P remains on the ground and Q does not reach the pulley.




Figure 1: Diagram showing string with P and Q on either end passing over a pulley.

Part A Acceleration of P

Calculate the acceleration of P while it is in motion to 2 significant figures.

Part B Tension in the string

Calculate the tension in the string to 2 significant figures.

Part C Mass of Q

Find the mass of Q .

Part D Height of Q

Calculate the greatest height of Q above the ground.

Part E Tension in the chain

It is given that the mass of the pulley is 0.50 kg.

State the magnitude of the tension in the chain which supports the pulley when P **is in motion**. Give your answer to 2 significant figures.

Part F Tension when Q moving upwards

State the magnitude of the tension in the chain which supports the pulley when P is at rest on the ground and Q is moving upwards. Give your answer to 2 significant figures.

Used with permission from UCLES, A Level Maths, June 2010, OCR M1, Question 7

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.