



Transcription Overview

Part A Transcription & Translation

A is a region of DNA that "codes for" a protein. The production of a protein happens in two main steps: transcription and translation.

Transcription is the process by which is produced from . In eukaryotic cells, this process happens inside .

Translation is the process by which is produced from . This process happens inside .

Items:

DNA

gene

the nucleus

a polypeptide chain

messenger RNA (mRNA)

a ribosome

Part B The process of transcription

Transcription begins when transcription factors and the enzyme bind to the promoter region of a gene. The enzyme then moves along the DNA, unwinding and unzipping it as it does so. The enzyme facilitates the binding of complementary RNA nucleotides to the strand (in the 5' to 3' direction), forming an mRNA strand that has the same base sequence as the strand of DNA (except that the DNA bases are replaced by RNA bases).

Items:

RNA polymerase

adenine

antisense/template

thymine

uracil

lagging

sense/coding

DNA polymerase

leading

guanine

cytosine

Part C Pre-mRNA to mRNA

The RNA strand produced by transcription is often called "pre-mRNA", and must undergo certain modifications in order to become mature mRNA.

A 5' cap is added to the end of the strand. This consists of a modified nucleotide. A poly(A) tail is added to the end of the strand. This consists of 100 to 250 nucleotides. These features allow the cell to identify the RNA as mRNA, which ensures that it will be exported from the nucleus (in eukaryotic cells) and bind to a ribosome.

Pre-mRNA also undergoes a process called splicing, during which the (non-coding regions) are removed so that only the (coding regions) remain.

Once the pre-mRNA has been capped, polyadenylated, and spliced, it is now a mature mRNA that can be translated.

Items:

uracil

5'

introns

guanine

3'

adenine

exons

cytosine




Transcription Diagrams


A Level




The images below represent DNA transcription. The labels (5' and 3') refer to the DNA strands (black). The DNA molecule is unzipping from bottom to top. The green arrows represent possible directions of nucleotide addition by RNA polymerase.




A




B




C



D



E



F

Which image above correctly illustrates the process of DNA transcription?

- ☐ A
- ☐ B
- ☐ C
- ☐ D
- ☐ E
- ☐ F




Physics. *You work it out.*

[Home](#) [Gameboard](#) [Biology](#) [Genetics](#) [Transcription](#) [DNA vs RNA](#)

DNA vs RNA

A Level



Part A DNA

Which of the following are true of DNA?

- ☐ is single-stranded in eukaryotic cells
- ☐ is double-stranded in eukaryotic cells
- ☐ contains thymine
- ☐ contains uracil
- ☐ contains ribose
- ☐ contains deoxyribose

Part B RNA

Which of the following are true of RNA?

- ☐ is single-stranded in eukaryotic cells
- ☐ is double-stranded in eukaryotic cells
- ☐ contains thymine
- ☐ contains uracil
- ☐ contains ribose
- ☐ contains deoxyribose

Transcription vs Replication

In the table below, identify which statements about transcription and DNA replication are correct, and which are incorrect. Fill in every box with either a tick (correct) or a cross (incorrect).

	Transcription	Replication
free nucleotides bind to both DNA strands	<input type="text"/>	<input type="text"/>
two new DNA molecules are produced	<input type="text"/>	<input type="text"/>
hydrogen bonds are broken between the two DNA strands	<input type="text"/>	<input type="text"/>
uracil nucleotides bind to adenine nucleotides	<input type="text"/>	<input type="text"/>
the process occurs along the entire length of the chromosome	<input type="text"/>	<input type="text"/>
the process only occurs at specific regions of the chromosome(s)	<input type="text"/>	<input type="text"/>

Items:

☐

☐

Adapted with permission from OCR A Level January 2003, Biology Foundation, Question 4

Gameboard:

STEM SMART Biology Week 7



Transcribe the Sequences

A Level



Part A Sense to RNA

A region of the sense/coding DNA strand contains the following base sequence (from 5' to 3'):

ATGCCGCAGTTC

Enter the sequence of the mRNA that would be transcribed from this gene region (from 5' to 3').

Enter your answer in all caps and without spaces.

Part B Antisense to RNA

A region of the antisense/template DNA strand contains the following base sequence (from 3' to 5'):

TACAGTCAGTCA

Enter the sequence of the mRNA that would be transcribed from this gene region (from 5' to 3').

Enter your answer in all caps and without spaces.

Part C RNA to sense

A region of mRNA contains the following base sequence (from 5' to 3'):

CAUGUCAAUGG

Enter the sense/coding strand sequence of DNA (from 5' to 3') that produced this mRNA sequence.

Enter your answer in all caps and without spaces.



Post-transcriptional Modifications




Figure 1: Post-transcriptional modification. A pre-mRNA transcript is modified to become a mature RNA through splicing, capping, and polyadenylation. UTR = untranslated region.

Match the labels from Figure 1 to the mRNA regions in the table below.

Label	Region
A	<div></div>
B	<div></div>
C	<div></div>
D	<div></div>

Items:

- 5' cap
- intron
- 3' cap
- C-terminus
- N-terminus
- exon
- poly(A) tail