



Exponential Equation 1

A Level Further A

P

P

P

P

P

P

Solve the following for y : $4^y = 8^{y+1}$

Created for isaacphysics.org by Julia Riley.

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.



Exponential Equation 2

A Level Further A

P

P

P

P

P

P

Solve the following for x : $3^x = \frac{1}{\left(9^{x-\frac{9}{4}}\right)}.$

Created for isaacphysics.org by David Lawrence

Gameboard:

STEM SMART Single Maths 8 - Logarithms

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.



Exponential Equation 3

A Level Further A

P

P

P

P

P

P

Solve the following for m : $\frac{1}{9^m} = 27^{1-m}$.

Created for isaacphysics.org by David Lawrence

Gameboard:
STEM SMART Single Maths 8 - Logarithms

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Log Laws 2ii](#)

Log Laws 2ii



Part A Express as a single log (i)

Express $\log_a 2 + \log_a 3$ as a single logarithm.

The following symbols may be useful: a , $\log()$

Part B Express as a single log (ii)

Express $2 \log_{10} x - 3 \log_{10} y$ as a single logarithm.

The following symbols may be useful: $\log()$, x , y

Used with permission from UCLES, A Level Maths, January 2008, OCR C2, Question 3

Gameboard:

STEM SMART Single Maths 8 - Logarithms

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Log Laws 1ii](#)

Log Laws 1ii



Part A Express in terms of $\log_2(x)$ (i)

Express $\log_2(x^2)$ in terms of $\log_2(x)$.

When you are entering your answer, note that $\log_a b$ can be written using $\log(b, a)$.

The following symbols may be useful: $\ln()$, $\log()$, \times

Part B Express in terms of $\log_2(x)$ (ii)

Express $\log_2(8x^2)$ in terms of $\log_2(x)$.

When you are entering your answer, note that $\log_a b$ can be written using $\log(b, a)$.

The following symbols may be useful: $\ln()$, $\log()$, \times

Part C Find $\log_3 y$

Given that $y^2 = 27$, find the value of $\log_3 y$.

The following symbols may be useful: $\ln()$, $\log()$, y

Used with permission from UCLES, A Level Maths, Specimen paper, OCR C2, Question 3

Gameboard:

STEM SMART Single Maths 8 - Logarithms

All materials on this site are licensed under the [**Creative Commons license**](#), unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Solving Equations & Logs 1ii](#)

Solving Equations & Logs 1ii



Part A Solve equation (i)

Use logarithms to solve the equation $5^{x-1} = 120$ for x , giving your answer correct to 3 significant figures.

Part B Solve equation (ii)

Solve the equation $\log_2 x + 2\log_2 3 = \log_2(x + 5)$. Give the value of x correct to 3 significant figures.

Used with permission from UCLES, A Level Maths, January 2011, OCR C2, Question 4

Gameboard:

STEM SMART Single Maths 8 - Logarithms

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Solving Equations & Logs 3ii](#)

Solving Equations & Logs 3ii

A Level



Part A Express log

Express $\log_3(4x + 7) - \log_3 x$ as a single logarithm.

When you are entering your answer, note that $\log_a b$ can be written using $\log(b, a)$.

The following symbols may be useful: $\ln()$, $\log()$, \times

Part B Solve equation

Hence solve the equation $\log_3(4x + 7) - \log_3 x = 2$. Give your answer in decimal form.

Part C Use logs

Use logarithms to solve the equation $7^x = 2^{x+1}$, giving the value of x correct to 3 significant figures.

Used with permission from UCLES, A Level Maths, January 2007, OCR C2, Question 5 and June 2009, OCR C2, Question 3

Gameboard:

STEM SMART Single Maths 8 - Logarithms

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Log Laws 1i](#)

Log Laws 1i

A Level



Part A Log laws

Given that $\log_a(x) = p$ and $\log_a(y) = q$, express $\log_a(xy)$ in terms of p and q .

The following symbols may be useful: p , q

Part B Simplify 1

Given that $\log_a(x) = p$ and $\log_a(y) = q$, express $\log_a\left(\frac{a^2x^3}{y}\right)$ in terms of p and q .

The following symbols may be useful: p , q

Part C Simplify 2

Express $\log_{10}(x^2 - 10) - \log_{10}x$ as a single logarithm.

When you are entering your answer, note that $\log_a b$ can be written using $\log(b, a)$.

The following symbols may be useful: $\log()$, x

Part D **Solve equation**

Solve the equation $\log_{10}(x^2 - 10) - \log_{10} x = 2 \log_{10} 3$. Write down the value of x .

Used with permission from UCLES, A Level Maths, January 2009, OCR C2, Question 8

Gameboard:

STEM SMART Single Maths 8 - Logarithms

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Log Laws 1i](#)

Log Laws 1i

A Level



It is given that $\ln x = p + 2$ and $\ln y = 3p$

Part A Log laws

Express $\ln(xy)$ in terms of p .

The following symbols may be useful: p

Part B Simplify 1

Express $\ln(x^3)$ in terms of p .

The following symbols may be useful: p

Part C Simplify 2

Express $\ln\left(\frac{y}{e}\right)$ in terms of p .

The following symbols may be useful: p

Part D Solve equation

Express y in terms of x and e , simplifying your answer.

The following symbols may be useful: e , x , y

Used with permission from UCLES, A Level Maths, January 2009, OCR C2, Question 8

Gameboard:

STEM SMART Single Maths 8 - Logarithms

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Solving Equations & Logs 3i](#)

Solving Equations & Logs 3i

A Level



Part A Solve equation

Solve the equation $2^{4x-1} = 3^{5-2x}$, giving your answer in the form $x = \frac{\log_{10} a}{\log_{10} b}$.

When you are entering your answer, note that $\log_a b$ can be written using $\log(b, a)$.

The following symbols may be useful: $\log()$, \times

Part B Find integer

Find the smallest integer n which satisfies the inequality $7^{2n} > e^{600}$.

Used with permission from UCLES, A Level Maths, June 2014, OCR C2, Question 5

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.