

<u>Home</u>

<u>Gameboard</u>

Maths

Friction 2i

Friction 2i






Figure 1: A particle P resting on a horizontal plane and attached to two light strings.

A particle P of weight $30\,\mathrm{N}$ rests on a horizontal plane. P is attached to two light strings making angles of $30\,^\circ$ and $50\,^\circ$ with the upward vertical, as shown in the **Figure 1**. The tension in each string is $15\,\mathrm{N}$, and the particle is in limiting equilibrium.

Part A Frictional force

I	Find the	e magnitude	e ot the	trictional	torce on I	P correct	to 3	significar	nt fiaures.
٠									


Does the frictional force on P point to the **left** or to the **right**?.

Part B Coefficient of friction

Find the coefficient of friction between P and the plane correct to 3 significant figures.

Used with permission from UCLES, A Level, June 2008, OCR M1, Question 4

All materials on this site are licensed under the ${\color{red} \underline{\textbf{Creative Commons license}}}$, unless stated otherwise.



Home Gameboard Maths Geometry Vectors Vectors in 3D 2

Vectors in 3D 2



The position vectors of three points A, B and C relative to the origin O are given by the vectors:

$$ec{OA} = egin{pmatrix} 4 \ 3 \ 3 \end{pmatrix} \qquad ec{OB} = egin{pmatrix} 8 \ 11 \ -1 \end{pmatrix} \qquad ec{OC} = egin{pmatrix} 8 \ 9 \ 5 \end{pmatrix}$$

Part A Midpoint of AB

Find the coordinates of point D, the midpoint of AB.

Give your answer in the form $a \underline{\pmb{i}} + b \underline{\pmb{j}} + c \underline{\pmb{k}}$.

Part B Find a unit vector

Find an exact expression for a unit vector with the same direction as \vec{AB} .

Give your answer in the form $a\underline{\pmb{i}} + b\underline{\pmb{j}} + c\underline{\pmb{k}}$, where $\underline{\pmb{i}}$, $\underline{\pmb{j}}$ and $\underline{\pmb{k}}$ are unit vectors in the x, y and z directions.

The following symbols may be useful: i, j, k

Part C The length of \vec{DC}

Find an exact expression for the length of \vec{DC} .

Part D Angle \hat{ACB}

Consider the lengths AB and DC. What can you conclude about the size of angle \hat{ACB} ?


- $\hat{ACB} < 45^{\circ}$
- $\hat{ACB} = 45^{\circ}$
- $\hat{ACB} > 45^{\circ}$
- $\hat{ACB} < 90^{\circ}$
- $\hat{ACB} = 90^{\circ}$
- $A\hat{C}B > 90^{\circ}$

Created for isaacphysics.org by Jonathan Waugh

Gameboard:

STEM SMART Single Maths 48 - Mechanics & Statistics

Revision



<u>Home</u> <u>Gameboard</u> Maths Statistics Probability Probabilities: Households

Probabilities: Households



The people living in 3 households are classified as children (C), parents (P) or grandparents (G). The numbers living in each house are shown in the table below.

House number 1	House number 2	House number 3
4C,1P,2G	2C,2P,3G	1C, 1G

Part A Scenario 1 - a grandparent

All the people in all 3 houses meet for a party. One person at the party is chosen at random. Calculate the probability of choosing a grandparent.

Part B Scenario 2 - a grandparent

A house is chosen at random. Then a person in that house is chosen at random. Using a tree diagram, or otherwise, calculate the probability that the person chosen is a grandparent.

Part C Scenario 2 - a parent


Given that the person chosen by the method in <u>Part B</u> is a grandparent, calculate the probability that there is also a parent living in the house.

Used with permission from UCLES, A level, June 2003, CIE, Paper 6 Probability & Statistics 1 (S1), Question 6

Gameboard:

STEM SMART Single Maths 48 - Mechanics & Statistics

Revision



<u>Home</u>

<u>Gameboard</u>

Maths

Projectiles: General 2i

Projectiles: General 2i



A child is trying to throw a small stone to hit a target painted on a vertical wall. The child and the wall are on horizontal ground. The child is standing a horizontal distance of $8\,\mathrm{m}$ from the base of the wall. The child throws the stone from a height of $1\,\mathrm{m}$ with speed $12\,\mathrm{m}\,\mathrm{s}^{-1}$ at an angle of $20\,^\circ$ above the horizontal.

Part A Finding the angle

Find the direction of motion of the stone when it hits the wall. Give your answer as an angle below the horizontal to 3 significant figures.

The child now throws the stone with a speed of $V\,\mathrm{m\,s^{-1}}$ from the same initial position and still at an angle of $20\,^\circ$ above the horizontal. This time the stone hits the target which is $2.5\,\mathrm{m}$ above the ground.


Find *V* to 3 significant figures.

Used with permission from UCLES, A Level, June 2014, OCR M2, Question 8

Gameboard:

STEM SMART Single Maths 48 - Mechanics & Statistics

Revision



<u>Home</u> <u>Gameboard</u> Physics Mechanics Statics Forces on a Supported Beam

Forces on a Supported Beam



A uniform beam AB of mass $15.0\,\mathrm{kg}$ and length $4.00\,\mathrm{m}$ is freely hinged to a vertical wall at A. The beam is held in equilibrium in a horizontal position by a light rod PQ of length $1.50\,\mathrm{m}$. P is fixed to the wall vertically below A and PQ makes an angle of 30.0° with the vertical. The force F exerted on the beam at Q by the rod is in the direction PQ. This force is shown in **Figure 1**.




Figure 1: The hinged beam AB supported by the light rod PQ.

Part A The force on the beam at Q

Find the magnitude of F, the force the rod exerts on the beam at Q.

Part B The force on the beam at A

Find the magnitude of the force exerted on the beam at A by the pivot.

Part C The angle of the force at ${\cal A}$


At A the force exerted on the beam is to the left. Find the angle the force makes to the horizontal. Use a positive sign if the force is directed above the horizontal and a negative sign if the force is directed below the horizontal.

Adapted from UCLES, A Level, June 2012, OCR M2, Question 3

Gameboard:

STEM SMART Single Maths 48 - Mechanics & Statistics

Revision



Home (

<u>Gameboard</u>

Maths Statistics

Probability

Normal Distribution 1

Normal Distribution 1



The random variable Y is normally distributed with mean μ and variance σ^2 , where μ and σ are integers. It is found that P(Y>150.0)=0.0228 and P(Y>143.0)=0.9332. Find the values of μ and σ .

Part A Value of μ

Enter the value of μ .

Part B Value of σ


Enter the value of σ .

Adapted with permission from UCLES, A level, Jun 2015, OCR, S2, Question 1

Gameboard:

STEM SMART Single Maths 48 - Mechanics & Statistics

Revision



<u>Home</u>

<u>Gameboard</u>

Maths

Statistics Hypothesis Tests Hypothesis Testing: Normal Distribution 3

Hypothesis Testing: Normal Distribution 3



Average Speed Cameras (ASC) are a way of encouraging motorists to reduce their speed. Modern systems use automated numberplate recognition to monitor how long it takes a car to travel between two points, from which the average speed of the car can be worked out.

A particular stretch of road is known to be an accident blackspot, and an ASC system is introduced with the aim of improving safety. Before the cameras are installed, the speed of the cars is monitored over a long period of time and it is found to obey a normal distribution with a mean of $75.0\,\mathrm{mph}$ and a standard deviation of $3.00\,\mathrm{mph}$. A few weeks after the cameras are installed the speed of the cars is monitored again. A sample of 400 cars is taken, and their speed is found to be normally distributed with a mean of $72.0 \,\mathrm{mph}$.

Null and alternative hypotheses Part A

A hypothesis test is carried out to determine whether the cameras have been effective in reducing driver
mean speed. Let μ represent the population mean speed. Drag and drop items into the boxes below to state
the null and alternative hypotheses for the test.

 H_0 : μ

 H_1 : μ

Items:

72.0

75.0

Part B Probability distributions

Drag and drop answers into the boxes below to describe the probability distributions involved in the hypothesis test. You may assume that the standard deviation of the speeds of the cars after the cameras are installed is the same as before the cameras were installed, $3.00\mathrm{mph}$.					
Let X be the speed of a car. If the null hypothesis is true, the speed of cars in the population is given by a normal distribution with a mean of $\begin{tabular}{ c c c c c c c c c c c c c c c c c c c$					
$X \sim \mathrm{N}($					
As X is normally distributed, the mean values of samples of X with size n will also be normally distributed. Let \overline{X} be the mean speed of a sample of 400 cars. Then the distribution of \overline{X} is					
$\overline{X} \sim \mathrm{N}(\bigcirc), \bigcirc)$					
Items:					
$egin{array}{cccccccccccccccccccccccccccccccccccc$					
Part C Carrying out the test					
Fill in the blanks to complete the description of the hypothesis test.					
Drag and drop items into the boxes below to carry out the hypothesis test at the 1% significance level.					
The hypothesis test is a $\overline{\hspace{1cm}}$ -tail test. At the 1% significance level, the critical region is \overline{X}					
The mean speed of the sample of 400 cars was found to be $72.0\mathrm{mph}$. This value is in the					
Therefore, the null hypothesis. There significant evidence that the mean speed of drivers is lower after the cameras were installed.					
Items:					
one two three $<$ \neq $=$ $>$ 74.65 74.95 75.05 75.35 acceptance region critical region reject do not reject is is not					

Created for isaacphysics.org by Jonathan Waugh

All materials on this site are licensed under the ${\color{red} \underline{\textbf{Creative Commons license}}}$, unless stated otherwise.