



# Binomial: All Rational n 2ii

A Level

P

P

P

Part A

Expansion

Expand  $(1 + 4x)^{\frac{1}{2}}$  in ascending powers of  $x$ , up to and including the term in  $x^2$ , simplifying the coefficients.

The following symbols may be useful:  $x$

Part B

Values of  $x$

State the set of values of  $x$  for which the expansion is valid. You may use the symbols  $<$   $>$   $=$   $x$  and  $|x|$  which can be entered as `abs(x)`.

The following symbols may be useful:  $<$ ,  $>$ , `abs()`,  $x$

Part C    Value of  $k$

In the expansion of

$$(1 + kx)(1 + 4x)^{\frac{1}{2}},$$

the coefficient of  $x$  is 7. Find the value of the constant  $k$  and hence the coefficient of  $x^2$ .

State the value of  $k$ .

The following symbols may be useful:  $k$


---

State the coefficient of  $x^2$ .

---

Used with permission from UCLES A-level Maths papers, 2003-2017.

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



# Binomial: All Rational n 4ii

A Level

P

P

P

Part A

Expansion

Expand  $(2 + x)^{-2}$  in ascending powers of  $x$  up to and including the term in  $x^3$ .

The following symbols may be useful:  $x$

Part B

Set of Values

State the set of values of  $x$  for which the expression is valid. You may use the symbols  $<$   $>$   $=$   $x$  and  $|x|$  which can be entered as `abs(x)`.

The following symbols may be useful:  $<$ ,  $>$ , `abs()`,  $x$

Part C

Coefficient

Hence find the coefficient of  $x^3$  in the expansion of  $\frac{1+x^2}{(2+x)^2}$ .


The following symbols may be useful:  $x$

Used with permission from UCLES A-level Maths papers, 2003-2017.

Gameboard:

**STEM SMART Single Maths 28 - Binomial Theorem - Rational n**

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



# Binomial: All Rational n 1i

A Level

P

P

P

Part A

Partial Fractions

Given that  $\frac{3x+4}{(1+x)(2+x)^2} \equiv \frac{A}{1+x} + \frac{B}{2+x} + \frac{C}{(2+x)^2}$ , find  $A$ ,  $B$ , and  $C$ .

Find  $A$ .

The following symbols may be useful: A

Find  $B$ .

The following symbols may be useful: B

Find  $C$ .

The following symbols may be useful: c

Part B

Expand

Hence or otherwise expand  $\frac{3x+4}{(1+x)(2+x)^2}$  in ascending powers of  $x$ , up to and including the term in  $x^2$ .

The following symbols may be useful: x

Part C Values of  $x$

State the set of values of  $x$  for which the expansion in the above part is valid.

Construct your answer from the items below.

Items:

$x$

$<$

$\leq$

$>$

$\geq$

$< x <$

$\leq x \leq$

$> x \text{ or } x >$

$\geq x \text{ or } x \geq$

$-4$

$-2$

$-\frac{3}{2}$

$-1$

$-\frac{1}{2}$

$0$

$\frac{1}{2}$

$1$

$\frac{3}{2}$

$2$


$4$

Used with permission from UCLES A-level Maths papers, 2003-2017.

Gameboard:

STEM SMART Single Maths 28 - Binomial Theorem - Rational n

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



# Binomial: All Rational n 3i

A Level

P

P

P

Find the coefficient of  $x^2$  in the expansion in ascending powers of  $x$  of

$$\sqrt{\frac{1+ax}{4-x}}$$


giving your answer in terms of  $a$ .

The following symbols may be useful: a

Used with permission from UCLES A-level Maths papers, 2003-2017.

Gameboard:  
[STEM SMART Single Maths 28 - Binomial Theorem - Rational n](#)

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.



# Binomial Expansion 1

## Pre-Uni Maths for Sciences 4.3.1

A Level

Further A

P

P

P

P

P

P

Rewrite each of the following expressions in the form  $a(1 + b)^n$ , where  $|b| < 1$ . Hence, using the appropriate binomial expansion, find the value of each of them correct to 4 dp.

Part A    $\sqrt{36.1}$

Rewrite  $\sqrt{36.1}$  in the form  $a(1 + b)^n$ , where  $|b| < 1$  and  $n = \frac{1}{2}$ . Hence, using the appropriate binomial expansion, find the value correct to 4 dp.

Give the value of  $a$ .

Give the value of  $b$  as a simplified fraction.

Using the binomial theorem, evaluate  $\sqrt{36.1}$  correct to 4 dp.



**Part B**  $\sqrt[3]{1.09}$ 

Rewrite  $\sqrt[3]{1.09}$  in the form  $a(1 + b)^n$ , where  $|b| < 1$  and  $n = \frac{1}{3}$ . Hence, using the appropriate binomial expansion, find the value correct to 4 dp.

Give the value of  $a$ .

---

Give the value of  $b$  as a simplified fraction.

---

Using the binomial theorem, evaluate  $\sqrt[3]{1.09}$  correct to 4 dp.

---

**Part C**  $1/\sqrt{1.04}$ 

Rewrite  $\frac{1}{\sqrt{1.04}}$  in the form  $a(1 + b)^n$ , where  $|b| < 1$  and  $n = -\frac{1}{2}$ . Hence, using the appropriate binomial expansion, find the value correct to 4 dp.

Give the value of  $a$ .

---

Give the value of  $b$  as a simplified fraction.

---

Using the binomial theorem, evaluate  $\frac{1}{\sqrt{1.04}}$  correct to 4 dp.

---

Part D  $\sqrt[3]{125.4}$

Rewrite  $\sqrt[3]{125.4}$  in the form  $a(1 + b)^n$ , where  $|b| < 1$  and  $n = \frac{1}{3}$ . Hence, using the appropriate binomial expansion, find the value correct to 4 dp.

Give the value of  $a$ .

---

Give the value of  $b$  as a simplified fraction.

---

Using the binomial theorem, evaluate  $\sqrt[3]{125.4}$  correct to 4 dp.

---

Created for isaacphysics.org by Julia Riley.

Gameboard:

**STEM SMART Single Maths 28 - Binomial Theorem - Rational n**

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Physics. *You work it out.*

# Binomial: All Rational n 5i

A Level

P

P

P

Part A

Expansion

Expand  $(1 - 3x)^{-\frac{1}{3}}$  in ascending powers of  $x$ , up to and including the term in  $x^3$ .

The following symbols may be useful:  $x$

Part B

Coefficient of  $x^3$

Hence find the coefficient of  $x^3$  in the expansion of  $\left(1 - 3(x + x^3)\right)^{-\frac{1}{3}}$

Used with permission from UCLES A-level Maths papers, 2003-2017.

Gameboard:

**STEM SMART Single Maths 28 - Binomial Theorem - Rational n**

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Algebra](#) [Series](#) [Maclaurin Series - Potential due to Dipole](#)

# Maclaurin Series - Potential due to Dipole

Further A University



Pre-Uni Maths for Sciences C3.9

## Part A Potential on the axis

An electric dipole consists of two charges  $+q$  and  $-q$  separated in the  $z$  direction by a very small distance  $a$ . The electric potential  $V(z)$  a distance  $z$  away from the centre of the dipole in a direction along the line joining the two charges is given by

$$V(z) = \frac{q}{4\pi\epsilon_0} \left( \frac{1}{z - \frac{a}{2}} - \frac{1}{z + \frac{a}{2}} \right).$$

Find an approximate expression for  $V(z)$  (assume that  $z \gg a$  and obtain the first non-zero term in the Maclaurin (or binomial) expansion of  $V(z)$ ).

The following symbols may be useful:  $a$ ,  $\epsilon_0$ ,  $\pi$ ,  $q$ ,  $z$

## Part B Potential in any direction

An electric dipole consists of two charges  $+q$  and  $-q$  separated by a distance  $a$ . The electric potential  $V$  a distance  $r$  from the centre of the dipole in a direction making an angle  $\theta$  to the line joining the two charges is given approximately by

$$V \approx \frac{q}{4\pi\epsilon_0} \left( \frac{1}{\sqrt{r^2 - ar \cos \theta}} - \frac{1}{\sqrt{r^2 + ar \cos \theta}} \right).$$

Assuming that  $r \gg a$  show that  $V \approx \frac{A \cos \theta}{r^2}$  and find an expression for  $A$ .

The following symbols may be useful:  $a$ ,  $\epsilon_0$ ,  $\pi$ ,  $q$ ,  $r$

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.