



Physics. *You work it out.*

[Home](#)   [Gameboard](#)   [Maths](#)   [Integration - Trig Manipulations 1ii](#)

# Integration - Trig Manipulations 1ii


A Level  


Use integration to find the exact value of  $\int_{\frac{\pi}{16}}^{\frac{\pi}{8}} (9 - 6 \cos^2 4x) \, dx$ .

The following symbols may be useful: pi

Used with permission from UCLES A-level Maths papers, 2003-2017.

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



[Home](#) [Gameboard](#) [Maths](#) [Integration - Trig Manipulations 3ii](#)

# Integration - Trig Manipulations 3ii

A Level




Find  $\int_0^{\frac{\pi}{4}} \frac{1 - 2 \sin^2 x}{1 + 2 \sin x \cos x} dx$ , giving your answer in the form  $a \ln b$ .

Used with permission from UCLES A-level Maths papers, 2003-2017.

Gameboard:

**STEM SMART Single Maths 36 - Integration by Parts & Differential Equations**

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



[Home](#)   [Gameboard](#)   Maths   Integration - Trig Manipulations 3i

## Integration - Trig Manipulations 3i

## A Level



## Part A Simplify

Simplify as far as possible  $\frac{1}{1-\tan x} - \frac{1}{1+\tan x}$ .

The following symbols may be useful:  $x$

## Part B Integrate

Hence evaluate  $\int_{\frac{\pi}{12}}^{\frac{\pi}{6}} \left( \frac{1}{1 - \tan x} - \frac{1}{1 + \tan x} \right) dx$ , giving your answer in the form  $a \ln(b)$ .


The following symbols may be useful:  $\pi$

Used with permission from UCLES A-level Maths papers, 2003-2017.

Gameboard:

## STEM SMART Single Maths 36 - Integration by Parts & Differential Equations

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



# Integration by Substitution 2i




**Part A**   The substitution  $x = a \sin \theta$

By using the substitution  $x = a \sin \theta$ , find the exact value of

$$\int_{\frac{1}{2}a}^a \sqrt{a^2 - x^2} \, dx$$

The following symbols may be useful: a, pi

---

**Part B**    **Area of a segment**

The diagram shows the circle  $x^2 + y^2 = a^2$  and the line  $x = \frac{1}{2}a$ . Find the area of the shaded region, giving your answer in an exact form.


The following symbols may be useful:  $a$ ,  $\pi$

Used with permission from UCLES A-level Maths papers, 2003-2017.

Gameboard:

**STEM SMART Single Maths 36 - Integration by Parts & Differential Equations**

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



# Integration by Parts 3ii



Evaluate  $\int_0^{\frac{\pi}{2}} x \cos x \, dx$ , giving your answer in an exact form.


The following symbols may be useful:  $\pi$

Used with permission from UCLES A-level Maths papers, 2003-2017.

Gameboard:

**STEM SMART Single Maths 36 - Integration by Parts & Differential Equations**

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



# Integration by Parts 5

Pre-Uni Maths for Sciences K4.5

A Level

P

P

P

Find, by integrating by parts twice,  $\int_0^{\pi/3} e^{-x} \sin x \, dx$ .


The following symbols may be useful: e, pi

Created for isaacphysics.org by Julia Riley

Gameboard:

[\*\*STEM SMART Single Maths 36 - Integration by Parts & Differential Equations\*\*](#)

All materials on this site are licensed under the [\*\*Creative Commons license\*\*](#), unless stated otherwise.



# Integration by Parts 2ii



Find the exact value of  $\int_1^8 \frac{1}{\sqrt[3]{x}} \ln(x) dx$ , giving your answer in the form  $A \ln(2) + B$ , where  $A$  and  $B$  are constants to be found.


Used with permission from UCLES A-level Maths papers, 2003-2017.

Gameboard:

**STEM SMART Single Maths 36 - Integration by Parts & Differential Equations**

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.





[Home](#) [Gameboard](#) [Maths](#) [Integration of Differential Equations 1ii](#)

# Integration of Differential Equations 1ii



The gradient of a curve at the point  $(x, y)$ , where  $x > -2$ , is given by

$$\frac{dy}{dx} = \frac{1}{3y^2(x+2)}$$


The points  $(1, 2)$  and  $(q, 1.5)$  lie on the curve. Find the value of  $q$ , giving your answer correct to 3 significant figures.

Used with permission from UCLES A-level Maths papers, 2003-2017.

Gameboard:

**STEM SMART Single Maths 36 - Integration by Parts & Differential Equations**

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



# Integration of Differential Equations 4i



## Part A   Derivative

If  $y = \operatorname{cosec} x$  then find an expression for  $\frac{dy}{dx}$ .

The following symbols may be useful: `Derivative(y, x)`, `arccos()`, `arccosec()`, `arccot()`, `arcsec()`, `arcsin()`, `arctan()`, `cos()`, `cosec()`, `cot()`, `ln()`, `log()`, `sec()`, `sin()`, `tan()`, `x`, `y`

## Part B   Solve

Solve the differential equation

$$\frac{dx}{dt} = -\sin x \tan x \cot t$$

given that  $x = \frac{\pi}{6}$  when  $t = \frac{\pi}{2}$ .

The following symbols may be useful: `arccos()`, `arccosec()`, `arccot()`, `arcsec()`, `arcsin()`, `arctan()`, `cos()`, `cosec()`, `cot()`, `ln()`, `log()`, `sec()`, `sin()`, `t`, `tan()`, `x`

Used with permission from UCLES A-level Maths papers, 2003-2017.

Gameboard:  
**STEM SMART Single Maths 36 - Integration by Parts & Differential Equations**

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.


Physics. *You work it out.*[Home](#) [Gameboard](#) [Maths](#) [Constructing Differential Equations 1i](#)

# Constructing Differential Equations 1i

A Level  
P P P

A cylindrical container has a height of 200 cm. The container was initially full of a chemical but there is a leak from a hole in the base (**Figure 1**). When the leak is noticed, the container is half-full and the level of the chemical is dropping at a rate of  $1 \text{ cm min}^{-1}$ .

It is required to find for how many minutes the container has been leaking. To model the situation it is assumed that, when the depth of the chemical remaining is  $x$  cm, the rate at which the level is dropping is proportional to  $\sqrt{x}$ .



**Figure 1:** Cylindrical container that is leaking from its base.

Part A Differential equation

State an appropriate differential equation for the rate of change of height of chemical in the tank.

The following symbols may be useful:  $\text{Derivative}(x, t)$ ,  $k$ ,  $t$ ,  $x$

---

---

Part B Solve

Solve this differential equation, giving  $x$  in terms of  $t$ , the time in minutes since the leak began.

The following symbols may be useful:  $t$ ,  $x$

---

---

Part C Time

Calculate the length of time that the container has been leaking for. Give your answer to 3 significant figures.

---

Used with permission from UCLES A-level Maths papers, 2003-2017.

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.