

[Home](#) [Gameboard](#) [Maths](#) [Trigonometry: Double Angles 1ii](#)

# Trigonometry: Double Angles 1ii

**A Level**

## Part A The form $a \sin^2 \theta + b \sin \theta + c = 0$

Express the equation  $(\operatorname{cosec} \theta)(3 \cos 2\theta + 7) + 11 = 0$  in the form  $a \sin^2 \theta + b \sin \theta + c = 0$ , where  $a$ ,  $b$ , and  $c$  are constants and  $a > 0$ .

Enter the values of  $a$ ,  $b$  and  $c$ :

$a =$

$b =$

$c =$

## Part B Solve

Hence solve, for  $-180^\circ < \theta < 180^\circ$ , the equation  $(\operatorname{cosec} \theta)(3 \cos 2\theta + 7) + 11 = 0$ .

Give your answers in degrees, to three significant figures.


Enter your answers in order from lowest value of  $\theta$  to highest.

(lowest value)

(highest value)

Used with permission from UCLES A-level Maths papers, 2003-2017.

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.



Trigonometry: Double Angles 2ii

A Level

c

c

c

Part A   sin Double Angle

Write down the identity expressing  $\sin 2\theta$  in terms of  $\sin \theta$  and  $\cos \theta$ .

The following symbols may be useful: theta

Part B    $\sin 2\alpha$

Given that  $\sin \alpha = \frac{1}{4}$  and  $\alpha$  is acute, find the exact value of  $\sin 2\alpha$ .

The following symbols may be useful: alpha

Part C   Solve


Solve in degrees, for  $0^\circ < \beta < 90^\circ$ , the equation  $5 \sin 2\beta \sec \beta = 3$ , giving your answer in degrees to three significant figures.

Used with permission from UCLES A-level Maths papers, 2003-2017.

Gameboard:

[STEM SMART Single Maths 31 - Compound and Double-Angle Formulae](#)

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



# Trigonometry: Combined Angles 5i



The value of  $\tan 10^\circ$  is denoted by  $p$ . Find, in terms of  $p$ , the value of:

**Part A**    $\tan 55^\circ$

$\tan 55^\circ$

The following symbols may be useful:  $p$

---

**Part B**    $\tan 5^\circ$

$\tan 5^\circ$

The following symbols may be useful:  $p$

---

**Part C**    $\tan \theta$

$\tan \theta$ , where  $\theta$  satisfies the equation  $3 \sin (\theta + 10^\circ) = 7 \cos (\theta - 10^\circ)$ .

The following symbols may be useful:  $p$ ,  $\theta$

---


Used with permission from UCLES A-level Maths papers, 2003-2017.

Gameboard:

**STEM SMART Single Maths 31 - Compound and Double-Angle Formulae**

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.





# Functions: Reciprocal Trig 2i

A Level

P

P

P

It is given that  $A$  and  $B$  are angles such that

$$\sec^2 A - \tan A = 13 \quad \text{and} \quad \sin B \sec^2 B = 27 \cos B \operatorname{cosec}^2 B.$$

**Part A**   Largest value of  $\tan(A - B)$

Find the largest possible exact value of  $\tan(A - B)$ .

The following symbols may be useful:  $A$ ,  $B$

---

---

**Part B**   Smallest value of  $\tan(A - B)$

Give the smallest possible value of  $\tan(A - B)$ .

The following symbols may be useful:  $A$ ,  $B$

---

---


Used with permission from UCLES A-level Maths papers, 2003-2017.

Gameboard:

**STEM SMART Single Maths 31 - Compound and Double-Angle Formulae**

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.





# Trigonometry and R-Form 1

A Level



**Part A**     $5 \cos x + 12 \sin x$

Express  $5 \cos x + 12 \sin x$  in the form  $R \cos(x - \alpha)$ , where  $R > 0$  and  $0^\circ < \alpha < 90^\circ$ .

State the value of  $R$ :

$R =$

Give the value of  $\alpha$  in degrees, to three significant figures:

$\alpha =$

Part B Transformations

Give details of a pair of transformations which transform the curve  $y = \cos x$  to the curve  $y = 5 \cos x + 12 \sin x$ .

Available items

Stretch parallel to the  $x$ -axis by a factor of  $\frac{1}{R}$ .

Stretch parallel to the  $y$ -axis by a factor of  $\frac{1}{R}$ .

Stretch parallel to the  $y$ -axis by a factor of  $R$ .

Translation in the positive  $y$  direction by  $\alpha$ .

Translation in the negative  $y$  direction by  $\alpha$ .

Stretch parallel to the  $x$ -axis by a factor of  $R$ .

Translation in the positive  $x$  direction by  $\alpha$ .

Translation in the negative  $x$  direction by  $\alpha$ .

Part C Solve

Solve, for  $0^\circ < x < 360^\circ$ , the equation  $5 \cos x + 12 \sin x = 2$ , giving your answers in degrees correct to the nearest  $0.1^\circ$ .

Enter your answers in order from lowest value of  $\theta$  to highest.

 (lowest value) (highest value)

Used with permission from UCLES A-level Maths papers, 2003-2017.



All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.




Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Trigonometry: Combined Angles 3i](#)

## Trigonometry: Combined Angles 3i

A Level  
P P P

In **Figure 1**, ABCD represents a rectangular table with sides 3.5 m and 1.5 m. It has been turned so it wedges in a passage of width 2.5 m.



**Figure 1:** The rectangular table ABCD.

**Part A** Value of  $7 \sin \theta + 3 \cos \theta$

Given that  $\theta$  is the acute angle between the longer side and the passage, as shown in the diagram, find the exact value of  $7 \sin \theta + 3 \cos \theta$ .

The following symbols may be useful:  $\cos()$ ,  $\sin()$ ,  $\tan()$ , theta

**Part B    The form  $R \sin (\theta^\circ + \alpha^\circ)$**

Express  $7 \sin \theta + 3 \cos \theta$  in the form  $R \sin (\theta + \alpha)$ , where  $R > 0$  and  $0^\circ < \alpha < 90^\circ$ .

Give the exact value of  $R$ .

The following symbols may be useful: R

Give the value of  $\alpha$  to 3 significant figures.

**Part C    Find  $\theta$**


Find  $\theta$ , to 3 significant figures.

Used with permission from UCLES A-level Maths papers, 2003-2017.

Gameboard:

**STEM SMART Single Maths 31 - Compound and Double-Angle Formulae**

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



# Wave Interference

Pre-Uni Maths for Sciences H4.10

A Level  
C C C

Two waves

$$\psi_1 = A \cos \left( 2\pi ft - \left( \frac{2\pi}{\lambda} \right) x + \phi \right)$$

and

$$\psi_2 = A \cos \left( 2\pi ft - \left( \frac{2\pi}{\lambda} \right) x - \phi \right)$$

interfere, such that the resultant wave is given by  $\psi = \psi_1 + \psi_2$ . Express  $\psi$  as the product of two trigonometric functions.

Express  $\psi$  as the product of two trigonometric functions.

The following symbols may be useful: A, f, lambda, phi, pi, t, x

Created for isaacphysics.org by Julia Riley.

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.