



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Integration \(powers of x\) 4ii](#)

Integration (powers of x) 4ii

A Level



Part A Cubic

Find $\int (x^3 + 8x - 5) \, dx$.

The following symbols may be useful: c , x

Part B Square root

Find $\int 12\sqrt{x} \, dx$.

The following symbols may be useful: c , x

Used with permission from UCLES, A Level Maths, January 2009, OCR C2, Question 1

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.



Indefinite Integrals 1

A Level Further A

P

P

P

P

P

P

Find the following indefinite integrals.

Part A Integrate $(3x - 1)(x + 1)$

Find $\int (3x - 1)(x + 1)dx$.

The following symbols may be useful: k , x

Part B Integrate $(\sqrt{p} - \frac{1}{p})^2$

Find $\int \left(\sqrt{p} - \frac{1}{p}\right)^2 dp$.

The following symbols may be useful: k , p

Created for isaacphysics.org by Julia Riley

Gameboard:

STEM SMART Single Maths 12 - Introducing Integration

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Indefinite Integrals 2

A Level

Further A

P

P

P

P

P

P

Find the following indefinite integrals.

Part A Integrate $\frac{q^2+3}{q^{\frac{5}{2}}}$

Find $\int \frac{q^2+3}{q^{\frac{5}{2}}}dq$.

The following symbols may be useful: k , q

Part B Integrate $2z(z^2-1)(z^2+1)$

Find $\int 2z(z^2-1)(z^2+1)dz$.


The following symbols may be useful: k , z

Created for isaacphysics.org by Julia Riley

Gameboard:

STEM SMART Single Maths 12 - Introducing Integration

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.



Integrating Powers 1

A Level Further A

P

P

P

P

P

P

Part A Integrate $4x^3$

Find the indefinite integral of $4x^3$.

The following symbols may be useful: k , x

Part B Integrate αx .

Find $\int_0^{x_0} \alpha x \, dx$, where α is a constant.

The following symbols may be useful: α , x , x_0

Created for isaacphysics.org by Julia Riley

Gameboard:

STEM SMART Single Maths 12 - Introducing Integration

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Integrating Powers 2

A Level

Further A

P

P

P

P

P

P

Part A Integrate ax^{-8}

Find the indefinite integral of ax^{-8} , where a is a constant.

The following symbols may be useful: a , k , x

Part B Integrate $\frac{4}{x^2}$

Find $\int_1^2 \frac{4}{x^2} dx$.


The following symbols may be useful: k , x

Created for isaacphysics.org by Julia Riley

Gameboard:

STEM SMART Single Maths 12 - Introducing Integration

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Integration (powers of x) 1ii

A Level

P

P

P

Part A Find integral

Find $\int (2x + 1)(x + 3)dx$

The following symbols may be useful: c , x

Part B Evaluate integral

Evaluate $\int_0^9 \frac{1}{\sqrt{x}}dx$.

Used with permission from UCLES, A Level Maths, June 2005, OCR C2, Question 3

Gameboard:

[STEM SMART Single Maths 12 - Introducing Integration](#)

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.



Definite Integrals 3

A Level

Further A

P

P

P

P

P

P

For the integrals given below, find the values of a and m respectively.

Part A $\int_{-1}^a x^{-\frac{4}{5}} \mathrm{d}x = 10$

Given that the definite integral $\int_{-1}^a x^{-\frac{4}{5}} \mathrm{d}x = 10$, find the value of a .

The following symbols may be useful: a

Part B $\int_0^4 \frac{m+1}{2} x^m \mathrm{d}x = 4$

Given that the integral $\int_0^4 \frac{m+1}{2} x^m \mathrm{d}x = 4$, find the value of m .


The following symbols may be useful: m

Created for isaacphysics.org by Julia Riley

Gameboard:

STEM SMART Single Maths 12 - Introducing Integration

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Definite Integrals 2

A Level

Further A

P

P

P

P

P

P

Find the following integrals.

Part A $\int_1^\infty \frac{3}{2x\sqrt{x}} \mathrm{d}x$

Find $\int_1^\infty \frac{3}{2x\sqrt{x}} \mathrm{d}x$.

Part B $\int_{-8}^0 \frac{1}{\sqrt[3]{x}} \mathrm{d}x$

Find $\int_{-8}^0 \frac{1}{\sqrt[3]{x}} \mathrm{d}x$.

Part C $\int_{-1}^1 \left(1 + x + \frac{x^2}{2} + \frac{x^3}{6}\right) \mathrm{d}x$

Find $\int_{-1}^1 \left(1 + x + \frac{x^2}{2} + \frac{x^3}{6}\right) \mathrm{d}x$. Give your answer as an improper fraction.

Created for isaacphysics.org by Julia Riley

Gameboard:

STEM SMART Single Maths 12 - Introducing Integration

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Calculus](#) [Integration](#) [Integrating Powers 4](#)

Integrating Powers 4

A Level Further A



Part A Integrate $\frac{A}{r^7} - \frac{B}{r^{13}}$

Find $\int_a^\infty \left(\frac{A}{r^7} - \frac{B}{r^{13}} \right) dr$.

(The force between, for example, two atoms of an inert gas, a distance r apart is given by $\left(\frac{A}{r^7} - \frac{B}{r^{13}} \right)$, where A and B are (negative) constants; the first term is the attractive force between them (the van der Waals interaction, due to their fluctuating induced dipoles) and the second is the repulsive force due to the overlap of their electron shells. The integral describes the potential energy of such a system i.e. the work done bringing one atom from infinity to within a distance a of the other atom.)

Find $\int_a^\infty \left(\frac{A}{r^7} - \frac{B}{r^{13}} \right) dr$.

The following symbols may be useful: A , B , a

Part B Integrate $\frac{C}{x^2} + D$

Find $\int_{x_1}^{x_2} \left(\frac{C}{x^2} + D \right) dx$.

(The function $\left(\frac{C}{x^2} + D \right)$, where C and D are constants, could describe the component of an electric field in the x -direction due to a combination of the field due to a point charge at the origin and a uniform field in the x -direction. The integral is then the potential difference between two points x_1 and x_2 on the x -axis.)

Find $\int_{x_1}^{x_2} \left(\frac{C}{x^2} + D \right) dx$.

The following symbols may be useful: C , D , x_1 , x_2

Created for isaacphysics.org by Julia Riley

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.