



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Physics](#) [Fields](#) [Electric Fields](#) [Essential Pre-Uni Physics H3.2](#)

## Essential Pre-Uni Physics H3.2



Physical constants which may be necessary to answer the problems on this page can be found within the hint tabs.

For electrons moving at a speed greater than 10% of the speed of light, you should only claim that your answer is approximate (unless you have used relativistic equations). If you reckon that the electron is travelling at a speed greater than 80% of the speed of light, you should decline to give your answer unless using relativity

Convert  $3.0 \times 10^{-11}$  J into electron volts.

---

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Physics](#) [Fields](#) [Electric Fields](#) [Essential Pre-Uni Physics H3.5](#)

# Essential Pre-Uni Physics H3.5

A Level

Physical constants which may be necessary to answer the problems on this page can be found within the hint tabs.

For electrons moving at a speed greater than 10% of the speed of light, you should only claim that your answer is approximate (unless you have used relativistic equations). If you reckon that the electron is travelling at a speed greater than 80% of the speed of light, you should decline to give your answer unless using relativity

How fast is an alpha particle going if it is accelerated by a 1.5 MV potential? Assume that the alpha particle has twice the charge and four times the mass of a proton.

---

Gameboard:

**STEM SMART Physics 40 - School of Fields - Accelerators**

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Physics](#) [Fields](#) [Electric Fields](#) [Energy and Fields - Accelerator 23.2](#)

## Energy and Fields - Accelerator 23.2

A Level  


Calculate the voltage needed to accelerate a proton to  $3.5 \times 10^6 \text{ m s}^{-1}$  from rest.

Gameboard:

**STEM SMART Physics 40 - School of Fields - Accelerators**

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.

*Physics. You work it out.*[Home](#) [Gameboard](#) [Physics](#) [Fields](#) [Electric Fields](#) [Essential Pre-Uni Physics H3.6](#)

# Essential Pre-Uni Physics H3.6



Physical constants which may be necessary to answer the problems on this page can be found within the hint tabs.

For electrons moving at a speed greater than 10% of the speed of light, you should only claim that your answer is approximate (unless you have used relativistic equations). If you reckon that the electron is travelling at a speed greater than 80% of the speed of light, you should decline to give your answer unless using relativity

To trigger a particular nuclear reaction, a deuterium nucleus (same charge as the proton, but twice the mass) needs to have a kinetic energy of  $4.0 \times 10^{-13}$  J. What accelerating voltage is needed?

Gameboard:

**STEM SMART Physics 40 - School of Fields - Accelerators**

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Physics. *You work it out.*

[Home](#)[Gameboard](#)[Physics](#)[Fields](#)[Combined Fields](#)[Vectors and Fields - Mass Spectrometer 30.2](#)

# Vectors and Fields - Mass Spectrometer 30.2

**A Level**

Calculate the speed electrons emerge from a 95 V accelerator. Assume that the electrons start from rest.

Gameboard:

**STEM SMART Physics 40 - School of Fields - Accelerators**

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Physics. *You work it out.*

[Home](#)[Gameboard](#)[Physics](#)[Fields](#)[Combined Fields](#)[Vectors and Fields - Mass Spectrometer 30.6](#)

# Vectors and Fields - Mass Spectrometer 30.6

**A Level**

Calculate the specific charge  $q/m$  of a particle travelling at  $2.0 \times 10^6 \text{ m s}^{-1}$  in a magnetic field if the path radius  $r = 11.9 \text{ mm}$  and the flux density  $B = 0.175 \text{ T}$ .

Gameboard:

**STEM SMART Physics 40 - School of Fields - Accelerators**

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Physics. *You work it out.*

[Home](#)[Gameboard](#)[Physics](#)[Fields](#)[Combined Fields](#)[Vectors and Fields - Mass Spectrometer 30.7](#)

# Vectors and Fields - Mass Spectrometer 30.7

**A Level**

Calculate the voltage  $V_s$  needed in a velocity selector to pass  $1.6 \times 10^6 \text{ m s}^{-1}$  electrons in a  $2.2 \text{ T}$  magnetic field if the velocity selector plate gap  $d = 6.5 \text{ cm}$ .

Gameboard:

**STEM SMART Physics 40 - School of Fields - Accelerators**

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Physics. *You work it out.*

[Home](#)[Gameboard](#)[Physics](#)[Fields](#)[Combined Fields](#)[Vectors and Fields - Mass Spectrometer 30.11](#)

# Vectors and Fields - Mass Spectrometer 30.11

**A Level**

A singly charged ion is accelerated by a 650 kV potential before passing into a region with a 1.25 T magnetic field. It curves with a radius of 0.322 m. Calculate its mass.

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.