

[Home](#) [Gameboard](#) [Chemistry](#) [Organic](#) [Reactions](#) [Isopentyl Bromide](#)

Isopentyl Bromide



When isopentyl bromide (**F**), $(\text{CH}_3)_2\text{CHCH}_2\text{CH}_2\text{Br}$, reacts with hot aqueous ethanolic KOH, two products are formed: compound **G**, $\text{C}_5\text{H}_{12}\text{O}$, and compound **H**, $\text{C}_7\text{H}_{16}\text{O}$.




Figure 1: Isopentyl bromide (**F**)

Part A Hydroxide with ethanol

The hydroxide ion and ethanol can take part in an acid-base reaction. Write an equation to represent this. State symbols are not required.

Part B **Compound G**

What is compound **G**?

Use the [structure editor](#) to generate a SMILES string.

In the editor, after drawing your structure, click on the round, yellow smiley face to generate a SMILES string. Copy the SMILES string and paste it in the answer box.

[Using the structure editor](#)

Part C **Compound H**

What is compound **H**?

Use the [structure editor](#) to generate a SMILES string.

In the editor, after drawing your structure, click on the round, yellow smiley face to generate a SMILES string. Copy the SMILES string and paste it in the answer box.

[Using the structure editor](#)

Part D **Type of reaction**

State the type of reaction undergone by compound **F**

Part E **F with ammonia**

Draw the structure of the product derived from compound **F** by reaction with concentrated aqueous ammonia.

Use the [structure editor](#) to generate a SMILES string.

In the editor, after drawing your structure, click on the round, yellow smiley face to generate a SMILES string. Copy the SMILES string and paste it in the answer box.

[Using the structure editor](#)

Part F **Reaction with potassium cyanide**

Draw the structure of the product derived from compound **F** by reaction with ethanolic potassium cyanide.

Use the [structure editor](#) to generate a SMILES string.

In the editor, after drawing your structure, click on the round, yellow smiley face to generate a SMILES string. Copy the SMILES string and paste it in the answer box.

[Using the structure editor](#)

Adapted with permission from UCLES, A-Level Chemistry, November 1994, Paper 1, Question 10

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.



Physics. You work it out.

[Home](#) [Gameboard](#) [Chemistry](#) [Organic](#) [Reactions](#) [Mechanism Types](#)

Mechanism Types



Part A Perspex intermediate

The following reaction is an intermediate stage in the manufacture of the important polymer known as *Perspex*.




Figure 1: Intermediate in manufacture of *Perspex*

What type of mechanism is involved in this reaction?

- ☐ free radical substitution
- ☐ electrophilic substitution
- ☐ electrophilic addition
- ☐ nucleophilic substitution
- ☐ nucleophilic addition

Part B **Reaction with chlorine**

A non-polar organic compound undergoes a reaction with chlorine [$A_r(\text{Cl}) = 35.5$] when light is shone upon the reaction mixture. The relative molecular mass of the product is 34.5 greater than that of the original compound.

The reaction is most likely to be:

- ☐ electrophilic substitution
 - ☐ electrophilic addition
 - ☐ nucleophilic substitution
 - ☐ free radical substitution
 - ☐ nucleophilic addition
-

Part A adapted with permission from UCLES, A-Level Chemistry, November 1991, Paper 1, Question 23 ;

Part B adapted with permission from OCSEB, A-Level Chemistry, June 1994, Paper 1, Question 21

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.


Physics. *You work it out.*[Home](#) [Gameboard](#) [Chemistry](#) [Organic](#) [Reactions](#) [Reactions of \$\text{NH}_3\$](#)

Reactions of NH_3



Part A With haloalkanes

An amine is produced in the following reaction.




What is the mechanism?

- ☐ nucleophilic substitution
- ☐ electrophilic addition
- ☐ nucleophilic addition
- ☐ electrophilic substitution

Part B With aldehydes

Ethanal can react with ammonia as shown.



Which kind of chemical reaction takes place?

- ☐ addition-elimination
 - ☐ electrophilic addition
 - ☐ nucleophilic addition
 - ☐ free-radical addition
-

Part A adapted with permission from UCLES, A-Level Chemistry, June 1996, Paper 3, Question 22;

Part B adapted with permission from OCR, A-Level Chemistry, June 1999, Paper 3, Question 21

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Chemistry](#) [Organic](#) [Reactions](#) [Organic Reaction Intermediates](#)

Organic Reaction Intermediates



Part A Hydrobromic acid with ethene

Hydrogen bromide reacts with ethene to form bromoethane. Which of the following is the best description of the organic intermediate?

- ☐ It has a negative charge
- ☐ It contains carbon, hydrogen and bromine
- ☐ Its structure is planar
- ☐ It is a free radical
- ☐ It is an electrophile

Part B Carbocation intermediate

In which of the following reactions is a cation an intermediate?

- ☐ $\text{CH}_3\text{CH}_2\text{Br} + \text{NaOH} \xrightarrow{\text{H}_2\text{O}} \text{CH}_3\text{CH}_2\text{OH} + \text{NaBr}$
- ☐ $\text{CH}_3\text{CH}_2\text{Cl} + 2 \text{NH}_3 \longrightarrow \text{CH}_3\text{CH}_2\text{NH}_2 + \text{NH}_4\text{Cl}$
- ☐ $\text{CH}_3\text{CH}_3 + \text{Cl}_2 \longrightarrow \text{CH}_3\text{CH}_2\text{Cl} + \text{HCl}$
- ☐ $\text{CH}_3\text{CHO} + \text{HCN} \xrightarrow{\text{CN}^-} \text{CH}_3\text{CH}(\text{OH})\text{CN}$
- ☐ $\text{CH}_2=\text{CH}_2 + \text{Br}_2 \longrightarrow \text{CH}_2\text{BrCH}_2\text{Br}$

Part A adapted with permission from UCLES, A-Level Chemistry, 1988, Paper 3, Question 23;

Part B adapted with permission from UCLES, A-Level Chemistry, June 1992, Paper 4, Question 22

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.



Physics. You work it out.


[Home](#) [Gameboard](#) [Chemistry](#) [Organic](#) [Reactions](#) [Compounds from Haloalkanes](#)

Compounds from Haloalkanes



Part A Compounds from C₃H₇Br

The compound C₃H₇Br undergoes a sequence of reactions as follows:



What could be the formulae for **X**, **Y** and **Z**?

	X	Y	Z
A	CH ₃ CH ₂ CH ₂ OH	CH ₃ CH ₂ COOH	CH ₃ CH ₂ CHO
B	CH ₃ CH ₂ CH ₂ OH	CH ₃ CH(OH)CH ₂ OH	CH ₃ COOH
C	CH ₃ CH ₂ CH ₂ OH	CH ₃ CH ₂ CHO	CH ₃ CH ₂ COOH
D	CH ₃ CH(OH)CH ₃	CH ₃ COCH ₃	CH ₃ COOH

- ☐ **A**
- ☐ **B**
- ☐ **C**
- ☐ **D**

Part B **Compounds from chloroethane**

Chloroethane is converted into a carboxylic acid containing one more carbon atom through a two-stage process.

Which of the following compounds could be the intermediate in the synthesis of the carboxylic acid?

- ☐ $\text{CH}_3\text{CH}_2\text{COOCH}_3$
 - ☐ $\text{CH}_3\text{CH}_2\text{OH}$
 - ☐ $\text{CH}_3\text{CH}_2\text{CN}$
 - ☐ $\text{CH}_3\text{CH}_2\text{CH}_2\text{NH}_2$
 - ☐ $\text{CH}_3\text{CH}_2\text{CH}_2\text{CN}$
-

Part A adapted with permission from UCLES, A-Level Chemistry, November 1994, Paper 4, Question 26;

Part B adapted with permission from UCLES, A-Level Chemistry, June 1993, Paper 3, Question 25

All materials on this site are licensed under the [Creative Commons license](https://creativecommons.org/licenses/by/4.0/), unless stated otherwise.



Physics. You work it out.

[Home](#) [Gameboard](#) [Chemistry](#) [Organic](#) [Reactions](#) [Compounds from But-2-ene](#)

Compounds from But-2-ene

A Level

Complete the reaction scheme shown below which starts with but-2-ene. In each of the boxes **A** to **D** give the principal organic product or intermediate compound.

Use the [structure editor](#) to generate a SMILES string.




Figure 1: Compounds from but-2-ene

In the editor, after drawing your structure, click on the round, yellow smiley face to generate a SMILES string. Copy the SMILES string and paste it in the answer box.

[Using the structure editor](#)

Part A **A**

A is:

Part B B

B is:

Part C C

C is:

Part D D

D is:

Adapted with permission from UCLES, A-Level Chemistry, June 1996, Chains and Rings, Question 1


All materials on this site are licensed under the [Creative Commons license](https://creativecommons.org/licenses/by/4.0/), unless stated otherwise.

Physics. *You work it out.*[Home](#) [Gameboard](#) [Chemistry](#) [Organic](#) [Reactions](#) [Ketones with HCN Further](#)

Ketones with HCN Further



The initial product of the reaction between HCN and propanone gives **A** which is then subjected to a dehydration reaction to produce **B**.



Part A Propanone with HCN

What is **A**?

Use the [structure editor](#) to generate a SMILES string.

In the editor, after drawing your structure, click on the round, yellow smiley face to generate a SMILES string. Copy the SMILES string and paste it in the answer box.

[Using the structure editor](#)

Part B Dehydration

What is **B**?

Use the [structure editor](#) to generate a SMILES string.

In the editor, after drawing your structure, click on the round, yellow smiley face to generate a SMILES string. Copy the SMILES string and paste it in the answer box.

[Using the structure editor](#)

Part C Hydrolysis

The product of the dehydration reaction **B** is hydrolysed under acidic conditions to give **C**.

What is **C**?

Use the [structure editor](#) to generate a SMILES string.

In the editor, after drawing your structure, click on the round, yellow smiley face to generate a SMILES string. Copy the SMILES string and paste it in the answer box.

[Using the structure editor](#)

Part A and B adapted with permission from UCLES, A-Level Chemistry, June 1992, Paper 4, Question 28;

Part C created for isaacphysics.org by R. Less

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.



Physics. You work it out.

[Home](#) [Gameboard](#) [Chemistry](#) [Organic](#) [Reactions](#) [More Nitriles](#)

More Nitriles



Part A Reaction with cyanide ions

Which of the following compounds could be the product of a reaction involving a nucleophilic attack by cyanide ions in aqueous ethanolic solution?

- ☐ $\text{CH}_2=\text{CHCN}$
- ☐ $(\text{CH}_3)_2\text{CHCN}$
- ☐ CH_3CONH_2
- ☐ $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}_2$

Part B Hydrolysis of CS

CS has the structure shown below, is an active component of 'tear gas' and is readily hydrolysed.




Figure 1: Structure of CS

Which of the following is a possible hydrolysis product of CS?




Figure 2: Possible hydrolysis products of CS

- ☐ **A**
- ☐ **B**
- ☐ **C**
- ☐ **D**

Part A adapted with permission from UCLES, A-Level Chemistry, November 1994, Paper 4, Question 23;

Part B adapted with permission from UCLES, A-Level Chemistry, June 1995, Paper 4, Question 25



Physics. You work it out.

[Home](#) [Gameboard](#) [Chemistry](#) [Organic](#) [Reactions](#) [Role of Reagent](#)

Role of Reagent



Part A Bradosol

Bradosol is a compound used for the relief of sore throats. It is produced in the following reaction.




Figure 1: *Bradosol* synthesis

What is the role of compound **Y** in this reaction?

- ☐ an electrophile
- ☐ a nucleophile
- ☐ a ligand
- ☐ a reducing agent

Part B Inorganic reagent

In which reaction does the inorganic reagent act as a nucleophile?

- ☐ $\text{CH}_3\text{CH}=\text{CH}_2 + \text{Br}_2 \longrightarrow \text{CH}_3\text{CHBrCH}_2\text{Br}$
- ☐ $\text{CH}_3\text{CH}_2\text{NH}_2 + \text{HCl} \longrightarrow [\text{CH}_3\text{CH}_2\text{NH}_3]^+\text{Cl}^-$
- ☐ $\text{CH}_3\text{CH}_2\text{Br} + \text{NaOH} \longrightarrow \text{CH}_3\text{CH}_2\text{OH} + \text{NaBr}$
- ☐ $\text{CH}_3\text{CH}_3 + \text{Cl}_2 \xrightarrow{h\nu \text{ (light)}} \text{CH}_3\text{CH}_2\text{Cl} + \text{HCl}$
-

Part A adapted with permission from UCLES, A-Level Chemistry, June 1998, Paper 3, Question 19;

Part B adapted with permission from UCLES, A-Level Chemistry, November 1998, Paper 3, Question 20

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Physics. You work it out.

[Home](#) [Gameboard](#) [Chemistry](#) [Organic](#) [Reactions](#) [Reaction Types](#)

Reaction Types

A Level



Part A Sunburn ointment

Many sunburn ointments contain benzocaine which relieves the pain caused by sunburn. It can be made in the laboratory by using the following reaction scheme.




Figure 1: Preparation of benzocaine

Which of the following statements about this reaction scheme are correct?

1 Step 1 is an oxidation.


2 Step 2 is an esterification.

3 Step 3 is a reduction.

- ☐ **1, 2 and 3** are correct
 - ☐ **1 and 2** only are correct
 - ☐ **1 and 3** only are correct
 - ☐ **2 and 3** only are correct
 - ☐ **1** only is correct
 - ☐ **2** only is correct
 - ☐ **3** only is correct
-

Part B Aldol

The Russian composer Borodin was also a research chemist who discovered a reaction in which two ethanal molecules combine to form a compound commonly known as aldol (reaction I). Aldol forms another compound on heating (reaction II).



Which of the following best describes reactions I and II?

	I	II
A	addition	elimination
B	addition	reduction
C	elimination	reduction
D	substitution	elimination

- ☐ **A**
- ☐ **B**
- ☐ **C**
- ☐ **D**

Part A adapted with permission from UCLES, A-Level Chemistry, November 1993, Paper 4, Question 36;

Part B adapted with permission from UCLES, A-Level Chemistry, November 1994, Paper 4, Question 21

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.