



Physics. *You work it out.*

[Home](#) [Gameboard](#) Maths Functions from Differential Equations 2ii

Functions from Differential Equations 2ii

A Level



The gradient of a curve is given by $\frac{dy}{dx} = 6x - 4$. The curve passes through the distinct points $(2, 5)$ and $(p, 5)$.

Part A Equation of curve

Find the equation of the curve.

The following symbols may be useful: x , y

Part B Find p

Find the value of p .

The following symbols may be useful: p

Used with permission from UCLES, A level, January 2010, Paper 4722, Question 2.

Gameboard:

STEM SMART Single Maths 13 - Using Integration

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) Maths Functions from Differential Equations 1ii

Functions from Differential Equations 1ii

A Level



The gradient of a curve is given by $\frac{dy}{dx} = 12\sqrt{x}$. The curve passes through the point (4, 50). Find the equation of the curve.

The following symbols may be useful: c , x , y

Used with permission from UCLES, A Level, June 2012, Paper 4722, Question 2

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.



Equation of Curve

A Level

P

P

P

A function $v(u)$ is such that $\frac{dv}{du} = \frac{1}{3}u^{\frac{1}{3}} \left(1 - \frac{1}{u}\right)$ and $v(8) = -1$.

Find the equation of the function $v(u)$.

The following symbols may be useful: u , v

Created for isaacphysics.org by Julia Riley

Gameboard:

STEM SMART Single Maths 13 - Using Integration

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Functions from Differential Equations 1i](#)

Functions from Differential Equations 1i

A Level



A curve has an equation which satisfies $\frac{dy}{dx} = kx(2x - 1)$ for all values of x . The point $P(2, 7)$ lies on the curve and the gradient of the curve at P is 9.

Part A Find k

Find the value of the constant k . Give your answer as an improper fraction.

The following symbols may be useful: k

Part B Equation of curve

Find the equation of the curve.

The following symbols may be useful: x , y

Used with permission from UCLES, A Level, January 2013, Paper 4722, Question 3

Gameboard:

[**STEM SMART Single Maths 13 - Using Integration**](#)

All materials on this site are licensed under the [Creative Commons license](#), unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Calculus](#) [Integration](#) [Integrating to Find An Area 1](#)

Integrating to Find An Area 1

A Level



This question is about the use of integration to find the area of one or more regions between a curve and the x -axis.

Part A Calculating the area under $y = x^2 + 1$




Figure 1: The graph of $y = x^2 + 1$

Figure 1 shows the curve $y = x^2 + 1$. The region between the curve and the x -axis, bounded by the lines $x = 0$ and $x = 3$, is labelled A . Calculate the area of A .


Part B Calculating the area above $y = x^2 - 1$ 

Figure 2: The graph of $y = x^2 - 1$

Figure 2 shows the curve $y = x^2 - 1$. The region between the curve and the x -axis, bounded by the lines $x = -1$ and $x = 1$, is labelled B . Calculate the area of B .

Part C Integrating $5x(x^2 - 1)$

Calculate the value of the integral $\int_{-1}^1 5x(x^2 - 1)dx$.

Part D The region enclosed between $y = 5x(x^2 - 1)$ and the x -axis




Figure 3: The graph of $y = 5x(x^2 - 1)$

Use **Figure 3** to explain why $\int_{-1}^1 5x(x^2 - 1)dx$ does not give the total area of the regions enclosed by the curve $y = 5x(x^2 - 1)$ and the x -axis between $x = -1$ and $x = +1$. Drag and drop options into the spaces provided to complete your answer.

The curve $y = 5x(x^2 - 1)$ intercepts the x -axis at $x = -1$, $x = 0$ and $x = 1$. For $-1 < x < 0$ the curve is the x -axis. Hence, the value of $\int_{-1}^0 5x(x^2 - 1)dx$ is and equal to the area of region . However, for $0 < x < 1$ the curve is the x -axis. Hence, the value of the integral $\int_0^1 5x(x^2 - 1)dx$ is . The area of the region labelled is given by $-\int_0^1 5x(x^2 - 1)dx$.

The total area of the shaded regions is $\int_{-1}^0 5x(x^2 - 1)dx - \int_0^1 5x(x^2 - 1)dx$. This is not the same as $\int_{-1}^1 5x(x^2 - 1)dx$, which has a value of 0 as the curve has symmetry and the contributions from the parts above and below the x -axis cancel out exactly.

Items:

Part E Calculating an area for $y = 5x(x^2 - 1)$

Calculate the area of the region enclosed between the curve $y = 5x(x^2 - 1)$ and the x -axis for x between -1 and 1 .

Created for isaacphysics.org by Jonathan Waugh

Gameboard:

STEM SMART Single Maths 13 - Using Integration

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Integration and Area 1ii](#)

Integration and Area 1ii

A Level



Figure 1 shows part of the curve $y = x^2 - 3x$ and the line $x = 5$.




Figure 1: A graph of $y = x^2 - 3x$ and the line $x = 5$

Part A Area of shaded regions

Which of the following expressions gives the total area of the regions shaded in **Figure 1**? Explain your choice.

- ☐ $\int_0^5 (x^2 - 3x) \, dx$
- ☐ $\int_3^5 (x^2 - 3x) \, dx - \int_0^3 (x^2 - 3x) \, dx$
- ☐ $\int_0^3 (x^2 - 3x) \, dx + \int_3^5 (x^2 - 3x) \, dx$

Part B Find area

Use integration to find the total area of the shaded regions. Give your answer to 3 significant figures.

Used with permission from UCLES, A Level Maths, January 2012, OCR C2, Question 7

Gameboard:

STEM SMART Single Maths 13 - Using Integration

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Calculus](#) [Integration](#) [Area Under a Curve 1](#)

Area Under a Curve 1

A Level



A graph of the functions $y = (x - 2)(x + 1)$ and $y = x + 1$ is shown in **Figure 1**. Find the areas of the shaded regions labelled A and B. A is the region between P and Q enclosed by the curve $y = (x - 2)(x + 1)$ and the x -axis; B is the region between Q and R below the curve $y = (x - 2)(x + 1)$ and above the x -axis.




Figure 1: A graph of the functions $y = (x - 2)(x + 1)$ and $y = x + 1$. A is the region between P and Q enclosed by the curve $y = (x - 2)(x + 1)$ and the x -axis; B is the region between Q and R below the curve $y = (x - 2)(x + 1)$ and above the x -axis.

Part A Region A

Find the area of the region A. Give your answer in the form of an improper fraction.

Part B Region B

Find the area of the region B. Give your answer in the form of an improper fraction.

Created for isaacphysics.org by Julia Riley

Gameboard:

STEM SMART Single Maths 13 - Using Integration

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Integration and Area 3i](#)

Integration and Area 3i

A Level



Figure 1 shows the graph of $y = 1 - 3x^{-\frac{1}{2}}$.




Figure 1: Graph of $y = 1 - 3x^{-\frac{1}{2}}$.

Part A Find intersection with x -axis

Find the x -coordinate of the intersection of that curve with the x -axis.

The following symbols may be useful: x

Part B Find a

The region enclosed by the curve, the x -axis and the line $x = a$ (where $a > 9$) has an area equal to 4 square units, find the value of a .

The following symbols may be useful: a

Gameboard:

STEM SMART Single Maths 13 - Using Integration

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Calculus](#) [Integration](#) [Area Under a Curve 3](#)

Area Under a Curve 3

A Level Further A



A graph of the functions $y = \frac{1}{2\sqrt{x}}$ and $y = 2x\sqrt{x}$ for $x \geq 0$ is shown in **Figure 1**. Find the area of the shaded region OPQR.




Figure 1: A graph of the functions $y = \frac{1}{2\sqrt{x}}$ and $y = 2x\sqrt{x}$ for $x \geq 0$. The shaded region OPQR is bounded by the line $x = 0$, the line $y = 1$, the curve $y = \frac{1}{2\sqrt{x}}$ and the curve $y = 2x\sqrt{x}$.

Part A The x coordinate of Q

Deduce the x coordinate of the point Q.

Part B The x coordinate of R

Find the x coordinate of the point R.

Part C The area of OPQR

Find the area of the shaded region OPQR, giving your answer in an exact form.

Created for isaacphysics.org by Julia Riley

All materials on this site are licensed under the [Creative Commons license](https://creativecommons.org/licenses/by/4.0/), unless stated otherwise.