

Electrochemistry Introduction

Subject & topics: Chemistry | Physical | Electrochemistry

Stage & difficulty: A Level P1

In electrochemistry, a potential difference between two half-cells is set up by having different redox couples and/or different concentrations of a given redox couple present.

Part A

Introduction

The potential of the side half-cell is measured relative to the side one, so the cell potential is given by subtracting the reduction potential of the side from the reduction potential of the side. Standard reduction potentials can be tabulated, which correspond to values recorded under standard conditions against the standard electrode. The conventional cell reaction consists of the right-hand side and the left-hand side (which can be thought of as subtracting the left-hand side), making sure that the number of electrons (n) transferred is the same for both sides.

Items:

Part B

Linking equations

Working out cell potentials can be useful in itself, but as a result of connections to other thermodynamic quantities, tabulated standard reduction potentials allow us to calculate, for example, equilibrium constants even of non-redox reactions or processes such as a salt dissolving.

Given that $\Delta G^\circ = -nFE^\circ = -RT \ln K$, rearrange the equation for K (the equilibrium constant) as a function of n (the number of electrons transferred), F (the Faraday constant), E° (the standard cell potential, for which you should use E° in your expression), R (the universal gas constant) and T (the temperature).

The following symbols may be useful: E° , F , K , R , T , e , $\ln()$, $\log()$, n

Electrode Potential & Cell Potential 1

Essential Pre-Uni Chemistry L1.1

Subject & topics: Chemistry | Physical | Electrochemistry

Stage & difficulty: A Level P2

Name the element whose reduction is used as a standard by which all electrode potentials are measured.

- ☐ Oxygen
- ☐ Silver
- ☐ Helium
- ☐ Hydrogen
- ☐ Iron
- ☐ Fluorine
- ☐ Lithium
- ☐ Platinum

Question deck:

STEM SMART Chemistry Week 37

Electrode Potential & Cell Potential 3

Essential Pre-Uni Chemistry L1.3

Subject & topics: Chemistry | Physical | Electrochemistry **Stage & difficulty:** A Level P2

The standard electrode potential, E^\ominus , for the reduction, $\text{Br}_2(\text{aq}) + 2\text{e}^- \longrightarrow 2\text{Br}^-(\text{aq})$ is 1.09 V. Give the E^\ominus value for the reduction, $\frac{1}{2}\text{Br}_2(\text{aq}) + \text{e}^- \longrightarrow \text{Br}^-(\text{aq})$.

Question deck:

STEM SMART Chemistry Week 37

Electrode Potential & Cell Potential 4

Essential Pre-Uni Chemistry L1.4

Subject & topics: Chemistry | Physical | Electrochemistry

Stage & difficulty: A Level P2

E^\ominus for the reaction, $\text{Ce}^{4+}(\text{aq}) + \text{e}^- \longrightarrow \text{Ce}^{3+}(\text{aq})$ is 1.70 V. Give the E^\ominus value for the oxidation half-reaction, $\text{Ce}^{3+}(\text{aq}) \longrightarrow \text{Ce}^{4+}(\text{aq}) + \text{e}^-$.

Question deck:

STEM SMART Chemistry Week 37

Electrode Potential & Cell Potential 5

Essential Pre-Uni Chemistry L1.5


Subject & topics: Chemistry | Physical | Electrochemistry **Stage & difficulty:** A Level P2

Reduction	E^\ominus / V
$\text{Zn}^{2+}(\text{aq}) + 2\text{e}^- \longrightarrow \text{Zn}(\text{s})$	-0.76
$\text{Cr}^{3+}(\text{aq}) + 3\text{e}^- \longrightarrow \text{Cr}(\text{s})$	-0.74
$\text{Fe}^{2+}(\text{aq}) + 2\text{e}^- \longrightarrow \text{Fe}(\text{s})$	-0.44
$\text{Cu}^{2+}(\text{aq}) + \text{e}^- \longrightarrow \text{Cu}^+(\text{aq})$	+0.16
$\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \longrightarrow \text{Cu}(\text{s})$	+0.34
$\text{Cu}^+(\text{aq}) + \text{e}^- \longrightarrow \text{Cu}(\text{s})$	+0.52
$\text{Fe}^{3+}(\text{aq}) + \text{e}^- \longrightarrow \text{Fe}^{2+}(\text{aq})$	+0.77
$\text{Ag}^+(\text{aq}) + \text{e}^- \longrightarrow \text{Ag}(\text{s})$	+0.80
$\text{Cr}_2\text{O}_7^{2-}(\text{aq}) + 6\text{e}^- + 14\text{H}^+(\text{aq}) \longrightarrow 2\text{Cr}^{3+}(\text{aq}) + 7\text{H}_2\text{O}(\text{l})$	+1.33

Use the standard electrode potentials tabulated above to calculate the standard cell potentials due to the following reactions, giving your answers to 2 decimal places throughout:


Part A

(a)




Part B

(b)




Part C

(c)



Part D

(d)



Question deck:


STEM SMART Chemistry Week 37

Free Energy Changes 9

Essential Pre-Uni Chemistry H2.9

Subject & topics: Chemistry | Physical | Entropy **Stage & difficulty:** A Level P2

The displacement of hydrogen from acid by iron,



has a standard cell potential of 0.44 V. Find the associated standard Gibbs free energy change. (Faraday constant = 96 485 C mol⁻¹)

Question deck:

STEM SMART Chemistry Week 37

Free Energy & Kc 1

Essential Pre-Uni Chemistry L2.1


Subject & topics: Chemistry | Physical | Electrochemistry **Stage & difficulty:** A Level P2

Reduction	E^\ominus / V
$\text{Zn}^{2+}(\text{aq}) + 2\text{e}^- \longrightarrow \text{Zn}(\text{s})$	-0.76
$\text{Cr}^{3+}(\text{aq}) + 3\text{e}^- \longrightarrow \text{Cr}(\text{s})$	-0.74
$\text{Fe}^{2+}(\text{aq}) + 2\text{e}^- \longrightarrow \text{Fe}(\text{s})$	-0.44
$\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \longrightarrow \text{Cu}(\text{s})$	+0.34
$\text{Cu}^+(\text{aq}) + \text{e}^- \longrightarrow \text{Cu}(\text{s})$	+0.52
$\text{Fe}^{3+}(\text{aq}) + \text{e}^- \longrightarrow \text{Fe}^{2+}(\text{aq})$	+0.77
$\text{Ag}^+(\text{aq}) + \text{e}^- \longrightarrow \text{Ag}(\text{s})$	+0.80
$\text{Cr}_2\text{O}_7^{2-}(\text{aq}) + 6\text{e}^- + 14\text{H}^+(\text{aq}) \longrightarrow 2\text{Cr}^{3+}(\text{aq}) + 7\text{H}_2\text{O}(\text{l})$	+1.33

Use the standard electrode potentials tabulated above to find ΔG^\ominus for the following reactions:


Part A

(a)




Part B

(b)



Part C

(c)



Question deck:

STEM SMART Chemistry Week 37

Free Energy & Kc 2

Essential Pre-Uni Chemistry L2.2

Subject & topics: Chemistry | Physical | Electrochemistry **Stage & difficulty:** A Level P2

Reduction	E^\ominus / V
$\text{Zn}^{2+}(\text{aq}) + 2\text{e}^- \longrightarrow \text{Zn}(\text{s})$	-0.76
$\text{Cr}^{3+}(\text{aq}) + 3\text{e}^- \longrightarrow \text{Cr}(\text{s})$	-0.74
$\text{Fe}^{2+}(\text{aq}) + 2\text{e}^- \longrightarrow \text{Fe}(\text{s})$	-0.44
$\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \longrightarrow \text{Cu}(\text{s})$	$+0.34$
$\text{Cu}^+(\text{aq}) + \text{e}^- \longrightarrow \text{Cu}(\text{s})$	$+0.52$
$\text{Fe}^{3+}(\text{aq}) + \text{e}^- \longrightarrow \text{Fe}^{2+}(\text{aq})$	$+0.77$
$\text{Ag}^+(\text{aq}) + \text{e}^- \longrightarrow \text{Ag}(\text{s})$	$+0.80$
$\text{Cr}_2\text{O}_7^{2-}(\text{aq}) + 6\text{e}^- + 14\text{H}^+(\text{aq}) \longrightarrow 2\text{Cr}^{3+}(\text{aq}) + 7\text{H}_2\text{O}(\text{l})$	$+1.33$

Use the standard electrode potentials tabulated above to find ΔG^\ominus for the following reactions:

Part A

(a)

$\text{Ag}^+(\text{aq}) + \text{Fe}^{2+}(\text{aq}) \longrightarrow \text{Fe}^{3+}(\text{aq}) + \text{Ag}(\text{s})$. Give your answer to 1 significant figure.

Part B

(b)

$3 \text{Zn(s)} + \text{Cr}_2\text{O}_7^{2-}(\text{aq}) + 14 \text{H}^+(\text{aq}) \longrightarrow 3 \text{Zn}^{2+}(\text{aq}) + 2 \text{Cr}^{3+}(\text{aq}) + 7 \text{H}_2\text{O(l)}$. Give your answer to 3 significant figures.

Part C

(c)

$2 \text{Cr(s)} + 3 \text{Cu}^{2+}(\text{aq}) \longrightarrow 2 \text{Cr}^{3+}(\text{aq}) + 3 \text{Cu(s)}$. Give your answer to 3 significant figures.

Question deck:

STEM SMART Chemistry Week 37

Electrode Potential & Cell Potential 7

Essential Pre-Uni Chemistry L1.7

Subject & topics: Chemistry | Physical | Electrochemistry **Stage & difficulty:** A Level P2

Reduction	E^\ominus / V
$\text{Zn}^{2+}(\text{aq}) + 2\text{e}^- \longrightarrow \text{Zn}(\text{s})$	-0.76
$\text{Cr}^{3+}(\text{aq}) + 3\text{e}^- \longrightarrow \text{Cr}(\text{s})$	-0.74
$\text{Fe}^{2+}(\text{aq}) + 2\text{e}^- \longrightarrow \text{Fe}(\text{s})$	-0.44
$\text{Cu}^{2+}(\text{aq}) + \text{e}^- \longrightarrow \text{Cu}^+(\text{aq})$	$+0.16$
$\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \longrightarrow \text{Cu}(\text{s})$	$+0.34$
$\text{Cu}^+(\text{aq}) + \text{e}^- \longrightarrow \text{Cu}(\text{s})$	$+0.52$
$\text{Fe}^{3+}(\text{aq}) + \text{e}^- \longrightarrow \text{Fe}^{2+}(\text{aq})$	$+0.77$
$\text{Ag}^+(\text{aq}) + \text{e}^- \longrightarrow \text{Ag}(\text{s})$	$+0.80$
$\text{Cr}_2\text{O}_7^{2-}(\text{aq}) + 6\text{e}^- + 14\text{H}^+(\text{aq}) \longrightarrow 2\text{Cr}^{3+}(\text{aq}) + 7\text{H}_2\text{O}(\text{l})$	$+1.33$

Using the data tabulated above, calculate the standard cell potential for:

Part A

(a)




Part B

(b)



Part C

(c)



Question deck:

STEM SMART Chemistry Week 37

Combining Potentials


Subject & topics: Chemistry | Physical | Electrochemistry **Stage & difficulty:** A Level C2

When half-cell potentials are combined to form a cell potential, the process is relatively straightforward: we balance the two half-equations to include the same number of electrons, and subtract the left-hand half-cell potential from the right-hand half-cell potential.

When two half-cell potentials need to be combined to instead form another half-cell potential, for a third half-reaction, the process is a little more complicated.

$A^{(n_1+n_2)+} + n_1e^- \longrightarrow A^{n_2+}$ has a half-cell potential of x and $A^{n_2+} + n_2e^- \longrightarrow A$ has a half-cell potential of y .

Derive an expression for the half-cell potential of the following reaction:



The following symbols may be useful: n_1 , n_2 , x , y